Enantioselective Gold-Catalyzed Pictet–Spengler Reaction

Nicolas Glinsky-Olivier, Shengwen Yang, Pascal Retailleau, Vincent Gandon,
Xavier Guinchard

► To cite this version:

Nicolas Glinsky-Olivier, Shengwen Yang, Pascal Retailleau, Vincent Gandon, Xavier Guinchard. Enantioselective Gold-Catalyzed Pictet–Spengler Reaction. *Organic Letters*, 2019, 21 (23), pp.9446 - 9451. 10.1021/acs.orglett.9b03656 . hal-03004206

HAL Id: hal-03004206

<https://hal.science/hal-03004206>

Submitted on 27 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The Enantioselective Gold-Catalyzed Pictet-Spengler Reaction

Nicolas Glinsky-Olivier,^a Shengwen Yang,^{b,c} Pascal Retailleau,^a Vincent Gandon,^{b,c*} Xavier Guinchard^{a*}

^a Institut de Chimie des Substances Naturelles, CNRS UPR 2301, Université Paris-Sud, Université Paris-Saclay, 1 avenue de la Terrasse, 91198 Gif-sur-Yvette, France.

^b Institut de Chimie Moléculaire et des Matériaux d'Orsay, CNRS UMR 8182, Université Paris-Sud, Université Paris-Saclay, Bâtiment 420, 91405 Orsay cedex, France.

^c Laboratoire de Chimie Moléculaire (LCM), CNRS UMR 9168, Ecole Polytechnique, IP Paris, route de Saclay, 91128 Palaiseau cedex, France.

Supporting Information Placeholder

ABSTRACT: Cationic chiral Au(I)-complexes catalyze asymmetric Pictet-Spengler reactions between tryptamines and arylaldehydes. The resulting tetrahydro- β -carboline are obtained with wide functional group tolerance in high yields and ee's (up to 95%). Aldehydes bearing polar and/or protic functions are well tolerated. The reaction features a hitherto unknown C2-auration of the indole as the key step, supported by both kinetic isotope effect (KIE) measurement and DFT calculations.

The Pictet-Spengler reaction is the acid-catalyzed condensation of β -arylethylamines with aldehydes, leading to tetrahydroisquinoline,¹ or tetrahydro- β -carboline (TH- β -Cs)² through the intermediacy of an iminium. With more than a century of intense research, the Pictet-Spengler reaction has now won its spurs as one of the most efficient and reliable method for the synthesis of tetrahydro- β -carboline, a major class of heterocyclic compounds (Scheme 1).³ Some of these derivatives are natural products, such as (*R*)-tryptargine or strictosidine. Numerous compounds embedding this structural unit are endowed with wide and interesting biological activities and are currently available as drugs or drug candidates.^{3b}

Scheme 1. Structure and Natural Occurrence of Tetrahydro- β -Carbolines

Numerous efforts have been made by synthetic chemists to control the absolute stereochemistry of C1.⁴ Enantioselective approaches to tetrahydro- β -carboline include the reduction of dihydro- β -carboline, using Ru or Rh chiral complexes via asymmetric transfer hydrogenation reactions⁵ or biocatalyzed reductions.⁶ However, the Pictet-Spengler re-

action is unarguably the leading synthetic strategy for the efficient synthesis of chiral tetrahydro- β -carboline.⁷ Beyond diastereoselective Pictet-Spengler reactions, enantioselective variants of the reaction can be catalyzed by Pictet-Spenglerase enzymes,⁸ or by Brønsted acid organocatalysis.⁹ Jacobsen, Seidel, List or Hiemstra have developed elegant enantioselective Pictet-Spengler reactions using chiral thioureas¹⁰ or chiral phosphoric acids¹¹ as catalysts, that have found numerous applications in total synthesis.^{11d, 12} Each of these methods have their strengths and weaknesses, either in the nature of the aldehyde used, the substituting pattern of the tryptamine or the electronic environment around the indole ring. However, in the context of such an important synthetic transformation, one must note that no enantioselective strategies using organometallic catalysis have been reported.

In the course of a project aiming at the synthesis of polycyclic indole derivatives,¹³ we used extensively the chiral phosphoric acid catalyzed Pictet-Spengler reaction to prepare key tetrahydro- β -carboline moieties that were cyclized to more complex polycyclic compounds using palladium.^{13a, 13b} or gold-catalysis.^{13c, 13d} During the study of the carboamination of *N*-allyltetrahydro- β -carboline to alkynes, we incidentally identified Au(I) complexes as efficient catalysts for the Pictet-Spengler reaction and developed a sequence combining this step with the carboamination cyclization (Scheme 2, eq. 1).^{13d} Precedence in the field was shown using achiral Au(I)¹⁴ or Au(III)¹⁵ complexes but it was however not clear from the literature the role that gold complexes could play in such a reaction, traditionally catalyzed by acids. In this paper, we demonstrate that chiral Au(I) complexes efficiently catalyze enantioselective Pictet-Spengler reactions (Scheme 2, eq. 2) and report our efforts in determining the scope of the reaction and understanding its mechanism.

Scheme 2. Gold-Catalyzed Synthesis of TH-β-Cs

We initiated this work by establishing a suitable catalytic system. After we checked that tryptamine itself did not react in these conditions,¹⁶ we prepared a series of tryptamines **1** with diverse substituents to test under different catalytic conditions. We first engaged *N*-benzyl tryptamine **1a** in a Pictet-Spengler reaction with benzaldehyde **2a** in the presence of the Gagos' catalyst $\text{Ph}_3\text{PAuNTf}_2$ **4**¹⁷ in DCM and molecular sieves,¹⁸ leading to the expected product **3aa** in 90% yield (Table 1, Entry 1). By anticipation that the use of chiral neutral Au(I) complexes necessitates a silver activation, we checked that the silver salt AgNTf_2 was not catalytically active (Entry 2). We next selected the DTBM-MeO-BIPHEP chiral ligand providing the digold complex $\text{L}_1(\text{AuCl})_2$ (3 mol%), using AgNTf_2 (5.8 mol%) as chloride abstractor and screened tryptamines **1a-f** (Entries 3-8). Tryptamines **1b** and **1c** bearing an allyl and a naphthylmethyl group led to moderate enantiomeric excesses (Entries 4 and 5), while all other tryptamines substituted by a benzyl-type protecting group led to an average 75% ee in good yields (Entries 3, 6-8). The best enantiomeric excess was obtained with a CH_2Mes group, leading to 83% ee (Entry 8). We further screened a series of chiral diphosphines (Entries 9-13),¹⁹ that identified (DM)-SEGPHOS L_4 as the best chiral ligand, delivering a 93% ee (Entry 11).²⁰

The optimal conditions were then applied to a series of aromatic aldehydes using $\text{L}_4(\text{AuCl})_2$ as the precatalyst and **1f** as the tryptamine partner on a 0.3 mmol scale (Scheme 3). Benzaldehyde afforded **3fa** in 97% yield and 92% ee. Aldehydes bearing a substituent in the *para* position led to excellent results with a good tolerance towards halide, alkyl and aryl substituents (**3fb** to **3ff**). Suitable crystals of compound **3fe** allowed to determine the absolute configuration of the stereogenic center as *R*.²¹ The presence of an electron-donating group led to a drop of the ee to 73% in **3fg**. Aldehyde **2h**, including a boronic acid function, led to **1fh** in 88% ee. The reaction performed from terephthalaldehyde **2i** led to **3fi** in an excellent 95% ee and 86% yield. Remarkably, there was no trace of difunctionalization in spite of the presence of a second aldehyde function. Finally, against all odds, the reaction tolerated the highly electrophilic nitro group, reputed reactive towards Au(I) complexes.²² The corresponding tetrahydro-β-carboline **3fj** was isolated in an excellent yield and with 72% ee. *Ortho*-substituted aromatic aldehydes were next screened. Halide substituents were found compatible with the established conditions, although a better ee was obtained with the less sterically demanding fluorine atom, leading to **3fl** in 87% ee. On the contrary, mesomeric electron-withdrawing groups such as cyanide or nitro resulted in lower ee's. Aldehydes substituted at the *meta* position by either halogens or trifluoromethyl groups resulted in excellent ee's, higher than 90% for **3fo-3fq**. Despite a significant drop of the ee to 61%, the reaction was compatible with an unprotected phenol function leading to

Table 1. Optimization of the Reaction^a

entry	1	catalyst	yield (%) ^b	ee ^c
1 ^d	1a	$\text{Ph}_3\text{PAuNTf}_2$ 4	3aa , 90	-
2	1a	AgNTf_2	3aa , 0	-
3	1a	$\text{L}_1(\text{AuCl})_2$	3aa , 99	75
4	1b	$\text{L}_1(\text{AuCl})_2$	3ba , 90	57
5	1c	$\text{L}_1(\text{AuCl})_2$	3ca , 50	52
6	1d	$\text{L}_1(\text{AuCl})_2$	3da , 88	74
7	1e	$\text{L}_1(\text{AuCl})_2$	3ea , 80	75
8	1f	$\text{L}_1(\text{AuCl})_2$	3fa , 84	83
9	1f	$\text{L}_2(\text{AuCl})_2$	3fa , 82	91
10	1f	$\text{L}_3(\text{AuCl})_2$	3fa , 32	80
11	1f	$\text{L}_4(\text{AuCl})_2$	3fa , quant.	93
12	1f	$\text{L}_5(\text{AuCl})_2$	3fa , 82	79
13	1f	$\text{L}_6(\text{AuCl})_2$	3fa , 99	78

^a Reactions were run using **1** (0.1 mmol), **2a** (0.2 mmol) in DCM (1 mL) at rt. ^b Isolated yields. ^c Determined by chiral HPLC. ^d No silver salt added.

compound **3fr** in 92% yield. When the reaction was performed with a dialdehyde (*m*-phthalaldehyde **2s**), **3fs** was isolated in an excellent 95% ee with, again, no trace of difunctionalization. In a similar manner, with 3-acetylbenzaldehyde **2t**, the gold-catalyzed functionalization proved totally selective to the aldehyde over the ketone function. Bicyclic aldehydes were next screened and it was shown that 2-naphthaldehyde **2u** or 6-quinolinecarboxaldehyde **2v** were well tolerated, leading to 89 and 74% ee's, respectively. Compound **3fw** was obtained with a moderate 51% ee from 1-naphthaldehyde. Interestingly though, its aza-analogous 4-quinolinecarboxaldehyde **2x** led to **2fx** in a moderate 34% yield but with an excellent 80% ee. Finally, the piperonaldehyde led to **3fy** in 73% ee, introducing 1,3-benzodioxole scaffold present in numerous indolic drugs such as tadalafil.²³ Finally aliphatic aldehydes were screened, pointing out the limitation of this methodology to aromatic aldehydes. Indeed, with 3-phenylpropanal **2z**, the corresponding tetrahydro-β-carboline **3fz** was obtained in a low 18% yield and 50% ee. Other tryptamines bearing either a substituent on the indole ring or another group on the nitrogen were next used under the optimized conditions (Scheme 4). The reaction tolerated a methyl and a methoxy group at position 5 of the indole ring using **2s** as the aldehyde leading to **3hs** and **3js** with good enantioselectivities.

Scheme 3. Scope of Aldehydes in the Gold-Catalyzed Enantioselective Pictet-Spengler Reaction with Tryptamine 1f

^a Reactions were run using 1f (0.3 mmol), 2 (0.6 mmol), powdered activated 3 Å molecular sieves (300 mg) in DCM (3 mL) at rt. ^b The product 3fa was obtained in 89% yield and 83% ee when the reaction was performed at a 1 mmol scale of tryptamine 1f.

Scheme 4. Additional Scope

When the reaction was performed from *N*-benzyltryptamine 3a, the product 3as was obtained in an excellent 86% ee which dropped to 65% ee using an allyl group. A *p*-methoxy substituent proved very similar to a benzyl group, leading to 3es in 86% ee and 67% yield.

The mechanism of the Pictet-Spengler reaction is controversial about the nature of the addition of the indole ring to the iminium *ii* (Scheme 5).²⁴ However, regardless of the exact mechanism for the cyclization, all known Pictet-Spengler reactions undergo initially the formation of an iminium *ii* from a hemiaminal *i* and a protic catalyst, this step generating water (Scheme 5, eq. 1). The mechanistic issue associated with the Au(I)-catalyzed version of this reaction stands in (1) the questionable formation of the iminium and water while the reaction is seemingly performed in the absence of protons and the role of the Au(I) complex during this step (Scheme 5, eq. 2) and (2) the nature of the intermediates. Our hypothesis is that coordination of the π -Lewis acid Au(I) complex to the indole ring from **A** via C2 modifies the acidity of the proton at C2, thereby allowing an intramolecular dehydration from intermediate **B** to **C** (Scheme 5, eq. 3).^{25,26} However very little is known about the actual auration of indoles, most of the examples being extremely specific.^{27,28} The catalytic cycle would then consist in the formation of the key iminium **C** from the tryptamine **1**, aldehyde **2** and the gold catalyst through intermediate **A**. The addition of the indole ring to the iminium would

occur via the direct addition of C2 to lead to **D** (or alternatively, the C3 addition). After a deauration step regenerating the catalytic species, the product would be obtained (Scheme 6).

Scheme 5. Putative and Proposed Mechanistic Steps for the Formation of the Key Iminium

Scheme 6. Mechanistic Proposal

Figure 1. Computed Dehydration Pathway and Final Steps to the Product

The Au(I)-catalyzed Pictet-Spengler reaction of tryptamine **1f** with benzaldehyde using AuPPh_3^+ as model catalyst was studied by means of DFT computations (see the Supporting Information for details and additional computations). The investigations started by the computational analysis of the initial steps of the reaction from the hemiaminal **B**. Here, we focus on the process of dehydration and the formation of the final product. Figure 1 shows the results obtained on the computed mechanism that was postulated in Scheme 6. All energies are Gibbs free energy at 298 K including solvent correction for dichloromethane. The hemiaminal complex **B**, located at -0.7 kcal/mol, displays the Au(I) fragment at the C2 position of the indole moiety. The dehydration process involving the C2-H bond to form iminium **C** requires 22.7 kcal/mol of free energy of activation, i.e. 22.0 kcal/mol from the reference system. Subsequently, intermediate **C** may form the spiro complex **E** via C3 attack to the iminium. This virtually thermoneutral spiroindolization requires only 3.3 kcal/mol of free energy of activation. It is thus easily reversible. Importantly, a C3 to C2 C-C bond migration that would form a 6-member ring **D** could not be modeled, which is consistent with previous observations in the field of organocatalyzed Pictet-Spengler reactions.²⁴ Alternatively, intermediate **C** can also be directly cyclized from the C2 position to give **D** at the expense of a moderate energy barrier of 4.8 kcal/mol. This step is appreciably exergonic by 16.6 kcal/mol. Finally, intermediate **D** releases the catalyst and the observed product, lying at -5.7 kcal/mol on the free energy surface.

In conclusion, we have developed a new route to chiral tetrahydro- β -carbolines with high enantioselectivities using Au(I) catalysis. Though limited to aromatic aldehydes, the reaction tolerates a large number of reactive functions with high chemoselectivity. The Au(I) complex serves as an activator by binding to the indole ring thereby allowing the formation of the key iminium with concomitant metalation at C2 of the indole ring. It is to bet that this new reactivity will pave the way to new strategies in the growing field of gold catalysis applied to heterocyclic chemistry.

ASSOCIATED CONTENT

Supporting Information

Details of the experimental procedures, NMR spectra, HPLC, spectroscopical and computational data. The Supporting Information is available free of charge on the ACS Publications website.

AUTHOR INFORMATION

Corresponding Authors

xavier.guinchard@cnsr.fr
vincent.gandon@u-psud.fr

ACKNOWLEDGMENT

N. Glinsky-Olivier thanks the MESRI and the CHARMMAT Laboratory of Excellence (ANR-11-LABX0039) for financial support. X. Guinchard thanks Dr. Angela Marinetti (ICSN) for her support. S. Yang thanks the CSC for PhD grant.

REFERENCES

- Pictet, A.; Spengler, T., Über die Bildung von Isochinolin derivaten durch Einwirkung von Methylal auf Phenyl-Ethylamin, Phenyl-alanin und Tyrosin. *Ber. Dtsch. Chem. Ges.* **1911**, *44*, 2030.
- Tatsui, G., Über die Synthese von Carbolinderivaten. *Yakugaku Zasshi* **1928**, *48*, 453-459.
- (a) Domínguez, G.; Pérez-Castells, J., Chemistry of β -Carbolines as Synthetic Intermediates. *Eur. J. Org. Chem.* **2011**, *2011*, 7243-7253. (b) Laine, A.; Lood, C.; Koskinen, A., Pharmacological Importance of Optically Active Tetrahydro- β -carbolines and Synthetic Approaches to Create the C1 Stereocenter. *Molecules* **2014**, *19*, 1544. (c) Cao, R.; Peng, W.; Wang, Z.; Xu, A., Beta-Carboline alkaloids: biochemical and pharmacological functions. *Curr. Med. Chem.* **2007**, *14*, 479-500.
- (a) Glinsky-Olivier, N.; Guinchard, X., Enantioselective Catalytic Methods for the Elaboration of Chiral Tetrahydro- β -carbolines and Related Scaffolds *Synthesis* **2017**, *49*, 2605-2620. (b) Rao, R. N.; Maiti, B.; Chanda, K., Application of Pictet-Spengler Reaction to Indole-Based Alkaloids Containing Tetrahydro- β -carboline Scaffold in Combinatorial Chemistry. *ACS Comb. Sci.* **2017**, *19*, 199-228.
- Uematsu, N.; Fujii, A.; Hashiguchi, S.; Ikariya, T.; Noyori, R., Asymmetric Transfer Hydrogenation of Imines. *J. Am. Chem. Soc.* **1996**, *118*, 4916-4917.
- (a) Espinoza-Moraga, M.; Petta, T.; Vasquez-Vasquez, M.; Laurie, V. F.; Moraes, L. A. B.; Santos, L. S., Bioreduction of β -carboline imines to amines employing *Saccharomyces bayanus*. *Tetrahedron: Asymmetry* **2010**, *21*, 1988-1992. (b) Mirabal-Gallardo, Y.; Soriano, M. d. P. C.; Santos, L. S., Stereoselective bioreduction of β -carboline imines through cell-free extracts from earthworms (*Eisenia foetida*). *Tetrahedron: Asymmetry* **2013**, *24*, 440-443. (c) Aleku, G. A.; Man, H.; France, S. P.; Leipold, F.; Hussain, S.; Toca-Gonzalez, L.; Marchington, R.; Hart, S.; Turkenburg, J. P.; Grogan, G.; Turner, N. J., Stereoselectivity and Structural Characterization of an Imine Reductase (IRED) from *Amicycolatopsis orientalis*. *ACS Catal.* **2016**, *6*, 3880-3889. (d) Leipold, F.; Hussain, S.; Ghislieri, D.; Turner, N. J., Asymmetric Reduction of Cyclic Imines Catalyzed by a Whole-

Cell Biocatalyst Containing an (S)-Imine Reductase. *ChemCatChem* **2013**, *5*, 3505-3508.

7. (a) Cox, E. D.; Cook, J. M., The Pictet-Spengler condensation - a new direction for an old reaction. *Chem. Rev.* **1995**, *95*, 1797-1842. (b) Stockigt, J.; Antonchick, A. P.; Wu, F. R.; Waldmann, H., The Pictet-Spengler Reaction in Nature and in Organic Chemistry. *Angew. Chem. Int. Ed.* **2011**, *50*, 8538-8564. (c) Lorenz, M.; Van Linn, M. L.; Cook, J. M., The Asymmetric Pictet-Spengler Reaction. *Curr. Org. Synth.* **2010**, *7*, 189.

8. Maresh, J. J.; Giddings, L. A.; Friedrich, A.; Loris, E. A.; Panjikar, S.; Trout, B. L.; Stockigt, J.; Peters, B.; O'Connor, S. E., Strictosidine Synthase: Mechanism of a Pictet-Spengler Catalyzing Enzyme. *J. Am. Chem. Soc.* **2008**, *130*, 710.

9. Kampen, D.; Reisinger, C. M.; List, B., Chiral Brønsted acids for asymmetric organocatalysis. *Top. Curr. Chem.* **2010**, *291*, 395-456.

10. (a) Taylor, M. S.; Jacobsen, E. N., Highly enantioselective catalytic acyl-Pictet-Spengler reactions. *J. Am. Chem. Soc.* **2004**, *126*, 10558-10559. (b) Klausen, R. S.; Jacobsen, E. N., Weak Brønsted Acid-Thiourea Co-catalysis: Enantioselective, Catalytic Protio-Pictet-Spengler Reactions. *Org. Lett.* **2009**, *11*, 887-890. (c) Mittal, N.; Sun, D. X.; Seidel, D., Conjugate-Base-Stabilized Brønsted Acids: Catalytic Enantioselective Pictet-Spengler Reactions with Unmodified Tryptamine. *Org. Lett.* **2014**, *16*, 1012-1015.

11. (a) Seayad, J.; Seayad, A. M.; List, B., Catalytic asymmetric Pictet-Spengler reaction. *J. Am. Chem. Soc.* **2006**, *128*, 1086-1087. (b) Wanner, M. J.; van der Haas, R. N. S.; de Cuba, K. R.; van Maarseveen, J. H.; Hiemstra, H., Catalytic asymmetric Pictet-Spengler reactions via sulphenyliminium ions. *Angew. Chem. Int. Ed.* **2007**, *46*, 7485-7487. (c) Sewgobind, N. V.; Wanner, M. J.; Ingemann, S.; de Gelder, R.; van Maarseveen, J. H.; Hiemstra, H., Enantioselective BINOL-phosphoric acid catalyzed Pictet-Spengler reactions of N-benzyltryptamine. *J. Org. Chem.* **2008**, *73*, 6405-6408. (d) Huang, D.; Xu, F.; Lin, X.; Wang, Y., Highly Enantioselective Pictet-Spengler Reaction Catalyzed by SPINOL-Phosphoric Acids. *Chem. Eur. J.* **2012**, *18*, 3148-3152.

12. (a) Mergott, D. J.; Zuend, S. J.; Jacobsen, E. N., Catalytic Asymmetric Total Synthesis of (+)-Yohimbine. *Org. Lett.* **2008**, *10*, 745. (b) Kerschgens, I. P.; Claveau, E.; Wanner, M. J.; Ingemann, S.; van Maarseveen, J. H.; Hiemstra, H., Total syntheses of mitragynine, paynantheine and speciogynine via an enantioselective thiourea-catalysed Pictet-Spengler reaction. *Chem. Commun.* **2012**, *48*, 12243-12245. (c) Piemontesi, C.; Wang, Q.; Zhu, J., Enantioselective Synthesis of (+)-Peganumine A. *J. Am. Chem. Soc.* **2016**, *138*, 11148-11151. (d) Raheem, I. T.; Thiara, P. S.; Peterson, E. A.; Jacobsen, E. N., Enantioselective Pictet-Spengler-Type Cyclizations of Hydroxylactams: H-Bond Donor Catalysis by Anion Binding. *J. Am. Chem. Soc.* **2007**, *129*, 13404-13405. (e) Badillo, J. J.; Silva-Garcia, A.; Shupe, B. H.; Fetting, J. C.; Franz, A. K., Enantioselective Pictet-Spengler reactions of isatins for the synthesis of spiroindolones. *Tetrahedron Lett.* **2011**, *52*, 5550-5553. (f) Duce, S.; Pescioli, F.; Gramigna, L.; Bernardi, L.; Mazzanti, A.; Ricci, A.; Bartoli, G.; Bencivenni, G., An Easy Entry to Optically Active Spiroindolinones: Chiral Brønsted Acid-Catalysed Pictet-Spengler Reactions of Isatins. *Adv. Synth. Catal.* **2011**, *353*, 860. (g) Wanner, M. J.; Boots, R. N. A.; Eradus, B.; de Gelder, R.; van Maarseveen, J. H.; Hiemstra, H., Organocatalytic Enantioselective Total Synthesis of (-)-Arboricine. *Org. Lett.* **2009**, *11*, 2579-2581. (h) Herle, B.; Wanner, M. J.; van Maarseveen, J. H.; Hiemstra, H., Total Synthesis of (+)-Yohimbine via an Enantioselective Organocatalytic Pictet-Spengler Reaction. *J. Org. Chem.* **2011**, *76*, 8907-8912.

13. (a) Gobé, V.; Guinchard, X., Stereoselective synthesis of chiral polycyclic indolic architectures through Pd0-catalyzed tandem deprotection/cyclization of tetrahydro-beta-carbolines on allenes. *Chem. Eur. J.* **2015**, *21*, 8511-8520. (b) Gobé, V.; Guinchard, X., Pd(0)-Catalyzed Tandem Deprotection/Cyclization of Tetrahydro-beta-carbolines on Allenes: Application to the Synthesis of Indolo[2,3-a]quinolizidines. *Org. Lett.* **2014**, *16*, 1924-1927. (c) Gobé, V.; Dousset, M.; Retaillieu, P.; Gandon, V.; Guinchard, X., Gold(I)-Catalyzed Carboaminations of Allenes by N-Allyl Tetrahydro-beta-carbolines: An Experimental and Theoretical study. *Adv. Synth. Catal.* **2016**, *358*, 3960-3965. (d) Gobé, V.; Retaillieu, P.; Guinchard, X., Self-Relay Gold(I)-Catalyzed Pictet-Spengler/Cyclization Cascade Reaction for the Rapid Elaboration of Pentacyclic Indole Derivatives. *Chem. Eur. J.* **2015**, *21*, 17587-17590. (e) Gobé, V.; Retaillieu, P.; Guinchard, X., Domino Formation of Enamines - Intramolecular Cyclizations to 1-Aminotetralins from gamma-Aryllallene Aldehydes and Amines. *Org. Lett.* **2014**, *16*, 5438-5441.

14. Subba Reddy, B. V.; Swain, M.; Reddy, S. M.; Yadav, J. S.; Sridhar, B., Gold-Catalyzed Domino Cycloisomerization/Pictet-Spengler Reaction of 2-(4-Aminobut-1-yn-1-yl)anilines with Aldehydes: Synthesis of Tetrahydropyrido[4,3-b]indole Scaffolds. *J. Org. Chem.* **2012**, *77*, 11355-11361.

15. Yoon, S. W., Development of the Pictet-Spengler Reaction Catalyzed by AuCl3/AgOTf. *J. Org. Chem.* **2006**, *71*, 2521-2523.

16. A tryptamine substituted by a 4-nitrobenzenesulfonyl group did not react either.

17. Mézailles, N.; Ricard, L.; Gagosz, F., Phosphine Gold(I) Bis-(trifluoromethanesulfonyl)imide Complexes as New Highly Efficient and Air-Stable Catalysts for the Cycloisomerization of Enynes. *Org. Lett.* **2005**, *7*, 4133-4136.

18. The addition of molecular sieves is necessary to ensure a high conversion. It was also checked that 3 Å molecular sieves leads to the best results throughout Table 1.

19. (a) Zi, W.; Toste, D. F., Recent advances in enantioselective gold catalysis. *Chem. Soc. Rev.* **2016**, *45*, 4567-4589. (b) Wang, Y.-M.; Lackner, A. D.; Toste, F. D., Development of Catalysts and Ligands for Enantioselective Gold Catalysis. *Acc. Chem. Res.* **2014**, *47*, 889-901. (c) Cera, G.; Bandini, M., Enantioselective Gold(I) Catalysis with Chiral Monodentate Ligands. *Isr. J. Chem.* **2013**, *53*, 848-855. (d) Pradal, A.; Toullec, P. Y.; Michelet, V., Recent Developments in Asymmetric Catalysis in the Presence of Chiral Gold Complexes. *Synthesis* **2011**, 1501-1514.

20. Further screening of the silver salt, the solvent, the stoichiometry of the aldehyde and of the catalyst did not improve this result. See the Supporting Information.

21. This R configuration was tentatively assigned similarly to all the compounds reported in this paper by structural similarity.

22. (a) Huplé, D. B.; Ghorpade, S.; Liu, R.-S., Recent Advances in Gold-Catalyzed N- and O-Functionalizations of Alkynes with Nitrones, Nitroso, Nitro and Nitroxy Species. *Adv. Synth. Catal.* **2016**, *358*, 1348-1367. (b) Liu, F.; Yu, Y.; Zhang, J., Highly Substituted Furo[3,4-d][1,2]oxazines: Gold-Catalyzed Regiospecific and Diastereoselective 1,3-Dipolar Cycloaddition of 2-(1-Alkynyl)-2-alken-1-ones with Nitrones. *Angew. Chem. Int. Ed.* **2009**, *48*, 5505-5508. (c) Jónsson, H. F.; Evjen, S.; Fiksdahl, A., Gold(I)-Catalyzed [2 + 2 + 2] Cyclotrimerization of 1,3-Diarylpargyl Acetals. *Org. Lett.* **2017**, *19*, 2202.

23. Daugan, A.; Grondin, P.; Ruault, C.; de Gouvello, A. C. L.; Coste, H.; Kirilovsky, J.; Hyafil, F.; Labaudiniere, R., The discovery of tadalafil: A novel and highly selective PDE5 inhibitor. 1: 5,6,11,11a-tetrahydro-1H-imidazo 1',5':1,6 pyrido 3,4-b indole-1,3(2H)-dione analogues. *J. Med. Chem.* **2003**, *46*, 4525-4532.

24. (a) Zheng, C.; Xia, Z.-L.; You, S.-L., Unified Mechanistic Understandings of Pictet-Spengler Reactions. *Chem* **2018**, *4*, 1952-1966. (b) Unsworth, W. P., Understanding the Role of Spiroindolenines in Pictet-Spengler Reactions. *Chem* **2018**, *4*, 1767-1770. (c) Gobé, V.; Gandon, V.; Guinchard, X., Reactions Involving Tryptamines and delta-Allenyl Aldehydes: Competition between Pictet-Spengler Reaction and Cyclization to 1-Aminotetralins. *Adv. Synth. Catal.* **2018**, *360*, 1280-1288.

25. A and B might well be resonance structures.

26. With the natural high nucleophilicity of the indole ring at C3, another hypothesis through the auration at C3 of the indole has been considered and is presented in the Supporting Information.

27. (a) Ahlsten, N.; Perry, G. J. P.; Cambeiro, X. C.; Boorman, T. C.; Larrosa, I., A silver-free system for the direct C-H auration of arenes and heteroarenes from gold chloride complexes. *Catal. Sci. Technol.* **2013**, *3*, 2892-2897. (b) Gimeno, A.; Rodríguez-Gimeno, A.; Cuenca, A. B.; Ramírez de Arellano, C.; Medio-Simón, M.; Asensio, G., Gold(I)-catalysed cascade reactions in the synthesis of 2,3-fused indole derivatives. *Chem. Commun.* **2015**, *51*, 12384-12387. (c) Preindl, J.; Jouvin, K.; Laurich, D.; Seidel, G.; Fürstner, A., Gold- or Silver-Catalyzed Syntheses of Pyrones and Pyridine Derivatives: Mechanistic and Synthetic Aspects. *Chem. Eur. J.* **2016**, *22*, 237-247. (d) Biasiolo, L.; Belpassi, L.; Gaggioli, C. A.; Macchioni, A.; Tarantelli, F.; Ciancaleoni, G.; Zuccaccia, D., Cyclization of 2-Alkynyldimethylaniline on Gold(I) Cationic and Neutral Complexes. *Organometallics* **2016**, *35*, 595-604. (e) Arcadi, A.; Bianchi, G.; Chiarini, M.; D'Anniballe, G.; Marinelli, F., Gold-Catalyzed Conjugate Addition Type Reaction of Indoles with alpha,beta-Unsaturated Ketones. *Synlett* **2004**, *2004*, 944-950. (f) Heffernan, S. J.; Tellam, J. P.; Queru, M. E.; Silvanus, A. C.; Benito, D.; Mahon, M. F.; Hennessy, A. J.; Andrews, B. L.; Carbery, D. R., Double Gold-Catalysed Annulation of Indoles by Enynes. *Adv.*

Synth. Catal. **2013**, 355, 1149-1159. (g) Cambeiro, X. C.; Ahlsten, N.; Larrosa, I., Au-Catalyzed Cross-Coupling of Arenes via Double C–H Activation. *J. Am. Chem. Soc.* **2015**, 137, 15636-15639. (h) Zeng, X.; Kinjo, R.; Donnadieu, B.; Bertrand, G., Serendipitous Discovery of the Catalytic Hydroammoniumation and Methylation of Alkynes. *Angew. Chem. Int. Ed.* **2010**, 49, 942-945.

28. Pirovano, V., Gold-Catalyzed Functionalization Reactions of Indole. *Eur. J. Org. Chem.* **2018**, 2018, 1925-1945.