1 **Assessing behavioural traits of benthic foraminifera: implications for**
2 **sediment mixing**

3 Noémie Deldicq^{1*}, Laurent Seuront^{2,3,4}, Dewi Langlet¹, Vincent M.P. Bouchet¹

4 ¹Univ. Lille, CNRS, Univ. Littoral Côte d'Opale, UMR 8187, LOG, Laboratoire d'Océanologie et de Géosciences,
5 F 62930, Wimereux, France.

6 ²CNRS, Univ. Lille, Univ. Littoral Côte d'Opale, UMR 8187, LOG, Laboratoire d'Océanologie et de Géosciences,
7 F 62930, Wimereux, France.

8 ³Department of Marine Resource and Energy, Tokyo University of Marine Science and Technology, 4-5-7 Konan,
9 Minato-ku, Tokyo 108-8477, Japan.

10 ⁴Department of Zoology and Entomology, Rhodes University, Grahamstown, 6140, South Africa.

11

12

13 *Corresponding author:

14 E-mail: noemie.deldicq@outlook.fr

15 Tel: 03 21 99 29 42

16

17

18

19

20

21

22

23

24

25 **Short title: Intertidal foraminifera behavioural traits**

26 **Abstract**

27 The assessment of behavioural traits of marine organisms is increasingly recognized as a key
28 issue to understand their role in ecosystem processes, such as bioturbation and nutrient cycling.
29 The movement ability of intertidal foraminifera suggest that they may have a role, yet to be
30 quantified, in benthic-pelagic coupling through their movement on the sediment, at the
31 sediment-water interface and within the sediment. In this context, we investigated the
32 behavioural traits of 5 benthic foraminiferal species typical of European temperate mudflats
33 under standardized trophic light and temperature conditions. Behavioural traits related to
34 motion of *Ammonia tepida*, *Haynesina germanica*, *Criboelphidium williamsoni*, *Miliammina*
35 *fusca* and *Quinqueloculina seminula* species were assessed through their travelled distance,
36 velocity, tortuosity of the path, position in the sediment and activity index. By analogy with
37 macrofauna bioturbation functional groups, we describe the studied foraminifera as
38 biodiffusors-species with 3 sub-groups defined according to their vertical position in the
39 sediment. *C. williamsoni* belongs to the epifaunal-biodiffusors group while *A. tepida* and *H.*
40 *germanica* belongs to surficial-biodiffusors. Finally, the species *Q. seminula* and *M. fusca* are
41 considered as gallery-biodiffusors species. Our results further suggest that features such as
42 velocity, activity and tortuosity may mediate sediment-mixing intensity. Therefore, *Q.*
43 *seminula*, *H. germanica* and *C. williamsoni* which are the most active species, would have
44 larger effect than the less active *A. tepida* and *M. fusca* on particles reworking rates. Taken
45 together, our results suggest that benthic foraminifera may play an underestimated role in
46 bioturbation processes.

47 **Keywords**

48 Benthic foraminifera – intertidal - motion behaviour – functional trait – bioturbators groups

49

50 1. Introduction

51 Trait-based studies have largely been implemented to describe ecosystem functioning,
52 especially over the last decade (Braeckman et al. 2010, Cardinale et al. 2011, Gothland et al.
53 2014). Specifically, traits — defined as “*the morphological, physiological or phenological*
54 *features measurable at the individual level, from the cell to the whole-organism level*” (Violle
55 et al. 2007) — can either be demographical (e.g. birth, mortality), biological (e.g. size, growth),
56 ecophysiological (e.g. nutrient assimilation, resource uptake) or behavioural (e.g. locomotion,
57 species interactions). These features determine the role of a species in the ecosystem; hence
58 allow defining functional traits (Violle et al. 2007, Gagic et al. 2015). For instance, a
59 comprehensive assessment of species-specific behavioural traits associated to locomotion is
60 one way to understand the role of species in the structure and functioning of coastal ecosystems.
61 Specifically in soft sediment environment, the behavioural traits related to faunal motion are
62 intrinsically considered as functional traits involved in bioturbation processes through the
63 displacement of sediments particles and the related enhancement of fluxes of both dissolved
64 and particulate materials (Mermillod-Blondin et al. 2004, Mermillod-Blondin & Rosenberg
65 2006, Kristensen et al. 2012, 2014). For instance, surface sediment displacement intensity is
66 directly linked to the type of activity (feeding, tube building) in the terebellid polychaete
67 *Eupolyornia nebulosa* (Maire et al. 2007c). Recent methodological improvements such as high-
68 frequency image analysis and automated acquisition help to quantify behavioural traits
69 associated to bioturbation, see Maire et al. (2008) for a review. Continuous observation of *Abra*
70 *ovata* showed that this bivalve’s behavioural activity is linked to sediment-mixing intensity
71 (Maire et al. 2007a). Depending on their behavioural traits, bioturbating species can induce
72 various changes on the benthic compartment such as the microbial community structure (Banta
73 et al. 1999, Marinelli et al. 2002, Papaspyrou et al. 2006) and in the biogeochemical reactions
74 occurring in the sedimentary column (Gutierrez & Jones 2006). Trait-based approaches hence

75 allow the description of different functional groups of macrofauna species such as gallery-
76 diffusors, biodiffusors, regenerator, upward- and downward-conveyors (François et al. 1997).
77 There is still, however, a limited knowledge on the role of the meiobenthic compartment in
78 bioturbation processes (see review in Schratzberger & Ingels 2017).

79 Meiofaunal taxa activity (nematodes and copepods) increase bacterial growth, oxygen
80 fluxes, denitrification and organic matter mineralisation (Aller & Aller 1992, Bradshaw et al.
81 2006, Middelburg & Meysman 2007, Nascimento et al. 2012, Piot et al. 2013, Bonaglia et al.
82 2014). Benthic foraminifera, despite their high abundance and ecological importance in the
83 marine meiobenthos (Murray 2006, Schönfeld et al. 2012), have essentially been ignored in
84 most studies dealing with meiofaunal bioturbation. Furthermore, little is known on
85 foraminiferal traits (e.g. habitats, metabolism, feeding modes, displacements), and only a few
86 studies quantified the motion behaviour of foraminifera (e.g. Kitazato 1981, 1988, Seuront &
87 Bouchet 2015, Jauffrais et al. 2016b). Hence, all benthic foraminiferal functions may not yet be
88 identified, particularly those involved in bioturbation processes. Benthic foraminifera can move
89 over relatively long distances (Seuront & Bouchet 2015) and their movements types (rotating,
90 spinning, crawling) are driven by both their number of pseudopods (Kitazato 1994) and the
91 morphology of their tests (i.e. spiral or quinqueloculine, Wetmore 1988). Foraminiferal motion
92 varies in intensity (Severin 1987, Gross 2002) and induces sediment displacements (Kitazato
93 1988, Hemleben & Kitazato 1995) that have so far only been quantified for deep-sea
94 communities (Gross 2002). These displacements create trails at the sediment surface (e.g.
95 *Quinqueloculina impressa*, *Criboelphidium excavatum*) and galleries (e.g. *Ammotium cassis*,
96 *Criboelphidium excavatum* subsp. *clavatum*) in the sediment (Richter 1964, Severin et al.
97 1982, Linke & Lutze 1993, Hemleben & Kitazato 1995, Bornmalm et al. 1997, Gross 2002).
98 Recently, Seuront and Bouchet (2015) showed a negative geotaxis in *Ammonia tepida* and a
99 positive geotaxis in both *C. excavatum* and *Haynesina germanica* suggesting that these species

100 move on and in the sedimentary column to colonize different microhabitats depending on their
101 ecological requirements (Wetmore 1988, Linke & Lutze 1993, Murray 2006). Hence, benthic
102 foraminiferal functional traits associated to motion behaviour may exist and be involved in
103 sediment mixing. This further suggests that the assessment of behavioural traits is an absolute
104 prerequisite to a comprehensive description of foraminiferal role in particulate fluxes at the
105 sediment-water interface. Yet, behavioural traits such as activity, motion intensity and vertical
106 position are poorly described in benthic foraminifera. In this context, the objectives of this study
107 are (i) to assess the horizontal and vertical dynamic of 5 intertidal foraminiferal species at the
108 sediment-water interface, (ii) to quantify the motion behaviour of these species and (iii) to use
109 the emergent vertical position and behavioural traits as a stepping stone to objectively classify
110 these species into functional groups. In intertidal mudflats from temperate environment, oxygen
111 penetration depth rarely reached 1 cm and foraminifera are mainly distributed in the 0-1 cm
112 layer (Geslin et al. 2011, Cesbron et al. 2016). This study hence specifically focused on
113 foraminifera living in the top 1st cm of sediment.

114 **2. Material and Methods**

115 *2.1. Sediment sampling and experimental set-up*

116 Surface sediment (0-1 cm) was collected from February to June 2018 at low tide in the
117 Authie Bay (50°22'20"N, 1°35'45"E), an intertidal mudflat located on the French coasts of the
118 English Channel for living benthic foraminifera. Sampled sediment was stored in plastic
119 containers (100 ml) and transported to the laboratory, where it was washed through a 125 µm
120 mesh-size sieve. Living benthic foraminifera were subsequently individually sorted with a
121 brush, identified and their pseudopodial activities checked with an inverted phase-contrast
122 microscope. Five intertidal species were considered in this study due to their high density at the
123 study site: the planispiral species *Haynesina germanica* and *Criboelphidium williamsoni*, the
124 trochospiral species *Ammonia tepida*, the agglutinated species *Miliammina fusca* and the

125 porcelainous species *Quinqueloculina seminula*. Recent molecular investigations showed that
126 the phylotypes *H. germanica* S16, *C. williamsoni* S1 and *A. tepida* T6 occur in the Authie Bay
127 (Schweizer, unpublished data). Individual sizes ranged from 300 to 800 µm in diameter.
128 Depending on the abundance at the time of sampling, 8 to 33 individuals per species were used
129 for the experimentation (Table 1). Before behavioural observations, individuals were kept
130 overnight 12h-long for acclimation under the experimental condition in a controlled-
131 temperature room at 18°C. Behavioural observations were performed in 300 ml aquaria filled
132 with 25-30ml defrost Authie-Bay sediment and 250 ml unfiltered eastern English-Channel
133 seawater (30 PSU). We used defrost sediment to make sure sediment was free of moving macro-
134 and meio-organisms (since nematodes, copepods and macrofaunal organisms are killed during
135 sediment freezing) and that only foraminiferal tracks could be observed at the sediment surface.
136 After the 12-h acclimation period, foraminifera were placed randomly at the surface of sediment
137 (Fig. 1). Note that the overlaying water was fully saturated in oxygen though a 10-min vigorous
138 air bubbling immediately before spreading living foraminifera on the sediment surface. At the
139 end of the 24h-experiment, dissolved-oxygen saturation was about 56% in the overlaying
140 seawater right above the sediment-water interface (HI9829 Multiparameter Meter, Hannah
141 instruments).

142 2.2. Quantification of behavioural traits

143 Individual displacements in and on the sediment were recorded by time-lapse photography
144 (1 image every 10 min for 24 hours) using a digital camera (Nikon V1 with a Nikkor 10–30
145 mm lens; Fig. 1). For each foraminifera, 144 images were combined in image-analysis software
146 Fiji to extract (x,y) coordinates using the Manual tracking plugin (Schindelin et al. 2019). The
147 behavioural traits of the above-mentioned species were investigated adapting the method
148 described in Seuront and Bouchet (2015). Five parameters were used to characterise the traits
149 of each study species.

150 First, the level of activity i.e. time allocated to motion by each individual, was estimated
 151 with the activity index A_i . This index is based on the ratio between the total time taken by one
 152 individual to move from its initial to its final position (t_{move}) and the time that the individual
 153 spent moving between these positions (t_{active}):

$$154 \quad A_i = 100 \times (t_{\text{active}} / t_{\text{move}})$$

155 The distance travelled by each individual between two images (*i.e.* 10 min) was calculated
 156 as:

$$157 \quad D_t = \sqrt{(x_t - x_{t+10})^2 + (y_t - y_{t+10})^2}$$

158 where (x_t, y_t) and (x_{t+10}, y_{t+10}) are the coordinates between two successive images taken at times
 159 t and $t+10$ min and the velocity (mm h^{-1}) of each individual was subsequently calculated
 160 considering the total distance travelled in 24 h.

161 The complexity (or tortuosity) of movement paths was assessed using fractal analysis. The
 162 fractal dimension measures the degree to which the trajectory fills the available space and is
 163 bounded between $D = 1$ for a line (*i.e.* the simplest instance of a trajectory) and $D = 2$ for a
 164 movement so complex that it actually fills the whole available space. The fractal dimensions of
 165 foraminifera trajectories were estimated using the box dimension method (see Seuront, 2010,
 166 2015 for reviews), which relies on the “ l cover” of the object, *i.e.* the number of boxes of length
 167 l required to cover the object. A more practical alternative is to superimpose a regular grid of
 168 boxes of length l on the object and count the number of boxes occupied by a subset of the object.
 169 This procedure is repeated using different values of l . The surface occupied by the trajectory of
 170 a foraminifera is then estimated using a series of counting boxes spanning a range of surfaces
 171 down to some small fraction of the entire surface. The number of occupied boxes increases with
 172 decreasing box size, leading to the following power-law relationship:

$$173 \quad N(l) \propto l^{-D_b}$$

174 where l is the box size, $N(l)$ is the number of boxes occupied by the trajectory, and D_b is the
175 box fractal dimension. The fractal dimension D_b is estimated from the slope of the linear trend
176 of the log-log plot of $N(l)$ versus l .

177 Finally, to assess the preferential position of the 5 studied species, the vertical position in
178 the sediment of each individual was determined for every picture based on a 3 depths categories
179 classification. When the test remained visible at the surface and the width of the path was
180 indistinguishable, the individual was considered to be crawling on the sediment surface (Fig.
181 2A,D). In turn, when the individual was burrowing into the sediment, its position was divided
182 in two categories: it was considered as moving at the sediment-water interface when half of the
183 test was visible (Fig. 2B,E) and to be burrowed in the sediment when a swelling at the sediment
184 surface was the only indication of the presence of the test in the sediment (Fig. 2C,F).

185 2.3. Statistical analyses

186 Because activity index, distance travelled, and velocity were non-normally distributed
187 (Kolmogorov-Smirnov test, $p < 0.05$), multiple comparisons between species were conducted
188 using the Kruskal-Wallis test, and a subsequent multiple comparison procedure based on the
189 Mann-Whitney test was eventually used to identify distinct group of measurements. The
190 presence of significant differences in fractal dimensions between species was assessed using an
191 analysis of covariance (Zar 2009). All statistical analyses were performed using © R.3.5.2.
192 software (R Core Team 2019).

193 3. Results

194 Among the 230 individuals picked for the experiment, 103 individuals were analysed as we
195 kept only those possible to track from the start to the end of the experiment (Table 1).

196 3.1. Activity index

197 *Criboelphidium williamsoni* was the most active species with a mean activity index at 89%
198 corresponding to 21/24h of displacement in the sediment (Fig. 3A, Table 1). This species

activity index is significantly larger ($p < 0.05$) than *Ammonia tepida*, *Haynesina germanica* and *Quinqueloculina seminula*, which are not significantly different from each other ($p < 0.05$). These 4 species have a significantly higher activity index than *Miliammina fusca* ($p < 0.05$) which was the less active species with a mean activity index at 50% (Fig. 3A).

3.2. Distance and velocity

Quinqueloculina seminula moved significantly faster ($p < 0.05$, Fig. 3C) and over longer distances (Fig. 3B) than the other. The four other species were discriminated in two other homogeneous groups with species moving slowly over short distances (*Miliammina fusca* and *Ammonia tepida*; Fig. 3B,C, Table 1) and species that moved at intermediate speed over intermediate distances (*Criboelphidium williamsoni* and *Haynesina germanica*; Fig. 3B,C, Table 1).

3.3. Tortuosity

Fractal dimensions D were in the range 1-1.3, indicating relatively linear trajectories. The analysis of covariance identified a group of 4 species (*Miliammina fusca*, *Ammonia tepida*, *Haynesina germanica* and *Quinqueloculina seminula*) moving significantly more tortuously than *Criboelphidium williamsoni* (Fig. 3D).

3.4. Vertical position

Miliammina fusca and *Quinqueloculina seminula* individuals were essentially observed (i.e. 100% and 70 to 90%, respectively) burrowed in the sediment throughout the experiment (Fig. 4A,B). Conversely, the vast majority (90-100%) of *Criboelphidium williamsoni* individuals remained on the sediment surface during the first three hours, before progressively moved to the sediment-water interface (10 to 100%) and were barely observed down into the sediment (Fig. 4C). *Haynesina germanica* gradually burrowed from the sediment surface down into the sediment with time to be essentially (75 to 80%) located into the sediment after 15 hours of observations (Fig. 4D). Finally, *Ammonia tepida* exhibited a slightly different temporal pattern.

First, a vast majority of them (95%) dug from the sediment surface down into the sediment in less than an hour, before progressively spreading out back up towards to sediment-water interface (Fig. 4E).

4. Discussion

This study is based on 24-h long observations assessing the behavioural traits of five species of benthic foraminifera to further understand their implication in sediment mixing. For instance, macrofaunal species behavioural traits affect the way they displace sediment particles (François et al. 1997, Maire et al. 2006, 2007a). It allowed for their classification in functional groups i.e. biodiffusors, regenerators or conveyors (François et al. 1997, Kristensen et al. 2012). The intensity of species activity is further directly linked to their life-mode which drives sediment mixing and bio-irrigation (Gérino et al. 2003, Gilbert et al. 2007, Maire et al. 2008, 2016, Kristensen et al. 2012). Considering that foraminifera fundamentally displace sediment over short distances due to their small size (Gross 2002, present work), we hereafter consider foraminifera as biodiffusors since they are “*organisms with activities that usually results in a constant and random local sediment bio-mixing over short distance*” (Kristensen et al. 2012).

More specifically, biodiffusors are typically divided in 3 subgroups (epifaunal-, gallery- and surficial-biodiffusors) depending on their life-mode in the sediment (Kristensen et al. 2012). In this context, our high-frequency image analysis of foraminiferal behaviour showed species-specific preferential depths of activity, that we used hereafter to classify the studied intertidal foraminiferal species in different biodiffusors functional subgroups (Fig. 5). Furthermore, motion-behaviour features (activity index, travelled distance, velocity and tortuosity) drive intra-functional group variability.

4.1. Foraminifera as biodiffusors

4.1.1. Epifaunal-biodiffusors

248 In our experiments, *Criboelphidium williamsoni* only moved on the sediment surface and
249 at the sediment-water interface. This observation is consistent with previous studies describing
250 this species as epifaunal (Allison et al. 2010) and reporting the highest density of the species in
251 the uppermost oxygenated sediment layers (Alve & Murray 2001, Bouchet et al. 2009). This
252 kleptoplastic species can host 10 times more active chloroplasts in its cytoplasm than other
253 temperate-water species (Lopez 1979). Kleptoplasty suggesting a preference for well-lighted
254 surface sediment, this species most likely have a surface-limited effect on the sediment mixing
255 and bio-irrigation. Consequently, *C. williamsoni* may be considered as an epifaunal-biodiffusor
256 (Fig. 5) which include “organisms that occur predominantly above the sediment–water
257 interface. Their activities are limited to near-surface sediments and generally redistribute fine
258 particles randomly over very short distances along the surface” (Kristensen et al. 2012).

259 4.1.2. Surficial-biodiffusors

260 *Haynesina germanica* and *Ammonia tepida* tended to avoid the surface sediment and both
261 preferred the sediment-water interface and burrowed position. Specifically, at the end of the
262 experiment, *A. tepida* individuals were evenly distributed between these two positions while *H.*
263 *germanica* ones preferred to be completely burrowed. These results are consistent with *in situ*
264 observations where *A. tepida* is found on and in the sediment (Goldstein et al. 1995, Bouchet et
265 al. 2009) while *H. germanica* mainly occur at the sediment surface (Alve & Murray 2001,
266 Bouchet et al. 2009). The latter can sequester photosynthetically active chloroplasts, which
267 might be used as food sources under low-light condition (Jauffrais et al. 2016a). Our results
268 suggest that both species could alternate between epifaunal and infaunal micro-habitats (Fig.
269 5). As a consequence, we classify these species in the surficial-biodiffusors group which are
270 “organisms with activities mostly restricted to the uppermost few centimetres of the sediment,
271 and these species rarely venture above the sediment–water interface” (Kristensen et al. 2012).

272 To be consistent with this definition, further assessments are needed to understand how deep
273 *H. germanica* and *A. tepida* can mix the sediment.

274 4.1.3. Gallery-biodiffusors

275 Quinqueloculine-shaped species (i.e. *Quinqueloculina seminula* and *Miliammina fusca*)
276 moved immediately from the surface down into the sediment layer and stayed buried during the
277 remaining time of experiment. These two species are hence clearly not restricted to the
278 sediment-water interface. *Quinqueloculina seminula* has, however, previously been described
279 as an epifaunal species (Di Bella et al. 2015, Martins et al. 2015) preferring oxic-zone (Moodley
280 et al. 1998), but is also known to be tolerant to low-oxygen events (Bernhard et al. 1997, Martins
281 et al. 2011, 2013, Langlet et al. 2014). Since in our 24-hours long observations specimens never
282 went back to the surface, these species may rather be considered as infaunal; with the ability to
283 create galleries in the sediment (Severin & Erskian 1981, Severin et al. 1982). As a
284 consequence, *Q. seminula* and *M. fusca* should be considered as gallery-biodiffusors (Fig. 5)
285 which are organisms “inducing diffusive local bio-mixing of particles and vertical transport of
286 particles from the upper regions of the sediment to the lower limit of burrow penetration”
287 (Kristensen et al. 2012).

288 4.2. Features explaining intra-functional group variability

289 Life traits such as individual size and foraging strategy affects role in species-specific
290 functional role in bioturbation processes (François et al. 1997, Gérino et al. 2003, Solan et al.
291 2004, Mermillod-Blondin 2011). For instance, macrofauna species displace a quantity of
292 sediment which is a direct proportion to their volumetric size (Dorgan et al. 2005). In our study,
293 quinqueloculine-shaped species (i.e. *Quinqueloculina seminula* and *Miliammina fusca*) are
294 species with the larger test volume suggesting that they would displace more sediment than the
295 other species investigated in the present work. However, effects of benthic fauna on fluxes at
296 the sediment-water interface depend on sediment reworking and bio-irrigation modes rather

297 than on the biogenic structure volume produced (Bouchet et al. 2009). Intensity of bioturbation
298 is hence a complex interplay between numerous life traits.

299 More specifically, the rate by which particles are physically moved also depends on feeding
300 strategies (Gérino et al. 2007). Tortuosity gives key information on that aspect (Pyke 1984; Bell
301 1991). In our study, the five species exhibited a relatively low tortuosity value. It implies that
302 they explored their environment extensively with close-to-linear trajectory. Such extensive
303 search strategy (or transecting e.g. Bell 1991) in the case of the 5 studied species, is optimal
304 under spatially-located food patches (Pyke 1984, Seuront & Stanley 2014, Seuront & Bouchet
305 2015). This foraging strategy is consistent with what is known of the trophic ecology of the five
306 species used in this study e.g. microphytobenthos, bacteria, metazoans (Nomaki et al. 2008,
307 Dupuy et al. 2010, Jauffrais et al. 2016b, Chronopoulou et al. 2019). For instance, *Haynesina*
308 *germanica* and *Ammonia tepida* are both herbivorous, feeding on benthic diatoms (Ward et al.
309 2003, Pascal et al. 2008). Recently, an in situ study showed that the feeding behaviour of
310 intertidal benthic foraminifera is more complex than what is observed experimentally
311 (Chronopoulou et al. 2019). Intertidal foraminifera exhibited clear varied and species-specific
312 trophic behaviour and were actually able to feed on different food sources. Specifically, *A.*
313 *tepida* may prefer to feed on algae but it is actually also able to feed on bacteria (Pascal et al.
314 2008). Since microphytobenthos, bacteria and metazoans microscale (i.e. centimetre-scale)
315 distributions are extremely patchy (Pinckney & Sandulli 1990, Danovaro et al. 2001, Spilmont
316 et al. 2011), all the studied species would benefit from the extensive search strategy to optimize
317 their likelihood to locate food patches. Such extensive displacement tends to maximize
318 sediment mixing (Seuront 2010, Viswanathan et al. 2011). However, species feeding strategy
319 has not been investigated throughout our experiment since we used homogenised defrost
320 surface-sediment therefore containing labile organic-matter, living bacteria, dead macro- and
321 meio-faunal organisms and the non-filtered overlaying contains microalgae. To further assess

a supprimé: is

a supprimé: This behavioural strategy is consistent with an optimal foraging strategy.

325 whether foraging-strategy would affect sediment-mixing rate, motion-behaviour of
326 foraminiferal species in a patchy and homogeneous controlled-food-condition needs to be
327 assess.

328 The five studied species showed significant differences between their respective activity
329 index, travelled distance and velocity. Within the functional groups above mentioned, species
330 may not have the same sediment mixing intensity and this intensity may most likely depend on
331 the detailed properties of species-specific motion behaviour. Within the gallery-biodiffusors,
332 *Quinqueloculina seminula* was the second most active species and travelled the longest
333 distances, while *Miliammina fusca* was the less active species and over lowest distances.
334 Similarly, the surficial-biodiffusor *Haynesina germanica* was more active and moved further
335 than *Ammonia tepida*. As a consequence, *Q. seminula* and *H. germanica* may rework a larger
336 volume of sediment. Furthermore, irrespective of their functional group, the most active species
337 (i.e. *Criboelphidium williamsoni*, *Q. seminula* and *H. germanica*) may contribute more to
338 sediment mixing than the less active species such as *A. tepida* and *M. fusca* (Fig. 5).

339 Conclusion

340 Based on the monitoring of species-specific behavioural traits this study assigned for the
341 first time intertidal benthic foraminiferal species into the biodiffusors-functional group. Hence,
342 the preferential depth of activity being fundamentally the prime feature allowing differentiating
343 between species. Secondary features like (i) size, (ii) feeding-modes, (iii) activity index, (iv)
344 travelled distance (v) velocity and (vi) tortuosity would most likely mediate the intensity of
345 bioturbation and explain the intra-functional group differences. To definitively validate our
346 classification, further work is needed on the quantification of sediment reworking rate of the
347 five studied species. Furthermore, our observations showed that foraminifera create physical
348 disturbances at the sediment surface, previously described as sediment pellets (Chandler 1989).
349 Such a pelletised surface layer created by benthic organisms activity can ease the resuspension

of sediments by tidal currents (Davis 1993, Willows et al. 1998, Orvain et al. 2003, 2004) and affect sediment bio-irrigation. These physical and chemical changes will affect microbial communities (Bertics & Ziebis 2009, Piot et al. 2013), organic matter mineralization and nutrient cycle (Gilbertson et al. 2012, Aller 2014). Meiofaunal species activity e.g. copepods, nematodes and foraminifera will consequently increase organic matter and NO_x fluxes which strongly affect benthic-pelagic coupling and therefore ecosystem functions (Danovaro et al. 2008, de Goeij et al. 2013). This suggest that foraminifera might play a underestimated role on sediment cohesiveness and dissolved-elements benthic fluxes (Schratzberger & Ingels 2017). This study supports the fact that, in a context of biodiversity change, assessing life traits of benthic foraminifera is critically needed to understand their role in ecosystem functioning.

360

361 **Acknowledgements**

The authors thank the Région Hauts-de-France, the Ministère de l'Enseignement Supérieur et de la Recherche and the European fund for regional economic development for their financial support through the attribution of the CPER research project CLIMIBIO. Noémie Deldicq PhD fellowship is funded by the Ministère de l'Enseignement Supérieur et de la Recherche. Dewi Langlet Post-doctoral fellowship is supported by the STARS research project COFFEE of the Région Hauts-de-France and the CPER research project CLIMIBIO. Molecular investigations of spiral-shaped foraminifera phylotype were done within the research project AMTEP, funded by the CNRS-INSU program EC2CO.

370 **References**

- 371 Aller RC (2014) Sedimentary diagenesis, depositional environments and benthic fluxes. In
372 Holland HG & Turekian KK (eds) Treatise on Geochemistry (2nd end). Elsevier p 293-
373 334.
374 Aller RC, Aller JY (1992) Meiofauna and solute transport in marine muds. *Limnol Oceanogr*
375 37:1018–1033.
376 Allison N, Austin W, Paterson D, Austin H (2010) Culture studies of the benthic foraminifera
377 *Elphidium williamsoni*: evaluating pH, $\Delta[\text{CO}_3^{2-}]$ and inter-individual effects on test
378 Mg/Ca. *Chem Geol* 274:87–93.

379 Alve E, Murray JW (2001) Temporal variability in vertical distributions of live (stained)
 380 intertidal foraminifera, Southern England. *J Foraminif Res* 31:12–24.
 381 Banta GT, Holmer M, Jensen MH, Kristensen E (1999) Effects of two polychaete worms,
 382 *Nereis diversicolor* and *Arenicola marina*, on decomposition in an organic-poor and an
 383 organic-enriched marine sediment. *Aquat Microb Ecol* 19:189–204.
 384 Bell WJ (1991) Searching behaviour: the behavioural ecology of finding resources. In:
 385 Springer, Berlin.
 386 Bernhard JM, Sen Gupta BK, Borne PF (1997) Benthic foraminiferal proxy to estimate dysoxic
 387 bottom-water oxygen concentrations: Santa Barbara basin, US Pacific continental margin.
 388 *J Foraminif Res* 27:301–310.
 389 Bertics VJ, Ziebis W (2009) Biodiversity of benthic microbial communities in bioturbated
 390 coastal sediments in controlled by geochemical microniches. *ISME J* 3:1269–1285.
 391 Bonaglia S, Nascimento FJA, Bartoli M, Klawonn I, Brüchert V (2014) Meiofauna increases
 392 bacterial denitrification in marine sediments. *Nat Commun* 5:5133.
 393 Bornmalm L, Corliss BH, Tedesco K (1997) Laboratory observations of rates and patterns of
 394 movement of continental margin benthic foraminifera. *Mar Micropaleontol* 29:175–184.
 395 Bouchet VMP, Sauriau PG, Debenay JP, Mermillod-Blondin F, Schmidt S, Amiard JC, Dupas
 396 B (2009) Influence of the mode of macrofauna-mediated bioturbation on the vertical
 397 distribution of living benthic foraminifera: first insight from axial tomodesitometry. *J*
 398 *Exp Mar Bio Ecol* 371:20–33.
 399 Bradshaw C, Kumblad L, Fagrell A (2006) The use of tracers to evaluate the importance of
 400 bioturbation in remobilising contaminants in Baltic sediments. *Estuar Coast Shelf Sci*
 401 66:123–134.
 402 Braeckman U, Provoost P, Gribsholt B, Gansbeke D Van, Middelburg JJ, Soetaert K, Vincx M,
 403 Vanaverbeke J (2010) Role of macrofauna functional traits and density in biogeochemical
 404 fluxes and bioturbation. *Mar Ecol Prog Ser* 399:173–186.
 405 Cardinale BJ, Matulich KL, Hooper DU, Byrnes JE, Duffy E, Gamfeldt L, Balvanera P, Connor
 406 MIO, Gonzalez A (2011) The functional role of producer diversity in ecosystems. *Am J*
 407 *Bot* 98:572–592.
 408 Cesbron F, Geslin E, Jorissen FJ, Delgard ML, Charrieau L, Deflandre B, Jézéquel D, Anschutz
 409 P, Metzger E (2016) Vertical distribution and respiration rates of benthic foraminifera:
 410 contribution to aerobic remineralization in intertidal mudflats covered by *Zostera noltei*
 411 meadows. *Estuar Coast Shelf Sci* 179:23–38.
 412 Chandler GT (1989) Foraminifera may structure meiobenthic communities. *Oecologia* 81:354–
 413 360.
 414 Chronopoulou PM, Salonen L, Bird C, Reichard GJ, Koho KA (2019) Metabarcoding insights
 415 into the tropic behaviour and identity of intertidal benthic foraminifera. *Front Microbiol*
 416 10:1169.
 417 Danovaro R, Armeni M, Dell’Anno A, Fabiano M, Manini E, Marrale D, Pusceddu A, Vanucci
 418 S (2001) Small-scale distribution of bacteria, enzymatic activities, and organic matter in
 419 coastal sediments. *Microb Ecol* 42: 177–185.
 420 Danovaro R, Gambi C, Dell’Anno A, Corinaldesi C, Fraschetti S, Vanreusel A, Vincx M,
 421 Gooday AJ (2008) Exponential decline of deep-sea ecosystem functioning linked to
 422 benthic biodiversity loss. *Curr Biol* 18:1–8.
 423 Davis R (1993) The role of bioturbation in sediment resuspension interaction with physical
 424 shearing. *J Exp Mar Bio Ecol* 171:187–200.
 425 de Goeij JM, van Oevelen D, Vermeij MJA, Osinga R, Middelburg JJ, de Goeij AFPM,
 426 Admiraal W (2013) Surviving in a marine desert: the sponge loop retains resources within
 427 coral reefs. *Science* 342:108–110.
 428 Di Bella L, Frezza V, Conte AM, Chiocci FL (2015) Benthic foraminiferal assemblages in

active volcanic area of the Azores Islands (North Atlantic Ocean). *Ital J Geosci* 134:50–59.

Dorgan KM, Jumars PA, Johnson B, Boudreau BP, Landis E (2005) Burrow extension by crack propagation. *Nature* 433:475.

Dupuy C, Rossignol L, Geslin E, Pascal PY (2010) Predation of mudflat meio-macrofaunal metazoans by a calcareous foraminifer, *Ammonia tepida* (Cushman, 1926). *J Foraminifer Res* 40:305–312.

François F, Poggiale JC, Durbec JP, Stora G (1997) A new approach for the modelling of sediment reworking induced by a macrobenthic community. *Acta Biotheor* 45:295–319.

Gagic V, Bartomeus I, Jonsson T, Taylor A, Winqvist C, Fischer C, Slade EM, Steffan-Dewenter I, Emmerson M, Potts SG, Tschardt T, Weisser W, Bommarco R (2015) Functional identity and diversity of animals predict ecosystem functioning better than species-based indices. *Proc R Soc B Biol Sci* 282:20142620.

Gérino M, Stora G, François-Carcaillet F, Gilbert F, Poggiale JC, Mermillod-Blondin F, Desrosiers G, Vervier P (2003) Macro-invertebrate functional groups in freshwater and marine sediments: a common mechanistic classification. *Vie Milieu* 53:221–231.

Gérino M, Frignani M, Mugnai C, Bellucci LG, Prevedelli D, Valentini A, Castelli A, Sauvage S (2007) Quantification of bioturbation processes and method to identify related functional groups. *Acta Oecol* 33:14–25.

Geslin E, Risgaard-Petersen N, Lombard F, Metzger E, Langlet D, Jorissen F (2011) Oxygen respiration rates of benthic foraminifera as measured with oxygen microensors. *J Exp Mar Biol Ecol* 396: 108–114.

Gilbert F, Hulth S, Grossi V, Poggiale JC, Desrosiers G, Rosenberg R, Gérino M, François-Carcaillet F, Michaud E, Stora G (2007) Sediment reworking by marine benthic species from the Gullmar Fjord (Western Sweden): importance of faunal biovolume. *J Exp Mar Bio Ecol* 348:133–144.

Gilbertson WW, Solan M, Prosser JI (2012) Differential effects of microorganism-invertebrate interactions in benthic nitrogen cycling. *FEMS Microbiol Ecol* 82:11–22.

Goldstein ST, Watkins GT, Kuhn RM (1995) Microhabitats of salt marsh foraminifera: St. Catherines Island, Georgia, USA. *Mar Micropaleontol* 26:17–29.

Gothland M, Dauvin JC, Denis L, Dufossé F, Jobert S (2014) Biological traits explain the distribution and colonisation ability of the invasive shore crab *Hemigrapsus takanoi*. *Estuar Coast Shelf Sci* 142:41–49.

Gross O (2002) Sediment interactions of foraminifera: implications for food degradation and bioturbation processes. *J Foraminifer Res* 32:414–424.

Gutierrez JL, Jones CG (2006) Physical ecosystem engineers as agent of biogeochemical heterogeneity. *Bioscience* 56:227–236.

Hemleben C, Kitazato H (1995) Deep-sea foraminifera under long time observation in the laboratory. *Deep Res I* 42:827–832.

Jauffrais T, Jesus B, Metzger E, Mouget JL, Jorissen F, Geslin E (2016a) Effect of light on photosynthetic efficiency of sequestered chloroplasts in intertidal benthic foraminifera (*Haynesina germanica* and *Ammonia tepida*). *Biogeosciences* 13:2715–2726.

Jauffrais T, Jesus B, Geslin E, Briand F, Martin V (2016b) Locomotion speed of the benthic foraminifer *Ammonia tepida* exposed to different nitrogen and carbon sources. *J Sea Res* 118:52–58.

Kitazato H (1981) Observation of behaviour and mode of life of benthic foraminifera in laboratory. *Geoscience Rep, Shizuoka University* 6:61–71.

Kitazato H (1988) Locomotion of some benthic foraminifera in and on sediments. *J Foraminifer Res* 18:344–349.

Kitazato H (1994) Foraminiferal microhabitats in four marine environments around Japan. *Mar*

479 Micropaleontol 24:29–41.

480 Kristensen E, Delefosse M, Quintana CO, Flindt MR, Valdemarsen T (2014) Influence of
481 benthic macrofauna community shifts on ecosystem functioning in shallow estuaries.
482 Front Mar Sci 1:1–14.

483 Kristensen E, Penha-Lopes G, Delefosse M, Valdemarsen T, Quintana CO, Banta GT (2012)
484 What is bioturbation? the need for a precise definition for fauna in aquatic sciences. Mar
485 Ecol Prog Ser 446:285–302.

486 Langlet D, Baal C, Geslin E, Metzger E, Zuschin M, Riedel B, Risgaard-Petersen N,
487 Stachowitsch M, Jorissen FJ (2014) Foraminiferal species responses to in situ,
488 experimentally induced anoxia in the Adriatic Sea. Biogeosciences 11:1775–1797.

489 Linke P, Lutze GF (1993) Microhabitat preferences of benthic foraminifera—a static concept or
490 a dynamic adaptation to optimize food acquisition? Mar Micropaleontol 20:215–234.

491 Lopez E (1979) Algal chloroplasts in the protoplasm of three species of benthic foraminifera:
492 taxonomic affinity, viability and persistence. Mar Biol 53:201–211.

493 Maire O, Duchêne JC, Rosenberg R, De Mendonça JB, Grémare A (2006) Effects of food
494 availability on sediment reworking in *Abra ovata* and *A. nitida*. Mar Ecol Prog Ser
495 319:135–153.

496 Maire O, Duchêne JC, Grémare A, Malyuga VS, Meysman FJR (2007a) A comparison of
497 sediment reworking rates by the surface deposit-feeding bivalve *Abra ovata* during
498 summertime and wintertime, with a comparison between two models of sediment
499 reworking. J Exp Mar Bio Ecol 343:21–36.

500 Maire O, Duchêne JC, Amouroux JM, Grémare A (2007c) Activity patterns in the terebellid
501 polychaete *Eupolyornia nebulosi* assessed using a new image analysis system. Mar Biol
502 151:737–749.

503 Maire O, Lecroart P, Meysman F, Rosenberg R, Duchêne JC, Grémare A (2008) Quantification
504 of sediment reworking rates in bioturbation research: a review. Aquat Biol 2:219–238.

505 Maire O, Barras C, Gestin T, Nardelli MP, Romero-Ramirez A, Duchêne JC, Geslin E (2016)
506 How does macrofaunal bioturbation influence the vertical distribution of living benthic
507 foraminifera? Mar Ecol Prog Ser 561:83–97.

508 Marinelli RL, Lovell CR, Wakeham SG, Ringelberg DB, White DC (2002) Experimental
509 investigation of the control of bacterial community composition in macrofaunal burrows.
510 Mar Ecol Prog Ser 235:1–13.

511 Martins MVA, Yamashita C, Sousa SHM, Martins P, Laut LLM, Figueira RCL, Mahiques MM,
512 Ferreira da Silva E, Alveirinho Dias JM, Rocha F (2011) The response of benthic
513 foraminifera to pollution and environmental stress in Ria de Aveiro (N Portugal). J Iber
514 Geol 37:231–246.

515 Martins MVA, Frontalini F, Tramonte KM, Figueira RCL, Miranda P, Sequeira C, Fernández-
516 Fernández S, Dias JA, Yamashita C, Renó R, Laut LLM, Silva FS, Rodrigues MAC,
517 Bernardes C, Nagai R, Sousa SHM, Mahiques M, Rubio B, Bernabeu A, Rey D, Rocha F
518 (2013) Assessment of the health quality of Ria de Aveiro (Portugal): heavy metals and
519 benthic foraminifera. Mar Pollut Bull 70:18–33.

520 Martins MVA, Zaaboub N, Aleya L, Frontalini F, Pereira E, Miranda P, Mane M, Rocha F,
521 Laut LLM, El Bour M (2015) Environmental quality assessment of Bizerte lagoon
522 (Tunisia) using living foraminifera assemblages and a multiproxy approach. PloS One
523 10:e0137250.

524 Mermillod-Blondin F (2011) The functional significance of bioturbation and biodeposition on
525 biogeochemical processes at the water–sediment interface in freshwater and marine
526 ecosystems. J North Am Benthol Soc 30:770–778.

527 Mermillod-Blondin F, Rosenberg R (2006) Ecosystem engineering: the impact of bioturbation
528 on biogeochemical processes in marine and freshwater benthic habitats. Aquat Sci 68:434–

442.

- Mermillod-Blondin F, Rosenberg R, François-Carcaillet F, Norling K, Mauclaire L (2004) Influence of bioturbation by three benthic infaunal species on microbial communities and biogeochemical processes in marine sediment. *Aquat Microb Ecol* 36:271–284.
- Middelburg JJ, Meysman FJR (2007) Burial at sea. *Science* 316:1294–1295.
- Moodley L, Van Der Zwaan GJ, Rutten GMW, Boom RCE, Kempers AJ (1998) Subsurface activity of benthic foraminifera in relation to porewater oxygen content: laboratory experiments. *Mar Micropaleontol* 34:91–106.
- Murray JW (2006) Ecology and applications of benthic foraminifera. In: Cambridge University Press, Cambridge.
- Nascimento FJA, Näslund J, Elmgren R (2012) Meiofauna enhances organic matter mineralization in soft sediment ecosystems. *Limnol Oceanogr* 57:338–346.
- Nomaki H, Ogawa NO, Ohkouchi N, Suga H, Toyofuku T, Shimanaga M, Nakatsuka T, Kitazato H (2008) Benthic foraminifera as trophic links between phytodetritus and benthic metazoans: carbon and nitrogen isotopic evidence. *Mar Ecol Prog Ser* 357:153–164.
- Orvain F, Le Hir P, Sauriau PG (2003) A model of fluff layer erosion and subsequent bed erosion in the presence of the bioturbator, *Hydrobia ulvae*. *J Mar Res* 61:823–851.
- Orvain F, Sauriau PG, Sygut A, Joassard L, Le Hir P (2004) Interacting effects of *Hydrobia ulvae* bioturbation and microphytobenthos on the erodibility of mudflat sediments. *Mar Ecol Prog Ser* 278:205–223.
- Papasprou S, Kristensen E, Cox RP (2006) Microbial reaction rates and bacterial communities in sediment surrounding burrows of two nereidid polychaetes (*Nereis diversicolor* and *N. virens*). *Mar Biol* 148:541–550.
- Pascal PY, Dupuy C, Richard P, Niquil N (2008) Bacterivory in the common foraminifer *Ammonia tepida*: isotope tracer experiment and the controlling factors. *J Exp Mar Biol Ecol* 359: 55–61.
- Pinckney J, Sandulli R (1990) Spatial autocorrelation analysis of meiofaunal and microalgal populations on an intertidal sandflat: scale linkage between consumers and resources. *Estuar Coast Shelf Sci* 30: 341–353.
- Piot A, Nozais C, Archambault P (2013) Meiofauna affect the macrobenthic biodiversity–ecosystem functioning relationship. *Oikos* 123:1–11.
- Pyke GH (1984) Optimal foraging theory: a critical review. *Annu Rev Ecol Syst* 15:523–575.
- R Core Team (2018) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- Richter G (1964) Zur ökologie der Foraminiferida II: lebensraum und lebenweise von Nonion depressulum, *Elphidium excavatum* und *Elphidium selseyense*. *Nat Mus* 94:421–430.
- Schindelin J, Arganda-carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A (2019) Fiji: an open-source platform for biological-image analysis. *Nat Methods* 9:676–682.
- Schönfeld J, Alve E, Geslin E, Jorissen F, Korsun S, Spezzaferri S, Abramovich S, Almogi-Labin A, Armynot du Châtelet E, Barras C, Bergamin L, Bicchi E, Bouchet V, Cearreta A, Di Bella L, Dijkstra N, Disaro ST, Ferraro L, Frontalini F, Gennari G, Golikova E, Haynert K, Hess S, Husum K, Martins V, McGann M, Oron S, Romano E, Sousa SM, Tsujimoto A (2012) The FOBIMO (FORaminiferal Blo-MONitoring) initiative-towards a standardised protocol for soft-bottom benthic foraminiferal monitoring studies. *Mar Micropaleontol* 94–95:1–13.
- Schratzberger M, Ingels J (2017) Meiofauna matters: the roles of meiofauna in benthic ecosystems. *J Exp Mar Bio Ecol* 502:1–14.
- Seuront, L (2010) Fractals and multifractals in ecology and aquatic science. In: CRC Press,

Boca Raton, Florida. Seuront L (2010) How does salinity influence the swimming speed
 of the estuarine calanoid copepod *Eurytemora affinis*? J Plankton Res 32:1223-1225.
 Seuront L (2015) On uses, misuses and potential abuses of fractal analysis in zooplankton
 behavioral studies: A review, a critique and a few recommendations. Physica A 432:410–
 434.
 Seuront L, Stanley HE (2014) Anomalous diffusion and multifractality optimises mating
 encounters in the ocean. P Natl Acad Sci USA 111:2206-2211.
 Seuront L, Bouchet VMP (2015) The devil lies in details: new insights into the behavioural
 ecology of intertidal foraminifera. J Foraminifer Res 45:390–401.
 Severin KP (1987) Laboratory observations of the rate of subsurface movement of a small
 miliolid foraminifer. J Foraminifer Res 17:110-116.
 Severin KP, Erskian MG (1981) Laboratory experiments on the vertical movement of
Quinqueloculina impressa Reuss through sand. J Foraminifer Res 11:133–136.
 Severin KP, Culver SJ, Blanpied C (1982) Burrows and trails produced by *Quinqueloculina*
impressa Reuss, a benthic foraminifer, in fine-grained sediment. Sedimentology 29:897–
 901.
 Solan M, Wigham BD, Hudson IR, Kennedy R, Coulon CH, Norling K, Nilsson HC, Rosenberg
 R (2004) In situ quantification of bioturbation using time-lapse fluorescent sediment
 profile imaging (f-SPI), luminophore tracers and model simulation. Mar Ecol Prog Ser
 271:1–12.
 Spilmont N, Seuront L, Meziane T, Welsh D (2011) There's more to the picture than meets the
 eye: sampling microphytobenthos in a heterogeneous environment. Estuar Coast Shelf Sci
 95: 470-476.
 Violle C, Navas M, Vile D, Kazakou E, Fortunel C (2007) Let the concept of trait be functional!
 Oikos 116:882–892.
 Viswanathan GM, da Luz MGE, Raposo EP, Stanley HE (2011) The physics of foraging: an
 introduction to random searching and biological encounters. In: Cambridge University
 Press, Cambridge.
 Ward JN, Pond DW, Murray JW (2003) Feeding of benthic foraminifera on diatoms and
 sewage-derived organic matter: an experimental application of lipid biomarker techniques.
 Mar Environ Research 56: 515-530.
 Wetmore K (1988) Burrowing and sediment movement by benthic foraminifera, as shown by
 time-lapse cinematography. Rev Paléobiologie 2:921–927.
 Willows RI, Widdows J, Wood RG (1998) Influence of an infaunal bivalve on the erosion of
 an intertidal cohesive sediment: a flume and modeling study. Limnol Oceanogr 43:1332–
 1343.
 Zar J (2009) Biostatistical analysis, fifth edition. In: Pearson, Upper Saddle River, New Jersey,
 USA.

619 **Table captions**

620 Table 1. Number N of individuals of each species used for experimentations and the subsequent
621 number n of individuals for which (x,y) coordinates were extracted with mean ($\bar{X}$) and standard
622 deviation (σ) for each parameters.

Species		<i>Q. seminula</i>	<i>H. germanica</i>	<i>C. williamsoni</i>	<i>A. tepida</i>	<i>M. fusca</i>
Total individual	N	30	90	30	60	20
Analysed individual	n	16	30	16	33	8
Distance (mm)	$\bar{X}$	88.25	51.51	44.18	17.29	14.06
	σ	32.95	19.63	13.50	5.94	4.67
Velocity (mm.h ⁻¹)	$\bar{X}$	3.67	2.15	1.84	0.72	0.59
	σ	1.37	0.82	0.56	0.25	0.19
Activity index (%)	$\bar{X}$	76.33	83.31	89.80	78.49	50.77
	σ	17.41	12.21	7.08	13.17	15.52
Tortuosity	$\bar{X}$	1.16	1.15	1.10	1.14	1.12
	σ	0.05	0.03	0.05	0.03	0.03

623

624 **Figures captions**

625

626

627 Figure 1. Experimental set-up for time-lapse assessment of foraminiferal life trait

628

Figure 2. Schematic representation (A, B and C) and images (D, E and F) of the three vertical position categories, which can be taken by a foraminifer. (A and D) Surface – (B and E) Sediment-water interface – (C and F) Burrowed. Examples from *Haynesina germanica* (D and E) and *Quinqueloculina seminula* (F). Scale bars = 0.2 mm.

Figure 3. Box and whiskers plots of the four calculated parameters (A) Activity index (%) – (B) Distance (mm) – (C) Velocity (mm.h⁻¹) – (D) Tortuosity for all species. The letters ‘a’, ‘b’, ‘c’ indicate significant distinct groups of measurements (Mann-Whitney test, $p < 0.05$). The box represents the first, second and third quartiles and the whiskers extend to 1.5 times the interquartile range values outside this range are represented by open circles.

650 Figure 4. Temporal evolution of the vertical position of each studied species

Species	Activity	Distance	Position
<i>Quinqueloculina seminula</i>	Intensive	High	Burrowed
<i>Criboelphidium williamsoni</i>		Medium	Surface
<i>Haynesina germanica</i>		Low	Surface → Burrowed
<i>Ammonia tepida</i>	Burrowed		
<i>Milliammina fusca</i>

651
652 Figure 5. Conceptual schema summarizing the difference between studied species and their
653 classification into functional groups with *Q. seminula* and *M. fusca* in gallery-biodiffusor group,
654 *A. tepida* and *H. germanica* in surficial-biodiffusor and *C. williamsoni* in epifaunal-
655 biodiffusors.

656
657