Dear Author,

Here are the proofs of your article.

- You can submit your corrections **online**, via **e-mail** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- You can also insert your corrections in the proof PDF and **email** the annotated PDF.
- For fax submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the **journal title**, **article number**, and **your name** when sending your response via e-mail or fax.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/ corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style. Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.
- Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**
- The **printed version** will follow in a forthcoming issue.

Please note

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL: [http://dx.doi.org/\[DOI\]](http://dx.doi.org/[DOI]).

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information go to: <http://www.link.springer.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us if you would like to have these documents returned.

Metadata of the article that will be visualized in OnlineFirst

ArticleTitle	Modelling mitochondrial ROS production by the respiratory chain	
Article Sub-Title		
Article CopyRight	Springer Nature Switzerland AG (This will be the copyright line in the final PDF)	
Journal Name	Cellular and Molecular Life Sciences	
Corresponding Author	Family Name	Mazat
	Particle	
	Given Name	Jean-Pierre
	Suffix	
	Division	UMR 5095
	Organization	IBGC CNRS
	Address	1 Rue Camille Saint-Saëns 33077, Bordeaux Cedex, France
	Division	
	Organization	Université de Bordeaux
	Address	146 Rue Léo-Saignat, 33076, Bordeaux Cedex, France
	Phone	+33-556-999-041
	Fax	
	Email	jean-pierre.mazat@u-bordeaux.fr
	URL	
	ORCID	http://orcid.org/0000-0002-7832-7455
Author	Family Name	Devin
	Particle	
	Given Name	Anne
	Suffix	
	Division	UMR 5095
	Organization	IBGC CNRS
	Address	1 Rue Camille Saint-Saëns 33077, Bordeaux Cedex, France
	Phone	
	Fax	
	Email	
	URL	
	ORCID	
Author	Family Name	Ransac
	Particle	
	Given Name	Stéphane
	Suffix	
	Division	UMR 5095
	Organization	IBGC CNRS
	Address	1 Rue Camille Saint-Saëns 33077, Bordeaux Cedex, France
	Division	

Organization	Université de Bordeaux
Address	146 Rue Léo-Saignat, 33076, Bordeaux Cedex, France
Phone	
Fax	
Email	
URL	
ORCID	

Schedule	Received	4 November 2019
	Revised	4 November 2019
	Accepted	12 November 2019
Abstract	<p>ROS (superoxide and oxygen peroxide in this paper) play a dual role as signalling molecules and strong oxidizing agents leading to oxidative stress. Their production mainly occurs in mitochondria although they may have other locations (such as NADPH oxidase in particular cell types). Mitochondrial ROS production depends in an interweaving way upon many factors such as the membrane potential, the cell type and the respiratory substrates. Moreover, it is experimentally difficult to quantitatively assess the contribution of each potential site in the respiratory chain. To overcome these difficulties, mathematical models have been developed with different degrees of complexity in order to analyse different physiological questions ranging from a simple reproduction/simulation of experimental results to a detailed model of the possible mechanisms leading to ROS production. Here, we analyse experimental results concerning ROS production including results still under discussion. We then critically review the three models of ROS production in the whole respiratory chain available in the literature and propose some direction for future modelling work.</p>	
Keywords (separated by '-')	ROS - Superoxide - Oxygen peroxide - Respiratory chain - Modelling	
Footnote Information

REVIEW

Modelling mitochondrial ROS production by the respiratory chain

Jean-Pierre Mazat^{1,2} · Anne Devin¹ · Stéphane Ransac^{1,2}

Received: 4 November 2019 / Revised: 4 November 2019 / Accepted: 12 November 2019

© Springer Nature Switzerland AG 2019

Abstract

ROS (superoxide and oxygen peroxide in this paper) play a dual role as signalling molecules and strong oxidizing agents leading to oxidative stress. Their production mainly occurs in mitochondria although they may have other locations (such as NADPH oxidase in particular cell types). Mitochondrial ROS production depends in an interweaving way upon many factors such as the membrane potential, the cell type and the respiratory substrates. Moreover, it is experimentally difficult to quantitatively assess the contribution of each potential site in the respiratory chain. To overcome these difficulties, mathematical models have been developed with different degrees of complexity in order to analyse different physiological questions ranging from a simple reproduction/simulation of experimental results to a detailed model of the possible mechanisms leading to ROS production. Here, we analyse experimental results concerning ROS production including results still under discussion. We then critically review the three models of ROS production in the whole respiratory chain available in the literature and propose some direction for future modelling work.

Keywords ROS · Superoxide · Oxygen peroxide · Respiratory chain · Modelling

Introduction

Accumulating evidence has suggested that reactive oxygen species (ROS), such as superoxide, hydrogen peroxide and other reactive forms of oxygen, play an important role in a broad range of cellular signalling processes [1–4]. However, at high concentrations, ROS damage proteins, lipid membranes, DNA and triggers PTP opening [5, 6] generating what is called oxidative stress. Oxidative stress is defined as a perturbation in the balance between the production of reactive oxygen species (free radicals) and antioxidant defences and contributes to pathologies such as cancer, ischemic cardiac injury and stroke, neurodegenerative diseases and other age-related degenerative conditions [7, 8]. Given their deleterious effects, ROS production is usually finely tuned by ROS-scavenging systems.

The mitochondrial electron transport chain is one of the major providers of ROS in most cells. In C2C12 myoblasts,

Wong et al. [9] show that 45% of ROS comes from mitochondria and 40% from NADPH oxidase. Work by many investigators (see [10] for a review) has largely established that complexes I and III of the mitochondrial respiratory chain are the major sources of reactive oxygen species (ROS), in the form of superoxide (O_2^-) and hydrogen peroxide H_2O_2 . However, despite intensive biochemical and biophysical studies of electron and proton transfer in the respiratory chain (for reviews, see [9, 11–15]) many questions about the mechanisms of O_2^- generation, particularly in physiological conditions, remain unsolved.

The production of superoxide/hydrogen peroxide is difficult to assess, particularly their site of production and their dependence upon the experimental conditions (respiratory substrate, inhibitors). Furthermore, when working with the whole respiratory chain in isolated mitochondria or in whole cells, it is difficult to assess the relative contribution of each separate site and to take into account contribution of the scavenging systems. This is why theoretical models of ROS generation can be useful to facilitate the quantitative analysis of the features controlling mitochondrial O_2^- production and help in the elucidation of experimental results and eventually predict new discriminant experimental protocols. Several theoretical models of ROS production by complex I and III exist [16–21] and also one involving complex II

✉ Jean-Pierre Mazat
jean-pierre.mazat@u-bordeaux.fr

¹ UMR 5095, IBGC CNRS, 1 Rue Camille Saint-Saëns 33077, Bordeaux Cedex, France

² Université de Bordeaux, 146 Rue Léo-Saignat, 33076 Bordeaux Cedex, France

[22]. However, we found only three theoretical models in the literature aiming at describing ROS production by the whole respiratory chain under different conditions. First, we will describe the main experimental results concerning ROS production for which a large consensus exists, and that theoretical models should reproduce, as well as experimental results leading to contradictory hypotheses, between which theoretical models might help to decide. We will emphasize the main points that a theoretical model must explain/simulate and finally we will proceed to the critical description of the three theoretical models.

Experimental data: the different sites of ROS production and the role of inhibitors

Using specific inhibitors of different sites of ROS production (superoxide and/or hydrogen peroxide), particularly inhibitors that do not prevent electron flow and varying the respiratory substrates, the group of Martin Brand has finely dissected the different sites of mitochondrial ROS production [9, 10]. They identified eleven distinct sites associated with respiratory complexes or enzymes and they gave an estimation of the maximal ROS production flux for each site. If we limit our description to the respiratory chain complexes (Fig. 1), the main producers are: the Flavin site of complex I (I_F), the ubiquinone reducing site of complex I (I_Q), the Flavin site of complex II (II_F) and the ubiquinone oxidizing site of complex III (III_{Qo}), (see Fig. 1 in [10] and Fig. 1b in [23]). Site III_{Qo} has, at least, twice the capacity of any other site (see Fig. 2 in [10]).

ROS production by complex I

Site I_F has long been proposed as a site of superoxide production [24, 25]. However, Brand et al. [10, 26] show that much of the ROS production previously ascribed to site I_F truly arises from other dehydrogenases, particularly

sites O_F (2-oxoglutarate dehydrogenase) and P_F (pyruvate dehydrogenase).

Site I_Q during reverse electron transport (RET), the majority of ROS arise from site I_Q . In RET (Fig. 2a), electrons are forced back into complex I by the high QH_2/Q ratio and the high proton motive force generated by electron flow through complexes III and IV. In this process, the electron flow is driven backwards by the consumption of proton motive force. Thus, production of ROS at site I_Q during reverse electron transport has a strong dependence on proton motive force [27–32]. However, several authors showed that it is much more sensitive to the magnitude of the pH gradient than of the membrane potential, even at constant proton motive force [30, 31, 33]. These results suggest that site I_Q is linked to an electroneutral proton-translocating step in the proton-pumping mechanism of complex I [34, 35]. The localization at site I_Q is confirmed by the inhibition of ROS production at this site by rotenone. Then addition of antimycin stimulates ROS production at site III_{Qo} (Figs. 2b, 4).

ROS production by complex II

The Flavin site II_F (Fig. 3) is supposed to be the site of ROS production in complex II and displays a similar maximal capacity of ROS production as site I_Q [10]. However, Grivennikova et al. [22] claim that their data are indicative of the [3Fe–4S] centre, close to the ubiquinone reduction site, as the site of superoxide generation in this complex. Wild-type complex II makes little contribution to ROS production in isolated mammalian mitochondria under normal conditions [36]. However, mutations in this complex can lead to abundant ROS production and cause pathologies [37]. ROS production by the Flavin site of complex II, site II_F , in isolated mitochondria [23] requires two conditions: there must be a source of electrons to reduce the Flavin (succinate), and the site must be open, probably to allow access of oxygen [10].

This behaviour results in a bell-shaped response of ROS production to succinate concentration, with a maximum in the region of the K_M of complex II for succinate

Fig. 1 Possible sites of ROS production (red arrows) in respiratory chain. In the text, the site Qo and Qi of complex III are called III_{Qo} and III_{Qi} . All ROS are generated in the matrix except for ROS in III_{Qo} which are partly extruded in the IMS. *FET* forward electron transport (with NADH), *RET* reverse electron transport (with succinate)

Fig. 2 ROS production in the respiratory chain by complex I and III with succinate. **a** RET results from the addition of succinate alone at high concentration. The thickness of the red arrows corresponds to the first addition of succinate without any inhibitor. Rotenone inhibits

this ROS production. **b** High ROS production by complex III in the presence of antimycin A (AA) [a similar ROS production occurs at complex III with antimycin when electrons are supplied by NADH in complex I in the absence of succinate and rotenone (Rot)]

(100–500 μM) [22, 23, 38]. The capacity for ROS production at site II_F can be measured in two particular conditions illustrated in Fig. 3 in the presence of inhibitors of complexes I and III [10]. In rat skeletal muscle mitochondria, the maximum capacity for ROS production of site II_F is very high, of the same order as site I_Q (Fig. 2 in [10]).

ROS production by complex III (bc1 complex) at site III_{QO}

Complex III (Fig. 4) at site Q_0 (III_{QO} site) has the highest capacity of ROS production (Fig. 2 in [10]). It is commonly accepted that the complex III ROS production is due to the formation during the catalytic process of an instable semiquinone SQ in Q_0 . However, the mechanism of a semiquinone formation in Q_0 is still the matter of controversy, which has to be taken into account for modelling. It is widely accepted that the modified Q-cycle mechanism proposed by Mitchell [39, 40] and subsequently refined by Crofts [41, 42] correctly describes the bc₁ complex operation. It is based on a bifurcation of the two electrons

coming from the QH_2 molecule bound at the Q_0 site. The first electron is transferred to the iron sulphur protein (ISP) and the second to the lower potential heme b_L . The electron on heme b_L moves within the cytochrome b to reduce the higher potential heme b_H , which in turn reduces ubiquinone (Q) at a second ubiquinone binding site Q_i (Fig. 4). The transfer of an electron from quinol to cytochrome c is a complex process involving: (i) a first electron transfer from quinol bound at the catalytic Q_0 site to a [2Fe–2S] cluster situated in the head of the Rieske iron sulphur protein (ISP) anchored in the inner mitochondrial membrane (ii) a large-scale movement of the head of reduced ISP towards cytochrome c_1 , (iii) the reduction of cytochrome c_1 and eventually, (iv) the reduction of cytochrome c by cytochrome c_1 and the return of ISP head to site Q_0 . How the steps following the transfer of the first electron on ISP interweave with the transfer of the second electron on b_L then b_H , is still debated. Three main scenarios (Fig. 5) have been proposed which may have implications for the semiquinone formation/lifetime and its reaction with oxygen:

Fig. 3 ROS production at site II_F (a), complex II in the presence of rotenone, myxothiazol and antimycin A generates ROS in the forward reaction: succinate around its K_M (100–500 μM) supplies electrons to reduce Q to QH_2 which accumulates and blocks electron pathway generating ROS in II_F . Malonate or high succinate concentrations cancel ROS formations but not Atpenin A5 binding in II_Q . **b**

ROS production by complex II in the reverse reaction with electrons supplied by other dehydrogenases to accumulate QH_2 and provide electrons in II_F to form ROS. In these conditions, ROS formation is inhibited by Atpenin A5 or Malonate or high succinate concentrations (after [10, 23])

- The ISP leaves the Q_o site to reduce c_1 before the second electron jumps on b_L and then b_H [43, 44] (Fig. 5a).
- The ISP leaves the Q_o site to reduce c_1 after the second electron transfers from b_L to b_H [45, 46] (Fig. 5b).
- A bypass/short-circuit mechanism can occur when reduced b_L transfers its electron not on b_H but in the reverse direction on a quinone in Q_o , which can be either the product of the reaction or of a newly bound quinone [47–49] (Fig. 5c).

Summary of the main points to consider in a mathematical model of mitochondrial ROS generation

- The main site of ROS production by the respiratory chain are III_{QO} , I_Q and II_F with a decreased maximum capacity in this order.

- In III_{QO} and I_Q , ROS are produced by the semi-quinone Q^- . The ROS production in II_F is through the fully reduced FMNH^- species.
- All ROS species by respiratory chain are produced in the matrix except for III_{QO} site, which produces ROS both in the matrix and in the intermembrane space.
- ROS production can also occur in the matrix at the FMN site of complex II (II_F) at low succinate concentration around 100–500 μM , with a maximal capacity analogous to the one of I_Q .
- The relative contributions of distinct mitochondrial sites depend on many factors: the substrates being oxidized, the energetic demands of the cell, the transmembrane potential, the amount of QH_2 pool and ultimately the cell type.

Fig. 4 Modified Q-cycle and ROS generation in complex III. The mechanism of modified Q-cycle [41, 42] involves the bifurcation of electrons in Qo. The first electron (green pathway) reduces a FeS centre in the head of the Rieske protein which moves towards c_1 (dotted green arrows). An unstable semiquinone (SQ) can be formed before

the second electron (blue pathway) is transferred to a Q molecule in Q_i to form a stable semiquinone (bold SQ). A second cycle of QH_2 oxidation in Q_o is necessary to completely reduce the SQ in Q_i in a new QH_2 molecule. After the two cycles, two Cyt_{ox} have been reduced in two Cyt_{red} and a net QH_2 has been oxidized to Q

Mathematical models of ROS production through the whole respiratory chain

Models of the group of Aon and Cortassa

In [50], Kembro et al. extended previous mathematical models of the mitochondrial respiratory chain [51, 52] to account for ROS production. Their ROS production model is purely phenomenological, with a function called ‘shunt’ which is a small percentage of the rate of respiration (VO_2) and depends on the state 3 or 4 of the respiration rate (Table 2 in [50]). In addition, the model involves the important contribution of ROS scavenging systems to study the balance between ROS production and scavenging in different redox environments. Model simulations were compared with experiments from isolated heart mitochondria reported in the same paper. However, in their conclusions the authors note that their model “is unable to simulate the increase in ROS levels when mitochondria evolve into state 4 respiration” [50].

Shortly after, the same authors published a new model [53] which is, above all, a detailed respiratory chain model including variables describing the concentrations of ubiquinone, ubiquinol, and ubisemiquinone, along with the oxidation states of cytochrome c and the redox centres in complex III, i.e., the high- and low-potential b-hemes (b_H and b_L) and cytochrome c_1 . They use the forward and reverse rate constants for electron transfer of complexes II–IV from the model of Demin et al. [54, 55]. A model of complex I was based on the non-equilibrium thermodynamic description

from Magnus and Keizer’s mitochondrial model [56]. First, the authors fit the experimental results in which the variables such as b_L and b_H reduction states [57, 58] are measured because these variables play an important role in semiquinone concentration at the Q_o site of complex III. The production of ROS in complex I is supposed to occur from $FMNH^-$ according to Pryde and Hirst [59], although the authors test the possibility of ROS production in site I_Q (but in the absence of ROS production by $FMNH^-$); they claim that they obtain similar results (not shown) and conclude that “Given that the modelling results for the two different hypotheses were identical, this modelling experiment was unable to distinguish between the two mechanisms” [53]. They also reproduce the pH dependence of ROS production by complex I observed in [33] (Fig. 5 in [53]). The ROS production from complex III is modelled from Demin and involves the semiquinone in Q_o , a highly reduced quinone pool and a high proton motive force [54, 55]. The authors were aware of other hypotheses [47, 60] but did not take them into account in their model.

In addition, they added to their model of ROS production a previous minimal model of ROS scavenging [61], which allowed them to draw a U-shaped dependence of the ROS balance between production and scavenging as a function of mitochondrial redox environment (sum of redox potentials for NADH, NADPH and GSH weighted by their respective concentrations [62]): the measured ROS production is high in reducing environment which favours ROS production which cannot be destroyed at limited scavenging mechanisms, but are also high in oxidizing environment,

Fig. 5 ISP movement, transfer of the second electron and ROS formation. The dotted red arrows indicate the possible formation of O_2^- from the semiquinone SQ in Qo **a** 'early dissociation hypothesis': O_2^- formation can occur in the same time as reduced ISP leaves Qo site, i.e. before the second electron is transferred to heme b_L ; **b** 'late dissociation hypothesis': reduced ISP leaves Qo site after the second electron is transferred from b_L to b_H and **c** return of the electron of reduced b_L on a Q molecule in Qo forming the semiquinone which can react with oxygen

as, despite a lower ROS production, the scavenging mechanisms are also less efficient (less regenerated in oxidizing environment). The intermediate environment corresponds to a minimum in the ROS production.

More recently, the same authors proposed an experimental and theoretical approach to assess the effects of β -oxidation in the heart on redox and energy metabolism [63]. They described the antagonist effects of fatty acids as respiratory substrates but also as uncouplers on the respiration rate (VO_2) and on the ROS production rate in relation with the level of the antioxidant system.

Markevich's model [64]

In order to study the mitochondrial production of ROS in different conditions of respiratory substrates and membrane potential ($\Delta\psi$), Markevich and Hoek developed an elaborate computational model of the whole respiratory chain. Complex II, modelled by only one simple rate equation, is not taken into account in their work for ROS production. Electron transfers inside complex I and III are detailed with several sites of ROS production. In accordance with Kussmaul and Hirst [24], they proposed that O_2^- is formed by the transfer of one electron from the fully reduced Flavin FMNH⁻ to O_2 . In addition, they assumed that the semiquinone Q^- in site I_Q is a second site of O_2^- formation in Complex I. Regarding complex III, they considered, as generally accepted, that the unstable semiquinone Q^- in site Qo is the site of O_2^- formation. According to the various mechanisms of electron transfer that have been proposed in Qo (see above), Markevich and Hoek compare three variations of their model. The first one considers that O_2^- formation occurs at the same time as the reduced Rieske protein (Iron Sulphur Protein or ISP) leaves the site Q_o to transfer its electron to c_1 ('early dissociation of ISP', Fig. 5a) i.e. before the second electron is transferred to the heme b_L [43, 65]. The second variation suggests that the Rieske protein leaves the Qo site after the transfer of the second electron to b_L and then to b_H ('late dissociation of ISP', Fig. 5b) [46]. In a third variation proposed by Dröse and Brandt [47] they considered that the oxidized quinone can leave the Qo site before ISP and before b_L transfers its electron to b_H . This may allow the return of the reduced b_L electron on an oxidized Q molecule (bypass/short-circuit mechanism also proposed in [48,

49]) by the reversion of the reactions to form a semiquinone which can react with O_2 and explain the activating role of the oxidized quinone in O_2^- formation by complex III (Fig. 5c).

The values of the kinetic parameters of superoxide production were chosen such that the computer-simulated rates of ROS generation were close to those observed in liver mitochondria [66, 67].

Their model is detailed enough to dissect ROS production at each site in different experimental conditions, particularly as a function of the transmembrane potential $\Delta\psi$ with different respiratory substrates (NADH alone, succinate alone, NADH⁺ succinate and NADH⁺ Rotenone (ROT), Fig. 3 in [64]). The difference in ROS production when pH or $\Delta\psi$ were changed is noteworthy (Fig. 4 in [64]). Furthermore, comparing the different variations of electron transfer in site Qo of complex III, they proposed that the scenario with 'late dissociation of ISP' is more likely. Finally, their third variation of the model (with 'late dissociation of ISP' and with binding of Q when $cytb_L$ is reduced) qualitatively reproduced the results of Dröse and Brand [47] and of Quinlan et al. [19] (see also Fig. 12 in [64]). They dissected this behaviour by analysing the amount of individual species, oxidized Q, reduced b_L and the b_L^- .Q.ISP complex (Fig. 12B in [64]). Their results stressed the necessity of a detailed modelling in the multifaceted field of ROS production by the respiratory chain. Furthermore Markevich and Hoek showed another advantage of modelling, i.e. generating testable hypotheses, some of them were confirmed later on, such as the ROS production by RET in I_Q [10]. Incidentally, they offer a careful model of the calculation of the different volumes and concentrations of the different mitochondrial compartments (membrane, matrix, etc.). In fact, this very detailed model is not far from a complete model of the respiratory chain, i.e. the expression of VO_2 as a function of different substrates (Fig. 3E in their paper) and can certainly be used to test other hypotheses.

Bazil and Vinnakota's model [68]

Bazil et al. [68] integrated their previous models of superoxide and hydrogen peroxide production by complexes I [20] and III [69] into an updated Beard's model of oxidative phosphorylation [70] that can simulate both the respiratory dynamics associated with ATP production and the kinetics of ROS production in a single integrated system. The authors distinguished hydrogen peroxide and superoxide generation.

The authors showed that the kinetic control is distributed and depends upon the experimental conditions as experimentally reported by many authors [71–73] and they fitted experimental results obtained using isolated rat heart mitochondria at low (1 mM) and high (5 mM) Pi concentrations.

They showed that ROS production depends indirectly on Pi concentration through changes in pH and $\Delta\psi$.

Model simulations predicted that complex III is responsible for more ROS production during physiological working conditions relative to complex I in the condition of forward electron transport, where electrons are transferred along the Electron Transfer Chain (ETC) from NADH and succinate to O_2 . However, simulating ischemia/reperfusion mechanism, they showed that an accumulation of succinate leading to a highly reduced quinone pool can explain a burst of ROS generated in complex I corroborating the experimental results of Chouchani et al. [8] (who also used a mathematical model of ROS production in complex I [74]). Of note the authors explored the bistability behaviour reported in [16–18] characterized by different rates of ROS production in the same conditions. This is mainly due to their model of ROS production by complex III [69] as emphasized in several complex III models aimed to simulate this phenomenon [16–18, 21].

Conclusion and future prospective

We have analysed in this review the three models currently available for ROS production by the whole respiratory chain. Two of them focused on ROS production in heart [50, 53, 68] in which ROS production during ischemia–reperfusion is a medical concern. The third one by Markevich and Hoek [64] estimated parameters from liver mitochondria but is in fact rather general. It is difficult to compare these models because their aims were different. The models of Aon and Cortassa and of Bazil and Vinnakota tried to simulate in the simplest way possible the experimental results in the heart. Furthermore, Aon and Cortassa studied the physiological interplay of the scavenging mechanisms with ROS production. On the contrary, the Markevich and Hoek model [64] aimed at understanding the intimate mechanisms of ROS production in different conditions. Their model is more detailed than the two others and analysing the consequences of different scenarios for the complex reaction of complex III, proposed a mechanism of the controversial mode of bifurcation of electrons in the Q_o site of bc_1 complex. This illustrates two purposes of modelling, either to derive a phenomenological simple model which can be used to study physiological question such as the balance between ROS production and scavenging in different redox situations or to test different mechanistic hypotheses of ROS production as done by Markevich and Hoek [64]. However, even in this latter case some simplifications were made. For instance, they bring together almost all FeS centres redox reactions of complex I in one reaction and concerning complex III, they assume the release of reduced ISP from Q_o in the same time as the second electron transfer on b_L , which is far from acknowledged.

Another hypothesis, made in all models, is the localization of ROS production. A general consensus is emerging that I_F , I_Q , II_F and III_{Q_o} are the main sites of ROS production in the respiratory chain and there are good evaluations of their maximal capacities [10]. All models take these sites as ROS production sites, excluding site II_F . In addition, some indications exist of the possibility of ROS production at other sites. In principle, any reduced redox centre with midpoint potential close to that of the O_2/O_2^- couple (-160 mV), should be able to produce O_2^- when O_2 is (spatially) close enough to accept an electron. The respiratory chain complexes contain iron–sulphur centres and hemes which are, in principle, able to transfer their electrons to O_2 . The limiting factor will be the distance, the O_2 molecule being at the closest at the surface of the protein. This is perhaps the reason why the binding sites, opened on the external medium and allowing O_2 to diffuse near the redox active site, are favoured for ROS generation. Although left aside by a number of experimenters, there have been published experiments proposing such redox centres as producing superoxide. For complex I, the iron–sulphur centre N_2 which is close to the quinone binding site has been proposed in [75], as well as the iron–sulphur centre $N1a$ which is not very far from the FMN in [76]. The Fe_3S_4 iron–sulphur centre of complex II which is close to the quinone binding site was proposed as a superoxide producing site [22]. Similarly, it has been proposed that reduced heme b_L is able to generate ROS in complex III [77]. One way of approaching this problem without any a priori would be to calculate the probability of reacting with the oxygen of all the redox centres of the respiratory complexes by using the equations developed by Moser and Dutton [78], that considers the distance and the difference in redox potential between the redox centres. Such a stochastic treatment takes into account all possible reactions (i.e. the reactions with a reasonable probability) and all the different oxidized/reduced species, but only when they are produced contrary to what occurs using differential equations (see for instance [79, 80]).

Of note, all models listed above are studied at steady-state. Let us also emphasize that experimental studies of transient phases of ROS production might be more informative on the intimate mechanisms than the simple consideration of steady-states.

To sum up, it appears that a complete model of ROS production by the respiratory chain still remains to be developed by incorporating at least the generation of ROS by the dehydrogenases of the Krebs cycle (including the II_F site of the succinate dehydrogenase) in a deterministic (differential equations) or/and a stochastic approach.

Acknowledgements This work was supported by the Agence Nationale de la Recherche and the Conseil National de la Recherche Scientifique

(CNRS). The authors wish to thank Prof. Michel Rigoulet for careful proofreading of the manuscript and constructive discussions.

Author contributions JPM conceived and coordinated the study and wrote the paper. SR and AD contributed to the writing of the paper, the design of the figures and the analysis of the literature. All authors approved the final version of the manuscript.

Compliance with ethical standards

Conflict of interest The corresponding author declares no conflict of interests on behalf of all authors.

References

- Dröge W (2002) Free radicals in the physiological control of cell function. *Physiol Rev* 82:47–95. <https://doi.org/10.1152/physrev.00018.2001>
- Sies H, Berndt C, Jones DP (2017) Oxidative stress. *Annu Rev Biochem* 86:715–748. <https://doi.org/10.1146/annurev-biochem-061516-045037>
- Sies H (2018) On the history of oxidative stress: concept and some aspects of current development. *Curr Opin Toxicol* 7:122–126. <https://doi.org/10.1016/j.cotox.2018.01.002>
- Boveris A, Chance B (1973) The mitochondrial generation of hydrogen peroxide. General properties and effect of hyperbaric oxygen. *Biochem J* 134:707–716
- Petronilli V, Costantini P, Scorrano L et al (1994) The voltage sensor of the mitochondrial permeability transition pore is tuned by the oxidation-reduction state of vicinal thiols. Increase of the gating potential by oxidants and its reversal by reducing agents. *J Biol Chem* 269:16638–16642
- Schieber M, Chandel NS (2014) ROS function in redox signaling and oxidative stress. *Curr Biol* 24:R453–R462. <https://doi.org/10.1016/j.cub.2014.03.034>
- Harman D (1956) Aging: a theory based on free radical and radiation chemistry. *J Gerontol* 11:298–300
- Chouchani ET, Pell VR, Gaude E et al (2014) Ischaemic accumulation of succinate controls reperfusion injury through mitochondrial ROS. *Nature* 515:431–435. <https://doi.org/10.1038/nature13909>
- Wong H-S, Benoit B, Brand MD (2019) Mitochondrial and cytosolic sources of hydrogen peroxide in resting C2C12 myoblasts. *Free Radic Biol Med* 130:140–150. <https://doi.org/10.1016/j.freeradbiomed.2018.10.448>
- Brand MD (2016) Mitochondrial generation of superoxide and hydrogen peroxide as the source of mitochondrial redox signaling. *Free Radic Biol Med* 100:14–31. <https://doi.org/10.1016/j.freeradbiomed.2016.04.001>
- Murphy MP (2009) How mitochondria produce reactive oxygen species. *Biochem J* 417:1–13. <https://doi.org/10.1042/BJ20081386>
- Detaille D, Pasdois P, Sémont A et al (2019) An old medicine as a new drug to prevent mitochondrial complex I from producing oxygen radicals. *PLoS ONE* 14:e0216385. <https://doi.org/10.1371/journal.pone.0216385>
- Andreyev AY, Kushnareva YE, Murphy AN, Starkov AA (2015) Mitochondrial ROS metabolism: 10 years later. *Biochem Mosc* 80:517–531. <https://doi.org/10.1134/S0006297915050028>
- Bouchez C, Devin A (2019) Mitochondrial biogenesis and mitochondrial reactive oxygen species (ROS): a complex relationship regulated by the cAMP/PKA signaling pathway. *Cells* 8:287. <https://doi.org/10.3390/cells8040287>
- Dubouchaud H, Walter L, Rigoulet M, Batandier C (2018) Mitochondrial NADH redox potential impacts the reactive oxygen species production of reverse electron transfer through complex I. *J Bioenerg Biomembr* 50:367–377. <https://doi.org/10.1007/s10863-018-9767-7>
- Selivanov VA, Votyakova TV, Pivtoraiko VN et al (2011) Reactive oxygen species production by forward and reverse electron fluxes in the mitochondrial respiratory chain. *PLoS Comput Biol* 7:e1001115. <https://doi.org/10.1371/journal.pcbi.1001115>
- Selivanov VA, Votyakova TV, Zeak JA et al (2009) Bistability of mitochondrial respiration underlies paradoxical reactive oxygen species generation induced by anoxia. *PLoS Comput Biol* 5:e1000619. <https://doi.org/10.1371/journal.pcbi.1000619>
- Selivanov VA, Cascante M, Friedman M et al (2012) Multistationary and oscillatory modes of free radicals generation by the mitochondrial respiratory chain revealed by a bifurcation analysis. *PLoS Comput Biol* 8:e1002700. <https://doi.org/10.1371/journal.pcbi.1002700>
- Quinlan CL, Gerencsér AA, Treberg JR, Brand MD (2011) The mechanism of superoxide production by the antimycin-inhibited mitochondrial Q-cycle. *J Biol Chem* 286:31361–31372. <https://doi.org/10.1074/jbc.M111.267898>
- Bazil JN, Pannala VR, Dash RK, Beard DA (2014) Determining the origins of superoxide and hydrogen peroxide in the mammalian NADH: ubiquinone oxidoreductase. *Free Radic Biol Med*. <https://doi.org/10.1016/j.freeradbiomed.2014.08.023>
- Guillaud F, Dröse S, Kowald A et al (2014) Superoxide production by cytochrome bc1 complex: a mathematical model. *Biochim Biophys Acta (BBA) Bioenerg* 1837:1643–1652. <https://doi.org/10.1016/j.bbabi.2014.05.358>
- Grivennikova VG, Kozlovsky VS, Vinogradov AD (2017) Respiratory complex II: ROS production and the kinetics of ubiquinone reduction. *Biochim Biophys Acta (BBA) Bioenerg* 1858:109–117. <https://doi.org/10.1016/j.bbabi.2016.10.008>
- Quinlan CL, Orr AL, Perevoshchikova IV et al (2012) Mitochondrial complex II can generate reactive oxygen species at high rates in both the forward and reverse reactions. *J Biol Chem* 287:27255–27264. <https://doi.org/10.1074/jbc.M112.374629>
- Kussmaul L, Hirst J (2006) The mechanism of superoxide production by NADH:ubiquinone oxidoreductase (complex I) from bovine heart mitochondria. *PNAS* 103:7607–7612. <https://doi.org/10.1073/pnas.0510977103>
- Hirst J, King MS, Pryde KR (2008) The production of reactive oxygen species by complex I. *Biochem Soc Trans* 36:976–980. <https://doi.org/10.1042/BST0360976>
- Quinlan CL, Goncalves RLS, Hey-Mogensen M et al (2014) The 2-oxoacid dehydrogenase complexes in mitochondria can produce superoxide/hydrogen peroxide at much higher rates than complex I. *J Biol Chem* 289:8312–8325. <https://doi.org/10.1074/jbc.M113.545301>
- Hansford R, Hogue B, Mildažienė V (1997) Dependence of H₂O₂ formation by rat heart mitochondria on substrate availability and donor age. In: Hansford RG, Hogue BA, Mildažienė V (eds)
- Miwa S, St-Pierre J, Partridge L, Brand MD (2003) Superoxide and hydrogen peroxide production by *Drosophila mitochondria*. *Free Radic Biol Med* 35:938–948. [https://doi.org/10.1016/S0891-5849\(03\)00464-7](https://doi.org/10.1016/S0891-5849(03)00464-7)
- Votyakova TV, Reynolds IJ (2001) $\Delta\psi$ m-dependent and -independent production of reactive oxygen species by rat brain mitochondria. *J Neurochem* 79:266–277. <https://doi.org/10.1046/j.1471-4159.2001.00548.x>
- Lambert AJ, Brand MD (2004) Inhibitors of the quinone-binding site allow rapid superoxide production from mitochondrial NADH: ubiquinone oxidoreductase (complex I). *J Biol Chem* 279:39414–39420. <https://doi.org/10.1074/jbc.M406576200>

31. Liu S (1997) Generating, partitioning, targeting and functioning of superoxide in mitochondria. *Biosci Rep* 17:259–272. <https://doi.org/10.1023/A:1027328510931>
32. Korshunov SS, Skulachev VP, Starkov AA (1997) High protonic potential actuates a mechanism of production of reactive oxygen species in mitochondria. *FEBS Lett* 416:15–18. [https://doi.org/10.1016/S0014-5793\(97\)01159-9](https://doi.org/10.1016/S0014-5793(97)01159-9)
33. Lambert AJ, Brand MD (2004) Superoxide production by NADH: ubiquinone oxidoreductase (complex I) depends on the pH gradient across the mitochondrial inner membrane. *Biochem J* 382:511–517. <https://doi.org/10.1042/BJ20040485>
34. Treberg JR, Quinlan CL, Brand MD (2011) Evidence for two sites of superoxide production by mitochondrial NADH-ubiquinone oxidoreductase (complex I). *J Biol Chem* 286:27103–27110. <https://doi.org/10.1074/jbc.M111.252502>
35. Treberg JR, Brand MD (2011) A model of the proton translocation mechanism of complex I. *J Biol Chem* 286:17579–17584. <https://doi.org/10.1074/jbc.M111.227751>
36. St-Pierre J, Buckingham JA, Roebuck SJ, Brand MD (2002) Topology of superoxide production from different sites in the mitochondrial electron transport chain. *J Biol Chem* 277:44784–44790. <https://doi.org/10.1074/jbc.M207217200>
37. Ackrell BAC (2002) Cytopathies involving mitochondrial complex II. *Mol Aspects Med* 23:369–384. [https://doi.org/10.1016/S0098-2997\(02\)00012-2](https://doi.org/10.1016/S0098-2997(02)00012-2)
38. Siebels I, Dröse S (2013) Q-site inhibitor induced ROS production of mitochondrial complex II is attenuated by TCA cycle dicarboxylates. *Biochim Biophys Acta (BBA) Bioenerg* 1827:1156–1164. <https://doi.org/10.1016/j.bbabi.2013.06.005>
39. Mitchell P (1976) Possible molecular mechanisms of the proton-motive function of cytochrome systems. *J Theor Biol* 62:327–367. [https://doi.org/10.1016/0022-5193\(76\)90124-7](https://doi.org/10.1016/0022-5193(76)90124-7)
40. Trumpower BL (1990) The protonmotive Q cycle. Energy transduction by coupling of proton translocation to electron transfer by the cytochrome bc1 complex. *J Biol Chem* 265:11409–11412
41. Crofts AR, Meinhardt SW, Jones KR, Snozzi M (1983) The role of the quinone pool in the cyclic electron-transfer chain of *Rhodospseudomonas sphaeroides* a modified Q-cycle mechanism. *Biochim Biophys Acta (BBA) Bioenerg* 723:202–218. [https://doi.org/10.1016/0005-2728\(83\)90120-2](https://doi.org/10.1016/0005-2728(83)90120-2)
42. Crofts AR (2004) The Q-cycle—a personal perspective. *Photosynth Res* 80:223–243. <https://doi.org/10.1023/B:PRES.0000030444.52579.10>
43. Crofts AR, Hong S, Wilson C et al (2013) The mechanism of ubiquinol oxidation at the Qo-site of the cytochrome bc1 complex. *Biochim Biophys Acta (BBA) Bioenerg* 1827:1362–1377. <https://doi.org/10.1016/j.bbabi.2013.01.009>
44. Victoria D, Burton R, Crofts AR (2013) Role of the -PEWY-glutamate in catalysis at the Qo-site of the Cyt bc1 complex. *Biochim Biophys Acta (BBA) Bioenerg* 1827:365–386. <https://doi.org/10.1016/j.bbabi.2012.10.012>
45. Esser L, Gong X, Yang S et al (2006) Surface-modulated motion switch: capture and release of iron–sulfur protein in the cytochrome bc1 complex. *PNAS* 103:13045–13050. <https://doi.org/10.1073/pnas.0601149103>
46. Yu C-A, Cen X, Ma H-W et al (2008) Domain conformational switch of the iron–sulfur protein in cytochrome bc1 complex is induced by the electron transfer from cytochrome bL to bH. *Biochim Biophys Acta (BBA) Bioenerg* 1777:1038–1043. <https://doi.org/10.1016/j.bbabi.2008.03.033>
47. Dröse S, Brandt U (2008) The mechanism of mitochondrial superoxide production by the cytochrome bc1 complex. *J Biol Chem* 283:21649–21654. <https://doi.org/10.1074/jbc.M803236200>
48. Osyczka A, Moser CC, Daldal F, Dutton PL (2004) Reversible redox energy coupling in electron transfer chains. *Nature* 427:607. <https://doi.org/10.1038/nature02242>
49. Osyczka A, Moser CC, Dutton PL (2005) Fixing the Q cycle. *Trends Biochem Sci* 30:176–182. <https://doi.org/10.1016/j.tibs.2005.02.001>
50. Kembro JM, Aon MA, Winslow RL et al (2013) Integrating mitochondrial energetics, redox and ROS metabolic networks: a two-compartment model. *Biophys J* 104:332–343. <https://doi.org/10.1016/j.bpj.2012.11.3808>
51. Cortassa S, Aon MA, Winslow RL, O'Rourke B (2004) A mitochondrial oscillator dependent on reactive oxygen species. *Biophys J* 87:2060–2073. <https://doi.org/10.1529/biophysj.104.041749>
52. Wei A-C, Aon MA, O'Rourke B et al (2011) Mitochondrial energetics, pH regulation, and ion dynamics: a computational–experimental approach. *Biophys J* 100:2894–2903. <https://doi.org/10.1016/j.bpj.2011.05.027>
53. Gauthier LD, Greenstein JL, Cortassa S et al (2013) A computational model of reactive oxygen species and redox balance in cardiac mitochondria. *Biophys J* 105:1045–1056. <https://doi.org/10.1016/j.bpj.2013.07.006>
54. Demin OV, Kholodenko BN, Skulachev VP (1998) A model of O₂-generation in the complex III of the electron transport chain. *Mol Cell Biochem* 184:21–33
55. Demin OV, Gorianin II, Kholodenko BN, Westerhoff HV (2001) Kinetic modeling of energy metabolism and generation of active forms of oxygen in hepatocyte mitochondria. *Mol Biol (Mosk)* 35:1095–1104
56. Magnus G, Keizer J (1997) Minimal model of beta-cell mitochondrial Ca²⁺ handling. *Am J Physiol Cell Physiol* 273:C717–C733. <https://doi.org/10.1152/ajpcell.1997.273.2.C717>
57. Kim N, Ripple MO, Springett R (2012) Measurement of the mitochondrial membrane potential and pH gradient from the redox poise of the hemes of the bc1 complex. *Biophys J* 102:1194–1203. <https://doi.org/10.1016/j.bpj.2012.02.003>
58. Brown GC, Brand MD (1985) Thermodynamic control of electron flux through mitochondrial cytochrome bc1 complex. *Biochem J* 225:399–405. <https://doi.org/10.1042/bj2250399>
59. Pryde KR, Hirst J (2011) Superoxide is produced by the reduced flavin in mitochondrial complex I a single, unified mechanism that applies during both forward and reverse electron transfer. *J Biol Chem* 286:18056–18065. <https://doi.org/10.1074/jbc.M110.186841>
60. Borek A, Sarewicz M, Osyczka A (2008) Movement of the iron–sulfur head domain of cytochrome bc1 transiently opens the catalytic Qo site for reaction with oxygen. *Biochemistry* 47:12365–12370. <https://doi.org/10.1021/bi801207f>
61. Aon MA, Stanley BA, Sivakumaran V et al (2012) Glutathione/thioredoxin systems modulate mitochondrial H₂O₂ emission: an experimental–computational study. *J Gen Physiol* 139:479–491. <https://doi.org/10.1085/jgp.201210772>
62. Schafer FQ, Buettner GR (2001) Redox environment of the cell as viewed through the redox state of the glutathione disulfide/glutathione couple. *Free Radic Biol Med* 30:1191–1212. [https://doi.org/10.1016/S0891-5849\(01\)00480-4](https://doi.org/10.1016/S0891-5849(01)00480-4)
63. Cortassa S, Sollott SJ, Aon MA (2017) Mitochondrial respiration and ROS emission during β -oxidation in the heart: an experimental–computational study. *PLoS Comput Biol* 13:e1005588. <https://doi.org/10.1371/journal.pcbi.1005588>
64. Markevich NI, Hoek JB (2015) Computational modeling analysis of mitochondrial superoxide production under varying substrate conditions and upon inhibition of different segments of the electron transport chain. *Biochim Biophys Acta* 1847:656–679. <https://doi.org/10.1016/j.bbabi.2015.04.005>
65. Crofts AR, Lhee S, Crofts SB et al (2006) Proton pumping in the bc1 complex: a new gating mechanism that prevents short circuits. *Biochim Biophys Acta (BBA) Bioenerg* 1757:1019–1034. <https://doi.org/10.1016/j.bbabi.2006.02.009>

66. Batandier C, Guigas B, Demaille D et al (2006) The ROS production induced by a reverse-electron flux at respiratory-chain complex I is hampered by metformin. *J Bioenerg Biomembr* 38:33–42. <https://doi.org/10.1007/s10863-006-9003-8>
67. Hoffman DL, Brookes PS (2009) Oxygen sensitivity of mitochondrial reactive oxygen species generation depends on metabolic conditions. *J Biol Chem* 284:16236–16245. <https://doi.org/10.1074/jbc.M809512200>
68. Bazil JN, Beard DA, Vinnakota KC (2016) Catalytic coupling of oxidative phosphorylation, ATP demand, and reactive oxygen species generation. *Biophys J* 110:962–971. <https://doi.org/10.1016/j.bpj.2015.09.036>
69. Bazil JN, Vinnakota KC, Wu F, Beard DA (2013) Analysis of the kinetics and bistability of ubiquinol: cytochrome c oxidoreductase. *Biophys J* 105:343–355. <https://doi.org/10.1016/j.bpj.2013.05.033>
70. Beard DA (2005) A biophysical model of the mitochondrial respiratory system and oxidative phosphorylation. *PLoS Comput Biol* 1:e36. <https://doi.org/10.1371/journal.pcbi.0010036>
71. Groen AK, Wanders RJ, Westerhoff HV et al (1982) Quantification of the contribution of various steps to the control of mitochondrial respiration. *J Biol Chem* 257:2754–2757
72. Gellerich FN, Bohnsack R, Kunz W (1983) Control of mitochondrial respiration. The contribution of the adenine nucleotide translocator depends on the ATP- and ADP-consuming enzymes. *Biochim Biophys Acta* 722:381–391
73. Letellier T, Malgat M, Mazat JP (1993) Control of oxidative phosphorylation in rat muscle mitochondria: implications for mitochondrial myopathies. *Biochim Biophys Acta* 1141:58–64
74. Smith AC, Robinson AJ (2011) A metabolic model of the mitochondrion and its use in modelling diseases of the tricarboxylic acid cycle. *BMC Syst Biol* 5:102. <https://doi.org/10.1186/1752-0509-5-102>
75. Genova ML, Ventura B, Giuliano G et al (2001) The site of production of superoxide radical in mitochondrial complex I is not a bound ubiquinone but presumably iron–sulfur cluster N2. *FEBS Lett* 505:364–368. [https://doi.org/10.1016/S0014-5793\(01\)02850-2](https://doi.org/10.1016/S0014-5793(01)02850-2)
76. Kushnareva Y, Murphy AN, Andreyev A (2002) Complex I-mediated reactive oxygen species generation: modulation by cytochrome c and NAD(P)⁺ oxidation–reduction state. *Biochem J* 368:545–553. <https://doi.org/10.1042/bj20021121>
77. Nohl H, Jordan W (1986) The mitochondrial site of superoxide formation. *Biochem Biophys Res Commun* 138:533–539. [https://doi.org/10.1016/S0006-291X\(86\)80529-0](https://doi.org/10.1016/S0006-291X(86)80529-0)
78. Moser CC, Keske JM, Warncke K et al (1992) Nature of biological electron transfer. *Nature* 355:796. <https://doi.org/10.1038/355796a0>
79. Ransac S, Parisey N, Mazat J-P (2008) The loneliness of the electrons in the bc1 complex. *Biochim Biophys Acta (BBA) Bioenerg* 1777:1053–1059. <https://doi.org/10.1016/j.bbabi.2008.05.003>
80. Ransac S, Mazat J-P (2010) How does antimycin inhibit the bc1 complex? A part-time twin. *Biochim Biophys Acta (BBA) Bioenerg* 1797:1849–1857. <https://doi.org/10.1016/j.bbabi.2010.05.014>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Journal:	18
Article:	3381

Author Query Form

Please ensure you fill out your response to the queries raised below and return this form along with your corrections

Dear Author

During the process of typesetting your article, the following queries have arisen. Please check your typeset proof carefully against the queries listed below and mark the necessary changes either directly on the proof/online grid or in the 'Author's response' area provided below

Query	Details Required	Author's Response
AQ1	Kindly check and confirm the edit made in the title.	
AQ2	Kindly check and confirm the corresponding affiliation is correctly identified.	
AQ3	Kindly check and update ref. [27] with complete details.	