Visible Light-Mediated C–H Alkylation of Pyridine Derivatives

Fatima Rammal,[†] Di Gao,[†] Sondes Boujnah,[†] Annie–Claude Gaumont,[†] Aqeel, A. Hussein,[‡] and Sami Lakhdar^{§*}

[†] Normandie Univ., LCMT, ENSICAEN, UNICAEN, CNRS, 6, Boulevard Maréchal Juin, Caen 14000 France

[‡] College of Dentistry, University of Al-Ameed, Karbala, Iraq

[§] Université Paul Sabatier, Laboratoire Hétérochimie Fondamentale et Appliquée (LHFA, UMR 5069), 118 Route de Narbonne, 31062, Toulouse Cedex 09, France

Supporting Information Placeholder

ABSTRACT: We report herein a visible light-mediated C–H alkylation of pyridine derivatives that proceeds by simple combination of a large variety of N-alkoxypyridinium ions with alkanes in the presence of 2 mol% of fac-Ir(ppy)₃ under blue illumination. The mild reaction conditions together with the high group functional tolerance make of this process a useful synthetic platform for the construction of structurally strained heterocycles. Detailed mechanistic investigations, including Density Functional Theory (DFT) calculations and quantum yield measurement, allowed to understand factors controlling the reactivity and the selectivity of the reaction.

Alkylated heteroarenes are ubiquitous chemical motifs present in pharmaceuticals, natural products, and ligand scaffolds.^{1,2} While access to such molecules can be achieved with different synthetic methodologies, the oxidative cross-coupling between two C–H compounds is evidently one of the most elegant approaches as it provides a direct construction of the C–C bond in a step- and atom-economical fashion.^{3–7} For instance, the generation of an alkyl radical through a hydrogen atom transfer (HAT) event and its subsequent addition to a heteroarene ring followed by a formal hydrogen atom loss can provide a rapid and direct method for heterocycle C–H alkylation. This approach, known as the Minisci reaction,^{8–11} gained much attention during the last decade with the spectacular renaissance of the field of photoredox catalysis that provides very mild conditions for the generation of alkyl radicals.^{12–19} While various elegant photoredox Minisci reactions employing HAT strategy for the generation of the alkyl radical have been developed, there is in most cases need to use an external oxidant and stoichiometric amounts of Brønsted acids to activate the heteroarene ring towards radical addition.^{20–27}

Scheme 1. (A) Reported Approaches for C–H Alkylation of Heteroarenes. (B) Current work. HAT denotes Hydrogen Atom Transfer.

A Common Photocatalytic Minisci Reactions

B This work

Based on their abilities to generate strong hydrogen atom abstractors (alkoxy radicals) under visible light photocatalytic conditions,^{28,29} as well as their high electrophilicity, *N*-alkoxypyridinium ions (NAPs) are unique scaffolds to achieve photocatalytic Minisci reaction. This approach has nicely been demonstrated by the Hong group in a certain number of examples, including acylation and intramolecular alkylations.^{30–34} However, the direct association of NAPs with alkanes to achieve Minisci reaction has not been reported so far. Furthermore, it is well known that pyridines are less reactive than quinolines and isoquinolines in various Minisci reactions.^{35–37} We therefore decided to investigate the reactions of NAPs with alkanes under mild photoredox conditions.

Table 1. Optimization of the Photocatalytic Alkylation Reaction between the *N*-alkoxypyridinium 1a and Cyclohexane 2a.^a

entry ^a	photocatalyst (PC)	base	solvent	3a, yield [%] ^b
1	Ru(bpy) ₃	NaHCO ₃	MeCN	traces
2	Eosin Y	NaHCO ₃	MeCN	traces
3	Hong's PC	NaHCO ₃	MeCN	64
4	<i>fac</i> -Ir(ppy) ₃	NaHCO ₃	MeCN	95
5	<i>fac</i> -Ir(ppy) ₃	NaHCO ₃	THF	traces
6	<i>fac</i> -Ir(ppy) ₃	NaHCO ₃	DMF	traces
7	<i>fac</i> -Ir(ppy) ₃	NaHCO ₃	DCM	traces
8	<i>fac</i> -Ir(ppy) ₃	NaHCO ₃	AcOEt	traces
9	<i>fac</i> -Ir(ppy) ₃	NaHCO ₃	DMSO	traces
10	<i>fac</i> -Ir(ppy) ₃	NaHCO ₃	CHCl ₃	traces
11	<i>fac</i> -Ir(ppy) ₃	Na ₂ CO ₃	MeCN	30
12	<i>fac</i> -Ir(ppy) ₃	K ₂ HPO ₄	MeCN	55

13	<i>fac</i> -Ir(ppy) ₃	K ₂ CO ₃	MeCN	41
14	–	NaHCO ₃	MeCN	traces
15	<i>fac</i> -Ir(ppy) ₃ ^c	NaHCO ₃	MeCN	0
16	<i>fac</i> -Ir(ppy) ₃	–	MeCN	33

^a Reaction conditions: *N*-methoxy-4-methylpyridinium methyl sulfate **1a** (0.21 mmol, 1 equiv), cyclohexane **2a** (1.1 mmol, 5 equiv), solvent (4.2 mL), blue LEDs (5W), 1 h. ^b NMR yields are determined from ¹H NMR spectroscopy using 1,1,2,2-tetrachloroethane as internal standard. ^c reaction performed in the dark.

We initially explored the reaction conditions by investigating the effect of the photocatalyst, solvent, and the base through the reaction of the *N*-methoxypyridinium (**1a**) with **2a**. As shown in Table 1, *fac*-Ir(ppy)₃ gave excellent conversion of **3a** (Table 1, entry 4, 95%) when 5 equivalents of **2a** were mixed with **1a** in the presence of NaHCO₃ as a base in acetonitrile under blue light irradiation. While good conversion was obtained with the 3-phosphonated quinolinone (entry 3, 64%), known as the Hong photocatalyst, under the same conditions as with *fac*-Ir(ppy)₃, only traces of **3a** were observed with Ru(bpy)₃ and Eosin Y (entries 1 and 2). A solvent screening, including THF, DMF, DCM, AcOEt, DMSO, and CH₃Cl (entries 5–10) revealed no formation of the desired adduct. We next studied the effect of the base on the reaction and found modest conversion (30–55%) with the inorganic bases Na₂CO₃ (entry 11), K₂HPO₄ (entry 12) and K₂CO₃ (entry 13). The absence of photocatalyst (entry 14) and the visible-light irradiation (entry 15) were detrimental for the C–H alkylation as no product has been observed. Finally, 33% of **3a** was observed when the base was not employed under the optimized conditions (entry 16). The result might be attributed to the pyridinium counterion that played the role of the base in the process.

Next, we set out to investigate the scope of this visible-light C–H alkylation of heteroarenes with alkanes (Figure 1). The reaction of cyclohexane (**2a**) with 4-substituted *N*-alkoxypyridinium ions proceeded smoothly, giving the 2-alkylated pyridines (**3a–3f**) in 50–99% yields. Remarkably, the reaction works well with both electron-donating or electron-withdrawing groups, tolerating cyano (**3c**), trifluoromethyl (**3d**) and carbonyl (**3e**) groups.

The reaction works equally well with 2-substituted pyridiniums. Both regioisomers C2- and C4-alkylated pyridines (**3g–3j**) were obtained in equal proportion (1:1 ratio), except in the case of 2-benzylpyridinium ion where the C2 adduct was obtained as a major regioisomer (C2:C4 = 4:1). The reaction is not only restricted to pyridines as alkylated quinolines (**3k–3n**), 2,2-quinoline (**3o**) and phenanthridine (**3p**) were obtained in good to excellent yields (Figure 1).

Figure 1. Photocatalytic C–H Alkylation of Heteroarenes.^a

^a Reaction conditions: *N*-alkoxy-pyridinium methylsulfate **1** (1 equiv), **2a** (5 equiv), *fac*-Ir(ppy)₃ (2 mol%), NaHCO₃ (1.2 equiv) CH₃CN (0.05 M), blue LEDs (5W), 1 h.

We further extend the scope of the photocatalytic approach to cycloheptane (**2b**) and cyclopentane (**2c**) (Figure 2). Remarkably, a large variety of heteroarylated-alkanes (**3q–3zb**) were isolated in yields going from 34 to 99%. Here again, the reaction is compatible with quinolines derivatives.

The reaction mechanism of the visible-light-mediated alkylation of heteroarenes is depicted in Figure 2. It starts with the generation of the methoxyl radical through single-electron reduction of the *N*-alkoxy-pyridinium ion by the excited state of the photocatalyst (PC*). This event is thermodynamically viable and we have previously demonstrated the generation of (MeO•) by EPR spectroscopy.^{38,39,40,41} DFT calculations show plausible HAT between the methoxyl radical and cyclohexane (**2a**) to form the cyclohexyl radical (**Int 1**). The latter reacts at the

C4 of the pyridinium ion (**1**) position ($\Delta G^\ddagger = 14.4$ kcal/mol) to form the radical intermediate (**Int2**). The radical addition at the C2 position proceeds with an activation energy of 14.8 kcal/mol. This weak difference between both attacks, *i.e.* at C2 and C4 position, may rationalize the low regioselectivity observed. Next, a fast deprotonation of **Int2** by a base gives the **Int3** that upon N–O bond cleavage forms the final product **3** and release the methoxyl radical to start a new reaction cycle. This radical chain pathway explains the high measured quantum yield of the reaction $\Phi = 62.9$ (see Supporting Information for details). The calculated mechanism is in good agreement with the experimental reaction time.

Figure 2. Free energy profile for the photoinitiated C–C bond formation in the reaction of pyridinium ion and cyclohexane. Calculations were performed at the SMD-(ACN)-M06-2X/def2-TZVP//B3LYP/6-31+G(d) level of theory. Energies between brackets are reported for the addition at C2-position. See Supporting Information for computational details.

Although visible-light-mediated alkylation of heteroarenes has widely been reported during the last few years, the reaction scope has mainly been restricted to isoquinolines, quinolines. In this study, we showed that alkylation of pyridines can also be achieved efficiently by simply combining *N*-methoxypyridinium ions with alkanes in the presence of low loading (2 mol %) of the readily accessible *fac*-Ir(ppy)₃ under blue light illumination. The broad scope and functional group tolerance are excellent features of this method. The detailed experimental and computational mechanistic studies conducted in this study are not only allowed understanding factors controlling this photoalkylation reaction but will also help the design of new visible light-induced C–H functionalization of heteroarenes.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website. Quantum yield measurement, DFT calculations, experimental procedures, characterization data, and NMR spectra

AUTHOR INFORMATION

Corresponding Author

* Dr. S. Lakhdar

Normandie Univ., LCMT, ENSICAEN, UNICAEN, CNRS, 6, Boulevard Maréchal Juin, Caen 14000 France
Université Paul Sabatier, Laboratoire Hétérochimie Fondamentale et Appliquée (LHFA, UMR 5069), 118 Route de
Narbonne, 31062, Toulouse Cedex 09, France
lakhdar@chimie.ups-tlse.fr

ACKNOWLEDGMENT

The authors thank the CNRS, Normandie Université, Labex Synorg (ANR-11-LABX-0029) for financial support. F. R. is indebted to the French Ministry of Research for a doctoral fellowship. The authors acknowledge the computational resources from iridis4 supercomputers supported by the University of Southampton. The authors thank Ms. Alya Inial for their help with the preparation of compound 3a.

DEDICATION

This work is dedicated to Prof. Pierre. H. Dixneuf (University of Rennes) for his excellent contribution to organometallic chemistry and catalysis.

REFERENCES

- (1) Vitaku, E.; Smith, D. T.; Njardarson, J. T. Analysis of the Structural Diversity, Substitution Patterns, and Frequency of Nitrogen Heterocycles among U.S. FDA Approved Pharmaceuticals. *J. Med. Chem.* **2014**, *57* (24), 10257–10274. <https://doi.org/10.1021/jm501100b>.
- (2) Baumann, M.; Baxendale, I. R. An Overview of the Synthetic Routes to the Best Selling Drugs Containing 6-Membered Heterocycles. *Beilstein J. Org. Chem.* **2013**, *9*, 2265–2319. <https://doi.org/10.3762/bjoc.9.265>.
- (3) Chen, X.; Engle, K. M.; Wang, D. H.; Jin-Quan, Y. Palladium(II)-Catalyzed C-H Activation/C-C Cross-Coupling Reactions: Versatility and Practicality. *Angew. Chemie - Int. Ed.* **2009**, *48* (28), 5094–5115. <https://doi.org/10.1002/anie.200806273>.
- (4) Rouquet, G.; Chatani, N. Catalytic Functionalization of C(Sp²)-H and C(Sp³)-H Bonds by Using Bidentate Directing Groups. *Angew. Chemie - Int. Ed.* **2013**, *52* (45), 11726–11743. <https://doi.org/10.1002/anie.201301451>.
- (5) Park, S. J.; Price, J. R.; Todd, M. H. Oxidative Arylation of Isochroman. *J. Org. Chem.* **2012**, *77* (2), 949–955. <https://doi.org/10.1021/jo2021373>.
- (6) Matcha, K.; Antonchick, A. P. Metal-Free Cross-Dehydrogenative Coupling of Heterocycles with Aldehydes. *Angew. Chemie - Int. Ed.* **2013**, *52* (7), 2082–2086. <https://doi.org/10.1002/anie.201208851>.
- (7) Liu, D.; Liu, C.; Lei, A. Copper-Catalysed Oxidative C–H/C–H Coupling between Olefins and Simple Ethers. *Chem. Commun.* **2014**, *50* (28), 3623–3626. <https://doi.org/10.1039/c4cc00867g>.
- (8) Proctor, R. S. J.; Phipps, R. J. Recent Advances in Minisci-Type Reactions. *Angew. Chemie - Int. Ed.* **2019**, *58* (39), 13666–13699. <https://doi.org/10.1002/anie.201900977>.
- (9) Caronna, T.; Gardini, G. P.; Minisci, F. Nucleophilic Character of Acyl Radicals: A New Selective Type of Aromatic Acylation. *J. Chem. Soc. D Chem. Commun.* **1969**, 201 (5), 201. <https://doi.org/10.1039/C29690000201>.
- (10) Minisci, F.; Vismara, E.; Fontana, F. Recent Developments of Free-Radical Substitutions of Heteroaromatic Bases. *Heterocycles* **1989**, *28* (1), 489–519. <https://doi.org/10.3987/REV-88-SR1>.
- (11) Minisci, F.; Bertini, F.; Galli, R.; Perchinijmmo, M. Character of Alkyl. *Tetrahedron* **1971**, *27*, 3575.
- (12) Xuan, J.; Xiao, W. J. Visible-Light Photoredox Catalysis. *Angew. Chemie - Int. Ed.* **2012**, *51* (28), 6828–6838. <https://doi.org/10.1002/anie.201200223>.
- (13) Marzo, L.; Pagire, S. K.; Reiser, O.; König, B. Visible-Light Photocatalysis: Does It Make a Difference in Organic Synthesis? *Angew. Chemie - Int. Ed.* **2018**, *57* (32), 10034–10072. <https://doi.org/10.1002/anie.201709766>.
- (14) Yoon, T. P.; Ischay, M. A.; Du, J. Visible Light Photocatalysis as a Greener Approach to Photochemical Synthesis. *Nat. Chem.* **2010**, *2* (7), 527–532. <https://doi.org/10.1038/nchem.687>.
- (15) Wimmer, A.; König, B. Photocatalytic Formation of Carbon-Sulfur Bonds. *Beilstein J. Org. Chem.* **2017**, *14* (Scheme 1), 54–83. <https://doi.org/10.3762/bjoc.14.4>.
- (16) Romero, N. A.; Nicewicz, D. A. Organic Photoredox Catalysis. *Chem. Rev.* **2016**, *116* (17), 10075–10166. <https://doi.org/10.1021/acs.chemrev.6b00057>.
- (17) Prier, C. K.; Rankic, D. A.; MacMillan, D. W. C. Visible Light Photoredox Catalysis with Transition Metal Complexes: Applications in Organic Synthesis. *Chem. Rev.* **2013**, *113* (7), 5322–5363. <https://doi.org/10.1021/cr300503r>.

- (18) Narayanam, J. M. R.; Stephenson, C. R. J. Visible Light Photoredox Catalysis: Applications in Organic Synthesis. *Chem. Soc. Rev.* **2011**, *40* (1), 102–113. <https://doi.org/10.1039/b913880n>.
- (19) Wang, C. S.; Dixneuf, P. H.; Soulé, J. F. Photoredox Catalysis for Building C-C Bonds from C(Sp²)-H Bonds. *Chem. Rev.* **2018**, *118* (16), 7532–7585. <https://doi.org/10.1021/acs.chemrev.8b00077>.
- (20) Bosset, C.; Beucher, H.; Bretel, G.; Pasquier, E.; Queguiner, L.; Henry, C.; Vos, A.; Edwards, J. P.; Meerpoel, L.; Berthelot, D. Minisci-Photoredox-Mediated α -Heteroarylation of N-Protected Secondary Amines: Remarkable Selectivity of Azetidines. *Org. Lett.* **2018**, *20* (19), 6003–6006. <https://doi.org/10.1021/acs.orglett.8b00991>.
- (21) Li, G. X.; Hu, X.; He, G.; Chen, G. Photoredox-Mediated Minisci-Type Alkylation of N-Heteroarenes with Alkanes with High Methylene Selectivity. *ACS Catal.* **2018**, *8* (12), 11847–11853. <https://doi.org/10.1021/acscatal.8b04079>.
- (22) Dong, J.; Xia, Q.; Lv, X.; Yan, C.; Song, H.; Liu, Y.; Wang, Q. Photoredox-Mediated Direct Cross-Dehydrogenative Coupling of Heteroarenes and Amines. *Org. Lett.* **2018**, *20* (18), 5661–5665. <https://doi.org/10.1021/acs.orglett.8b02389>.
- (23) Grainger, R.; Heightman, T. D.; Ley, S. V.; Lima, F.; Johnson, C. N. Enabling Synthesis in Fragment-Based Drug Discovery by Reactivity Mapping: Photoredox-Mediated Cross-Dehydrogenative Heteroarylation of Cyclic Amines. *Chem. Sci.* **2019**, *10* (8), 2264–2271. <https://doi.org/10.1039/C8SC04789H>.
- (24) Devari, S.; Shah, B. A. Visible Light-Promoted C-H Functionalization of Ethers and Electron-Deficient Arenes. *Chem. Commun.* **2016**, *52* (7), 1490–1493. <https://doi.org/10.1039/c5cc08817h>.
- (25) Jin, J.; MacMillan, D. W. C. Direct α -Arylation of Ethers through the Combination of Photoredox-Mediated C-H Functionalization and the Minisci Reaction. *Angew. Chemie - Int. Ed.* **2015**, *54* (5), 1565–1569. <https://doi.org/10.1002/anie.201410432>.
- (26) Quattrini, M. C.; Fujii, S.; Yamada, K.; Fukuyama, T.; Ravelli, D.; Fagnoni, M.; Ryu, I. Versatile Cross-Dehydrogenative Coupling of Heteroaromatics and Hydrogen Donors via Decatungstate Photocatalysis. *Chem. Commun.* **2017**, *53* (15), 2335–2338. <https://doi.org/10.1039/c6cc09725a>.
- (27) Okugawa, N.; Moriyama, K.; Togo, H. Introduction of Ether Groups onto Electron-Deficient Nitrogen-Containing Heteroaromatics Using Radical Chemistry under Transition-Metal-Free Conditions. *European J. Org. Chem.* **2015**, *2015* (22), 4973–4981. <https://doi.org/10.1002/ejoc.201500584>.
- (28) He, F. S.; Ye, S.; Wu, J. Recent Advances in Pyridinium Salts as Radical Reservoirs in Organic Synthesis. *ACS Catal.* **2019**, *9* (10), 8943–8960. <https://doi.org/10.1021/acscatal.9b03084>.
- (29) Rössler, S. L.; Jelier, B. J.; Magnier, E.; Dagousset, G.; Carreira, E. M.; Togni, A. Pyridinium Salts as Redox-Active Functional Group Transfer Reagents. *Angew. Chemie - Int. Ed.* **2020**, *59* (24), 9264–9280. <https://doi.org/10.1002/anie.201911660>.
- (30) Kim, I.; Kang, G.; Lee, K.; Park, B.; Kang, D.; Jung, H.; He, Y. T.; Baik, M. H.; Hong, S. Site-Selective Functionalization of Pyridinium Derivatives via Visible-Light-Driven Photocatalysis with Quinolinone. *J. Am. Chem. Soc.* **2019**, *141* (23), 9239–9248. <https://doi.org/10.1021/jacs.9b02013>.
- (31) Jeon, J.; He, Y. T.; Shin, S.; Hong, S. Visible-Light-Induced Ortho-Selective Migration on Pyridyl Ring: Trifluoromethylative Pyridylation of Unactivated Alkenes. *Angew. Chemie - Int. Ed.* **2020**, *59* (1), 281–285. <https://doi.org/10.1002/anie.201912746>.
- (32) Jung, S.; Lee, H.; Moon, Y.; Jung, H. Y.; Hong, S. Site-Selective c-h Acylation of Pyridinium Derivatives by Photoredox Catalysis. *ACS Catal.* **2019**, *9* (11), 9891–9896. <https://doi.org/10.1021/acscatal.9b03367>.
- (33) Kim, I.; Park, B.; Kang, G.; Kim, J.; Jung, H.; Lee, H.; Baik, M. H.; Hong, S. Visible-Light-Induced Pyridylation of Remote C(Sp³)-H Bonds by Radical Translocation of N-Alkoxy-pyridinium Salts. *Angew. Chemie - Int. Ed.* **2018**, *57* (47), 15517–15522. <https://doi.org/10.1002/anie.201809879>.
- (34) Kim, Y.; Lee, K.; Mathi, G. R.; Kim, I.; Hong, S. Visible-Light-Induced Cascade Radical Ring-Closure and Pyridylation for the Synthesis of Tetrahydrofurans. *Green Chem.* **2019**, *21* (8), 2082–2087. <https://doi.org/10.1039/c9gc00414a>.
- (35) Liang, X. A.; Niu, L.; Wang, S.; Liu, J.; Lei, A. Visible-Light-Induced C(Sp³)-H Oxidative Arylation with Heteroarenes. *Org. Lett.* **2019**, *21* (7), 2441–2444. <https://doi.org/10.1021/acs.orglett.9b00744>.
- (36) Niu, L.; Liu, J.; Liang, X. A.; Wang, S.; Lei, A. Visible Light-Induced Direct α C-H Functionalization of Alcohols. *Nat. Commun.* **2019**, *10* (1), 1–7. <https://doi.org/10.1038/s41467-019-08413-9>.
- (37) Bieszczad, B.; Perego, L. A.; Melchiorre, P. Photochemical C-H Hydroxyalkylation of Quinolines and Isoquinolines. *Angew. Chemie - Int. Ed.* **2019**, *58* (47), 16878–16883. <https://doi.org/10.1002/anie.201910641>.
- (38) Quint, V.; Morlet-Savary, F.; Lohier, J. F.; Lalevée, J.; Gaumont, A. C.; Lakhdar, S. Metal-Free, Visible Light-Photocatalyzed Synthesis of Benzo[b]Phosphole Oxides: Synthetic and Mechanistic Investigations. *J. Am. Chem. Soc.* **2016**, *138* (23), 7436–7441. <https://doi.org/10.1021/jacs.6b04069>.
- (39) Inial, A.; Morlet-Savary, F.; Lalevée, J.; Gaumont, A. C.; Lakhdar, S.; Lakhdar, S. Visible-Light-Mediated Access to Phosphate

Esters. *Org. Lett.* **2020**, *22* (11), 4404–4407. <https://doi.org/10.1021/acs.orglett.0c01409>.

- (40) Quint, V.; Chouchène, N.; Askri, M.; Lalevée, J.; Gaumont, A. C.; Lakhdar, S. Visible-Light-Mediated α -Phosphorylation of: N - Aryl Tertiary Amines through the Formation of Electron-Donor-Acceptor Complexes: Synthetic and Mechanistic Studies. *Org. Chem. Front.* **2019**, *6* (1), 41–44. <https://doi.org/10.1039/c8qo00985f>.
- (41) Quint, V.; Noël-Duchesneau, L.; Lagadic, E.; Morlet-Savary, F.; Lalevée, J.; Gaumont, A.-C.; Lakhdar, S. Metal-Free Generation of Phosphorus-Centered Radicals for the Synthesis of Phosphorus-Based Heterocycles: A Personal Account. *Synth.* **2017**, *49* (15). <https://doi.org/10.1055/s-0036-1588499>.