Unexpected formation of 2-alkynylpyridines from iodo-selenopyridines by selective activation of C Se over C I bonds

Christine Tran, Guillaume Lefèvre, Mansour Haddad, Virginie Ratovelomanana-Vidal

► To cite this version:

Christine Tran, Guillaume Lefèvre, Mansour Haddad, Virginie Ratovelomanana-Vidal. Unexpected formation of 2-alkynylpyridines from iodo-selenopyridines by selective activation of C Se over C I bonds. *Catalysis Communications*, 2019, 130, pp.105757. 10.1016/j.catcom.2019.105757. hal-03011778

HAL Id: hal-03011778

<https://hal.science/hal-03011778>

Submitted on 18 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Short communication

Unexpected formation of 2-alkynylpyridines from iodo-selenopyridines by selective activation of C–Se over C–I bonds

Christine Tran, Guillaume Lefèvre, Mansour Haddad, Virginie Ratovelomanana-Vidal *

PSL Research University, Chimie ParisTech, CNRS, Institute of Chemistry for Life and Health Sciences, CSB2D 11 rue Pierre et Marie Curie, Paris 75005, France

ARTICLE INFO

Keywords:

Selenopyridines
Cross-coupling
Alkynylpyridines
Palladium
Copper
Alkynes

ABSTRACT

The synthesis of bicyclic 2-alkynylpyridines is performed in the presence of iodo-selenopyridines and terminal alkynes, using a palladium(II) catalyst, a copper(I) co-catalyst, and triethylamine as a base. This facile cross-coupling reaction leads to a wide variety of 2-alkynylpyridines in good yields. Mechanistic investigations provided by DFT calculations reveal that the reaction should involve a key selective activation of the C–Se bond of the 2-selenopyridyl moiety by the *in situ* generated Pd(0) catalyst, preferentially to the C–I bond activation. A transmetalation step with copper acetylide, followed by a reductive elimination, affords the 2-alkynylpyridine product.

1. Introduction

2-Alkynylpyridines exhibit interesting biological properties as second mitochondria-derived activator of caspases (SMAC) mimetics. These molecules could be applied for the treatment and prevention of abnormal cell proliferation in cancer diseases [1,2]. Additionally, 2-alkynylpyridines represent synthetically relevant intermediates in organic synthesis, because their triple bonds could readily undergo diverse transformations, such as annulation [3], cyclization [4], or 1,3-dipolar cycloaddition [5]. In this context, these compounds contribute to the construction of a large array of heterocyclic frameworks, including azaindoles [6,7], furo[3,4-*b*]pyridines [8], indolizines [9], isoquinolines [10], aryl-substituted pyridines [11], and *N*-metallacycles [12]. As a result, the construction of functionalized 2-alkynylpyridines constitutes an attractive challenge, considering the synthetic and biological aspects of this scaffold.

We have recently developed a practical route to access iodo-selenopyridine derivatives [13]. We envisaged the functionalization of these selenopyridines, through a Sonogashira cross-coupling reaction. First, a reaction was performed with diarylselenium **1a** and 1-hexyne (**2a**). Surprisingly, the desired product **4** was not obtained, but 2-alkynylpyridine **3a** was isolated with 91% yield (Scheme 1).

In view of this unexpected result (Scheme 1), we focused our studies around the construction of 2-alkynylpyridines derivatives [14–25]. In addition, some DFT calculations were performed to enlighten the

origin of this result, thus providing mechanistic insights into this access to the alkynylpyridine pattern.

2. Results and discussion

In light of the preliminary result, we thus investigated the scope of this reaction in the presence of diverse diarylselenium compounds. As depicted in Table 1, this method allows the access to a wide range of 2-alkynylpyridines from a large array of selenopyridines and diverse aryl/alkyl terminal alkynes. From selenopyridine **1a**, alkynylpyridines **3a–e** were obtained in good to excellent yields (71–91%, Table 1, entries 1 to 5). The reaction was also well-tolerated in the presence of 2-ethynylpyridine, with 75% yield (entry 6). Cross-coupling reactions [26], performed in the presence of **1b** [27], provided the desired compounds in 50 to 86% yields (Table 1, entries 7 to 11). The structure of **3i** was confirmed by 2D NMR (NOESY experiment, *cf.* Supporting Information). It is noticeable that the reaction involving phenylacetylene led to pyridine derivatives with lower yields, probably due to steric effects (Table 1, entry 10). A good tolerance was also noted with the selenopyridines **1c** and **1d** [27], bearing an oxygen atom or a diketone group, affording the expected alkynylpyridines **3l–o** with high yields (68 to 94%). Moreover, the preparation of unsymmetrical alkynylpyridines was envisaged with selenium derivatives **1e** and **1f** [27], furnishing the corresponding products **3p** and **3q** in 97 and 54% yield, respectively.

* Corresponding author.

Email address: virginie.vidal@chimie-paristech.fr (V. Ratovelomanana-Vidal)

In summary, the unexpected formation of 2-alkynylpyridines obtained from iodo-selenopyridines through a palladium/copper co-catalytic system was reported. This transformation provides a broad array of substrates in good to excellent yields. Mechanistic studies demonstrated that the key step of this transformation is a selective activation of the C—Se bond of the starting iodo-selenopyridine, which undergoes oxidative addition of the palladium(0) catalyst, whereas the C—I bond remains unreactive. DFT calculations suggest that this selectivity

Table 2
Coupling reaction for mechanistic insights with selenopyridine 1a.

Entry	Reaction conditions	Yield (%)
1	Without CuI and PdCl ₂ (PPh ₃) ₂	nr
2	With CuI only	nr
3	With PdCl ₂ (PPh ₃) ₂ only	nr
4	With CuI and PdCl ₂ (PPh ₃) ₂ , without NEt ₃	nr

a) Cu(I)/Cu(III) oxidative addition pathway

b) Pd(0)/Pd(II) C-I oxidative addition

c) Pd(0)/Pd(II) C-Se oxidative addition

Scheme 2. Competitive pathways investigated by DFT computation.

Scheme 3. DFT-computed pathway for the formation of I.

4. Experimental

4.1. General procedure for the synthesis of 2-alkynylpyridines

In a round-bottom flask, PdCl₂(PPh₃)₂ (20 mol%) and CuI (10 mol%) were solubilized in THF (1.5 mL) and NEt₃ (1.5 mL). The yellow solution was stirred for 10 min. A mixture of selenopyridine 1 (1.0 eq) and alkyne 2 (4.0 eq) in THF (1.5 mL) and NEt₃ (1.5 mL) was then added. The reaction was stirred at 50 °C for 8 h. The crude mixture was then quenched with a saturated aqueous solution of ammo-

nium chloride. The organic layer was extracted with ethyl acetate (x3), washed with water, dried over MgSO₄, filtered and concentrated under reduced pressure. The residue was purified by flash chromatography over silica gel to afford the corresponding alkynylpyridines.

4.2. DFT calculations

DFT calculations were performed using the Gaussian 09 code. See Gaussian 09, Revision D.01, M. J. Frisch et al., Gaussian, Inc., Wallingford CT, 2013, and see SI for additional details.

Scheme 4. Postulated reaction mechanism.

Acknowledgements

We thank the Ministère de l'Enseignement Supérieur de la Recherche et de l'Innovation, Chimie ParisTech (C.T.) and the CNRS for financial support. We are grateful to Dr. M.-N. Rager (Chimie ParisTech) for 2D NMR experiments.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.catcom.2019.105757>.

References

- [1] U. Reiser, G. Bader, W. Spevak, A. Steffen, A.L. Parkes, US Pat. 0225567 A1, 2013.
- [2] U. Reiser, G. Bader, W. Spevak, A. Steffen, A.L. Parkes, US Pat. 8859541 B2, 2014.
- [3] F. Roschangar, J. Liu, E. Estanove, M. Dufour, S. Rodríguez, V. Farina, E. Hickey, A. Hossain, P.-J. Jones, H. Lee, B.Z. Lu, R. Varsolona, J. Schröder, P. Beaulieu, J. Gillard, C.H. Senanayake, Tetrahedron Lett. 49 (2008) 363–366.
- [4] M. Dell'Acqua, G. Abbiati, A. Arcadi, E. Rossi, Org. Biomol. Chem. 9 (2011) 7836–7848.
- [5] K.A. Johnston, R.W. Allcock, Z. Jiang, I.D. Collier, H. Blakli, G.M. Rosair, P.D. Bailey, K.M. Morgan, Y. Kohno, D.R. Adams, Org. Biomol. Chem. 6 (2008) 175–186.
- [6] S. Cacchi, G. Fabrizi, L.M. Parisi, J. Comb. Chem. 7 (2005) 510–512.
- [7] C. Harcken, Y. Ward, D. Thomson, D. Riether, Synlett (20) (2005) 3121–3125.
- [8] G.P. Jana, B.K. Ghorai, Tetrahedron 63 (2007) 12015–12025.
- [9] D. Chernyak, S.B. Gadamssetty, V. Gevorgyan, Org. Lett. 10 (2008) 2307–2310.
- [10] S. Mukherjee, G.P. Jana, B.K. Ghorai, J. Organomet. Chem. 694 (2009) 4100–4106.
- [11] O.R. Abid, M. Nawaz, M.F. Ibad, R.A. Khera, V. Iaroshenko, P. Langer, Org. Biomol. Chem. 9 (2011) 2185–2191.
- [12] J.W. Teo, V.S. Sridevi, W.K. Leong, Organometallics 33 (2014) 1174–1180.
- [13] C. Tran, M. Haddad, V. Ratovelomanana-Vidal, Synthesis 51 (2019) 2532–2541.
- [14] T. León, P. Quinio, Q. Chen, P. Knochel, Synthesis 46 (2014) 1374–1379.
- [15] S. Roy, H. Plenio, Adv. Synth. Catal. 352 (2010) 1014–1022.
- [16] H. Kim, P.H. Lee, Adv. Synth. Catal. 351 (2009) 2827–2832.
- [17] G. Fabrizi, A. Goggiani, A. Sferazza, S. Cacchi, Angew. Chem. Int. Ed. 49 (2010) 4067–4070.
- [18] G. Úr, T. Kálai, M. Balog, B. Bognár, G. Gulyás-Fekete, K. Hideg, Synth. Commun. 45 (2015) 2122–2129.
- [19] W. Zhang, J. Cheng, Z. Huang, Q. Zhou, X. Chen, J. Qian, J. Chem. Res. 12 (2006) 781–782.
- [20] Y. Kelgokmen, M. Zora, RSC Adv. 6 (2016) 4608–4621.
- [21] K.C. Nicolaou, K. Namoto, A. Ritzén, T. Ulven, M. Shoji, J. Li, G. D'Amico, D. Liotta, C.T. French, M. Wartmann, K.-H. Altmann, P. Giannakakou, J. Am. Chem. Soc. 123 (2001) 9313–9323.
- [22] C.S. Consorti, F.R. Flores, F. Rominger, J. Dupont, Adv. Synth. Catal. 348 (2006) 133–141.
- [23] M. Zhang, T. Jia, C.Y. Wang, P.J. Walsh, J. Am. Chem. Soc. 137 (2015) 10346–10350.
- [24] D. Bhatt, N. Patel, H. Chowdhury, P.V. Bharatam, A. Goswami, Adv. Synth. Catal. 360 (2018) 1876–1882.
- [25] P.J. Hamzik, A.-S. Goutierre, T. Sakai, R.L. Danheiser, J. Organomet. Chem. 82 (2017) 12975–12991.
- [26] Iodo-selenopyridine derivatives **1** (Table 1) were also evaluated in additional cross-coupling reactions. The Suzuki coupling was attempted with **1** (X = indene-1,3-dione) with phenyl boronic acid (2.0 equiv), Pd(OAc)₂ (5 mol%), K₂CO₃ (3.0 equiv), CataCXium A (0.1 equiv) in DMF (0.3 M) at 90 °C for 48 h, affording the recovered starting material. The Heck reaction was performed with **1** (X = 2,2-dimethyl-1,3-dioxane) in standard conditions (*tert*-butyl acrylate (3.5 equiv), Pd(OAc)₂ (20 mol%), K₂CO₃ (1.0 equiv), P^tBu₃ (2.0 equiv), DMA, 80 °C, 48 h), providing mainly degradation products.
- [27] For the synthesis of **1b–1f**, see ref. 13.
- [28] C. Glaser, Ber. Dtsch. Chem. Ges. 2 (1869) 422–424.
- [29] P. Siemsen, R.C. Livingston, F. Diederich, Angew. Chem. Int. Ed. 39 (2000) 2632–2657.
- [30] M. Font, T. Parella, M. Costas, X. Ribas, For recent examples of Cu(I)/Cu(III) catalytic cycles involving the formation of C—Se bonds, Organometallics 31 (2012) 7976–7982.
- [31] K. Osakada, M. Maeda, Y. Nakamura, T. Yamamoto, A. Yamamoto, J. Chem. Soc. Chem. Commun. (6) (1986) 442–443.