Existence, uniqueness and stability of fractional impulsive functional differential inclusions

José Vanterler, Kishor Kucche

► To cite this version:

José Vanterler, Kishor Kucche. Existence, uniqueness and stability of fractional impulsive functional differential inclusions. 2020. hal-03012073

HAL Id: hal-03012073

<https://hal.science/hal-03012073>

Preprint submitted on 18 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Existence, uniqueness and stability of fractional impulsive functional differential inclusions

J. Vanterler da C. Sousa · Kishor D. Kucche

Received: date / Accepted: date

Abstract In the paper, we discuss necessary and sufficient conditions to obtain the existence, uniqueness and stability of solutions of fractional impulsive functional differential equations towards the ψ -Caputo fractional derivative, through fixed point theorem, Arzela-Ascoli theorem and multivalued analysis theory.

Keywords Fractional Differential inclusions, existence and uniqueness, stability, Banach fixed point, multivalued analysis theory

Mathematics Subject Classification (2010) MSC 26A33, 34A08, 34A12, 34A60, 49J24, 49K24.

1 Introduction

Fractional differential equations and inclusions arise naturally in various fields of science, with numerous applications [1, 35, 8, 36, 12, 25, 18, 21, 23, 24, 39]. Over the years, the search for discussing properties of solutions of fractional differential equations, namely: existence, uniqueness, attractiveness, controllability, observability, among others, has been of great community attention [26–29, 6, 4, 9, 34, 38]. Some authors dedicate to discuss more analytical properties of solutions, others already direct to applications.

In recent years, Benchohra et al. [24] and Chang et al. [11], have investigated the existence of solutions for impulsive functional inclusions Eq.(1) with $p(t) = 1$ and

J. Vanterler da C. Sousa
Centro de Matemática, Computação e Cognição, Universidade Federal do ABC,
Avenida dos Estados, 5001, Bairro Bangu, 09.210-580, Santo André, SP - Brazil
E-mail: vanterlermatematico@hotmail.com, jose.vanterler@edu.ufabc.br

Kishor D. Kucche
Department of Mathematics, Shivaji University, Kolhapur-416 004, Maharashtra, India.

Eq.(1), respectively

$$\begin{cases} (p(t)y'(t))' \in F(t, u_t), t \in J = [0, a], t \neq t_k, k = 1, 2, \dots, m, \\ \delta y(t)|_{t=t_k} = I_k(y(t_k^-)), k = 1, 2, \dots, m, \\ \delta y'(t)|_{t=t_k} = J_k(y(t_k^-)), k = 1, 2, \dots, m, \\ y(t) = \phi(t), t \in [-r, 0], y'(0) = \eta. \end{cases} \quad (1)$$

Other works on existence, uniqueness, controllability involving differential equations of a integer order, can be obtained at [17, 3, 5, 7, 11, 14, 20].

Em 2018 Wang et al. [35] discuss the controllability of non-instantaneous impulsive semilinear differential inclusions involving integer and fractional order derivatives i.e.,

$$\begin{cases} x'(t) \in A(t)x(t) + F(t, x_t) + B(u(t)), a.e.t \in \bigcup_{i=1}^{i=m} (s_i, t_{i+1}] \\ x(t_i^+) = g_i(t_i, x(t_i^-)) & i = 1, \dots, m \\ x(t) = g_i(t, x(t_i^-)) & , t \in \bigcup_{i=1}^{i=m} (t_i, s_i] \\ x(t) = \Psi(t) & , t \in [-r, 0] \end{cases}$$

and

$$\begin{cases} {}^C D_{s_i, t}^\alpha x(t) \in A(t)x(t) + F(t, x_t) + B(z(t)), a.e.t \in \bigcup_{i=1}^{i=m} (s_i, t_{i+1}] \\ x(t_i^+) = g_i(t_i, x(t_i^-)) & i = 1, \dots, m \\ x(t) = g_i(t, x(t_i^-)) & , t \in \bigcup_{i=1}^{i=m} (t_i, s_i] \\ x(t) = \Psi(t) & , t \in (-r, 0] \end{cases}$$

where $\alpha \in (0, 1)$, ${}^C D_{s_i, t}^\alpha(\cdot)$ is the Caputo fractional derivative with lower limit s_i of order α , A is the infinitesimal generator of C_0 -semigroup $\{T(t) : t \leq 0\}$, and the control function z is given in $L^p(J, X)$, $p > \frac{1}{\alpha}$, a Banach space of admissible control functions.

In the present paper, we highlight the advances in the theory of fractional differential equations involving fractional derivatives with respect to another function, since they are more general and global operators that allow the discussion and analysis of a wide class of particular cases. The discussions of these problems are motivated by the fractional operators, ψ -Hilfer and ψ -Caputo. Some works in this segment can be obtained at [19, 22, 15, 16, 2]. Although there is an interesting range of published works, many open questions and problems, as well as new tools need to be presented, in this sense, one of the main purpose of this work, is to provide new results for the area and consequently contribute to its strengthening.

Let $0 = t_1 < t_1 < \dots < t_p < t_{p+1} = T$, where T is a positive constant. Consider $F : [0, T] \times D \rightarrow \mathcal{P}(\mathbb{R}^n)$ is a multivalued map, where $D = \{\xi : [-r, 0] \rightarrow \mathbb{R}^n, \xi \text{ is continuous everywhere except for a finite number of points } \bar{t} \text{ at which } \xi(\bar{t}^-) \text{ and } \xi(\bar{t}^+) \text{ exist with } \xi(\bar{t}^-) = \xi(\bar{t})\}$, and $\mathcal{P}(\mathbb{R}^n)$ is the family of all nonempty subsets of $\mathbb{R}^n$.

Motivated by the above work and the innumerable open questions, in this paper, we consider the α^{th} order impulsive fractional functional differential equations

$$\begin{cases} {}^C\mathbb{D}_{0+}^{\alpha,\psi}u(t) \in F(t, u_t), \quad t \in J = [0, T], \quad t \neq t_k, \quad k = 1, 2, \dots, p, \\ \delta u^{(i)}(t_k) = \mathcal{I}_{ik}(u(t_k^-)), \quad i = 0, 1, \dots, m-1, \quad k = 1, 2, \dots, p, \\ u(t) = \phi(t), \quad t \in [-r, 0], \\ u^{(i)}(0) = \eta_i, \quad i = 1, 2, \dots, m-1, \end{cases} \quad (2)$$

where ${}^C\mathbb{D}_{0+}^{\alpha,\psi}(\cdot)$ is the ψ -Caputo fractional derivative of order $m-1 < \alpha \leq m$, $m \in \mathbb{N}$, $\phi \in D$ and $\mathcal{I}_{ik} \in C(\mathbb{R}^n, \mathbb{R}^n)$ ($i = 0, 1, \dots, m-1, k = 1, 2, \dots, p$). Moreover, $u^{(i)} : [0, T] \rightarrow \mathbb{R}^n$ which are piecewise continuous in $[0, T]$ with points of discontinuity of the first kind at the points $t_k \in [0, T]$, i.e., there exists the limits $u^{(i)}(t_k^+) < \infty$ and $u^{(i)}(t_k^-) = u^{(i)}(t_k) < \infty$, $u^{(m)} : [0, T] \rightarrow \mathbb{R}^n$, and $\delta u^{(i)}(t_k) = u^{(i)}(t_k^+) - u^{(i)}(t_k)$ ($i = 0, 1, \dots, m-1, k = 1, 2, \dots, p$). For any continuous function u defined on $[-r, T] \setminus \{t_1, \dots, t_p\}$ and any $t \in [0, T]$, we denote by u_t the element of D defined by $u_t(\theta) = u(t + \theta)$, $\theta \in [-r, 0]$. Here $u_t(\cdot)$ represents the history of the state from $t-r$, up to the present time t .

For the discussion of the main results of this paper, we will consider the following conditions:

- (A1) $F : [0, T] \times D \rightarrow \mathcal{P}(\mathbb{R}^n)$ has the property that $F(\cdot, \xi) : [0, T] \rightarrow P_{cp}(\mathbb{R}^n)$ is measurable for each $\xi \in D$.
- (A2) There exists nonnegative constants c_{ik} ($i = 0, 1, \dots, m-1, k = 1, \dots, p$) such that $|I_{ik}(u(t_k)) - I_{ik}(v(t_k))|, I_{ik}(0) = 0, i = 0, 1, \dots, m-1, k = 1, \dots, p$ and for all $u, v \in \Omega$.
- (A3) There exists a function $l \in L^1([0, T], \mathbb{R}_+)$ such that $H_d(F(t, \xi), F(t, \varphi)) \leq l(t) \|\xi - \varphi\|_D$, for a.e. $t \in [0, T]$ and any $\xi, \varphi \in D$, and $E_d(0, F(t, 0)) \leq l(t)$, for a.e. $t \in [0, T]$.
- (A4) Let $F : [0, T] \times D \rightarrow \mathcal{P}(\mathbb{R}^n)$ be a nonempty, compact valued multivalued map such that $(t, \xi) \rightarrow F(t, \xi)$ is $\mathcal{L} \times \mathcal{B}$ measurable, and $\xi \rightarrow F(t, \xi)$ is l.s.c for a.e. $t \in [0, T]$.
- (A5) There exists a function $M \in L^1([0, T], \mathbb{R}_+)$ such that $\|F(t, \xi)\| = \sup \{ \|v(t)\| : v(t) \in F(t, \xi) \} \leq M(t)$ for each $t \in [0, T]$.

We will here highlight the main contributions of this article explicitly and clearly. The first main contribution is to discuss the existence of solutions for the Eq.(2), that is, the following results:

Theorem 1 Assume that (A1), (A2) and (A3) are satisfied. Then Eq.(2) has at least one solution on $[-r, T]$, provided

$$\mathcal{L} = \frac{(\psi(T) - \psi(0))^{\alpha-1}}{\Gamma(\alpha)} \psi(T) \|l\|_{L^1} + \sum_{k=1}^p \sum_{i=0}^{m-1} \frac{(\psi(T) - \psi(0))^i}{i!} C_{ik} < 1.$$

Theorem 2 In addition to (A4) and (A5), assume that the following condition hold:

- (A6) There exists constants d_{ik} ($i = 0, 1, \dots, m-1, k = 1, 2, \dots, p$), such that $|\mathcal{I}_{ik}(u(t_k))| \leq d_{ik}|u(t_k)|$ for each $u \in \Omega$. Then Eq.(2) has at least one solution on $[-r, T]$, provided

$$\mathcal{L} = \sum_{k=1}^p \sum_{i=0}^{m-1} \frac{(\psi(T) - \psi(t_k))^i}{i!} d_{ik} < 1.$$

The second and third main contributions, is to guarantee the uniqueness and stability of solutions for Eq.(2), in other words, is to discuss the following results:

Theorem 3 *In addition to (A2), (A4) and (A5), assume that the following condition hold:*

(A7) *There exists non-negative constant b , such that $|f(u)(t) - f(v)(t)| \leq b\|u - v\|_\Omega$ for any $u, v \in \Omega$ where f is deduced in the proof of Theorem 2. Then Eq.(2) has a unique solution on $[-r, T]$, provided*

$$\mathcal{L} = \frac{b(\psi(T) - \psi(0))^\alpha}{\Gamma(\alpha + 1)} + \sum_{k=1}^p \sum_{i=0}^{m-1} C_{ik} \frac{\psi(T)^i}{i!} < 1.$$

Theorem 4 *Assume that all the conditions of Theorem 3 hold. Then the solution of the Eq.(2) is uniformly stable.*

In the rest, the paper is organized as follows. In section 2, we present definitions and essential results in obtaining the main results of this paper. Section 3 is divided into 3 subsections, each of which discusses a contribution from this paper, that is, the existence, uniqueness and uniform stability of solutions from Eq.(2).

2 Preliminaries

In this section, we present some fundamental concepts and results for the development of this paper.

Let (X, d) be a metric space and $N : X \rightarrow \mathcal{P}(X)$ be a multivalued map with nonempty values. We use the notations $P(X) = \{Y \in \mathcal{P}(X) : Y \neq \emptyset\}$, $P_{cl}(X) = \{Y \in \mathcal{P}(X) : Y \text{ closed}\}$, $P_b(X) = \{Y \in \mathcal{P}(X) : Y \text{ bounded}\}$, $P_c = \{Y \in \mathcal{P}(X) : Y \text{ convex}\}$, and $P_{cp}(X) = \{Y \in \mathcal{P}(X) : Y \text{ compact}\}$.

Let $\alpha \in (0, 1)$ and $f \in L^1[a, b]$, $[a, b]$ be a finite or infinite interval of the real line $\mathbb{R}$ ($-\infty \leq a < b \leq \infty$). Also let $\psi(\cdot)$ be an increasing and positive continuous function on $[a, b]$, having a continuous derivatives $\psi'(x) \neq 0$ on (a, b) . The left-sided and right-sided fractional integrals of a function f with respect to another function ψ on $[a, b]$, is defined [30–32]

$$\mathcal{J}_{a+}^{\alpha; \psi} f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x \psi'(s) (\psi(t) - \psi(s))^{\alpha-1} f(s) ds. \quad (3)$$

On the other hand, let $n - 1 < \alpha < n$, with $n \in \mathbb{N}$, $I = [a, b]$, is the interval such that $-\infty \leq a < b \leq \infty$ and there exist two function $f, \psi \in C^n([a, b], \mathbb{R})$ such that ψ is increasing and $\psi'(x) \neq 0$, for all $x \in I$. The ψ -Hilfer fractional derivatives left-sided and right-sided ${}^H\mathbb{D}_{a+,x}^{\alpha,\beta;\psi}(\cdot)$ function or order α and type $0 \leq \beta \leq 1$ is defined by [30–32]

$${}^H\mathbb{D}_{a+,x}^{\alpha,\beta;\psi} f(x) = \mathcal{J}_{a+,x}^{\beta(n-\alpha);\psi} \left(\frac{1}{\psi'(x)} \frac{d}{dx} \right)^n \mathcal{J}_{a+,x}^{(1-\beta)(n-\alpha);\psi} f(x). \quad (4)$$

The ψ -Hilfer fractional derivatives defined as above can be written in the following form

$${}^H\mathbb{D}_{a+,x}^{\alpha,\beta;\psi} f(x) = \mathcal{I}_{a+,x}^{\gamma-\alpha;\psi} \mathcal{D}_{a+,x}^{\gamma;\psi} f(x)$$

where $\mathcal{D}_{a+,x}^{\gamma;\psi}(\cdot)$ is left-sided ψ -Riemann-Liouville fractional derivative.

Taking limit $\beta \rightarrow 1$ on both sides of Eq.(4), we have the Caputo fractional derivative with respect to another function, given by [30]

$${}^C\mathbb{D}_{a+,x}^{\alpha;\psi} f(x) = \mathcal{I}_{a+,x}^{(n-\alpha);\psi} \left(\frac{1}{\psi'(x)} \frac{d}{dx} \right)^n f(x). \quad (5)$$

For the formulation of the main result of this paper, we use the Caputo fractional derivative with respect to another function given by Eq.(5).

For the definitions and lemmas presented below, we use the following references [37, 1, 8, 10, 13, 33].

Definition 1 A multivalued map $N : [0, T] \rightarrow P_{cl}(X)$ is said to be measurable if for each $x \in X$ the function $g : [0, T] \rightarrow \mathbb{R}_+$, defined by

$$g(t) = E_d(x, N(t)) = \inf \{ |x - z| : z \in N(t) \}$$

belongs to $L^1([0, T], \mathbb{R})$.

Definition 2 Let X be a nonempty closed subset of $\mathbb{R}^n$, and $N : X \rightarrow \mathcal{P}(\mathbb{R}^n)$ be a multivalued map with nonempty closed values. N is lower semicontinuous (l.s.c) on X if the set $\{x \in X : N(x) \cap C \neq \emptyset\}$ is open for each open set C in $\mathbb{R}^n$.

Definition 3 Let $A_{12} \subseteq [0, T] \times D$. A_{12} is $\mathcal{L} \otimes \mathcal{B}$ if A_{12} belongs to the σ -algebra generated by all sets of the form $\mathcal{J} \times \mathcal{D}$ where $\mathcal{J}$ is Lebesgue measurable in $[0, T]$ and $\mathcal{D}$ is Borel measurable in D .

Definition 4 Let X be a separable metric space and $\mathcal{N} : X \rightarrow \mathcal{P}(L^1([0, Y], \mathbb{R}^n))$ be a multivalued operator. We say $\mathcal{N}$ has property (BC) if:

1. $\mathcal{N}$ is (l.s.c).
2. $\mathcal{N}$ has nonempty closed and decomposable values.

In order to define the solutions of Eq.(2), we consider the following spaces:

$$PC = \left\{ u : [0, T] \rightarrow \mathbb{R}^n \mid u_k \in C((t_k, t_{k+1}], \mathbb{R}^n), k = 0, \dots, p \right. \\ \left. \text{and there exist } u(t_k^-) \text{ and } u(t_k^+) \text{ with } u(t_k^-) = u(t_k), k = 0, \dots, p \right\}$$

,which is a Banach space with the norm $\|u\|_{PC} = \max \left\{ \|u_k\|_{(t_k, t_{k+1}]}, k = 0, \dots, p \right\}$ where u_k is the restriction of u to $(t_k, t_{k+1}]$, $k = 0, \dots, p$.

Definition 5 Let $F : [0, T] \times D \rightarrow \mathcal{P}(\mathbb{R}^n)$ be a multivalued map with nonempty compact values. Assign to F the multivalued operator $\mathcal{F} : \Omega \rightarrow \mathcal{P}(L^1([0, T], \mathbb{R}^n))$ by letting $\mathcal{F}(u) = \{v \in L^1([0, T], \mathbb{R}^n) \mid v(t) \in F(t, u_t) \text{ for a.e } t \in [0, T]\}$. The operator $\mathcal{F}$ is called the Niemytzki operator associated with F .

Definition 6 A function $u \in \Omega$ is said to be a solution of Eq.(2) if u satisfies Eq.(2).

Definition 7 A multivalued operator $N : X \rightarrow P_{cl}(X)$ is called:

1. γ -Lipschitz if and only if there exists $\gamma > 0$ such that $H_d(N(x), N(y)) \leq \gamma d(x, y)$ for each $x, y \in X$.
2. contraction if and only if it is γ -Lipschitz with $\gamma < 1$.

Definition 8 Let $F : [0, T] \times D \rightarrow \mathcal{P}(\mathbb{R}^n)$ be a multivalued map with nonempty compact values. We say F is of lower semicontinuous type (l.s.c type) if its associated Niemytzki operator $\mathcal{F}$ is l.s.c and has nonempty closed and decomposable values.

Definition 9 The multivalued map N has a fixed point if there exists $x \in X$ such that $x \in N(x)$. The set of fixed points of the multivalued map N will be denoted by $\text{Fix } N$.

Definition 10 For a function $u : [-r, T] \rightarrow \mathbb{R}^n$, the set

$$S_{F,u} = \{v \in L^1([0, T], \mathbb{R}^n) | v(t) \in F(t, u_t)\}$$

is known as the set of selection functions.

Definition 11 F has a measurable selection if there exists a measurable function (single-valued) $h : [0, T] \rightarrow \mathbb{R}^n$ such that $h(t) \in S_{F,u}$ for each $t \in [0, T]$.

We will present some essential results from [37, 8, 10, 13, 33] to discuss the main results of this paper.

Lemma 1 Let $H_d : P(X) \times P(X) \rightarrow \mathbb{R}_+ \cup \{\infty\}$ by

$$H_d(A, B) = \max \{ \sup_{a \in A} E_d(a, B), \sup_{b \in B} E_d(A, b) \}$$

. Then $(P_{b,cl}(X), H_d)$ is a metric space and $(P_{cl}(X), H_d)$ is a complete metric space.

Lemma 2 Let $\Omega = D \cup PC$. Then Ω is a Banach space with norm

$$\|u\|_{\Omega} = \max \{ \|u\|_D, \|u\|_{PC} \}$$

Lemma 3 Let X be a separable metric space and $N : X \rightarrow \mathcal{P}(L^1([0, T], \mathbb{R}^n))$ be a multivalued operator which has property (BC). Then N has a continuous selection, i.e., there exists a continuous function $f : X \rightarrow L^1([0, T], \mathbb{R}^n)$ such that $f(x) \in N(x)$ for each $x \in X$.

Lemma 4 Let X be a normed linear space with $S \subset X$ convex and $0 \in S$. Assume $H : S \rightarrow S$ is a completely continuous operator. If the set

$$\varepsilon(H) = \{x \in S | x = \lambda H(x) \text{ for some } \lambda \in (0, 1) \text{ is bounded}\}$$

is bounded, then H has at least one fixed point in S .

Lemma 5 Let (X, d) be a complete metric space. If $N : X \rightarrow P_{cl}(X)$ is a contraction, then $\text{Fix } N \neq \emptyset$.

Lemma 6 $H \subseteq \Omega$ is a relatively compact set if and only if $H \subseteq \Omega$ is uniformly bounded and equicontinuous on each J_k ($k = 0, \dots, p$), where $J_0 = [-0, r]$, $J_k = (t_k, t_{k+1}]$ ($k = 0, \dots, p$).

Lemma 7 Let $F : [0, T] \times D \rightarrow \mathcal{P}(\mathbb{R}^n)$ be a multivalued map with nonempty, compact values. Assume (A4) and (A5) hold. Then F is of l.s.c type.

3 Mains results

In this section, we will discuss the main results of this paper, i.e., first we will investigate the existence of solutions through the fixed point technique and the Arzela-Ascoli theorem and through the necessary and sufficient conditions (A1)-(A6), as presented in the introduction. In this sense, assuming the conditions (A2), (A4) and (A5) imposing the condition a new condition (A7), we obtain the uniqueness of solutions via the Banach fixed point theorem. Finally, we discuss the uniform stability of solutions.

3.1 Existence of solutions

Theorem 5 Assume that (A1), (A2) and (A3) are satisfied. Then Eq.(2) has at least one solution on $[-r, T]$, provided

$$\mathcal{L} = \frac{(\psi(T) - \psi(0))^{\alpha-1}}{\Gamma(\alpha)} \|\psi(T)\|_{L^1} + \sum_{k=1}^p \sum_{i=0}^{m-1} \frac{(\psi(T) - \psi(0))^i}{i!} C_{ik} < 1.$$

Proof First, we consider multivalued map $\mathcal{G} : \Omega \rightarrow \mathcal{P}(\Omega)$ defined by $\mathcal{G}(u) = \{g \in \Omega\}$, where

$$g(t) = \begin{cases} \phi(t), & t \in [-r, 0] \\ \phi(0) + \sum_{i=1}^{m-1} \frac{\eta_i}{i!} (\psi(t) - \psi(0))^i + \frac{1}{\Gamma(\alpha)} \int_0^t \psi'(s) (\psi(t) - \psi(s))^{\alpha-1} h(s) ds \\ + \sum_{0 < t_k < t} \sum_{i=0}^{m-1} \frac{(\psi(t) - \psi(t_k))^i}{i!} \mathcal{J}_{ik}(u(t_k)), & t \in [0, T] \text{ and } h \in S_{F,u}. \end{cases}$$

It is clear that the fixed points of $\mathcal{G}$ are solutions of Eq.(2). For each $u \in \Omega$, the set $S_{F,u}$ is nonempty since by (A1), f has measurable selection.

We shall show that, $\mathcal{G}$ satisfies the assumptions of Lemma 5. The proof will be divided into two steps.

Step 1: $\mathcal{G}(u) \subseteq \mathcal{P}_{cl}(\Omega)$ for each $u \in \Omega$.

Indeed, let $\{u_n\} \in \mathcal{G}(u)$ such that $u_n \rightarrow u_*$. Then there exists $h_n \in S_{F,u}$, such that for each $t \in [0, T]$,

$$\begin{aligned} u_n(t) &= \phi(0) + \sum_{i=1}^{m-1} \frac{\eta_i}{i!} (\psi(t) - \psi(0))^i + \frac{1}{\Gamma(\alpha)} \int_0^t \psi'(s) (\psi(t) - \psi(s))^{\alpha-1} h_n(s) ds \\ &+ \sum_{0 < t_k < t} \sum_{i=0}^{m-1} \frac{(\psi(t) - \psi(t_k))^i}{i!} \mathcal{J}_{ik}(u(t_k)). \end{aligned}$$

Since $F(0, \psi)$ has compact values and (A5) hold, we may pass to a subsequence if necessary to get that h_n converges to h in $L^1([0, T], \mathbb{R}^n)$ and hence $h \in S_{F,u}$. Then, for each $t \in [0, T]$

$$u_n(t) \rightarrow u_*(t) = \phi(0) + \sum_{i=1}^{m-1} \frac{\eta_i}{i!} (\psi(t) - \psi(0))^i + \frac{1}{\Gamma(\alpha)} \int_0^t \psi'(s) (\psi(t) - \psi(s))^{\alpha-1} h(s) ds$$

$$+ \sum_{0 < t_k < t} \sum_{i=0}^{m-1} \frac{(\psi(t) - \psi(t_k))^i}{i!} \mathcal{J}_{ik}(u(t_k)).$$

So $u_* \in \mathcal{G}(u)$, and in particular, $\mathcal{G}(u) \subseteq \mathcal{P}_{cl}(\Omega)$.

Step 2. It can be shown that there exists $\mathcal{L} < 1$ such that $H_d(\mathcal{G}(u), \mathcal{G}(\bar{u})) \leq \mathcal{L}\|u - \bar{u}\|_\Omega$ for all $u, \bar{u} \in \Omega$.

Let $u, \bar{u} \in \Omega$ and $g \in \mathcal{G}(u)$. Then there exists $h(t) \in F(t, u_t)$ such that for each $t \in [0, T]$

$$\begin{aligned} g(t) &= \phi(0) + \sum_{i=1}^{m-1} \frac{\eta_i}{i!} (\psi(t) - \psi(0))^i + \frac{1}{\Gamma(\alpha)} \int_0^t \psi'(s) (\psi(t) - \psi(s))^{\alpha-1} h(s) ds \\ &\quad + \sum_{0 < t_k < t} \sum_{i=0}^{m-1} \frac{(\psi(t) - \psi(t_k))^i}{i!} \mathcal{J}_{ik}(u(t_k)). \end{aligned}$$

From **(A3)** it follows that, for each $t \in [0, T]$,

$$H_d(F(t, u_t), F(t, \bar{u}_t)) \leq l(t)\|u_t - \bar{u}_t\|_D.$$

Hence there exists $w(t) \in F(t, \bar{u}_t)$ such that

$$|h(t) - w(t)| \leq l(t)\|u_t - \bar{u}_t\|_D, \quad t \in [0, T].$$

Consider $U : [0, T] \rightarrow \mathcal{P}_{cl}(\mathbb{R}^n)$, given by $U(t) = \{w(t) : |h(t) - w(t)| \leq l(t)\|u_t - \bar{u}_t\|_D\}$. Since, the multivalued operator $V(t) = U(t) \cap F(t, \bar{u}_t)$ is measurable, there exists a function $\bar{h}(t)$ which is a measurable selection for V . So, $\bar{h}(t) \in F(t, \bar{u}_t)$ and $|h(t) - \bar{h}(t)| \leq l(t)\|u_t - \bar{u}_t\|_D$ for each $t \in [0, T]$.

We define, for each $t \in [0, T]$,

$$\begin{aligned} g(t) &= \phi(0) + \sum_{i=1}^{m-1} \frac{\eta_i}{i!} (\psi(t) - \psi(0))^i + \frac{1}{\Gamma(\alpha)} \int_0^t \psi'(s) (\psi(t) - \psi(s))^{\alpha-1} \bar{h}(s) ds \\ &\quad + \sum_{0 < t_k < t} \sum_{i=0}^{m-1} \frac{(\psi(t) - \psi(t_k))^i}{i!} \mathcal{J}_{ik}(\bar{u}(t_k)). \end{aligned}$$

Then, we have

$$\begin{aligned} &|g(t) - \bar{g}(t)| \\ &= \left| \frac{1}{\Gamma(\alpha)} \int_0^t \psi'(s) (\psi(t) - \psi(s))^{\alpha-1} h(s) ds - \frac{1}{\Gamma(\alpha)} \int_0^t \psi'(s) (\psi(t) - \psi(s))^{\alpha-1} \bar{h}(s) ds \right. \\ &\quad \left. + \sum_{0 < t_k < t} \sum_{i=0}^{m-1} \frac{(\psi(t) - \psi(t_k))^i}{i!} \mathcal{J}_{ik}(u(t_k)) - \sum_{0 < t_k < t} \sum_{i=0}^{m-1} \frac{(\psi(t) - \psi(t_k))^i}{i!} \mathcal{J}_{ik}(\bar{u}(t_k)) \right| \\ &\leq \frac{1}{\Gamma(\alpha)} \int_0^t \psi'(s) (\psi(t) - \psi(s))^{\alpha-1} |h(s) - \bar{h}(s)| ds \\ &\quad + \sum_{0 < t_k < t} \sum_{i=0}^{m-1} \frac{(\psi(t) - \psi(t_k))^i}{i!} |\mathcal{J}_{ik}(u(t_k)) - \mathcal{J}_{ik}(\bar{u}(t_k))| \end{aligned}$$

$$\begin{aligned}
&\leq \frac{(\psi(T) - \psi(0))^{\alpha-1}}{\Gamma(\alpha)} \psi'(T) \int_0^t |h(s) - \bar{h}(s)| ds \\
&\quad + \sum_{0 < t_k < t} \sum_{i=0}^{m-1} \frac{(\psi(T) - \psi(t_k))^i}{i!} C_{ik} |u(t_k) - \bar{u}(t_k)| \\
&\leq \frac{(\psi(T) - \psi(0))^{\alpha-1}}{\Gamma(\alpha)} \psi'(T) \int_0^t l(s) \|u_s - \bar{u}_s\|_D ds \\
&\quad + \sum_{k=1}^p \sum_{i=0}^{m-1} \frac{(\psi(T) - \psi(t_k))^i}{i!} C_{ik} |u(t_k) - \bar{u}(t_k)| \\
&\leq \frac{(\psi(T) - \psi(0))^{\alpha-1}}{\Gamma(\alpha)} \psi'(T) \|u - \bar{u}\|_{\Omega} \int_0^t l(s) ds \\
&\quad + \sum_{k=1}^p \sum_{i=0}^{m-1} \frac{(\psi(T) - \psi(t_k))^i}{i!} C_{ik} \|u - \bar{u}\|_{\Omega} \\
&\leq \left\{ \frac{(\psi(T) - \psi(0))^{\alpha-1}}{\Gamma(\alpha)} \psi'(T) \|l\|_{L^1} + \sum_{k=1}^p \sum_{i=0}^{m-1} \frac{(\psi(T) - \psi(t_k))^i}{i!} C_{ik} \right\} \|u - \bar{u}\|_{\Omega}. \quad (6)
\end{aligned}$$

So, $\|g(t) - \bar{g}(t)\|_{\Omega} \leq \mathcal{L} \|u - \bar{u}\|_{\Omega}$. By an analogous reasoning, obtained by interchanging the roles of u and $\bar{u}$, it follows that $H_d(G(u), G(\bar{u})) \leq L \|u - \bar{u}\|_{\Omega}$. Therefore $\mathcal{G}$ is contraction. By Lemma 5, $\mathcal{G}$ has a fixed point which is solution of Eq.(2).

Theorem 6 *In addition to (A4) and (A5), assume that the following condition holds:*

(A6) *There exists constants $d_{ik} (i = 0, 1, \dots, m-1, k = 1, 2, \dots, p)$, such that $|\mathcal{J}_{ik}(u(t_k))| \leq d_{ik} |u(t_k)|$ for each $u \in \Omega$. Then Eq.(2) has at least one solution on $[-r, T]$, provided*

$$\mathcal{L} = \sum_{k=1}^p \sum_{i=0}^{m-1} \frac{(\psi(T) - \psi(t_k))^i}{i!} d_{ik} < 1.$$

Proof Note that (A4), (A5) and Lemma 7 imply that f is of L. S. C. type. Then, from Lemma 3, there exists a continuous function $f : \Omega \rightarrow L^1([0, T], \mathbb{R}^n)$ such that $f(u) \in F(u)$, for each $u \in \Omega$.

We consider

$$\begin{cases} {}^C \mathbb{D}_{0+}^{\alpha, \psi} u(t) = f(u)(t), & t \in J = [0, T], t \neq t_k, k = 1, 2, \dots, p, \\ \delta u^{(i)}(t_k) = \mathcal{J}_{ik}(u(t_k)), & i = 0, 1, \dots, m-1, k = 1, 2, \dots, p, \\ u(t) = \phi(t), & t \in [-r, 0], \\ u^{(i)}(0) = \eta_i, & i = 1, 2, \dots, m-1. \end{cases} \quad (7)$$

We shall show that T satisfies all assumptions of Lemma 4. The proof will be given in four steps.

Step 1: T is continuous.

Since the functions f and $\mathcal{J}_{ik}$ are continuous, this conclusion can be easily obtained.

Step 2: T maps arbitrary bounded subset of Ω into on bounded set in Ω .

Let $B_a = \{u \in \Omega : \|u\|_\Omega \leq a\}$ be arbitrary bounded subset of Ω and $u \in B_a$, then there exists $f \in F(u)$ such that for $t \in [0, T]$,

$$\begin{aligned} T(u)(t) &= \phi(0) + \sum_{i=1}^{m-1} \frac{\eta_i}{i!} (\psi(t) - \psi(0))^i + \frac{1}{\Gamma(\alpha)} \int_0^t \psi'(s) (\psi(t) - \psi(s))^{\alpha-1} f(u)(s) ds \\ &\quad + \sum_{0 < t_k < t} \sum_{i=0}^{m-1} \frac{(\psi(t) - \psi(t_k))^i}{i!} \mathcal{J}_{ik}(u(t_k)). \end{aligned} \quad (8)$$

Using the conditions (A5) and (A6), we have

$$\begin{aligned} |T(u)(t)| &\leq |\phi(0)| + \sum_{i=1}^{m-1} \frac{|\eta_i|}{i!} (\psi(T) - \psi(0))^i + \frac{(\psi(T) - \psi(0))^{\alpha-1}}{\Gamma(\alpha)} \psi'(T) \int_0^T |f(u)(s)| ds \\ &\quad + \sum_{k=1}^p \sum_{i=0}^{m-1} \frac{(\psi(T) - \psi(0))^i}{i!} |\mathcal{J}_{ik}(u(t_k))| \\ &\leq |\phi(0)| + \sum_{i=1}^{m-1} \frac{|\eta_i|}{i!} (\psi(T) - \psi(0))^i + \frac{(\psi(T) - \psi(0))^{\alpha-1}}{\Gamma(\alpha)} \psi'(T) \int_0^T |M(s)| ds \\ &\quad + \sum_{k=1}^p \sum_{i=0}^{m-1} \frac{(\psi(t) - \psi(t_k))^i}{i!} d_{ik} |u(t_k)| \\ &\leq |\phi(0)| + \sum_{i=1}^{m-1} \frac{|\eta_i|}{i!} (\psi(T) - \psi(0))^i + \frac{(\psi(T) - \psi(0))^{\alpha-1}}{\Gamma(\alpha)} \psi'(T) \|M\|_{L^1} \\ &\quad + \sum_{k=1}^p \sum_{i=0}^{m-1} \frac{(\psi(t) - \psi(t_k))^i}{i!} d_{ik} \|u\|_\Omega \end{aligned}$$

for each $t \in [0, T]$.

Therefore, for each $u \in B_a$, we have

$$\begin{aligned} \|T(u)\|_\Omega &\leq |\phi(0)| + \sum_{i=1}^{m-1} \frac{|\eta_i|}{i!} (\psi(T) - \psi(0))^i + \frac{(\psi(T) - \psi(0))^{\alpha-1}}{\Gamma(\alpha)} \psi'(T) \|M\|_{L^1} + \mathcal{L} \|u\|_\Omega \\ &\leq |\phi(0)| + \sum_{i=1}^{m-1} \frac{|\eta_i|}{i!} (\psi(T) - \psi(0))^i + \frac{(\psi(T) - \psi(0))^{\alpha-1}}{\Gamma(\alpha)} \psi'(T) \|M\|_{L^1} + \mathcal{L} a. \end{aligned} \quad (9)$$

Therefore, $T(B_a)$ is bounded.

Step 3. T maps arbitrary bounded set of points into equicontinuous set in Ω .

Let $\tau_1, \tau_2 \in (t_k, t_{k+1})$, $k = 1, 2, \dots, p$, $\tau_1 < \tau_2$, and $u \in B_a$ be arbitrary bounded subset of Ω . By (8) we get

$$\begin{aligned} &|T(u)(\tau_2) - T(u)(\tau_1)| \\ &\leq \sum_{i=1}^{m-1} \frac{|\eta_i|}{i!} \left[(\psi(\tau_2) - \psi(t_k))^i - (\psi(\tau_1) - \psi(t_k))^i \right] \\ &\quad + \left| \frac{1}{\Gamma(\alpha)} \int_{\tau_1}^{\tau_2} \psi'(s) (\psi(\tau_2) - \psi(s))^{\alpha-1} f(u)(s) ds \right| \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{\Gamma(\alpha)} \int_0^{\tau_1} \psi'(s) \left((\psi(\tau_2) - \psi(s))^{\alpha-1} + (\psi(\tau_1) - \psi(s))^{\alpha-1} \right) f(u)(s) ds \Big| \\
& + \sum_{k=1}^p \sum_{i=0}^{m-1} \frac{(\psi(\tau_2) - \psi(t_k))^i}{i!} |\mathcal{J}_{ik}(u(t_k))| + \sum_{k=1}^p \sum_{i=0}^{m-1} \frac{(\psi(\tau_1) - \psi(t_k))^i}{i!} |\mathcal{J}_{ik}(u(t_k))| \\
& \leq \sum_{i=1}^{m-1} \frac{|\eta_i|}{i!} (\psi(\tau_2)^i - \psi(\tau_1)^i) + \left| \frac{1}{\Gamma(\alpha)} \int_{\tau_1}^{\tau_2} \sum_{i=0}^{m-1} C_{m-1}^i \psi'(s) \psi(s)^{\alpha-1-i} \psi(\tau_2)^i f(u)(s) ds \right. \\
& \quad \left. + \frac{1}{\Gamma(\alpha)} \int_0^{\tau_1} \sum_{i=0}^{m-1} C_{m-1}^i \psi'(s) \psi(s)^{\alpha-1-i} (\psi(\tau_2)^i - \psi(\tau_1)^i) f(u)(s) ds \right| \\
& \quad + \sum_{k=1}^p \sum_{i=0}^{m-1} \frac{(\psi(\tau_2) - \psi(t_k))^i + (\psi(\tau_1) - \psi(t_k))^i}{i!} |\mathcal{J}_{ik}(u(t_k))| \\
& \leq \sum_{i=1}^{m-1} \frac{|\eta_i|}{i!} (\psi(\tau_2)^i - \psi(\tau_1)^i) + \left| \frac{1}{\Gamma(\alpha)} \int_{\tau_1}^{\tau_2} \sum_{i=0}^{m-1} C_{m-1}^i \psi'(s) \psi(s)^{\alpha-1-i} \psi(\tau_2)^i f(u)(s) ds \right. \\
& \quad \left. + \frac{1}{\Gamma(\alpha)} \int_0^{\tau_1} \sum_{i=0}^{m-1} C_{m-1}^i \psi'(s) \psi(s)^{\alpha-1-i} (\psi(\tau_2)^i - \psi(\tau_1)^i) f(u)(s) ds \right| \\
& \quad + \sum_{k=1}^p \sum_{i=0}^{m-1} \frac{1}{i!} \sum_{j=0}^i C_i^j d_{ik} (\psi(\tau_2)^j - \psi(\tau_1)^j) \psi(\tau_k)^{i-j} \\
& \leq (\psi(\tau_2) - \psi(\tau_1)) \sum_{i=1}^{m-1} \frac{|\eta_i|}{i!} \frac{\psi(T)^{i-1}}{(i-1)!} + \frac{1}{\Gamma(\alpha)} \left[\psi'(T) \psi(T)^{\alpha-1} 2^{m-1} \int_{\tau_1}^{\tau_2} f(u)(s) ds \right. \\
& \quad \left. + \psi'(T) \psi(T)^{\alpha-2} (\psi(\tau_2) - \psi(\tau_1)) \int_0^{\tau_1} |f(u)(s)| ds \sum_{i=1}^{m-1} i C_{m-1}^i \right] \\
& \quad + a \sum_{k=1}^p \sum_{i=0}^{m-1} \frac{1}{i!} \sum_{j=0}^i d_{ik} (\psi(\tau_2)^j - \psi(\tau_1)^j) \psi(\tau_k)^{i-j} \\
& \leq (\psi(\tau_2) - \psi(\tau_1)) \sum_{i=1}^{m-1} \frac{|\eta_i|}{i!} \frac{\psi(T)^{i-1}}{(i-1)!} + \frac{1}{\Gamma(\alpha)} \left[\psi'(T) \psi(T)^{\alpha-1} 2^{m-1} \int_{\tau_1}^{\tau_2} M(s) ds \right. \\
& \quad \left. + \psi'(T) \psi(T)^{\alpha-2} (\psi(\tau_2) - \psi(\tau_1)) \|f\|_{L^1} \sum_{i=1}^{m-1} i C_{m-1}^i \right] \\
& \quad + a (\psi(\tau_2) - \psi(\tau_1)) \sum_{i=1}^{m-1} \frac{2^i \psi(T)^{i-1}}{(i-1)!} d_{ik} \\
& \leq (\psi(\tau_2) - \psi(\tau_1)) \sum_{i=1}^{m-1} \frac{|\eta_i|}{i!} \frac{\psi(T)^{i-1}}{(i-1)!} + \frac{1}{\Gamma(\alpha)} \left[\psi'(T) \psi(T)^{\alpha-1} 2^{m-1} \int_{\tau_1}^{\tau_2} M(s) ds \right. \\
& \quad \left. + \psi'(T) \psi(T)^{\alpha-2} (\psi(\tau_2) - \psi(\tau_1)) m 2^{m-1} \|f\|_{L^1} \right] + a (\psi(\tau_2) - \psi(\tau_1)) \sum_{i=1}^{m-1} \frac{2^i \psi(T)^{i-1}}{(i-1)!} d_{ik}.
\end{aligned}$$

According to the completely continuity of integrable function M , the right hand side of the above inequality tends to zero as $\tau_2 \rightarrow \tau_1$. The convergence for the uses $\tau_2, \tau_1 \in (0, t_1]$ and $[-r, 0]$ is obvious. Then $T(B_a)$ is equicontinuous set in Ω .

As a consequences of **Step 1** to **Step 3** together with Lemma 6 and the Ascoli-Arzelà theorem, we conclude that $T : \Omega \rightarrow \Omega$ is completely continuous.

Step 4: The set $\varepsilon(T) = \{u \in \Omega : u = \lambda T(u), \text{ for some } 0 < \lambda < 1\}$ is bounded.

For each $u \in \varepsilon(T)$, by (9), we have

$$\begin{aligned} \|u\|_{\Omega} &= \lambda \|T(u)\|_{\Omega} \\ &\leq |\phi(0)| + \sum_{i=1}^{m-1} \frac{|\eta_i|}{i!} (\psi(T) - \psi(0))^i + \frac{(\psi(T) - \psi(0))^{\alpha-1}}{\Gamma(\alpha)} \psi'(T) \|M\|_{L^1} + \mathcal{L} \|u\|_{\Omega} \end{aligned}$$

Then

$$\|u\|_{\Omega} \leq \frac{|\phi(0)| + \sum_{i=1}^{m-1} \frac{|\eta_i|}{i!} (\psi(T) - \psi(0))^i + \frac{(\psi(T) - \psi(0))^{\alpha-1}}{\Gamma(\alpha)} \psi'(T) \|M\|_{L^1}}{1 - \mathcal{L}}$$

i.e. $\varepsilon(T)$ is bounded.

In view of Lemma 4, we deduce that T has a fixed which in turn is a solution of Eq.(2).

3.2 Uniqueness of solutions

Theorem 7 *In addition to (A2), (A4) and (A5), assume that the following condition hold:*

- (A7) *There exists non-negative constant b , such that $|f(u)(t) - f(v)(t)| \leq b \|u - v\|_{\Omega}$ for any $u, v \in \Omega$ where f is deduced in the proof of Theorem 2. Then Eq.(2) has a unique solution on $[-r, T]$, provided*

$$\mathcal{L} = \frac{b(\psi(T) - \psi(0))^{\alpha}}{\Gamma(\alpha + 1)} + \sum_{k=1}^p \sum_{i=0}^{m-1} C_{ik} \frac{\psi(T)^i}{i!} < 1.$$

Proof By (8), for any $u, v \in \Omega$, and each $t \in [0, T]$, we obtain

$$\begin{aligned} |T(u)(t) - T(v)(t)| &\leq \frac{1}{\Gamma(\alpha)} \int_0^t \psi'(s) (\psi(t) - \psi(s))^{\alpha-1} |f(u)(s) - f(v)(s)| ds \\ &\quad + \sum_{0 < t_k < t} \sum_{i=0}^{m-1} \frac{(\psi(t) - \psi(t_k))^i}{i!} |\mathcal{J}_{ik}(u(t_k)) - \mathcal{J}_{ik}(\bar{u}(t_k))| \\ &\leq \frac{b}{\Gamma(\alpha)} \|u - v\|_{\Omega} \int_0^t \psi'(s) (\psi(t) - \psi(s))^{\alpha-1} ds \\ &\quad + \sum_{0 < t_k < t} \sum_{i=0}^{m-1} \frac{(\psi(t) - \psi(t_k))^i}{i!} C_{ik} |u(t_k) - \bar{u}(t_k)| \end{aligned}$$

$$\begin{aligned}
&\leq \frac{b}{\Gamma(\alpha+1)} \|u-v\|_{\Omega} (\psi(T) - \psi(0))^{\alpha} + \|u-v\|_{\Omega} \sum_{0 < t_k < t} \sum_{i=0}^{m-1} \frac{(\psi(T))^i}{i!} C_{ik} \\
&\leq \left(\frac{b}{\Gamma(\alpha+1)} (\psi(T) - \psi(0))^{\alpha} + \sum_{k=1}^p \sum_{i=0}^{m-1} \frac{(\psi(T))^i}{i!} C_{ik} \right) \|u-v\|_{\Omega}.
\end{aligned}$$

So

$$\|T(u) - T(v)\|_{\Omega} \leq \mathcal{L} \|u-v\|_{\Omega}.$$

Thus all conditions of the fixed point theorem of Banach are satisfied. Then Eq.(2) has a unique solution on $[-r, T]$.

3.3 Uniformly stable of solutions

Theorem 8 Assume that all the conditions of Theorem 3 hold. Then the solution of the Eq.(2) is uniformly stable.

Proof Let $u(t)$ be a solution of Eq.(2), and let $\bar{u}(t)$ be a solution of Eq.(2) satisfying condition $\bar{X}(0) = \bar{\phi}(0)$, then

$$u(t) - \bar{u}(t) = \begin{cases} \phi(t) - \bar{\phi}(t), & t \in [-r, 0], \\ \phi(0) - \bar{\phi}(0) + \frac{1}{\Gamma(\alpha)} \int_0^t \psi'(s) (\psi(t) - \psi(s))^{\alpha-1} |f(u)(s) - f(\bar{u})(s)| ds \\ + \sum_{0 < t_k < t} \sum_{i=0}^{m-1} \frac{(\psi(t) - \psi(t_k))^i}{i!} [\mathcal{J}_{ik}(u(t_k)) - \mathcal{J}_{ik}(\bar{u}(t_k))], & t \in [0, T]. \end{cases}$$

So we get for each $t \in (0, T]$

$$\begin{aligned}
\|u(t) - \bar{u}(t)\| &\leq \|\phi(0) - \bar{\phi}(0)\| + \frac{1}{\Gamma(\alpha)} \int_0^t \psi'(s) (\psi(t) - \psi(s))^{\alpha-1} \|f(u)(s) - f(\bar{u})(s)\| ds \\
&\quad + \sum_{0 < t_k < t} \sum_{i=0}^{m-1} \frac{(\psi(t) - \psi(t_k))^i}{i!} \|\mathcal{J}_{ik}(u(t_k)) - \mathcal{J}_{ik}(\bar{u}(t_k))\| \\
&\leq \|\phi - \bar{\phi}\| + \frac{b}{\Gamma(\alpha)} \|u - \bar{u}\|_{\infty} \int_0^t \psi'(s) (\psi(t) - \psi(s))^{\alpha-1} ds \\
&\quad + \sum_{0 < t_k < t} \sum_{i=0}^{m-1} \frac{(\psi(t) - \psi(t_k))^i}{i!} C_{ik} \|u(t_k) - \bar{u}(t_k)\| \\
&\leq \|\phi - \bar{\phi}\| + \frac{b}{\Gamma(\alpha+1)} \|u - \bar{u}\|_{\infty} (\psi(T) - \psi(0))^{\alpha} \\
&\quad + \|u - \bar{u}\|_{\infty} \sum_{k=1}^p \sum_{i=0}^{m-1} \frac{(\psi(T) - \psi(t_k))^i}{i!} C_{ik} \\
&\leq \|\phi - \bar{\phi}\| + \left(\frac{b}{\Gamma(\alpha+1)} (\psi(T) - \psi(0))^{\alpha} + \sum_{k=1}^p \sum_{i=0}^{m-1} \frac{(\psi(T))^i}{i!} C_{ik} \right) \|u - \bar{u}\|_{\infty}.
\end{aligned}$$

So we have

$$\|u - \bar{u}\|_{\infty} \leq \|\phi - \bar{\phi}\| + \mathcal{L}\|u - \bar{u}\|_{\infty}.$$

This gives

$$\|u - \bar{u}\|_{\infty} \leq \frac{1}{1 - \mathcal{L}} \|\phi - \bar{\phi}\|.$$

Therefore, if $\|\phi - \bar{\phi}\| < \delta(\varepsilon)$, then $\|u - \bar{u}\| < \varepsilon$ which implies that the solution of Eq.(2) is uniformly stable.

References

1. Abada, N., M. Benchohra, and H Hammouche. Existence and controllability results for nondensely defined impulsive semilinear functional differential inclusions. *J. Diff. Equ.* 246(10) (2009) 3834–3863.
2. Abdo, M. S., Satish K. Panchal, and H. S. Hussien. Fractional integro-differential equations with non-local conditions and ψ -Hilfer fractional derivative. *Math. Modell. Anal.* 24(4) (2019) 564–584.
3. Aubin, J-P., and Arrigo Cellina. *Differential inclusions: set-valued maps and viability theory*. Vol. 264. Springer Science & Business Media, 2012.
4. Ahmad, B., Y. Alruwaily, A. Alsaedi, and Juan J. Nieto "Fractional integro-differential equations with dual anti-periodic boundary conditions." *Diff. Integral Equ.* 33, (2020): 181-206.
5. Balasubramaniam, P., and D. Vinayagam. Existence of solutions of nonlinear stochastic integrodifferential inclusions in a Hilbert space. *Comput. Math. Appl.* 50(5) (2005) 809–821.
6. Baghani, H., J. Alzabut, J. Farokhi-Ostad and Juan J. Nieto .Existence and uniqueness of solutions for a coupled system of sequential fractional differential equations with initial conditions. *J. Pseudo-Diff. Operators Appl.* (2020) 1–11.
7. Benchohra, M., and A. Ouahab. Initial boundary value problems for second order impulsive functional differential inclusions. *Elec. J. Qual. Theory Diff. Equ.* 2003(3) (2003) 1–10.
8. Benchohra, M., and A. Ouahab. Initial boundary value problems for second order impulsive functional differential inclusions. *Electronic J. Qual. Theory Diff. Equ.* 2003(3) (2003) 1–10.
9. Boudjerida, A., D. Seba, and G. M. N'Guérékata. "Controllability of coupled systems for impulsive ϕ -Hilfer fractional integro-differential inclusions." *Applicable Anal.* (2020) 1–18.
10. Bressan, A., G. Colombo, Existence and selections of maps with decomposable values, *Studia Math.* 90 (1988) 69–86.
11. Chang, Y.-K. and Wan-Tong Li. Existence results for second order impulsive functional differential inclusions. *J. Math. Anal. Appl.* 301(2) (2005) 477–490.
12. Du, J., W. Jiang, and A. Ullah Khan Niazi. Approximate controllability of impulsive Hilfer fractional differential inclusions. *J. Nonlinear Sci. Appl.* 10(2) (2017) 595–611.
13. Fu, X.L., B.Q. Yan, Y.S. Liu, *Theory of impulsive differential system*, Science Press, Beijing, 2005.
14. Frigon, M.. *Théorèmes d'existence de solutions d'inclusions différentielles*. Topological Methods in Differential Equations and Inclusions. Springer, Dordrecht, (1995) 51–87.
15. Hari Krishnan, S. Note on the solution of random differential equations via ψ -Hilfer fractional derivative. *Adv. Diff. Equ.* 2018(1) (2018) 224.
16. Hari Krishnan, S., E. M. Elsayed, and K. Kanagarajan. Existence and uniqueness results for fractional pantograph equations involving ψ -Hilfer fractional derivative. *Dyn. Cont. Disc. Impulsive Sys.* 25 (2018) 319–328.
17. Hu, L., and Y. Ren. Existence results for impulsive neutral stochastic functional integro-differential equations with infinite delays. *Acta Appl. Math.* 111(3) (2010) 303–317.
18. Kilbas, A. A., H. M. Srivastava, J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, North-Holland Math. Stud., Vol. 204, Elsevier, Amsterdam, 2006.
19. Kucche, Kishor D., Ashwini D. Mali, and J. Vanterler da C. Sousa. On the nonlinear ψ -Hilfer fractional differential equations. *Comput. Appl. Math.* 38(2) (2019) 73.
20. Kryszewski, W., and S. Plaskacz. Periodic solutions to impulsive differential inclusions with constraints. *Nonlinear Analysis: Theory, Methods & Applications* 65(9) (2006) 1794–1804.
21. Liang, J., H. Yang, Controllability of fractional integro-differential evolution equation with nonlocal conditions, *Appl. Math. Comput.*, 254 (2015) 20–29.

22. Liu, K., J. Wang, and D. O'Regan. Ulam–Hyers–Mittag-Leffler stability for ψ -Hilfer fractional-order delay differential equations. *Adv. Diff. Equ.* 2019.1 (2019) 1–12.
23. Liu, Z., B. Zeng, Existence and controllability for fractional evolution inclusions of Clark's subdifferential type, *Appl. Math. Comput.*, 257 (2015) 178–189.
24. Ren, L., J. Wang, M. Feckan, Asymptotically periodic solutions for caputo type fractional evolution equations, *Fract Calc. Appl. Anal.*, 21(5) 2018 1294–1312.
25. Rudolf, H.. Applications of fractional calculus in physics. world scientific, 2000.
26. Sousa, J. Vanterler da C., and E. Capelas De Oliveira. "Ulam–Hyers stability of a nonlinear fractional Volterra integro-differential equation." *Appl. Math. Lett.* 81 (2018) 50–56.
27. Sousa, J. Vanterler da C., and E. Capelas de Oliveira. On the Ulam–Hyers–Rassias stability for nonlinear fractional differential equations using the ψ -Hilfer operator. *J. Fixed Point Theory Appl.* 20(3) (2018) 96.
28. Sousa, J. Vanterler da C., Gastão S. F. Frederico, and E. Capelas de Oliveira. ψ -Hilfer pseudo-fractional operator: new results about fractional calculus. *Comput. Appl. Math.* 39(4) (2020): 1–33.
29. Sousa, J. Vanterler da C., M. Benchohra, and Gaston M. N'Guérékata. Attractivity for differential equations of fractional order and ψ -Hilfer type. *Frac. Cal. Appl. Anal.* 23.4 (2020) 1188–1207.
30. Sousa, J. Vanterler da C., and E. Capelas de Oliveira. On the ψ -Hilfer fractional derivative. *Commun. Nonlinear Sci. Numer. Simul.* 60 (2018) 72–91.
31. Sousa, J. Vanterler da C., and E. Capelas de Oliveira. Leibniz type rule: ψ -Hilfer fractional operator. *Commun. Nonlinear Sci. Numer. Simul.* 77 (2019) 305–311.
32. Sousa, J. Vanterler da C., J. A. T. Machado, and E. Capelas de Oliveira. The ψ -Hilfer fractional calculus of variable order and its applications. *Comput. Appl. Math.* 39(4) (2020) 1–35.
33. Smart, D.R., *Fixed Point Theorems*, Cambridge Univ. Press, Cambridge, 1974.
34. Wang, JinRong, A. G. Ibrahim, and D. O'Regan. Topological structure of the solution set for fractional non-instantaneous impulsive evolution inclusions. *J. Fixed Point Theory Appl.* 20.2 (2018) 59.
35. Wang, J., A. G Ibrahim, D. O'Regan, Y. Zhou . Controllability for noninstantaneous impulsive semilinear functional differential inclusions without compactness. *Indagationes Math.* 29.5 (2018) 1362–1392.
36. Wang, J., G. Ibrahim, and D. O'Regan. Controllability of Hilfer fractional noninstantaneous impulsive semilinear differential inclusions with nonlocal conditions. *Nonlinear Analysis: Modelling and Control* 24(6) (2019) 958–984.
37. Ye, G., J. Shen, and J. Li. Existence results for m th-order impulsive functional differential inclusions. *Indagationes Math.* 22(1) (2011) 1–11.
38. Yu, X., A. Debbouche, and J. Wang. On the iterative learning control of fractional impulsive evolution equations in Banach spaces. *Math. Meth. Appl. Sci.* 40(17) (2017) 6061–6069.
39. Zhou, Y., *Fractional Evolution Equations and Inclusions: Analysis and Control*, Elsevier, Amsterdam, 2015.