Localized Afterslip at Geometrical Complexities Revealed by InSAR After the 2016 Central Italy Seismic Sequence

Lea Pousse-Beltran, Anne Socquet, Lucilla Benedetti, Marie-Pierre Doin,
Magali Rizza, Nicola d'Agostino

► To cite this version:

Lea Pousse-Beltran, Anne Socquet, Lucilla Benedetti, Marie-Pierre Doin, Magali Rizza, et al.. Localized Afterslip at Geometrical Complexities Revealed by InSAR After the 2016 Central Italy Seismic Sequence. *Journal of Geophysical Research: Solid Earth*, 2020, 125 (11), 10.1029/2019JB019065 . hal-03027096

HAL Id: hal-03027096

<https://hal.science/hal-03027096>

Submitted on 27 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1
2 Title:

3 **Localized afterslip at geometrical complexities revealed by InSAR after the 2016 Central Italy**
4 **seismic sequence**

5 Authors:

6 Léa Pousse-Beltran¹, Anne Socquet², Lucilla Benedetti¹, Marie-Pierre Doin², Magali Rizza¹,
7 Nicola D'Agostino³

8
9 ¹ Aix Marseille Université, CNRS, IRD, Collège de France, CEREGE, Aix-en-Provence, France

10 ² Université Grenoble-Alpes, Université de Savoie Mont-Blanc, CNRS, IRD, IFSTTAR, ISTerre,
11 38000 Grenoble, France

12 ³ Istituto Nazionale di Geofisica e Vulcanologia (INGV), Centro Nazionale Terremoti, via di Vigna
13 Murata 605, 00143, Rome, Italy

14
15
16 Email list:

17 Léa Pousse-Beltran: pousse@cerege.fr

18 Anne Socquet : anne.socquet@univ-grenoble-alpes.fr

19 Lucilla Benedetti : benedetti@cerege.fr

20 Marie-Pierre Doin : marie-pierre.doin@univ-grenoble-alpes.fr

21 Magali Rizza : rizza@cerege.fr

22 Nicola D'Agostino : nicola.dagostino@ingv.it

25 Keywords:

- 26 - Postseismic
- 27 - InSAR time-series
- 28 - 2016-2017 Amatrice-Norcia Seismic Sequence
- 29 - Geometrical complexity

30

31 Key points:

- 32 - We monitor pre and post-seismic deformation of the 2016 seismic sequence using two-
33 year InSAR time-series
- 34 - Centimetre scale post-seismic surface displacements are detected after October 30,
35 2016 Mw 6.5 mainshock (Norcia earthquake)
- 36 - Localized shallow afterslip occurred at structural complexity that may have hindered
37 the propagation of seismic ruptures

38

39

40 Abstract :

41 The Mw 6.5 Norcia earthquake occurred on October 30, 2016, along the Mt Vettore
42 fault (Central Apennines, Italy), it was the largest earthquake of the 2016-2017 seismic
43 sequence that started two months earlier with the Mw 6.0 Amatrice earthquake (August, 24).
44 To detect potential slow slip during the sequence, we produced Interferometric Synthetic
45 Aperture Radar (InSAR) time-series using 12 to 6-day repeat cycles of Sentinel-1A/1B images.
46 Time-series indicates that centimetre-scale surface displacements took place during the 10
47 weeks following the Norcia earthquake. Two areas of subsidence are detected: one in the
48 Castelluccio basin (hanging wall of the Mt Vettore fault), and one in the southern extent of the
49 Norcia earthquake surface rupture, near an inherited thrust. Poroelastic and viscoelastic
50 models are unable to explain these displacements. In the Castelluccio basin, the displacement
51 reaches 13.2 ± 1.4 mm in the ascending line of sight (LOS) on January 06, 2017. South of the
52 Norcia earthquake surface rupture (a zone between the Norcia and Amatrice earthquakes),
53 the post-seismic surface displacements affect a smaller area, but reach 35.5 ± 1.7 mm in
54 ascending LOS by January 2017 and follow a logarithmic temporal decay consistent with post-
55 seismic afterslip. Our analysis suggests that the structurally complex area located south of the
56 Norcia rupture (30 October) is characterized by a conditionally stable frictional regime. This
57 geometrical and frictional barrier likely halted rupture propagation during the Amatrice
58 (August 24) and Norcia (October 30) earthquakes at shallow depth (<3-4 km).

1 Introduction

Monitoring the spatial and temporal variations of the slip on a fault enables researchers to better assess stress build-up on seismic asperities and slip released during seismic cycle (Avouac, 2015; Bürgmann, 2018; Chen & Bürgmann, 2017; Harris, 2017). In the rate and state formulation, rupture propagation can be hindered by rate-strengthening sections of a fault, which tend to slip via creep rather than in seismic rupture (e.g., H. Perfettini et al., 2010; Hirose et al., 2010). Such barriers are also often associated with structural complexities – such as a change of strike, secondary faulting or interaction with inherited faults (e.g., King & Nabelek, 1985; King, 1986; Wesnousky, 1988). These structural complexities can act as a geometrical barrier, and are characterized by an increased equivalent strength (Nielsen & Knopoff, 1998). Locating aseismic slip on the fault and comparing these locations with those of seismic slip and fault segmentation is of pivotal importance to better characterize the frictional behavior of a fault system and its relation with structural complexities.

A notable seismic sequence occurred in the Central Apennines (Italy) in 2016-2017 (Chiarabba et al., 2018; Perouse et al., 2018; Cheloni et al., 2017; Huang et al., 2017; Cirella et al., 2018; Civico et al., 2018; Scognamiglio et al., 2018; Ragon et al., 2019), with four main events: the 24th August 2016 M_w 6.0 Amatrice event, the 26th October 2016 M_w 5.9 Visso event, the 30th October 2016 M_w 6.5 Norcia event, and the 18th January 2017 M_w 5.5 Campotosto event (Figure 1 and Table S1). This seismic sequence ruptured the complex Mt Vettore fault system (in red in Figure 1) (Pizzi et al., 2017; Porreca et al., 2018; Villani, Pucci, et al., 2018), and the adjacent Amatrice-Campotosto fault (in orange in Figure 1). During the Norcia earthquake, the rupturing of an antithetic fault on the opposite side of the Castelluccio basin seems necessary to fit geodetic data and is supported by alignments of relocated aftershocks (Chiaraluce et al., 2017; Walters et al., 2018; Cheloni et al., 2019). In addition, the role of an inherited west-dipping thrust called OAS (Olevano-Antrdoco-Sibillini thrust) in the Norcia earthquake coseismic rupture geometry has been widely discussed. While some studies suggested that only the Mt Vettore fault system was activated (Chiaraluce et al., 2017; Liu et al., 2017; Papadopoulos et al., 2017; Pavlides et al., 2017; Pizzi et al., 2017; Wang et al., 2018; Xu et al., 2017), others suggested that also the OAS thrust ruptured as a reactivated high-angle normal fault during the event, as suggested by geodetic and seismological observations (Cheloni et al., 2017; Scognamiglio et al., 2018; Walters et al., 2018). Although the reactivation of the thrust is not clearly demonstrated (Cheloni et al., 2019), the OAS appears to have played

a role in the aftershock distribution (Chiarabba et al., 2018; Chiaraluce et al., 2017; Pizzi et al., 2017). The earthquakes of the 2016 sequence appear to have nucleated near crosscutting structures that seem to have been loaded by previous ruptures in the sequence (Chiaraluce et al., 2017; Pino et al., 2019). This seismic sequence is thus an excellent case study to better understand the link between structural segmentation, aseismic slip and frictional properties that might control this rupture propagation.

Post-seismic processes during this sequence have been observed in the seismicity (e.g., Albano et al., 2018; Tung & Masterlark, 2018), but no aseismic slip has been detected with geodetic data so far. Using Interferometric Synthetic Aperture Radar (InSAR) time-series from Sentinel-1 data, we document a small but significant post-seismic deformation transient that we find is best explained by aseismic slip on faults associated with this sequence. Surface displacements are presented and analysed. We also explore simple modelling schemes that provide a framework for our interpretation and discussion.

2 Geological setting

The Central Apennines were affected by an extensional phase during the Jurassic, followed by a compressive phase during the Neogene (e.g., Calamita et al., 2011). The OAS (Olevano-Antrodoco-Sibillini thrust) is one the main thrusts resulting from the Neogene compressive phase (Calamita et al., 1994) and has been interpreted as a transpressive ramp (Di Domenica et al., 2012). In the area affected by the 2016-2017 seismic sequence, the OAS thrust is expressed by parallel splays associated with fault-bend folds characteristic of structural ramps (Calamita et al., 2012) (Figure 1). The ongoing ENE oriented extension of 2 to 4 mm/yr (D'Agostino, 2014; Carafa & Bird, 2016; Devoti et al., 2017), which probably began in the Early Pleistocene (e.g., Galadini & Galli, 2000), is currently accommodated through normal fault systems such as the Monte Vettore fault system that hosted the 2016-2017 seismic sequence.

3 Surface displacements during the seismic sequence

3.1 InSAR processing

Synthetic radar interferometry (InSAR) is now systematically used to constrain deformation fields (Elliott, Walters, et al., 2016) and can document centimetre to millimetre scale slow aseismic ground deformation using an adapted processing chain (Hussain et al.,

123 2018; Aslan et al., 2019). We used C band (5.5 cm wavelength) images from Sentinel-1A/B
 124 images (Figure 1-A) spanning almost two years (July 28, 2015, to June 11, 2017) for the
 125 ascending track (A117, subswath IW3). To confirm the main observations made on the
 126 ascending track, we processed descending track (D22, subswaths IW2 and IW3) images from
 127 October 26, 2016 to February 11, 2017. SAR images were processed in VV polarization. We
 128 used the NSBAS processing chain (Doin et al., 2011, 2015) modified for Sentinel data by
 129 Grandin (2015) to generate differential interferograms. The interferogram network, and
 130 examples of unfiltered and uncorrected interferograms, are provided in Figures S1 and S2. The
 131 Shuttle Radar Topography Mission digital elevation model (DEM) at 3 arc sec resolution (Rabus
 132 et al., 2003), resampled at 45m resolution, has been used to accurately coregister the focused
 133 SAR images and to correct interferograms from the topographic contribution to the
 134 interferometric phase. We removed a ramp in range and in azimuth for each interferogram
 135 using the methodology of Cavalié et al., (2007) included in NSBAS. We de-noised the
 136 interferograms before unwrapping using collinearity, a criterion to characterize at each pixel
 137 the local spatial variability of the phase (Pinel-Puysségur et al., 2012). The collinearity was used
 138 to adapt the strength of the filter. We filtered for a window of 12 pixels (400m in range and
 139 700m in azimuth); the filter is described in Doin et al. (2011) and is based on the collinearity
 140 value which weights the complex phase in a sliding window. For the filter we had the option
 141 of adapting the weighting of the phase within windows of different sizes. This size adaptation
 142 depends on the collinearity within the windows. Unwrapping was performed in 2D with the
 143 NSBAS chain (Grandin et al., 2012; Doin et al., 2015). After unwrapping, to account for errors
 144 associated with stratified troposphere, we removed a quadratic cross-function of elevation (z)
 145 and azimuth to ramps in azimuth (y) and in range (x) estimation following the function
 146 $ax+by+c+ez+fz \times az+g \times (z \times az)^2$ using a least-square approach (Daout et al., 2019). Time-series
 147 were then calculated following the NSBAS method (Doin et al., 2011; Daout et al., 2016) using
 148 an approach based on the Small Baseline Subset time-series Analysis (SBAS) of López-Quiroz
 149 et al., (2009)'s algorithm. The smoothing of the pixel time-series is performed by minimizing
 150 the Laplacian of the temporal evolution of the deformation (Cavalié et al., 2007). The final pixel
 151 size is 62 m in azimuth and 37 m in range. We removed pixels with an RMS value greater than
 152 0.7. For the ascending track, we build a time series spanning the 2 years (from 28/07/2015 to
 153 11/06/2017) (Figure S1 and see missing links in the time series in Figure S3). For this complete
 154 2-year time-series, we encounter problems in unwrapping the co-seismic interferograms due

to the large deformation with respect to the Sentinel wavelength in near field (aliasing). Fringes are too close in space and cannot be unwrapped. At pixels that are incoherent in the co-seismic interferogram (i.e. in near field), this causes gaps in the complete (2 yrs) time-series spanning the main earthquakes (Norcia, Amatrice and Campotosto earthquakes) (Figure S3), or lead to an underestimation of the coseismic displacement (brown circles in Figure 2-D). In addition, we also build three time-series in between earthquakes in order to avoid possible bias: before the 24th August (Amatrice earthquake), between 27th August and 26th October (between the Amatrice and Visso earthquake), and after the 30th October (Norcia earthquake). We excluded the SAR data from January 18, 2017, which produced noised interferograms. For the descending track, we build two time-series: between 27th August and 26th October (Visso earthquake), and after the 30th October (Norcia earthquake).

3.2 Time-series results and description of the main features

3.2.1 Ascending Track

The ascending time-series built before the seismic sequence (from July 28th 2015 to August 21th 2016) does not show significant nor localized surface displacements along the main faults (Figure S4-A,B). After October 30th, the (post-Norcia) time-series shows centimetre-scale displacements going away from the satellite in the LOS direction in three areas (Figure 2-B):

- South of Amatrice: In the area affected by the Campotosto earthquake (January 18th, 2017 Mw 5.0 -5.5 EQ) the ground surface moved away from the satellite by more than 60 mm in LOS (Figure 2-B). This coseismic displacement results in a step function in the time series, and can therefore be easily separated from any gradual post-seismic deformation.
- Near Arquata del Tronto: At the southern extremity of surface rupture of the Mw 6.5 October 30th Norcia earthquake (red faults in Figure 2), surface displacements are detected over an area of ~12 km², and follow logarithmic evolution (in Figure 2-D and Figure 3-A see time-series at point 1 where the cumulative post-Norcia displacements in LOS on January 06, 2017 is in average $\sim 35.5 \pm 1.7$ mm and reach 50.5 ± 2.1 mm on 30 April 2017)
- Castelluccio Basin: On the hanging wall of the Mt Vettore fault, slow deformation affects an area of ~50 km² (Figure 2), and is associated with displacements in the

186 LOS direction that are 13.2 ± 1.4 mm in average at Point 2 on 6 January and reach
187 37.9 ± 1.3 mm on 30 April 2017 (Figure 2-D and Figure 3-A).

188 To rule out possible bias due to the Campotosto earthquakes affecting the area near
189 Castelluccio and near Arquata del Tronto, we confirmed these previous observations with a
190 shorter time-series calculated between the Norcia and Campotosto earthquakes (November
191 1st – January 12th) (Figure 4). We prefer to use, for the rest of the manuscript, the longer post-
192 Norcia time-series (November 1st – June 11th) that shows a better signal-noise ratio.

193 In the post Amatrice earthquake time series (August 27 to October 26) we did not observe
194 any localized surface displacements similar to the pattern observed after October 30 (Norcia
195 earthquake) (Figure S4-B). Yet near the town of Amatrice, we observe diffuse surface
196 displacements ($< \sim 2.5$ cm) moving away from the satellite. However, this time-series is
197 constrained by only 9 scenes and 20 interferograms, which prevents from properly (i)
198 constraining a low amplitude signal and (ii) correcting for atmosphere and topography. The
199 surface displacements here have a low signal to noise ratio. The variance of the uncorrelated
200 noise is ~ 40 mm² (see Supplementary Text S1 and Figure S5-A), and the standard deviation
201 sigma of the noise is thus 6.3 mm. We take $3 \times \text{sigma} = 20$ mm to set our limit of detection. The
202 characteristic length scale of correlated noise is 5.0 km and there is autocovariance for
203 distances smaller than ~ 15 km (see Figure S5-A). Observed patterns cannot be differentiated
204 from noise, analyses on SAR images from other satellites should be carried out to confirm or
205 not the post-Amatrice (August 24 earthquake) surface displacements.

206 3.2.2 Descending Track

207 To confirm the post-October 30 (Norcia earthquake) observations we processed
208 descending interferograms (Figure 2-C). We used a 3-month dataset for the descending track
209 (November 1st, 2016 to February 11, 2017). Time-series calculated for the descending track
210 also indicate slow deformation after Norcia earthquake (October, 30), that reaches on average
211 20.5 ± 2.7 mm on January 24 in the LOS direction near Arquata del Tronto (point 1). The
212 deformation reached on average 10.8 ± 1.5 mm in the Castelluccio basin (in point 2). Assuming
213 negligible north-south displacements, by combining the results from ascending and
214 descending tracks, the displacement is dominated by subsidence in this area (Figure S6).

215 In the first order, deformation observed in the descending track is compatible with
216 results inferred from the ascending track. Noise and unwrapping issues led us to mask noisy
217 areas and resulted in more blank pixels in the descending track picture which should be used

with caution. The relief projected in the descending LOS geometry masked a ridge (Mt Bove - Mt Vettore – Mt Gorzano high massifs) (Figure S7). In addition to noise, this resulted in an incoherent area and made unwrapping difficult in that area. The temporal evolution of the features slightly differs from the one in the ascending track. The removal of the 7th November SAR images (affected by strong atmosphere conditions) and the poor constraint on the 19th November SAR images (see the network in Figure S1) could affect the time-series in November. Falcucci et al. (2018) had similar difficulties in using descending SAR images to survey the January 2017 Campotosto seismic event. This might be due to the early morning acquisition time, which amplifies decorrelation due to change of moisture level or thawing.

In addition, we produced time-series calculated between the Amatrice and Norcia earthquakes. This time-series is also affected by a short-wavelength atmospheric turbulence that makes it difficult to interpret (Figure S5-B-C). The variance of noise is $\sim 21 \text{ mm}^2$. The characteristic length scale of correlated noise is 5.0 km and there is autocovariance for distances smaller than $\sim 15 \text{ km}$ (see Figure S5). Again, displacement less than 19 mm for the ascending track and under 14 mm for the descending track, cannot be considered as detectable signal, therefore constraining smaller surface displacements between the Amatrice and Norcia events demands complementary time-series with SAR images from other satellite.

3.2.3 Comparison with GNSS

Few GNSS stations exist near the studied area (ARQT, LNSS see station locations in Figure 2-A). Those GNSS time-series agree with InSAR time-series sampled at the same locations (Figure S8).

3.3 Pattern temporal evolution

By sampling the time-series, the temporal evolution is shown by averaging pixel displacement in area. To enhance the precision of the displacement temporal evolution without using time-series inversion, we temporally track a stable pattern in our unwrapped interferograms (Grandin, 2009). The method is detailed in Supplementary Text S2 and is illustrated by Figures S9-S10.

This method can only be used on areas where the displacement pattern is well defined, and works well in our case to track the evolution of the area near Arquata del Tronto (pattern in Figure 4 which encompasses the point 1). After the Norcia earthquake, the amplitude ratio is higher, and a logarithmic decay is clearly identified there (Figure 3-B). The pattern tracking shows a temporal evolution similar to the averaging surface displacement method. We are

therefore confident that pixel averaging adequately captures the evolution of displacement that we want to describe. For the descending interferograms, the pattern is not precisely trackable and its time evolution is noisy (grey curve in Figure S10-C). This could be due to a poor correlation for the pattern in some interferograms. The evolution of the pattern around Castelluccio Basin (pattern in Figure 4) is poorly resolved too.

4 Modelling 3D deformation field

4.1 Comparison with topography and geology

To interpret the displacement pattern near Arquata del Tronto we compared it with the topography and the geology. There is no correlation with either the slope or the relief (Figure S11), thus we can exclude a gravitational process (i.e. landslide) as an explanation for the observed displacements. Concerning geology, the OAS thrusts fractured Meso-Cenozoic carbonate rocks over Neogene pelagic sediments; the carbonates host some of the largest aquifers in the Apennines while the Neogene sediments are thought to be an aquiclude (Figure S12) (Boni et al., 2010). The OAS delimiting these two units therefore acts as an impermeable boundary (Boni et al., 2010). Petitta et al., (2018) and Valigi et al., (2019) show that the 2016-2017 Italian seismic sequence had effects on spring discharge, water-table levels, and streamflow of those aquifers. In our results, the extent of the Maiolica unit, which hosts a shallow aquifer (Boni et al., 2010), corresponds well with the deformation area near Arquata del Tronto (Figure S11-A-B). The long pre-seismic InSAR time-series did not show any seasonal deformation in this aquifer, which means that groundwater seasonal processes can be excluded (point 1 in Figure 2-D). In addition, it seems difficult to get more than 1 cm of subsidence from winter rain in this area (i.e., Silverii et al., 2016).

4.2 Poro-elastic modelling

Poro-elastic effects have been studied during the 2016-2017 seismic sequence to explore the aftershocks and earthquakes triggering. Tung & Masterlark (2018) suggest that fluid migration caused by the Amatrice earthquake (August 24, 2016) could have triggered the Visso earthquake (October 26, 2016), as well as some associated aftershocks. They inferred that this poro-elastic triggering should have occurred in an intermediately fractured crust. They also estimated that afterslip and viscoelastic-relaxation were quite negligible with respect to poroelastic effects. Albano et al. (2018) also suggest that post-seismic fluid diffusion after the Amatrice earthquake is related to aftershocks. According to their pore fluid diffusion model, there could be some associated afterslip (~ 10 cm) after the Amatrice earthquake.

The Maiolica karst aquifer is shallow and unconfined but Roeloffs (1996) postulates that every aquifer reacts as a confined aquifer to a disturbance at short timescale. Here, to compare with the displacement pattern near Arquata del Tronto, we thus test a forward model of poro-elastic rebound using Relax software (Barbot & Fialko, 2010). We perform a simple poro-elastic model with no lateral variation of diffusivity, that does not account for the spatially heterogeneous distribution of local aquifers. We use the slip distribution models of the Norcia and Visso mainshocks (Maubant et al., 2017), (Figure S13 and Figure 5–A). We do not take into account the Amatrice earthquake since poroelastic rebound from the Amatrice earthquake would likely have stabilized by the time of the Visso and Norcia earthquakes (Figure 4-f in Albano et al., (2018)). We tested several diffusivities for a shallow layer (0-5 km depth), from 1.5 m².s proposed by the Tung and Masterlark (2018)’s aftershocks analysis to 10⁴ m²/s (maximum diffusivity value for karst) (Roeloffs, 1996). Other parameters are described in detail in Figure 5. This model predicts uplift in the Castelluccio Basin and near Arquata del Tronto (Figure 5-D) where we observe subsidence (Figure 5-C). This discrepancy allows us to rule out poro-elastic rebound as the main driver of observed deformation. These poroelastic models are not exhaustive, we only explore simple geometrical configurations using parameter values informed by the local geology, however, other karstic configurations would likely predict uplift as well.

4.3 Viscoelastic modelling

Viscoelastic relaxation in the crust and in the mantle is also an important aseismic process, and has been inferred to be the driver of post-seismic deformation following many earthquakes (e.g., Pollitz et al., 2001; Zhao et al., 2017). To test whether viscoelastic relaxation is a plausible driving mechanism for post-Norcia deformation, we used Relax to perform forward modelling in a simple layered framework. Our model uses the stress perturbation from the three mainshocks: the Amatrice earthquake using the slip distribution of Ragon et al., (2019) added to the Visso and Norcia earthquakes using the slip distribution model of Maubant et al., (2017). We set the upper, middle and lower crust thickness to the values proposed in Laske et al., (2013) and Verdecchia et., al. (2018) (see Table 1). We tested a viscoelastic relaxation governed by a Newtonian rheology with $\dot{\gamma} = \frac{\tau}{\eta}$ where $\dot{\gamma}$ is the viscous strain rate, τ is the deviatoric stress and η is the Newtonian viscosity. We used viscosity values η inferred in studies of viscoelastic relaxation modelling using GPS measurements following 1997 Umbria-Marche earthquakes, or using levelling line measurements following the 1915

Fucino earthquake (Amoruso et al., 2005; Aoudia et al., 2003; Riva et al., 2007). These studies inferred that the ductile structure of the Central Apennines consists of an elastic upper crust overlaying a middle crust (10^{18} - 10^{19} Pa s), a viscous lower crust (10^{17} - 10^{18} Pa s) and the upper mantle (10^{21} Pa s) (Table 1).

Although those models are geometrically simple and do not take into account earthquakes prior 2016, they predict at a long-wavelength uplift (>50 km) in the area where we observe subsidence (Figure 6). This discrepancy in both sign and wavelength allows us to rule out a simple visco-elastic relaxation as the main driver of observed deformation. Visco-elastic models are not exhaustive here: nonlinear viscoelastic rheologies (e.g., power law creep) in lower crust (Freed & Bürgmann, 2004) could also be investigated; however finite element models with creeping lower crust seem also to predict uplift in similar settings of normal fault systems (e.g., Thompson & Parsons, 2016). It is worth also mentioning that the observed logarithmic decay could also potentially be associated with Burgers rheology or shear zone (e.g., Hetland & Zhang, 2014) and could also be investigated in future work. Although, as with the poroelastic modeling, it seems unlikely that a different geometry or rheology could completely reverse the sign of uplift and subsidence (for example, a different rheology would only affect the temporal evolution/spatial distribution).

4.4 Modelling the temporal evolution of afterslip

The logarithmic-like temporal evolution of displacement (Figure 3-A), and the disagreement between the observations and the predictions of poroelastic and viscoelastic models, suggest that afterslip may have been the main driver of postseismic deformation (Marone et al., 1991). Thus to characterize the temporal evolution of the deformation, we fitted the deformation decay (of the raw time-series and of the pattern tracking evolution) with the logarithmic function (Figure 3-A-B respectively) from the Marone et al. (1991) model and Zhou et al. (2018) reformulation for rate-strengthening afterslip:

$$U(t) \approx \alpha \times \ln(1 + c \times t)$$

$$c = \frac{\beta \times V_i}{\alpha}$$

In which U: afterslip in time t, α : characteristic length scale, $\beta \times V_i$: initial rate at the beginning of the post-seismic period, with β a scaling vector by which the sliding rate evolves in response to the stress and V_i the pre-seismic slip rate.

At point 1 (near Arquata del Tronto), the good fit of the afterslip model suggests that the main post-seismic process is likely an afterslip phenomenon. Since we found similar results for c value (describing the temporal decay) using the raw pixel time series and using the pattern tracking (independent for the time-series inversion), we are confident in our fitting (Table 2). The first post-Norcia Sentinel image has been acquired 2.6 days after the Norcia earthquake; therefore those 2.6 days of early afterslip are missing in our data. We compare the temporal decay of the deformation with the cumulative moment and number of aftershocks (blue and green curves in Figure S14). Although the completeness magnitude is high (ML 3 in Figure S5 in Chiaraluce et al., (2017)), those two curves obtained with less than 50 events are not following the same trends, aftershocks do not seem to be induced by afterslip. At point 2 (near Castelluccio di Norcia), the temporal decay does not fit with the afterslip law as the rate at the beginning of the post-seismic period is close to zero (c value for the purple curve fit in Figure 3-A). Other post-seismic processes could be a complementary driver of the deformation here and should be taken into account in further modelling (i.e. finite element forward model accounting for afterslip superimposed with poro-elastic rebound and viscoelastic relaxation).

4.5 Afterslip modelling on faults

4.5.1 Inversion strategy and fit to the data

To obtain the afterslip distribution we use the Classic Slip Inversion (CSI) Python tools (Elliott, Jolivet, et al., 2016; Jolivet et al., 2015) to invert for the slip. We use the constrained least-squares formula of Tarantola (2005) to solve the inverse problem (see details in Supplementary Text S3 and Figures S15 to S18). We chose a dip of 40° for the Mt Vettore fault (as Cheloni et al., (2017) see Figure S15) and project the fault down dip from the mapped fault at the surface (modeled fault geometry in yellow Figure 2-A). We discretize the fault into 88 rectangular patches. The smoothed cumulative surface displacement on February 11, 2017, measured in both ascending and descending tracks are inverted to obtain the afterslip distribution. The resolution is good for short-wavelength features at shallow depths (<5 km depth) (Figure S16). As several geometries have been used to model the mainshock rupture (OAS reactivation, antithetic fault), we explored three cases: (1) slip only on the Mt Vettore Fault, (2) slip on the Mt Vettore Fault and on the OAS and (3) slip on the Mt Vettore Fault and an antithetic fault with a geometry similar to the one proposed by Cheloni et al., (2019) and Maubant et al., (2017) (dip 65°) (Figures S17 to S20). We also explored the rake: (1) dip-slip only and (2) variable rake. A rigorous statistical comparison between the cases is difficult due

to variation in model parameters and number of degrees of freedom, as explained by Cheloni et al., (2019). We thus used the RMS values to compare the different cases. Here the case that reduces the residuals the most is the inversion with an antithetic fault. This case is in addition consistent with geological and seismological observations (i.e. Cheloni et al., 2019). The different cases were compared using the RMS (root mean square), and allows us to choose the inversion with an antithetic fault (Table S2).

Both ascending and descending tracks are well modelled (Figure 7). It is worth noting that the resolution is low at depth (Figure S16). The total geodetic moment released by the afterslip model is equivalent to $M_w \sim 5.78$ (without taking into account the Campotosto earthquake) which corresponds to $\sim 8.7\%$ of the geodetic moment released during the Norcia earthquake ($M_w 6.5$).

4.5.2 Afterslip distribution

4.5.2.1 Near Arquata del Tronto

We obtain a maximum afterslip of ~ 100 mm below Arquata del Tronto at shallow depth (0- 2 km depth) (Figure 7 Figure 8). The shallow part depth $< 3-4$ km toward the south is associated with very low coseismic slip for both the Norcia and Amatrice earthquakes and is located at the edge of the coseismic asperities. This is also consistent with an afterslip process: the afterslip is located where the slip gradient increased the shear stress on the unruptured portions. This feature has also been observed after the L'Aquila earthquake (D'Agostino et al., 2012). We calculate a geodetic moment released equivalent to $M_w \sim 5.1$ in this area (patches above 2km depth and south of Arquata del Tronto). The cumulative moment released by the seismicity (Figure S14-D) at this date corresponds to $\sim 2.0\%$ of the moment released by the afterslip model. The deformation is thus mainly aseismic.

Brozzetti et al., (2019) found some surface ruptures south of the 30th October Norcia rupture ($42^\circ 47'N$ in their Fig. 1-C) mapped by Villani et al., (2018). The ruptures mapped by Brozzetti et al., (2019) are subtle and according to us may correspond to post-seismic deformation, as was observed after the L'Aquila earthquake (D'Agostino et al., 2012).

4.5.2.2 Castelluccio Basin

We observe a maximum slip of ~ 170 mm below Castelluccio at ~ 5 km depth. Some afterslip overlaps with the coseismic rupture area, but not in the area of maximum coseismic slip (Figure 7). As also the shape of the measured time-series did not fit with the afterslip's law

(Figure 3-A), a sole afterslip process is unlikely. It is possible that poroelastic and fluid flow processes could be at work here since there is a large basal aquifer (Boni et al., 2010). Modelling accounting for both processes could be performed, for example fully coupled poroelastic finite element numerical modelling with spatially variable material properties (e.g., Albano et al., 2017).

5 Discussion

5.1 Afterslip mechanism near Arquata del Tronto

Based on Dietrich (1979), Ruina (1983) and Marone et al., (1991) equations, we estimated α (the characteristic length scale over which the elastic stress changes by order of the frictional stress) in Figure 3 and Table 2. We can now estimate the friction parameter $(a - b)$ since according to Perfettini and Avouac (2004) and Zhou et al., (2018):

$$\alpha = \frac{(a - b) \times \sigma}{k}$$

with k : the effective stiffness, and σ the effective normal stress. This relation can be applied to our case since the duration of our analysis (~200 days) is much shorter than the characteristic time t_d (Gualandi et al., 2014). Where $t_d = \frac{\alpha}{V_{pl}}$ (>12 years) where V_{pl} is the local plate loading rate (<2.1 mm/yr (Puliti et al., 2020)). The fit to the displacement of the time-series sampled in point 1 by an afterslip law (Figure 3) leads to $\alpha = 30.9 \pm 4.0$ mm. To estimate σ , we assume that the mean normal stress may vary from the hydrostatic to lithostatic pressure, with a rock density of 2.5 kg.m^{-3} (Albano et al., 2018) and a depth of the slipping area of 3-5 km. This leads to $\sigma = 44\text{-}124$ MPa. For $k = G/h$ with $G = 30$ GPa (the shear modulus near the surface) and $h = 8$ km (rate-strengthening depth based on the coseismic slip models), we obtain $(a-b) = 3 \times 10^{-3} - 7.8 \times 10^{-4}$. More complex models (e.g. finite element model, heterogeneous $(a-b)$ values) could be performed by future studies in order to reproduce the surface displacements and to propose a more precise value of $(a-b)$. However, our estimation is in agreement with experiments on carbonates. Scuderi and Collettini (2016) found values of $(a - b)$ evolve from velocity strengthening behaviour ($a - b \approx 0.005$) at fluid pressure condition of sub-hydrostatic to a velocity neutral behaviour ($a - b$ approaching 0), when the fault is at near lithostatic fluid pressure. Pluymakers et al., (2016) found that wet anhydrite and dolomite gouges at depths <6 km, exhibit $(a-b)$ values ranging from 10^{-2} to 10^{-4} .

In Arquata del Tronto, the obtained (a-b) positive value (0.0026) is in agreement with a velocity-strengthening area. The obtained (a-b) value is also very low and corresponds to a quasi-neutral rate-dependency of friction implying a high sensitivity to stress perturbations. Potential stress perturbations needed to reach strengthening may involve either a decrease in effective normal stress and / or an increase in shear stress (Scholz, 1998). Walters et al., (2018) calculated the Coulomb stress change (CFF) after the Norcia earthquake. At shallow depths, they found a positive CFF change below Arquata del Tronto, which might have triggered the observed afterslip in this area. An increase in pore fluid pressure would have a similar effect. The afterslip is located near to the Olevano-Antrodoco-Sibillini (OAS) thrust. Chiarabba et al., (2018) detected a high V_p/V_s near the OAS thrust (around 3 km depth), indicating high pore pressure that could have been further increased by nearby major shocks (Amatrice and Norcia earthquake). This high pore pressure could have induced the observed afterslip allowing for the release of the residual accumulated stress. This scenario is also supported by the fact that pore fluid pressure variation is common in this area, where the inherited structures seem to control the rupture pattern and could also concentrate high shear stress. Future work could examine detailed Coulomb Stress modelling with high spatial resolution taking all fault complexity and pore fluid effects into account. These works could be useful, despite some limitations related to the resolution of the coseismic slip and by the ambiguity about which faults were involved.

If this overlapping of coseismic slip and afterslip is real below 4-5 km depth (below Arquata del Tronto) could be an artefact caused by the smoothing used to regularize the inversion of the slip distribution, combined with the fact that the resolution at depth is limited. If this overlapping of co-seismic and slow slip was confirmed, a simple interpretation of rate and state friction cannot be proposed here. Overlaps between coseismic slip and afterslip have been observed after several earthquakes. For example after 2004 M_w 6 Parkfield earthquake, afterslip was inferred to overlap the coseismic slip (Freed, 2007). Johnson et al., (2006) proposed frictional spatial heterogeneity to explain the afterslip distribution, and proposed (a-b) values on the order of $10^{-4} - 10^{-3}$. After the 2015 Ilaapel M_w 8 megathrust earthquake, Barnhart et al. (2016) also observed afterslip and coseismic slip overlapping, and advocated that stress heterogeneities likely provide the primary control on the afterslip distribution. After the 1978 M_w 7.3 Tabas-e-Golshan earthquake, Zhou et al., (2018) also observe such overlapping and found $(a-b) \sim 3 \cdot 10^{-3}$. They proposed that the fault is creeping during the whole

interseismic period and that the earthquake propagates through the rate-strengthening region although they also stated that a change in frictional properties is also possible due to shear heating. Thomas et al., (2017), after the 2003 M_w 6.8 Chengkung earthquake estimated that (a-b) varies with depth from 0.018 near the surface to less than 0.001 at depth larger than 19 km, to explain areas showing both seismic and aseismic behavior they advocated shear heating processes. Using numerical modelling, Noda and Lapusta (2013) argued that coseismic slip could propagate into velocity-strengthening regions of a fault, and that these regions can appear locked or creeping during the interseismic period. They observed this behavior due to rapid shear heating. For our case it is difficult to pinpoint the particular fault zone mechanism here, but we can propose a switch to (a-b) positive value below 4-5 km depth, due to coseismic shear heating processes (Thomas et al., 2017) requiring temperature changes of several hundred degrees (e.g., Rice, 2006). This could be a potential explanation since Smeraglia et al., (2017) suggested that nanostructures from the Mt Vettoreto Fault rocks were generated by coseismic shear heating and grain comminution (reduction of particle sizes).

5.2 Barrier of rupture propagation

Near Arquata del Tronto, at shallow depth (< 3-4 km) we observe that the aseismic slip is localized on an area associated with limited coseismic slip, for both the Amatrice and Norcia earthquakes. The observed aseismic slip seems located in an area where coseismic ruptures have not propagated during the Norcia or Amatrice earthquakes. This slowly slipping zone is located near a structural complexity (inherited thrust OAS), that Chiaraluce et al., (2017), Pizzi et al., (2017) and Puliti et al. (2020) interpreted as a barrier that concentrated stress after the Amatrice earthquake. We propose that maybe as a result of this geometric complexity (that may concentrate high stress and high porosity resulting from fracturation), this zone might be also characterized by frictional properties at shallow depth that could have favoured the arrest of the ruptures (e.g., Kaneko et al., 2010). Scholz (1998) shows that when an earthquake propagates into a velocity strengthening field it will produce a negative stress drop, that rapidly terminates propagation. Our simple calculations yield here a slightly positive (a-b) value, consistent with this model.

6 Conclusion

The Sentinel-1 InSAR time-series show cm-scale post-seismic displacements after the Norcia earthquake (October 30). The displacements are more obvious in the ascending track than in the descending track, and correspond to ~1 to 5 cm of subsidence in the Castelluccio basin and at the southern tip of the Norcia surface rupture. In the area of the Castelluccio basin, whether the deformation can be explained by afterslip only is less clear and would require additional modelling to include the effects of poro-elastic release or shear zone deformation. At the southern edge of the Norcia coseismic rupture, deformation evolves with time following a logarithmic decay consistent with afterslip triggered by the Norcia earthquake. Although good observations of the 2.6 first days of afterslip are not available, after 3 months the slow deformation evidenced in this study released a geodetic moment of $M_w \sim 5.78$, which corresponds to ~8.7 % of the geodetic moment released during the Norcia earthquake. Cumulated moment released by aftershocks in the southern tip of the Norcia rupture accounts for ~2.0 % of the moment released by the modelled deformation, the deformation is thus here mainly aseismic. This afterslip takes place at the southern tip of the Norcia rupturing patch, and seems to have acted as a barrier to the propagation of Norcia and Amatrice ruptures at shallow depth (<3-4 km). Our results suggest that the observed post-Norcia earthquake afterslip might have been triggered in response to heterogeneities of pore fluid pressure, potentially facilitated by structural complexity and intense faulting in the area. Such small and localized slow deformation due to afterslip may be best detected with InSAR time-series in regions of sparse GNSS coverage. It might actually be quite common and could be an underestimated phenomenon. This likely points toward a bias in the literature in favour of high and wider afterslip, that is easier to detect. However detection of small afterslip transient is crucial to further understand the physics of earthquakes, and the link between slow slip and seismic rupture (e.g., Kaneko et al., 2010; Rousset et al., 2017).

7 Tables

7.1 Table 1

	Depths (km)	Model 1 η	Model 2 η	Model 3 η	Model 4 η
Upper crust	0-10.7	Elastic	Elastic	Elastic	Elastic
Middle crust	10.7-21.7	Elastic	10^{19} Pa s	10^{18} Pa s	10^{18} Pa s
Lower crust	21.7-33.1	10^{18} Pa s	10^{18} Pa s	10^{18} Pa s	10^{17} Pa s
Upper mantle	33.1-below	10^{21} Pa s	10^{21} Pa s	10^{21} Pa s	10^{21} Pa s

Table 1 : Crust and mantle viscosities (η) tested in Figure 6.

7.2 Table 2

Fit	α	C (days ⁻¹)	γ	R2
Point 1 near Arquata del Tronto	30.9 ± 4.0	0.26 ± 0.26	-6.4 ± 6.2	0.87
Point 2 near Castelluccio di Norcia	156.4 ± 230.7	0.003 ± 0.007	2.54 ± 3.5	0.81
Pattern tracking	1.6 ± 0.1	0.24 ± 0.1	-0.37 ± 0.21	0.95

Table 2: Parameters obtained from the fit of the afterslip model in Figure 3 using a non-linear least squares method (Moré, 1978). R2 is the coefficient of determination of the regression. The pattern encompasses the point 1 (see Figure 4)

8 Figures
8.1 Figure 1

Figure 1: Neotectonic framework of the Central Apennines. Black lines are main normal active faults compiled from Benedetti (1999), Tesson et al., (2016) and references therein (VBFS: Mt Vettore – Mt Bove Fault System, A-C F: Amatrice-Campotosto Fault). In red, coseismic surface ruptures observed in the field after the mainshocks of the Central Italy 2016-2017 seismic sequence (Villani, Civico, et al., 2018). **(A)** Blue and red rectangles are, respectively, the descending (D22 subswaths IW2 and IW3) and ascending (A117, subswath IW3) Sentinel tracks processed in this study. **(B)** Black dashed line represents the Neogene OAS thrust (Olevano-AnTRODoco-Sibillini thrust) (Di Domenica et al., 2012). In orange the Amatrice-Campotosto fault which was activated during the 2016—2017 seismic sequence. ARQT and LNSS (in B) are GNSS stations available in this region from http://geodesy.unr.edu/NGLStationPages/gpsnetmap/GPSNetMap_MAG.html (Blewitt et al., 2018).

Figure 2: (A) Map of the studied area, green squares are the GPS stations, points 1-2 are the points of sampling. The modeled geometries of the Mt Vettore and antithetic faults are plotted (see section 4.5). Circles are the earthquakes (see Figure 1). (B-C): Time-series calculated cumulative post-Norcia (30 oct) displacements for ascending (B) and descending track (C), respectively. In panels A,B,C,E : Red lines are surface ruptures associated with 30th October 2016 Norcia earthquake (Villani, Civico, et al., 2018), and black lines are active faults. Black dashed line is the OAS thrust. (VBFS: Mt Vettore – Mt Bove Fault System, A-C F: Amatrice-Campotosto Fault, OAS: Olevano-Antròdoco-Sibillini thrust). (D) Displacement through time averaged over a circle of 20 pixels (dashed circles in panel E) in diameter around points 1-2. (E) Cumulative displacement on June 11, 2017, with respect to the 1st November 2016 date.

Figure 3: **(A)** Temporal variation of displacement averaged over a circle of 20 pixels in diameter through time of point 1 (black curve) and 2 (purple dashed curve) localized in Figure 2-E. We fitted the curve with afterslip model (see results in Table 2). **(B)** Temporal variation of pattern amplitude. Pattern is indicated in Figure 4. In red, the fit with an afterslip model (see results in Table 2).

580
581 8.4 Figure 4

582
583 Figure 4: Mean velocity map between Mw 6.5 Norcia and Mw 5.7 Campotosto earthquakes
584 (November 1st – January 12th). This map is computed using 35 ascending interferograms. The
585 swath A-B is used in Figure S11 to compare InSAR displacements and topography. Circles
586 represent the earthquakes (see color code in Figure 1).
587
588

590

591 Figure 5: Poro-elastic rebound modelled with Relax software (Barbot & Fialko, 2010). We used
 592 the coseismic slip models of the Norcia and Visso earthquakes from Maubant et al., (2017).
 593 **(A)** Mapview of the coseismic slip distribution. **(B)** Coseismic $-\sigma_{kk}$ modelled at 500m depth,
 594 calculated in an undrained homogenous elastic medium (Poisson $\nu=0.34$; Lamé $\lambda=6.36E+04$
 595 MPa, shear modulus $G=30$ GPa, Gravity wavelength $\gamma=5.39E-04$ km⁻¹). To remind, the pore
 596 pressure change (Δp) is equal to $-B \cdot \sigma_{kk}/3$, with (B the Skempton's coefficient). **(C)** Post-Norcia
 597 cumulative displacement map (11th February 2017) observed by InSAR (uplift component). **(D)**

Model-predicted poro-elastic rebound as of 25th February 2017, obtained using the difference between drained and undrained conditions. Poisson drained is 0.26 (cf. layer 1 in Albano et al. 2018), and for undrained conditions is 0.34 (from $\beta=0.4$ in the table 1 and equation 15 in Barbot et al., (2010)). The layer between (0 and 5 km) is characterized by a diffusivity equals to $1.5 \text{ m}^2/\text{s}$ (Tung & Masterlark, 2018). **(E)** Modelled poro-elastic rebound time-series at point A using several diffusivities, from $1.5 \text{ m}^2/\text{s}$ (Tung & Masterlark, 2018) to $10^4 \text{ m}^2/\text{s}$ (Roeloffs, 1996). Because the delay time is inversely proportional to the assumed diffusivity, the response for high diffusivity is shorter. (VBFS: Mt Vettore – Mt Bove Fault System, A-C F: Amatrice-Campotosto Fault, OAS: Olevano-Antrodoco-Sibillini thrust).

8.6 Figure 6

Figure 6: Surface displacements predicted by visco-elastic models, we used the Amatrice, Norcia and Visso earthquakes coseismic slip distribution modelled by Ragon et al., (2019) and Maubant et al., (2017). **(A)** Coseismic slip distribution of the Amatrice earthquake from Ragon et al., (2019) **(B-C-E-F)** Surface displacements predicted by four visco-elastic models on 24th February 2017 using Relax software (Barbot & Fialko, 2010). Mantle viscosity is fixed for all models at 10^{21} Pas. **(D)** Post-Norcia earthquake (30th October) cumulative displacement map (11th February 2017) observed by InSAR (uplift component). (VBFS: Mt Vettore – Mt Bove Fault System, A-C F: Amatrice-Campotosto Fault, OAS: Olevano-Antrodoco-Sibillini thrust)

Figure 7: Result of afterslip inversion using the CSI software (see Supplementary text S3 for details). **(A)** Subsampled displacement maps for the ascending (left column) and descending (right column) tracks. Top row shows the input subsampled data, middle row shows the synthetic displacements predicted by the model, and bottom row shows the residuals (i.e. model - data). The modeled fault geometries used for the inversion are plotted in Figure 2-A. The input maps are the smoothed cumulative displacement maps on February 11th (calculated in the ascending and descending post-30th October time-series). **(B)** Afterslip distribution inverted from the data on the main fault and the antithetic fault. Black arrows in patches show the rake. Our model has low resolution at depth (Figure S16). Coseismic slip distributions for

632 the Amatrice and Norcia earthquakes (Scognamiglio et al., (2018) are superimposed in white
633 and red , respectively. Comparison with the Cheloni et al.,(2019) model in Figure S21 shows
634 similar patterns.

635

636

638
639 Figure 8: Afterslip distribution model plotted on mapview. The slip distribution is inverted from
640 the data on the main fault and the antithetic fault (see Figure 7). (VBFS: Mt Vettore – Mt Bove
641 Fault System, A-C F: Amatrice-Campotosto Fault). Circles are the main earthquakes of the
642 seismic sequence.
643

9 Acknowledgments

We would like to thank the Editor, Associate Editor, Mong-Han Huang and two anonymous reviewers for their constructive suggestions, which helped to improve the manuscript substantially. We acknowledge the French Spatial Agency CNES (Centre National d'Etudes Spatiales) for funding this study (Program THEIA and CNES post-doctorate fellowship for L.Pousse-Beltran). This work was also funded through the TELLUS-ALEAS program from Institut des Sciences de l'Univers (INSU). CNES is warmly acknowledged from providing us with the Pléiades satellite images through ISIS and CEOS_seismic pilot (ESA) programs. Nicola D'Agostino was under an invited researcher position at Univ. Grenoble Alpes - OSUG - ISTERre when most of the work was performed. Most of the computations presented in this paper were performed using the Luke platform of the CIMENT infrastructure (<https://ciment.ujf-grenoble.fr>), which is supported by the Rhône-Alpes region (grant CPER07_13 CIRA), the OSUG@2020 labex (reference ANR10 LABX56), and the Equip@Meso project (reference ANR-10-EQPX-29-01) of the programme Investissements d'Avenir supervised by the Agence Nationale pour la Recherche. Franck Thollard, Christophe Laurent, Erwan Pathier, Louise Maubant and Simon Daout are warmly acknowledged for help and advice on NSBAS processing chain. We are also grateful to Théa Ragon and Romain Jolivet for support and advice in slip distribution modelling, the CSI software will be soon released online. We thank Louise Maubant and James Hollingsworth for sharing the Norcia slip distribution model. Alberto Pizzi, Bruno Pace, Paolo Boncio and Irene Puliti are also warmly acknowledged for fruitful discussions. The European Space Agency (ESA) Copernicus program has provided free and high-quality SAR data from Sentinel-1A and 1B (ESA, <https://scihub.copernicus.eu>) through the PEPS platform of CNES (Copernicus 2018 for Sentinel data. We thank Simone Tarquini for the supply of 10 m resolution TINITALY DEM. The afterslip distribution model will be provided in [10.5281/zenodo.3755127](https://doi.org/10.5281/zenodo.3755127). Seismicity used for this research is included in (Chiaraluce et al., 2017) and in <http://cnt.rm.ingv.it/>. The GNSS stations are downloaded from <http://geodesy.unr.edu/NGLStationPages/stations/>

10 References

- Albano, M., Barba, S., Solaro, G., Pepe, A., Christian, B., Moro, M., et al. (2017). Aftershocks, groundwater changes and postseismic ground displacements related to pore pressure gradients: Insights from the 2012 Emilia-Romagna earthquake. *Journal of Geophysical Research: Solid Earth*, 122(7), 5622–5638. <https://doi.org/10.1002/2017JB014009>
- Albano, M., Barba, S., Saroli, M., Polcari, M., Bignami, C., Moro, M., et al. (2018). Aftershock rate and pore fluid diffusion: Insights from the Amatrice-Visso-Norcia (Italy) 2016 seismic sequence. *Journal of Geophysical Research: Solid Earth*, 0(ja). <https://doi.org/10.1029/2018JB015677>
- Amoruso, A., Crescentini, L., D’Anastasio, E., & Martini, P. M. D. (2005). Clues of postseismic relaxation for the 1915 Fucino earthquake (central Italy) from modeling of leveling data. *Geophysical Research Letters*, 32(22). <https://doi.org/10.1029/2005GL024139>
- Aoudia, A., Borghi, A., Riva, R., Barzaghi, R., Ambrosius, B. a. C., Sabadini, R., et al. (2003). Postseismic deformation following the 1997 Umbria-Marche (Italy) moderate normal faulting earthquakes. *Geophysical Research Letters*, 30(7). <https://doi.org/10.1029/2002GL016339>
- Aslan, G., Lasserre, C., Cakir, Z., Ergintav, S., Özarpaci, S., Dogan, U., et al. (2019). Shallow Creep Along the 1999 Izmit Earthquake Rupture (Turkey) From GPS and High Temporal Resolution Interferometric Synthetic Aperture Radar Data (2011–2017). *Journal of Geophysical Research: Solid Earth*, 124(2), 2218–2236. <https://doi.org/10.1029/2018JB017022>
- Avouac, J.-P. (2015). From Geodetic Imaging of Seismic and Aseismic Fault Slip to Dynamic Modeling of the Seismic Cycle. *Annual Review of Earth and Planetary Sciences*, 43(1), 233–271. <https://doi.org/10.1146/annurev-earth-060614-105302>
- Barbot, S., & Fialko, Y. (2010). A unified continuum representation of post-seismic relaxation mechanisms: semi-analytic models of afterslip, poroelastic rebound and viscoelastic flow: Semi-analytic models of postseismic transient. *Geophysical Journal International*, 182(3), 1124–1140. <https://doi.org/10.1111/j.1365-246X.2010.04678.x>

699 Barnhart, W. D., Murray, J. R., Briggs, R. W., Gomez, F., Miles, C. P. J., Svarc, J., et al. (2016).
700 Coseismic slip and early afterslip of the 2015 Illapel, Chile, earthquake: Implications for
701 frictional heterogeneity and coastal uplift. *Journal of Geophysical Research: Solid Earth*,
702 121(8), 6172–6191. <https://doi.org/10.1002/2016JB013124>

703 Benedetti, L. (1999). *Sismotectonique de l'Italie et des régions adjacentes: fragmentation du*
704 *promontoire adriatique* (PhD Thesis). Paris 7.

705 Blewitt, G., Hammond, W., & Kreemer, C. (2018). Harnessing the GPS Data Explosion for
706 Interdisciplinary Science. *Eos*, 99. <https://doi.org/10.1029/2018EO104623>

707 Boni, C. F., TARRAGONI, C., MARTARELLI, L., & PIERDOMINICI, S. (2010). STUDIO
708 IDROGEOLOGICO NEL SETTORE NORD-OCCIDENTALE DEI MONTI SIBILLINI: UN
709 CONTRIBUTO ALLA CARTOGRAFIA IDROGEOLOGICA UFFICIALE. *Italian Journal of*
710 *Engineering Geology and Environment*, 2, 16. <https://doi.org/10.4408/IJEGE.2010->
711 02.O-02

712 Brozzetti, F., Boncio, P., Cirillo, D., Ferrarini, F., Nardis, R. de, Testa, A., et al. (2019). High-
713 Resolution Field Mapping and Analysis of the August–October 2016 Coseismic Surface
714 Faulting (Central Italy Earthquakes): Slip Distribution, Parameterization, and
715 Comparison With Global Earthquakes. *Tectonics*, 0(0).
716 <https://doi.org/10.1029/2018TC005305>

717 Bürgmann, R. (2018). The geophysics, geology and mechanics of slow fault slip. *Earth and*
718 *Planetary Science Letters*, 495, 112–134. <https://doi.org/10.1016/j.epsl.2018.04.062>

719 Calamita, F., Cello, G., Deiana, G., & Paltrinieri, W. (1994). Structural styles, chronology rates of
720 deformation, and time-space relationships in the Umbria-Marche thrust system
721 (central Apennines, Italy). *Tectonics*, 13(4), 873–881.
722 <https://doi.org/10.1029/94TC00276>

723 Calamita, F., Satolli, S., Scisciani, V., Eserice, P., & Pace, P. (2011). Contrasting styles of fault
724 reactivation in curved orogenic belts: Examples from the Central Apennines (Italy).
725 *Geological Society of America Bulletin*, 123(5–6), 1097–1111.
726 <https://doi.org/10.1130/B30276.1>

727 Calamita, F., Satolli, S., & Turtù, A. (2012). Analysis of thrust shear zones in curve-shaped belts:
728 Deformation mode and timing of the Olevano-AnTRODoco-Sibillini thrust
729 (Central/Northern Apennines of Italy). *Journal of Structural Geology*, 44, 179–187.
730 <https://doi.org/10.1016/j.jsg.2012.07.007>

731 Carafa, M. M. C., & Bird, P. (2016). Improving deformation models by discounting transient
732 signals in geodetic data: 2. Geodetic data, stress directions, and long-term strain rates
733 in Italy. *Journal of Geophysical Research: Solid Earth*, 121(7), 5557–5575.
734 <https://doi.org/10.1002/2016JB013038>

735 Cavalié, O., Doin, M.-P., Lasserre, C., & Briole, P. (2007). Ground motion measurement in the
736 Lake Mead area, Nevada, by differential synthetic aperture radar interferometry time
737 series analysis: Probing the lithosphere rheological structure. *Journal of Geophysical*
738 *Research: Solid Earth*, 112(B3), n/a–n/a. <https://doi.org/10.1029/2006JB004344>

739 Cheloni, D., De Novellis, V., Albano, M., Antonioli, A., Anzidei, M., Atzori, S., et al. (2017).
740 Geodetic model of the 2016 Central Italy earthquake sequence inferred from InSAR
741 and GPS data. *Geophysical Research Letters*, 44(13), 2017GL073580.
742 <https://doi.org/10.1002/2017GL073580>

743 Cheloni, D., Falcucci, E., & Gori, S. (2019). Half-graben rupture geometry of the 30 October
744 2016 MW 6.6 Mt. Vettore-Mt. Bove earthquake, central Italy. *Journal of Geophysical*
745 *Research: Solid Earth*, 0(ja). <https://doi.org/10.1029/2018JB015851>

746 Chen, K. H., & Bürgmann, R. (2017). Creeping faults: Good news, bad news? *Reviews of*
747 *Geophysics*, 55(2), 2017RG000565. <https://doi.org/10.1002/2017RG000565>

748 Chiarabba, C., Gori, P. D., Cattaneo, M., Spallarossa, D., & Segou, M. (2018). Faults geometry
749 and the role of fluids in the 2016-2017 Central Italy seismic sequence. *Geophysical*
750 *Research Letters*, 0(ja). <https://doi.org/10.1029/2018GL077485>

751 Chiaraluce, L., Stefano, R. D., Tinti, E., Scognamiglio, L., Michele, M., Casarotti, E., et al. (2017).
752 The 2016 Central Italy Seismic Sequence: A First Look at the Mainshocks, Aftershocks,
753 and Source Models. *Seismological Research Letters*, 88(3), 757–771.
754 <https://doi.org/10.1785/0220160221>

755 Cirella, A., Pezzo, G., & Piatanesi, A. (2018). Rupture Kinematics and Structural - Rheological
756 Control of the 2016 Mw6.1 Amatrice (Central Italy) Earthquake from Joint Inversion of
757 Seismic and Geodetic Data. *Geophysical Research Letters*, 0(ja).
758 <https://doi.org/10.1029/2018GL080894>

759 Civico, R., Pucci, S., Villani, F., Pizzimenti, L., Martini, P. M. D., Nappi, R., & Group, the O. E.
760 W. (2018). Surface ruptures following the 30 October 2016 Mw 6.5 Norcia earthquake,
761 central Italy. *Journal of Maps*, 14(2), 151–160.
762 <https://doi.org/10.1080/17445647.2018.1441756>

763 D’Agostino, N. (2014). Complete seismic release of tectonic strain and earthquake recurrence
764 in the Apennines (Italy). *Geophysical Research Letters*, 41(4), 1155–1162.
765 <https://doi.org/10.1002/2014GL059230>

766 D’Agostino, N., Cheloni, D., Fornaro, G., Giuliani, R., & Reale, D. (2012). Space-time distribution
767 of afterslip following the 2009 L’Aquila earthquake. *Journal of Geophysical Research:*
768 *Solid Earth*, 117, n/a-n/a. <https://doi.org/10.1029/2011jb008523>

769 Daout, S., Jolivet, R., Lasserre, C., Doin, M.-P., Barbot, S., Tapponnier, P., et al. (2016). Along-
770 strike variations of the partitioning of convergence across the Haiyuan fault system
771 detected by InSAR. *Geophysical Journal International*, 205(1), 536–547.

772 Daout, S., Sudhaus, H., Kausch, T., Steinberg, A., & Dini, B. (2019). Interseismic and Postseismic
773 Shallow Creep of the North Qaidam Thrust Faults Detected with a Multitemporal InSAR
774 Analysis. *Journal of Geophysical Research: Solid Earth*, 124(7), 7259–7279.
775 <https://doi.org/10.1029/2019JB017692>

776 Devoti, R., D’Agostino, N., Serpelloni, E., Pietrantonio, G., Riguzzi, F., Avallone, A., et al. (2017).
777 A Combined Velocity Field of the Mediterranean Region. *Annals of Geophysics*, 60(2).
778 <https://doi.org/10.4401/ag-7059>

779 Di Domenica, A., Turtù, A., Satolli, S., & Calamita, F. (2012). Relationships between thrusts and
780 normal faults in curved belts: New insight in the inversion tectonics of the Central-
781 Northern Apennines (Italy). *Journal of Structural Geology*, 42, 104–117.
782 <https://doi.org/10.1016/j.jsg.2012.06.008>

Dieterich, J. H. (1979). Modeling of rock friction: 1. Experimental results and constitutive equations. *Journal of Geophysical Research: Solid Earth*, 84(B5), 2161–2168. <https://doi.org/10.1029/JB084iB05p02161>

Doin, M.-P., Lodge, F., Guillaso, S., Jolivet, R., Lasserre, C., Ducret, G., et al. (2011). Presentation of the small baseline NSBAS processing chain on a case example: the Etna deformation monitoring from 2003 to 2010 using ENVISAT data. In *Proceedings of the Fringe symposium, Frascati, Italy, ESA SP-697*.

Doin, M.-P., Twardzik, C., Ducret, G., Lasserre, C., Guillaso, S., & Jianbao, S. (2015). InSAR measurement of the deformation around Siling Co Lake: Inferences on the lower crust viscosity in central Tibet. *Journal of Geophysical Research: Solid Earth*, 120(7), 5290–5310. <https://doi.org/10.1002/2014JB011768>

Elliott, J. R., Jolivet, R., González, P. J., Avouac, J.-P., Hollingsworth, J., Searle, M. P., & Stevens, V. L. (2016). Himalayan megathrust geometry and relation to topography revealed by the Gorkha earthquake. *Nature Geoscience*, 9(2), 174–180. <https://doi.org/10.1038/ngeo2623>

Elliott, J. R., Walters, R. J., & Wright, T. J. (2016). The role of space-based observation in understanding and responding to active tectonics and earthquakes. *Nature Communications*, 7, 13844. <https://doi.org/10.1038/ncomms13844>

Falcucci, E., Gori, S., Bignami, C., Pietrantonio, G., Melini, D., Moro, M., et al. (2018). The Campotosto seismic gap in between the 2009 and 2016-2017 seismic sequences of central Italy and the role of inherited lithospheric faults in regional seismotectonic settings. *Tectonics*, 0(ja). <https://doi.org/10.1029/2017TC004844>

Freed, A. M. (2007). Afterslip (and only afterslip) following the 2004 Parkfield, California, earthquake. *Geophysical Research Letters*, 34(6), L06312. <https://doi.org/10.1029/2006GL029155>

Freed, A. M., & Bürgmann, R. (2004). Evidence of power-law flow in the Mojave desert mantle. *Nature*, 430(6999), 548–551. <https://doi.org/10.1038/nature02784>

810 Galadini, F., & Galli, P. (2000). Active Tectonics in the Central Apennines (Italy) –Input Data for
811 Seismic Hazard Assessment. *Natural Hazards*, 22(3), 225–268.
812 <https://doi.org/10.1023/A:1008149531980>

813 Grandin, R. (2009). *L'apport de la géodésie spatiale dans la compréhension du processus de*
814 *rifting magmatique : l'exemple de l'épisode en cours en Afar Ethiopien (2005-2009)*.
815 Institut de physique du globe (Paris). Retrieved from
816 <http://www.theses.fr/2009GLOB0013>

817 Grandin, R. (2015). Interferometric Processing of SLC Sentinel-1 TOPS Data. In *FRINGE'15:*
818 *Advances in the Science and Applications of SAR Interferometry and Sentinel-1 InSAR*
819 *Workshop, Frascati, Italy, 23-27 March 2015*. Frascati, Italy.
820 <https://doi.org/10.5270/Fringe2015.pp116>

821 Grandin, R., Doin, M.-P., Bollinger, L., Pinel-Puysségur, B., Ducret, G., Jolivet, R., & Sapkota, S.
822 N. (2012). Long-term growth of the Himalaya inferred from interseismic InSAR
823 measurement. *Geology*, 40(12), 1059–1062. <https://doi.org/10.1130/G33154.1>

824 Gualandi, A., Serpelloni, E., & Belardinelli, M. E. (2014). Space–time evolution of crustal
825 deformation related to the Mw 6.3, 2009 L'Aquila earthquake (central Italy) from
826 principal component analysis inversion of GPS position time-series. *Geophysical*
827 *Journal International*, 197(1), 174–191. <https://doi.org/10.1093/gji/ggt522>

828 Harris, R. A. (2017). Large earthquakes and creeping faults. *Reviews of Geophysics*, 55(1),
829 2016RG000539. <https://doi.org/10.1002/2016RG000539>

830 Hetland, E. A., & Zhang, G. (2014). Effect of shear zones on post-seismic deformation with
831 application to the 1997 Mw 7.6 Manyi earthquake. *Geophysical Journal International*,
832 198(1), 259–269. <https://doi.org/10.1093/gji/ggu127>

833 Hirose, H., Asano, Y., Obara, K., Kimura, T., Matsuzawa, T., Tanaka, S., & Maeda, T. (2010). Slow
834 Earthquakes Linked Along Dip in the Nankai Subduction Zone. *Science*, 330(6010),
835 1502–1502. <https://doi.org/10.1126/science.1197102>

836 Huang, M.-H., Fielding, E. J., Liang, C., Milillo, P., Bekaert, D., Dreger, D., & Salzer, J. (2017).
837 Coseismic deformation and triggered landslides of the 2016 Mw 6.2 Amatrice

838 earthquake in Italy. *Geophysical Research Letters*.
839 <https://doi.org/10.1002/2016GL071687>

840 Hussain, E., Wright, T. J., Walters, R. J., Bekaert, D. P. S., Lloyd, R., & Hooper, A. (2018). Constant
841 strain accumulation rate between major earthquakes on the North Anatolian Fault.
842 *Nature Communications*, 9(1), 1392. <https://doi.org/10.1038/s41467-018-03739-2>

843 Johnson, K. M. (2006). Frictional Properties on the San Andreas Fault near Parkfield, California,
844 Inferred from Models of Afterslip following the 2004 Earthquake. *Bulletin of the*
845 *Seismological Society of America*, 96(4B), S321–S338.
846 <https://doi.org/10.1785/0120050808>

847 Jolivet, R., Simons, M., Agram, P. S., Duputel, Z., & Shen, Z.-K. (2015). Aseismic slip and
848 seismogenic coupling along the central San Andreas Fault. *Geophysical Research*
849 *Letters*, 42(2). <https://doi.org/10.1002/2014GL062222>

850 Kaneko, Y., Avouac, J.-P., & Lapusta, N. (2010). Towards inferring earthquake patterns from
851 geodetic observations of interseismic coupling. *Nature Geoscience*, 3(5), 363–369.
852 <https://doi.org/10.1038/ngeo843>

853 King. (1986). Speculations on the geometry of the initiation and termination processes of
854 earthquake rupture and its relation to morphology and geological structure. *Pure and*
855 *Applied Geophysics*, 124, 19. <https://doi.org/10.1007/BF00877216>

856 King, & Nabelek. (1985). Role of Fault Bends in the Initiation and Termination of Earthquake
857 Rupture. *Science*, 228(4702), 984–987. <https://doi.org/10.1126/science.228.4702.984>

858 Laske, G., Masters, G., Ma, Z., & Pasyanos, M. (2013). Update on CRUST1.0 - A 1-degree Global
859 Model of Earth's Crust, 15, EGU2013-2658. Presented at the EGU General Assembly
860 Conference Abstracts.

861 Liu, C., Zheng, Y., Xie, Z., & Xiong, X. (2017). Rupture features of the 2016 Mw 6.2 Norcia
862 earthquake and its possible relationship with strong seismic hazards. *Geophysical*
863 *Research Letters*, 44(3), 1320–1328. <https://doi.org/10.1002/2016GL071958>

864 López-Quiroz, P., Doin, M.-P., Tupin, F., Briole, P., & Nicolas, J.-M. (2009). Time series analysis of
865 Mexico City subsidence constrained by radar interferometry. *Journal of Applied*
866 *Geophysics*, 69(1), 1–15. <https://doi.org/10.1016/j.jappgeo.2009.02.006>

867 Marone, C. J., Scholtz, C. H., & Bilham, R. (1991). On the mechanics of earthquake afterslip.
 868 *Journal of Geophysical Research: Solid Earth*, 96(B5), 8441–8452.
 869 <https://doi.org/10.1029/91JB00275>

870 Maubant, L., Socquet, A., Hollingsworth, J., Pathier, E., & Pousse-Beltrán, L. (2017). The Seismic
 871 Sequence of the Norcia Earthquake, Italy 2016, seen by geodesy. In *Cargese, 2nd of*
 872 *October – 6th of October 2017*. France.

873 Moré, J. J. (1978). The Levenberg-Marquardt algorithm: Implementation and theory. In G. A.
 874 Watson (Ed.), *Numerical Analysis* (pp. 105–116). Springer Berlin Heidelberg.

875 Nielsen, S. B., & Knopoff, L. (1998). The equivalent strength of geometrical barriers to
 876 earthquakes. *Journal of Geophysical Research: Solid Earth*, 103(B5), 9953–9965.
 877 <https://doi.org/10.1029/97JB03293>

878 Noda, H., & Lapusta, N. (2013). Stable creeping fault segments can become destructive as a
 879 result of dynamic weakening. *Nature*, 493(7433), 518–521.
 880 <https://doi.org/10.1038/nature11703>

881 Papadopoulos, G. A., Ganas, A., Agalos, A., Papageorgiou, A., Triantafyllou, I., Kontoes, C., et
 882 al. (2017). Earthquake Triggering Inferred from Rupture Histories, DInSAR Ground
 883 Deformation and Stress-Transfer Modelling: The Case of Central Italy During August
 884 2016–January 2017. *Pure and Applied Geophysics*, 174(10), 3689–3711.
 885 <https://doi.org/10.1007/s00024-017-1609-8>

886 Pavlides, S., Chatzipetros, A., Papathanasiou, G., Georgiadis, G., Sboras, S., & Valkaniotis, S.
 887 (2017). Ground deformation and fault modeling of the 2016 sequence (24 Aug. – 30
 888 Oct.) in central Apennines (Central Italy). *Bulletin of the Geological Society of Greece*,
 889 51(0), 76–112. <https://doi.org/10.12681/bgsg.14334>

890 Perfettini, H., & Avouac, J.-P. (2004). Postseismic relaxation driven by brittle creep: A possible
 891 mechanism to reconcile geodetic measurements and the decay rate of aftershocks,
 892 application to the Chi-Chi earthquake, Taiwan. *Journal of Geophysical Research: Solid*
 893 *Earth*, 109(B2). <https://doi.org/10.1029/2003JB002488>

894 Perfettini, H., Avouac, J.-P., Tavera, H., Kositsky, A., Nocquet, J.-M., Bondoux, F., et al. (2010).
895 Seismic and aseismic slip on the Central Peru megathrust. *Nature*, 465(7294), 78–81.
896 <https://doi.org/10.1038/nature09062>

897 Perouse, E., Benedetti, L., Fleury, J., Rizza, M., Puliti, I., Billant, J., et al. (2018). Coseismic Slip
898 Vectors of 24 August and 30 October 2016 Earthquakes in Central Italy: Oblique Slip
899 and Regional Kinematic Implications. *Tectonics*, 37(10), 3760–3781.
900 <https://doi.org/10.1029/2018TC005083>

901 Petitta, M., Mastrorillo, L., Preziosi, E., Banzato, F., Barberio, M. D., Billi, A., et al. (2018). Water-
902 table and discharge changes associated with the 2016–2017 seismic sequence in
903 central Italy: hydrogeological data and a conceptual model for fractured carbonate
904 aquifers. *Hydrogeology Journal*, 1–18. <https://doi.org/10.1007/s10040-017-1717-7>

905 Pinel-Puységur, B., Michel, R., & Avouac, J.-P. (2012). Multi-link InSAR time series:
906 Enhancement of a wrapped interferometric database. *IEEE Journal of Selected Topics*
907 *in Applied Earth Observations and Remote Sensing*, 5(3), 784–794.
908 <https://doi.org/10.1109/JSTARS.2012.2196758>

909 Pino, N. A., Convertito, V., & Madariaga, R. (2019). Clock advance and magnitude limitation
910 through fault interaction: the case of the 2016 central Italy earthquake sequence.
911 *Scientific Reports*, 9(1), 5005. <https://doi.org/10.1038/s41598-019-41453-1>

912 Pizzi, A., Di Domenica, A., Gallovič, F., Luzi, L., & Puglia, R. (2017). Fault segmentation as
913 constraint to the occurrence of the main shocks of the 2016 Central Italy seismic
914 sequence. *Tectonics*, 2017TC004652. <https://doi.org/10.1002/2017TC004652>

915 Pluymakers, A. M. H., Niemeijer, A. R., & Spiers, C. J. (2016). Frictional properties of simulated
916 anhydrite-dolomite fault gouge and implications for seismogenic potential. *Journal of*
917 *Structural Geology*, 84, 31–46. <https://doi.org/10.1016/j.jsg.2015.11.008>

918 Pollitz, F. F., Wicks, C., & Thatcher, W. (2001). Mantle Flow Beneath a Continental Strike-Slip
919 Fault: Postseismic Deformation After the 1999 Hector Mine Earthquake. *Science*,
920 293(5536), 1814–1818. <https://doi.org/10.1126/science.1061361>

921 Porreca, M., Minelli, G., Ercoli, M., Brobia, A., Mancinelli, P., Cruciani, F., et al. (2018). Seismic
922 reflection profiles and subsurface geology of the area interested by the 2016-2017

923 earthquake sequence (Central Italy). *Tectonics*.
 924 <https://doi.org/10.1002/2017TC004915>

925 Puliti, I., Pizzi, A., Benedetti, L., Domenica, A. D., & Fleury, J. (2020). Comparing slip distribution
 926 of an active fault system at various timescales: insights for the evolution of the Mt.
 927 Vettore- Mt. Bove fault system in Central Apennines. *Tectonics*, *n/a*(*n/a*),
 928 e2020TC006200. <https://doi.org/10.1029/2020TC006200>

929 Rabus, B., Eineder, M., Roth, A., & Bamler, R. (2003). The shuttle radar topography mission—a
 930 new class of digital elevation models acquired by spaceborne radar. *ISPRS Journal of*
 931 *Photogrammetry and Remote Sensing*, *57*(4), 241–262.
 932 [https://doi.org/10.1016/S0924-2716\(02\)00124-7](https://doi.org/10.1016/S0924-2716(02)00124-7)

933 Ragon, T., Sladen, A., & Simons, M. (2019). Accounting for uncertain fault geometry in
 934 earthquake source inversions – II: application to the Mw 6.2 Amatrice earthquake,
 935 Central Italy. *Geophysical Journal International*. <https://doi.org/10.1093/gji/ggz180>

936 Rice, J. R. (2006). Heating and weakening of faults during earthquake slip. *Journal of*
 937 *Geophysical Research: Solid Earth*, *111*, n/a-n/a.
 938 <https://doi.org/10.1029/2005jb004006>

939 Riva, R. E. M., Borghi, A., Aoudia, A., Barzaghi, R., Sabadini, R., & Panza, G. F. (2007). Viscoelastic
 940 relaxation and long-lasting after-slip following the 1997 Umbria-Marche (Central Italy)
 941 earthquakes. *Geophysical Journal International*, *169*(2), 534–546.
 942 <https://doi.org/10.1111/j.1365-246X.2007.03315.x>

943 Roeloffs, E. (1996). Poroelastic Techniques in the Study of Earthquake-Related Hydrologic
 944 Phenomena. In *Advances in Geophysics* (Vol. 37, pp. 135–195). Elsevier.
 945 [https://doi.org/10.1016/S0065-2687\(08\)60270-8](https://doi.org/10.1016/S0065-2687(08)60270-8)

946 Rousset, B., Campillo, M., Lasserre, C., Frank, W. B., Cotte, N., Walpersdorf, A., et al. (2017). A
 947 geodetic matched filter search for slow slip with application to the Mexico subduction
 948 zone: GEODETIC MATCHED FILTER FOR SLOW SLIP. *Journal of Geophysical Research:*
 949 *Solid Earth*, *122*(12), 10,498–10,514. <https://doi.org/10.1002/2017JB014448>

950 Ruina, A. (1983). Slip instability and state variable friction laws. *Journal of Geophysical*
 951 *Research: Solid Earth*, 88(B12), 10359–10370.
 952 <https://doi.org/10.1029/JB088iB12p10359>
 953 Scholz, C. H. (1998). Earthquakes and friction laws. *Nature*, 391(6662), 37–42.
 954 <https://doi.org/10.1038/34097>
 955 Scognamiglio, L., Tinti, E., Casarotti, E., Pucci, S., Villani, F., Cocco, M., et al. (2018). Complex
 956 fault geometry and rupture dynamics of the Mw 6.5, 2016, October 30th central Italy
 957 earthquake. *Journal of Geophysical Research: Solid Earth*.
 958 <https://doi.org/10.1002/2018JB015603>
 959 Scuderi, M. M., & Collettini, C. (2016). The role of fluid pressure in induced vs. triggered
 960 seismicity: insights from rock deformation experiments on carbonates. *Scientific*
 961 *Reports*, 6(1), 24852. <https://doi.org/10.1038/srep24852>
 962 Silverii, F., D'Agostino, N., Métois, M., Fiorillo, F., & Ventafridda, G. (2016). Transient
 963 deformation of karst aquifers due to seasonal and multiyear groundwater variations
 964 observed by GPS in southern Apennines (Italy). *Journal of Geophysical Research: Solid*
 965 *Earth*, 121(11), 8315–8337. <https://doi.org/10.1002/2016JB013361>
 966 Smeraglia, L., Billi, A., Carminati, E., Cavallo, A., & Doglioni, C. (2017). Field- to nano-scale
 967 evidence for weakening mechanisms along the fault of the 2016 Amatrice and Norcia
 968 earthquakes, Italy. *Tectonophysics*. <https://doi.org/10.1016/j.tecto.2017.05.014>
 969 Tarantola, A. (2005). *Inverse Problem Theory and Methods for Model Parameter Estimation*.
 970 Society for Industrial and Applied Mathematics.
 971 <https://doi.org/10.1137/1.9780898717921>
 972 Tesson, J., Pace, B., Benedetti, L., Visini, F., Delli Rocoli, M., Arnold, M., et al. (2016). Seismic
 973 slip history of the Pizzalto fault (central Apennines, Italy) using in situ-produced ³⁶Cl
 974 cosmic ray exposure dating and rare earth element concentrations. *Journal of*
 975 *Geophysical Research: Solid Earth*, 121(3), 2015JB012565.
 976 <https://doi.org/10.1002/2015JB012565>
 977 Thomas, M. Y., Avouac, J.-P., & Lapusta, N. (2017). Rate-and-state friction properties of the
 978 Longitudinal Valley Fault from kinematic and dynamic modeling of seismic and aseismic

979 slip. *Journal of Geophysical Research: Solid Earth*, 2016JB013615.
 980 <https://doi.org/10.1002/2016JB013615>

981 Thompson, G. A., & Parsons, T. (2016). Vertical deformation associated with normal fault
 982 systems evolved over coseismic, postseismic, and multiseismic periods. *Journal of*
 983 *Geophysical Research: Solid Earth*, 121(3), 2153–2173.
 984 <https://doi.org/10.1002/2015JB012240>

985 Tung, S., & Masterlark, T. (2018). Delayed poroelastic triggering of the 2016 October Visso
 986 earthquake by the August Amatrice earthquake, Italy. *Geophysical Research Letters*.
 987 <https://doi.org/10.1002/2017GL076453>

988 Valigi, D., Mastroiello, L., Cardellini, C., Checcucci, R., Di Matteo, L., Frondini, F., et al. (2019).
 989 Springs discharge variations induced by strong earthquakes: the Mw 6.5 Norcia event
 990 (Italy, October 30th 2016). *Rendiconti Online Della Società Geologica Italiana*, 47, 141–
 991 146. <https://doi.org/10.3301/ROL.2019.25>

992 Verdecchia, A., Pace, B., Visini, F., Scotti, O., Peruzza, L., & Benedetti, L. (2018). The Role of
 993 Viscoelastic Stress Transfer in Long-Term Earthquake Cascades: Insights After the
 994 Central Italy 2016–2017 Seismic Sequence. *Tectonics*, 37(10), 3411–3428.
 995 <https://doi.org/10.1029/2018TC005110>

996 Villani, F., Civico, R., Pucci, S., Pizzimenti, L., Nappi, R., De Martini, P. M., et al. (2018). A
 997 database of the coseismic effects following the 30 October 2016 Norcia earthquake in
 998 Central Italy. *Scientific Data*, 5, 180049. <https://doi.org/10.1038/sdata.2018.49>

999 Villani, F., Pucci, S., Civico, R., Martini, P. M. D., Cinti, F. R., & Pantosti, D. (2018). Surface faulting
 1000 of the 30 October 2016 Mw 6.5 central Italy earthquake: detailed analysis of a complex
 1001 coseismic rupture. *Tectonics*, 0(ja). <https://doi.org/10.1029/2018TC005175>

1002 Walters, R. J., Gregory, L. C., Wedmore, L. N. J., Craig, T. J., McCaffrey, K., Wilkinson, M., et al.
 1003 (2018). Dual control of fault intersections on stop-start rupture in the 2016 Central Italy
 1004 seismic sequence. *Earth and Planetary Science Letters*, 500, 1–14.
 1005 <https://doi.org/10.1016/j.epsl.2018.07.043>

1006 Wang, L., Gao, H., Feng, G., & Xu, W. (2018). Source parameters and triggering links of the
 1007 earthquake sequence in central Italy from 2009 to 2016 analyzed with GPS and InSAR
 1008 data. *Tectonophysics*. <https://doi.org/10.1016/j.tecto.2018.07.013>

1009 Wesnousky, S. G. (1988). Seismological and structural evolution of strike-slip faults. *Nature*,
 1010 335(6188), 340. <https://doi.org/10.1038/335340a0>

1011 Xu, G., Xu, C., Wen, Y., & Jiang, G. (2017). Source Parameters of the 2016–2017 Central Italy
 1012 Earthquake Sequence from the Sentinel-1, ALOS-2 and GPS Data. *Remote Sensing*,
 1013 9(11), 1182. <https://doi.org/10.3390/rs9111182>

1014 Zhao, B., Bürgmann, R., Wang, D., Tan, K., Du, R., & Zhang, R. (2017). Dominant Controls of
 1015 Downdip Afterslip and Viscous Relaxation on the Postseismic Displacements Following
 1016 the Mw7.9 Gorkha, Nepal, Earthquake. *Journal of Geophysical Research: Solid Earth*,
 1017 122(10), 8376–8401. <https://doi.org/10.1002/2017JB014366>

1018 Zhou, Y., Thomas, M. Y., Parsons, B., & Walker, R. T. (2018). Time-dependent postseismic slip
 1019 following the 1978 Mw 7.3 Tabas-e-Golshan, Iran earthquake revealed by over 20 years
 1020 of ESA InSAR observations. *Earth and Planetary Science Letters*, 483, 64–75.
 1021 <https://doi.org/10.1016/j.epsl.2017.12.005>
 1022

1023 11 References for supplementary materials

1024 Blewitt, G, WC Hammond, and C Kreemer. 2018. “Harnessing the GPS Data Explosion for
 1025 Interdisciplinary Science.” *Eos* 99. <https://doi.org/10.1029/2018EO104623>.

1026 Boni, CARLO FELICE, CLAUDIA TARRAGONI, LUCIO MARTARELLI, and SIMONA PIERDOMINICI.
 1027 2010. “STUDIO IDROGEOLOGICO NEL SETTORE NORD-OCCIDENTALE DEI MONTI
 1028 SIBILLINI: UN CONTRIBUTO ALLA CARTOGRAFIA IDROGEOLOGICA UFFICIALE.” *Italian*
 1029 *Journal of Engineering Geology and Environment* 2: 16.
 1030 <https://doi.org/10.4408/IJEGE.2010-02.O-02>.

1031 Cheloni, D., E. Falcucci, and S. Gori. 2019. “Half-Graben Rupture Geometry of the 30 October
 1032 2016 MW 6.6 Mt. Vettore-Mt. Bove Earthquake, Central Italy.” *Journal of Geophysical*
 1033 *Research: Solid Earth* 0 (ja). <https://doi.org/10.1029/2018JB015851>.

Chiaraluce, L., R. Di Stefano, E. Tinti, L. Scognamiglio, M. Michele, E. Casarotti, M. Cattaneo, et al. 2017. "The 2016 Central Italy Seismic Sequence: A First Look at the Mainshocks, Aftershocks, and Source Models." *Seismological Research Letters* 88 (3): 757–71. <https://doi.org/10.1785/0220160221>.

Chiles, Jean-Paul, and Pierre Delfiner. 2009. *Geostatistics: Modeling Spatial Uncertainty*. Vol. 497. John Wiley & Sons.

Grandin, R. 2009. *L'Apport de La Géodésie Spatiale Dans La Compréhension Du Processus de Rifting Magmatique : L'exemple de l'épisode En Cours En Afar Ethiopien (2005-2009)*. Institut de physique du globe (Paris). <http://www.theses.fr/2009GLOB0013>.

Hansen, P. 1992. "Analysis of Discrete Ill-Posed Problems by Means of the L-Curve." *SIAM Review* 34 (4): 561–80. <https://doi.org/10.1137/1034115>.

Improta, Luigi, Diana Latorre, Lucia Margheriti, Anna Nardi, Alessandro Marchetti, Anna Maria Lombardi, Barbara Castello, et al. 2019. "Multi-Segment Rupture of the 2016 Amatrice-Visso-Norcia Seismic Sequence (Central Italy) Constrained by the First High-Quality Catalog of Early Aftershocks." *Scientific Reports* 9 (1): 6921. <https://doi.org/10.1038/s41598-019-43393-2>.

Jolivet, R., C. Lasserre, M.-P. Doin, S. Guillaso, G. Peltzer, R. Dailu, J. Sun, Z.-K. Shen, and X. Xu. 2012. "Shallow Creep on the Haiyuan Fault (Gansu, China) Revealed by SAR Interferometry." *Journal of Geophysical Research: Solid Earth* 117 (B6): B06401. <https://doi.org/10.1029/2011JB008732>.

Jolivet, R., M. Simons, P. S. Agram, Z. Duputel, and Z.-K. Shen. 2015. "Aseismic Slip and Seismogenic Coupling along the Central San Andreas Fault." *Geophysical Research Letters* 42 (2). <https://doi.org/10.1002/2014GL062222>.

Lohman, Rowena B., and Mark Simons. 2005. "Some Thoughts on the Use of InSAR Data to Constrain Models of Surface Deformation: Noise Structure and Data Downsampling." *Geochemistry, Geophysics, Geosystems* 6 (1): Q01007. <https://doi.org/10.1029/2004GC000841>.

Maubant, Louise, Anne Socquet, James Hollingsworth, Erwan Pathier, and Pousse Lea. 2017. "The Seismic Sequence of the Norcia Earthquake, Italy 2016, Seen by Geodesy." In

1063 *Colloque G2 2017 Géodésie - Rhéologie, 13-15 Nov. 2017 Nice. France.* <https://g2->
1064 [2017.sciencesconf.org/program](https://g2-2017.sciencesconf.org/program).

1065 Petitta, Marco, Lucia Mastrorillo, Elisabetta Preziosi, Francesca Banzato, Marino Domenico
1066 Barberio, Andrea Billi, Costanza Cambi, et al. 2018. "Water-Table and Discharge
1067 Changes Associated with the 2016–2017 Seismic Sequence in Central Italy:
1068 Hydrogeological Data and a Conceptual Model for Fractured Carbonate Aquifers."
1069 *Hydrogeology Journal*, January, 1–18. <https://doi.org/10.1007/s10040-017-1717-7>.

1070 Pierantoni, Pietro, Giovanni Deiana, and Sandro Galdenzi. 2013. "Stratigraphic and Structural
1071 Features of the Sibillini Mountains (Umbria-Marche Apennines, Italy)." *Italian Journal*
1072 *of Geosciences* 132 (3): 497–520. <https://doi.org/10.3301/IJG.2013.08>.

1073 Puliti, I., A. Pizzi, L. Benedetti, A. Di Domenica, and J. Fleury. 2020. "Comparing Slip Distribution
1074 of an Active Fault System at Various Timescales: Insights for the Evolution of the Mt.
1075 Vettore- Mt. Bove Fault System in Central Apennines." *Tectonics* n/a (n/a):
1076 e2020TC006200. <https://doi.org/10.1029/2020TC006200>.

1077 Radiguet, M., F. Cotton, M. Vergnolle, M. Campillo, B. Valette, V. Kostoglodov, and N. Cotte.
1078 2011. "Spatial and Temporal Evolution of a Long Term Slow Slip Event: The 2006
1079 Guerrero Slow Slip Event." *Geophysical Journal International* 184 (2): 816–28.
1080 <https://doi.org/10.1111/j.1365-246X.2010.04866.x>.

1081 Ragon, Théa, Anthony Sladen, and Mark Simons. 2019. "Accounting for Uncertain Fault
1082 Geometry in Earthquake Source Inversions – II: Application to the Mw 6.2 Amatrice
1083 Earthquake, Central Italy." *Geophysical Journal International*.
1084 <https://doi.org/10.1093/gji/ggz180>.

1085 Scognamiglio, L., E. Tinti, E. Casarotti, S. Pucci, F. Villani, M. Cocco, F. Magnoni, A. Michelini,
1086 and D. Dreger. 2018. "Complex Fault Geometry and Rupture Dynamics of the Mw 6.5,
1087 2016, October 30th Central Italy Earthquake." *Journal of Geophysical Research: Solid*
1088 *Earth*. <https://doi.org/10.1002/2018JB015603>.

1089 Sudhaus, Henriette, and Sigurjón Jonsson. 2009. "Improved Source Modelling through
1090 Combined Use of InSAR and GPS under Consideration of Correlated Data Errors:

1091 Application to the June 2000 Kleifarvatn Earthquake, Iceland.” *Geophysical Journal*
 1092 *International* 176 (2): 389–404. <https://doi.org/10.1111/j.1365-246X.2008.03989.x>.
 1093 Tarantola, Albert. 2005. *Inverse Problem Theory and Methods for Model Parameter Estimation*.
 1094 Society for Industrial and Applied Mathematics.
 1095 <https://doi.org/10.1137/1.9780898717921>.
 1096 Tarantola, Albert, and Bernard Valette. 1982. “Generalized Nonlinear Inverse Problems Solved
 1097 Using the Least Squares Criterion.” *Reviews of Geophysics* 20 (2): 219.
 1098 <https://doi.org/10.1029/RG020i002p00219>.
 1099 Tarquini, Simone, Stefano Vinci, Massimiliano Favalli, Fawzi Doumaz, Alessandro Fornaciai, and
 1100 Luca Nannipieri. 2012. “Release of a 10-m-Resolution DEM for the Italian Territory:
 1101 Comparison with Global-Coverage DEMs and Anaglyph-Mode Exploration via the
 1102 Web.” *Computers & Geosciences* 38 (1): 168–70.
 1103 <https://doi.org/10.1016/j.cageo.2011.04.018>.
 1104 Valigi, Daniela, Lucia Mastroiello, Carlo Cardellini, Roberto Checcucci, Lucio Di Matteo,
 1105 Francesco Frondini, Francesco Mirabella, Stefano Viaroli, and Isotta Vispi. 2019.
 1106 “Springs Discharge Variations Induced by Strong Earthquakes: The Mw 6.5 Norcia Event
 1107 (Italy, October 30th 2016).” *Rendiconti Online Della Società Geologica Italiana* 47
 1108 (March): 141–46. <https://doi.org/10.3301/ROL.2019.25>.
 1109 Villani, Fabio, Riccardo Civico, Stefano Pucci, Luca Pizzimenti, Rosa Nappi, Paolo Marco De
 1110 Martini, the Open EMERGEO Working Group, et al. 2018. “A Database of the Coseismic
 1111 Effects Following the 30 October 2016 Norcia Earthquake in Central Italy.” *Scientific*
 1112 *Data* 5 (March): 180049. <https://doi.org/10.1038/sdata.2018.49>.
 1113 Wackernagel, Hans. 2003. *Multivariate Geostatistics: An Introduction with Applications*. 3rd
 1114 ed. Berlin Heidelberg: Springer-Verlag. <https://doi.org/10.1007/978-3-662-05294-5>.
 1115 Wright, Tim J., Barry E. Parsons, and Zhong Lu. 2004. “Toward Mapping Surface Deformation
 1116 in Three Dimensions Using InSAR.” *Geophysical Research Letters* 31 (1).
 1117 <https://doi.org/10.1029/2003GL018827>.
 1118