**Impacts of copper and lead exposure on prokaryotic communities
from contaminated contrasted coastal seawaters: the influence of
previous metal exposure**

Clément Coclet^{1,2}, Cédric Garnier¹, Gaël Durrieu¹, Sébastien D'onofrio¹, Nicolas
Layglon¹, Jean-François Briand², Benjamin Misson^{1,*}

¹ *Université de Toulon, Aix Marseille Université, CNRS, IRD, Mediterranean Institute
of Oceanography (MIO), UM110, France.*

² *Université de Toulon, MAPIEM, EA 4323, Toulon, France.*

*: corresponding author; email: mission@univ-tln.fr; tel: +33 494 146 746; Université
de Toulon, CS60584, 83041 Toulon Cedex 9

Running title: Impacts of copper on prokaryotic communities depends on metal
exposure history

17 **ABSTRACT**

18 Our understanding of environmental factors controlling prokaryotic community is
19 largely hampered by the large environmental variability across spatial scales (e.g.
20 trace metal contamination, nutrient enrichment, physico-chemical variations) and the
21 broad diversity of bacterial pre-exposure to environmental factors. In this article, we
22 investigated the specific influence of copper (Cu) and lead (Pb) on prokaryotic
23 communities from the uncontaminated site, using mesocosm experiments. In
24 addition, we studied how pre-exposure (i.e. life history) affects communities, with
25 reference to previous metal exposure on the response of three prokaryotic
26 communities to similar Cu exposition. This study showed a stronger influence of Cu
27 contamination than Pb contamination on prokaryotic diversity and structure. We
28 identified 12, and 34 bacterial families and genera, respectively, contributing to the
29 significant differences observed in community structure between control and spiked
30 conditions. Taken altogether, our results point towards a combination of direct
31 negative responses to Cu contamination and indirect responses mediated by
32 interaction with phytoplankton. These identified responses were largely conditioned
33 by the previous exposure of community to contaminants.

34

35 Keywords: prokaryotic communities; copper contamination; metal exposure history;
36 coastal seawater; mesocosm experiment; high throughput sequencing

37

38 INTRODUCTION

39 Marine coastal environments are especially vulnerable to anthropogenic pressures
40 such as chemical contamination (Halpern *et al.* 2008; The Mermex Group *et al.*
41 2011). For example, copper (Cu) contamination in coastal environments is mainly
42 caused by urban and industrial wastes (Levin *et al.* 2001; Oursel *et al.* 2013), or
43 antifouling coatings close to harbors (Turner 2010). Lead (Pb) contamination is
44 mainly associated with historical short-term events (Xu *et al.* 2014; Dang *et al.* 2015a,
45 2015b) or historical fuel consumption for nautical traffic (Callender 2003). Cu
46 contamination is mainly encountered in seawaters of harbor areas while Pb
47 contamination are found in sediments, probably due to past events (Dang *et al.*
48 2015b). In Toulon Bay, Cu and Pb concentrations can be enriched up to more than
49 100-fold above geochemical background levels depending on the compartment (e.g.
50 sediment or water column) (Casas *et al.* 2008; Tessier *et al.* 2011; Jean *et al.* 2012;
51 Cossa *et al.* 2014; Dang *et al.* 2015b; Twining *et al.* 2015; Coclet *et al.* 2018, 2019).
52 Thus, the unique feature of Toulon Bay relates to a gradient of multiple trace metal
53 elements (TME) with very low concentrations in open parts comparable to levels
54 measured in open Mediterranean Sea (Morley *et al.* 1997). By comparison, most
55 enclosed and anthropized areas exhibit very high TME concentrations (Dang *et al.*
56 2015a; Coclet *et al.* 2018, 2019). Therefore, Toulon bay is a model ecosystem for the
57 study of ecological consequences of TME contamination in the marine coastal
58 environment.

59 In Toulon Bay where TME gradients are wider than classical physico-chemical
60 gradients, TME contamination have been proposed to influence directly the

bacterioplankton diversity and structure (Coclet *et al.* 2019). Several field studies clearly suggested TMEs as structuring factors of abundance, diversity and composition of bacterioplanktonic communities by selecting metal resistant taxa (Sheeba *et al.* 2017; Coclet *et al.* 2018, 2019; Goni-Urriza *et al.* 2018). A microcosm-based study revealed that artificial mixture of contaminants (nutrients, metals, and herbicides) induced a strong inhibition of growth, production, respiration and significant modifications of the functional patterns of bacterioplankton community (Pringault *et al.* 2016). Furthermore, both industrial effluent discharge and the resuspension of metal-contaminated marine sediments have been shown to affect microbial diversity in a resuspension experiment (Zouch *et al.* 2018). TME contamination typically found in coastal environments have also been observed to affect phytoplanktonic communities (Mackey *et al.* 2012; Lafabrie *et al.* 2013; Coclet *et al.* 2018), which can have indirect consequences for bacterioplankton community (Goni-Urriza *et al.* 2018; Coclet *et al.* 2019).

Life history of a microbial community can be defined as its past exposure to a multitude of environmental changes, like top-down (e.g. grazing pressure, exposition to toxic substances) as well as bottom-up (e.g. phytoplanktonic production) control. The pre-exposure of communities to harmful contaminants like TME can affect the microbial ecology of the marine system and therefore is of major concern (Hughes-Martiny *et al.* 2006; Bissett *et al.* 2010; Bell *et al.* 2013; Sjöstedt *et al.* 2018; Ward *et al.* 2019). However, most studies exploring patterns in microbial community diversity and structure in response to environmental change have focused on direct effects, without consideration of life history. Previous TME exposure can lead to an increase

of metal tolerance in microbial communities by the elimination of the most sensitive individuals and the replacement of sensitive species by tolerant ones (Gustavson *et al.* 1999). Serra *et al.* (2009) found that copper pulse exposure may be toxic to periphyton communities while continuous copper exposure may lead to community adaptation, which is often related to changes in species composition. All of these studies are in agreement with the pollution-induced community tolerance (PICT) model (Blanck, Wangberg and Molander 1988; Blanck 2002). However, while numerous studies developed the PICT approach based on photosynthetic efficiency, extracellular enzyme activities or respiration activity (Carman, Fleeger and Pomarico 2000; Soldo *et al.* 2005; Serra, Corcoll and Guasch 2009; Tlili 2013), little is known about how metal exposure impacts diversity and composition of marine prokaryotic communities previously exposed to similar contamination.

Assuming that the life history of its planktonic communities in Toulon Bay would mainly reflect TME exposure history due to the strong and chronic Cu and Pb contamination gradients of the area (Layglon *et al.*, under revision), in this study we aimed to answer two main questions: (1) to what extent do Cu and Pb contamination affect prokaryotic communities from uncontaminated site at relevant concentrations; and (2) does prokaryotic TME exposure history play a role in shaping prokaryotic diversity after a Cu exposure?

MATERIALS AND METHODS

Study area, seawater collection, processing, and storage

106 The study was conducted in Toulon Bay, located in the northwestern part of the
107 Mediterranean Sea (Figure 1 and Supplementary Material). Seawater used for the
108 first mesocosm experiment was collected at 1-m depth from site 41p (low
109 contaminated site) (Figure 1) on May 10th 2016 and seawater used for the second
110 mesocosm experiment was collected at 1-m depth from both sites LAZ (intermediate
111 contaminated site) and 6ext (highly contaminated site) on May 2nd 2017 (Figure 1).
112 The 3 sampling seawaters used for mesocosm experiments were geochemically
113 contrasted. They were representative of the North-West to South-East decreasing
114 gradient of trace metal contamination (mainly Cu and Pb) previously observed in
115 Toulon Bay (Figure 1). Minimal concentrations were recorded in the open-sea
116 sampling site (41p), while maximal concentrations were recorded in the northern and
117 enclosed part of the little bay (6ext) where the TME contamination is chronic (Coclet
118 et al., 2019, 2017; Layglon et al., under revision). Details on sample's information are
119 given in the supplementary Table 1. Seawater was collected using an air-
120 compressed Teflon pump (AstiPure™ II, Saint-Gobin) connected with Teflon tubing. It
121 was filtered on site through a 5-10 µm cellulose Polycap filter (Whatman) connected
122 to the Teflon tubing after the pump in order to remove the largest grazers. Filtered
123 seawater was collected in 50-L low-density polyethylene (LDPE) tanks (Nalgene®).
124 Back to the lab, each mesocosm was filled with 20-L of filtered seawater. The
125 remaining seawater was filtered through 0.2 µm Teflon Polycap filters (Whatman) to
126 eliminate all organisms, then stored at 4°C and subsequently used to re-fill
127 mesocosms during the experiment. All the sampling material, bottles, tanks, and
128 filters were pre-cleaned following rigorous protocols, detailed in supplementary

information and in Coclet *et al.* (2018), in order to keep metal concentrations as close as possible to their initial levels.

Mesocosm setup, and experimental design

For both mesocosm studies, twelve 20-L LDPE tanks (Nalgene®) were enclosed in a large box covered of plastics in order to avoid contamination in TMEs from ambient particles. Fresh air was continuously provided inside the box by a filtered-equipped fan in order to avoid humidity accumulation and to equilibrate temperature with the surrounding room. The tanks were filled with 5-10 µm filtered water, and thus included a natural marine microbial community mainly composed of free-living organisms. Seawater in each tank was continuously mixed with a Teflon magnetic stir bar used at low speed and oxygenized with 0.2 µm-filtered and humidified ambient air. The 12 tanks were submitted to a daily light/dark cycle of 14/10h, using full solar spectrum aquariophilly neon tubes and a surface light irradiance of 50 µmol.m².s¹ of white light. In the first year, the temperature of the room and of the mesocosms' box were kept constant at 19 ± 1.6°C, and 23 ± 1.5°C, respectively. In the second year, the room temperature and the mesocosm interior were kept constant at 19 ± 2.2°C and, 22 ± 2.0°C, respectively. To maintain resource availability during the experiments, each tank had a continuous opened water circulation system, using a peristaltic pump (Minipuls Evolution®, Gilson), to renew 50% of the total water volume in a week by 0.2 µm filtered seawater sampled at the same site.

The first mesocosm experiment was performed from May 10th to June 17th 2016 over a 6-week period. Twelve pre-cleaned 20-L LDPE (Nalgene®) tanks were filled with 5-10 µm filtered water from site 41p (Figure 1). A 2-week acclimation period

was carried out to establish relatively stable conditions in each tank. After the acclimation period, seawater from the 12 tanks was pooled, homogenized and then tanks were re-filled. Following the acclimation period, seawater in tanks from the first mesocosm study were assigned to four treatments: (1) site control (41p CTRL, i.e. no TME added to 41p seawater), (2) Cu addition (150 nM) (41p Cu), Pb addition (7 nM) (41p Pb), and (4) Cu and Pb addition (150 nM + 7 nM, respectively) (41p Cu+Pb) (Supplementary Figure 1). The Cu treatments were spiked with a stock solution of Cu nitrate (10^{-3} M) and the Pb treatments were spiked with a stock solution of Pb nitrate (10^{-3} M). According to the initial Cu and Pb concentrations measured in the 41p seawater used to fill the tanks, final concentrations of 150 nM and 7 nM for Cu and Pb respectively, were targeted. The water supplies were simultaneously spiked with similar concentrations of Cu, Pb, or both depending of treatment to maintain metal concentrations in tanks.

The second mesocosm was performed from May 2nd to June 12th 2017 over a 6-week period. Six 20-L LDPE (Nalgene®) tanks were filled with 5-10 μ m filtered water from each sampling site (LAZ and 6ext sites) (Figure 1). The treatments in the second mesocosm experiment were (1) site control (LAZ CTRL, i.e. no TME added to LAZ seawater), (2) Cu addition to LAZ seawater (150 nM) (LAZ Cu), (3) site control (6ext CTRL, i.e. no TME added to 6ext seawater), and (4) Cu addition to 6ext seawater (150 nM) (6ext Cu) (Supplementary Figure 2). The Cu treatments were spiked with a stock solution of Cu nitrate (10^{-3} M). According to the initial Cu concentrations measured in LAZ and 6ext seawaters used to fill the tanks, final concentration of 150 nM for Cu were targeted. The water supplies were simultaneous spiked with Cu depending of treatment to maintain metal concentrations in tanks.

177

178 **Experimental time plan and sampling**

179 After the acclimation period, seawater was sampled from the 12 tanks using a
180 peristaltic pump connected with Teflon tubing at different times over the course of the
181 first mesocosm experiment (0, 3, 7, 10, 14, 17, 21, and 25 days) after the Cu/Pb
182 spikes, and the second mesocosm experiment (0, 3, 7, 10, 14, 17, 21, 25 and 28
183 days) after the Cu spikes. At each sampling time, 333 mL of seawater was sampled
184 in each replicate of a given experimental condition and pooled in 1L FEP bottles
185 (Nalgene®) to obtain composite samples for each condition. It was then immediately
186 resampled for future trace metals (TM), dissolved organic carbon (DOC), and total
187 nitrogen (TN) quantifications as previously described (Omanović *et al.* 2006; Cindrić
188 *et al.* 2015), and fully explained in Supplementary Material. Additionally, seawater
189 (10-mL) for flow cytometry analysis were directly collected in 15-mL centrifuge tubes
190 (Falcon) from each of 12 tanks, fixed with 0.25% (final concentration) glutaraldehyde,
191 and stored at -80°C until analysis. Finally, at 0, 7, 14, 21, and 25 days, 1-L of
192 seawater from each of 12 tanks was collected in 1-L HDPE bottles and filtered
193 through 0.2 µm polycarbonate membranes (Millipore). Filters were stored at -80°C
194 until DNA extraction.

195

196 **Seawater physico-chemical measurements and analyses**

197 Temperature (°C), pH, dissolved oxygen concentration (mg.L⁻¹) and saturation (%)
198 were directly monitored daily into tanks, over the course of both mesocosm
199 experiments, using LDO10105 (Hach) multiparameter probe. Differential pulse
200 anodic stripping voltammetry (DPASV) was used to measure Cd, Cu, Pb, and Zn

dissolved concentrations. Details about instruments, analysis conditions, and data treatment are provided in Supplementary material and elsewhere (Oursel *et al.* 2013; Cindrić *et al.* 2015). DOC and TN concentrations were determined using a TOC-V_{CSH} analyzer (Shimadzu) (further details in Dang *et al.*, 2014; Coclet *et al.*, 2018, 2019).

Flow cytometry analysis

Autotrophic prokaryotes (*Synechococcus*-like, SYN), photosynthetic picoeukaryotes (PPE) and photosynthetic nanoeukaryotes (PNE) populations were characterized and enumerated using a BD Accury™ C6 (BD Biosciences) flow cytometer, as previously described (Coclet *et al.* 2018, 2019). Heterotrophic prokaryotes were enumerated after staining with SYBR Green as previously described (Cabrol, Quéméneur and Misson 2017).

DNA extraction, PCR amplification, and sequencing of 16S rRNA gene amplicons

DNA was extracted from the polycarbonate filters by a combination of enzymatic cell lysis (Ghiglione, Conan and Pujo-Pay 2009) and AllPrep DNA/RNA Mini Kit (QIAGEN) according to the manufacturer's instructions. Prokaryotic community structure was assessed by targeting the V4-V5 region of the 16S rRNA gene (Parada, Needham and Fuhrman 2016) and using Illumina Miseq 2 x 250pb paired-end sequencing (Genoscreen, France). The protocol for the DNA extraction and the library preparation is fully described in Coclet *et al.* (2019).

Bioinformatic analysis

Sequences were demultiplexed and assigned to corresponding samples using CASAVA (Illumina). Bioinformatic analysis were performed on sequences from two sequencing MiSeq runs in the same sequencing company were used for our two mesocosm experiments. The analysis of the raw sequences was done by following the standard pipeline of DADA2 (Callahan *et al.* 2016) in RStudio (R Core Team 2017) with the following parameters: maxN= 0, maxEE= c(2,2), truncQ= 2. Briefly, the package includes the following steps: filtering, dereplication, sample inference, chimera identification, and merging of paired-end reads. DADA2 infers exact amplicon sequence variants (ASVs) from sequencing data, instead of building operational taxonomic units from sequence similarity. The taxonomy assignments were done with the SILVA v.128 database (Pruesse *et al.* 2007; Quast *et al.* 2013) and the “assignTaxonomy” function in DADA2. For some ASVs, in order to obtain a finer taxonomical resolution, DADA2 package implements a method to make species level assignments based on exact matching between ASVs and sequenced reference strains in the SILVA v.128 database, using “addSpecies” function. Sequences classified as mitochondria or chloroplast were removed from the ASV table. The datasets from the samples were normalized by random subsampling to include an equal number of reads (n = 2078 reads). Samples with lower number of sequences than the number of reads chose to the normalization were discarded (Supplementary Table 1).

Statistical analysis

All plots and statistical analyses were performed with RStudio (R Core Team 2017). Alpha diversity and error estimates were performed using QIIME script

core_diversity_analyses.py (Caporaso *et al.* 2010), including Chao1, equitability, observed richness, Shannon and Simpson's diversity. Two-way analysis of variance (ANOVA) was applied to investigate the influences of Cu and Pb addition, experimental time, and their interaction on α -diversity metrics and the abundance heterotrophic prokaryotes' community. Nonmetric multidimensional scaling (NMDS) based on Bray-Curtis distance was applied to evaluate the overall differences in prokaryotic community structure, using the *vegan* package in R (Oksanen *et al.* 2016). Global and pairwise PERMANOVA tests, using *vegan* and *RVAideMemoire* packages, respectively were applied to test significant differences between treatments and sampling time. To find representative phylotypes associated with the treatments, similarity percentage (SIMPER) analyses were first applied to screen OTUs primarily responsible for the overall dissimilarity (i.e. contributing to more than 1% of total dissimilarity) in the prokaryotic communities between treatments at each time point using *vegan* package.

Results

Trace metal elements and additional physicochemical parameters

Initial seawater samples (Ti) used for the first (i.e. 41p) and the second (i.e. LAZ and 6ext) experiments were geochemically contrasted and presented variations in trace metal concentrations as expected (Supplementary Table 2 and Supplementary Table 3). Total Cu, Pb, and Zn concentrations for 41p were 2.44 nM, 0.13 nM, and 5.34 nM, respectively while the concentrations were 2.5-, 3-, and 3.6-fold higher than in LAZ respectively, representing 15 nM, 0.79 nM, and 40 nM. For initial 6ext seawater, total

272 Cu, Pb, and Zn concentrations were 15-, 18-, and 26-fold higher than in 41p
273 respectively, representing 36 nM, 2.32 nM, and 139 nM.

274 At the beginning of both mesocosm experiments (T0), trace metal
275 concentrations measured in control conditions were close to the ones encountered in
276 initial seawater samples (Ti) from 41p, LAZ, 6ext. Thus, this confirms that no
277 adsorption or contamination in metals occurred in our experiments. In both
278 mesocosms, metal concentrations added were close to the targeted concentrations
279 (150 nM of Cu and 7 nM of Pb) (Supplementary Table 2 and Supplementary Table
280 3). Finally, both Cu and Pb concentrations at the beginning (T0) were not different to
281 those measured at the end (TF) of each experiment, confirming no metal loss in
282 treatments where metals were added (Supplementary Table 2 and 3).

283 The DOC and TN concentrations were not significantly different between the 3
284 initial seawaters (Supplementary Table 2 and Supplementary Table 3). For both
285 mesocosm experiments, water temperature, pH, DO, and DOC did not exhibit
286 significant differences among conditions but all of these parameters experienced
287 temporal significant changes during the entire course of experiments (Kruskal-Wallis,
288 $p < 0.05$). Globally, for both mesocosm experiments, pH and water temperature
289 increased with time while DO saturation and DOC concentration decreased with time
290 (Supplementary Table 2 and Supplementary Table 3).

291

292 **Ultraplanktonic and heterotrophic prokaryotes' abundances**

293 During the first mesocosm experiment, total ultraplanktonic abundances were
294 determined by flow cytometry. Abundances were comparable between sampled

295 seawater (Ti) from 41p and both conditions 41p CTRL and 41p Pb ($1.1 \pm 0.1 \times 10^4$
296 cell.mL⁻¹ on average), with low temporal variability during the entire course of
297 experiment (Supplementary Figure 3A). Total ultraphytoplanktonic abundances in
298 conditions 41p Cu and 41p Cu+Pb, started to decrease 3 days after metal addition,
299 leading to significantly lower abundances than in conditions 41p CTRL and 41p Pb
300 until the end of the experiment (ANOVA, $P < 0.001$). At the beginning of the
301 experiment (T0), heterotrophic prokaryotes' abundance measured in all conditions
302 ($11 \pm 0.3 \times 10^4$ cell.mL⁻¹ on average) was significantly lower than in sampled
303 seawater (Ti) from the 41p (2.5×10^5 cell.mL⁻¹) (ANOVA, $P < 0.001$) (Supplementary
304 Figure 3B). From three days after metal addition to the end of the experiment,
305 heterotrophic prokaryotic abundance was significantly higher in conditions 41p Cu
306 and 41p Cu+Pb than in conditions 41p CTRL and 41p Pb (ANOVA, $P < 0.001$).

307 During the second mesocosm experiment, total ultraphytoplanktonic
308 abundance in LAZ (1.9×10^4 cell.mL⁻¹) was significantly higher than in 6ext seawater
309 (7.9×10^3 cell.mL⁻¹) (ANOVA, $P < 0.001$). After the acclimation period (T0), total
310 ultraphytoplanktonic abundance followed a similar pattern in all conditions,
311 decreasing 3 days after metal addition (Supplementary Figure 4A). From 3 days after
312 metal addition to the end of the experiment, total ultraphytoplanktonic abundance
313 was significantly lower in Cu-treated conditions than in LAZ and 6ext control
314 conditions (ANOVA, $P < 0.001$). Heterotrophic prokaryotic abundance was not
315 significantly different between LAZ (5.8×10^5 cell.mL⁻¹) and 6ext (5.0×10^5 cell.mL⁻¹)
316 initial seawaters (Supplementary Figure 4B). At the beginning of the experiment (T0),
317 this abundance was similar in all conditions ($41 \pm 1.0 \times 10^4$ cell.mL⁻¹), while from 7
318 days after metal addition to the day 17, it was higher in Cu-treated conditions than in

control conditions (ANOVA, $P < 0.001$). From 21 days after metal addition to the end of the experiment, heterotrophic prokaryotic abundance was similar in 6ext CTRL and 6ext Cu. Heterotrophic prokaryotic abundance from LAZ Cu was higher than in LAZ CTRL.

Prokaryotic alpha-diversity

For the first mesocosm experiment, all the α -diversity indices except equitability varied significantly over time (Table 1A and Supplementary Figure 5). Globally, prokaryotic α -diversity indices significantly decreased over time in all conditions (ANOVA, $P < 0.05$). After the metal addition (T0), the Chao1 index was significantly lower in Cu-treated conditions (i.e., 41p Cu and 41p Cu+Pb) relative to both initial seawaters from 41p and condition 41p CTRL (ANOVA, $P < 0.001$). Simpson's diversity was significantly higher in metal-treated conditions (i.e., 41p Cu, 41p Pb, and 41p Cu+Pb) relative to condition 41p CTRL (ANOVA, $P < 0.001$). No significant difference was observed in equitability, observed richness, and Shannon indices between conditions.

For the second mesocosm experiment, all the α -diversity indices except equitability varied significantly over time (Table 1B and Supplementary Figure 6). All prokaryotic α -diversity indices significantly decreased over time in all conditions, especially after the 7th day of the experiment (ANOVA, $P < 0.05$). Average Chao1, observed OTUs, and Shannon indices varied across conditions, and they were lower in condition LAZ Cu relative to LAZ CTRL (ANOVA, $P < 0.05$). No significant difference was observed in all α -diversity indices between conditions 6ext CTRL and

6ext Cu. Finally, all α -diversity indices measured in the second mesocosm experiment were lower compared to those of the first mesocosm experiment.

Prokaryotic community is strongly shaped by Cu in contrast to Pb

In the first mesocosm, the most abundant dominant phyla/classes (> 1% of all sequences across all samples) were *Alphaproteobacteria*, *Gammaproteobacteria*, *Flavobacteriia* and *Cyanobacteria*, representing 96% of the overall relative abundance. Archaea represented less than 1% of the prokaryotic community of all samples and thus were grouped with all the “other” rare taxa (< 1% of all sequences across all samples) in subsequent analysis.

Among the abundant families, 10 representative families, determined by SIMPER analysis, significantly differed treatments over time (Supplementary Table 5). Their relative abundances are shown in Figure 2. *Rhodobacteraceae*, *Family I* (*Cyanobacteria*), *Alteromonadaceae*, and *SAR11 Surface 1 clade* decreased in Cu-treated conditions compared to 41p CTRL or 41p Pb. These families contributed to $20 \pm 1.3\%$, $16 \pm 4.6\%$, $6.7 \pm 2.5\%$, and $3.0 \pm 0.76\%$ of the overall dissimilarity between these two conditions, respectively. Conversely, the relative abundance of *Hyphomonadaceae*, *Flavobacteriaceae*, and *Haliaceae* increased in Cu-treated conditions compared to 41p CTRL or 41p Pb and contributed to $34 \pm 8.1\%$, $17 \pm 4.4\%$, and $4.8 \pm 1.6\%$ respectively of the overall dissimilarity between these two groups of conditions.

Moreover, 19 genera showed overall significant differences between conditions at the end of the experiment (TF) (Supplementary Table 6). Among these genera, only 4 were responsible for the structural change in Pb-treated condition.

The relative abundance of *Alteromonas*, *Glaciecola*, and unknown *SAR11 Surface 2* clade significantly increased while *Erythrobacter* nearly disappeared when only Pb was added. *Synechococcus*, *Litorimicrobium*, and to a lesser extent, *CHAB XI 27* decreased in 41p Cu compared to 41p CTRL. Conversely, *Parvularcula*, *NS5 marine group* and *Hyphomonas* increased in 41p Cu compared to 41p CTRL (Figure 3).

Samples from the first mesocosm study were mainly clustered by conditions (PERMANOVA, $R^2 = 0.41$; $P < 0.001$) and in a lesser extent by sampling times (PERMANOVA, $R^2 = 0.34$; $P < 0.001$) or interaction (PERMANOVA, $R^2 = 0.17$; $P < 0.001$) (Figure 4 and Supplementary Table 4A). At the beginning of the experiment (T0), prokaryotic community structures were similar between all conditions, and significantly different from the initial seawater at 41p (Supplementary Table 4B). After the 7th day, Cu-treated conditions (41p Cu and 41p Cu+Pb) shown significantly different prokaryotic community structures from the other conditions (41p, 41p CTRL, and 41p Pb) at each sampling time, especially at the end of the experiment (TF). Finally, there was no significant difference in communities between 41p CTRL and 41p Pb, as well as between 41p Cu and 41p Cu+Pb at each sampling time (Supplementary Table 4).

Influence of TME exposure history on copper effects

Prokaryotic community composition and structure

Concerning prokaryotic communities sampled from 3 geochemically contrasted sites (a low contaminated site (41p), an intermediately contaminated site (LAZ), and a highly contaminated site (6ext)), the most abundant phyla/classes (> 1% of all sequences across all samples) were *Alphaproteobacteria*, *Gammaproteobacteria*,

Flavobacteriia and *Cyanobacteria*, representing 94% of the overall relative abundance. At the family taxonomic rank, *Rhodobacteraceae* ($25 \pm 1.5\%$), *Flavobacteriaceae* ($16 \pm 1.1\%$), *Hyphomonadaceae* ($11 \pm 1.3\%$), *SAR11 Surface 1 clade* ($6.3 \pm 0.89\%$), and *Halieaceae* ($6.0 \pm 0.59\%$) included the majority of the total sequences (Figure 5). As for the first mesocosm experiment, Archaea represented less than 1% of the prokaryotic community of all samples, and they were considered in the group “other” corresponding to the rare taxa in further analyses.

NMDS plot and PERMANOVA tests showed significant clusters corresponding to both sampling times and conditions (Figure 6 and Supplementary Table 7A). At all taxonomic rank, Cu addition (PERMANOVA, $R^2 = 0.58$; $P < 0.001$) showed a greater influence on community variations than time (PERMANOVA, $R^2 = 0.14$; $P < 0.001$) or interaction between time and experimental condition (PERMANOVA, $R^2 = 0.20$; $P < 0.001$) (Supplementary Table 7A). Prokaryotic community structures were significantly similar between initial seawaters from LAZ and 6ext and significantly different to the initial seawater from 41p (Supplementary Table 7B). At the beginning of the experiment (T0), similarly to what was observed for 41p in the first experiment, the prokaryotic community structures in each control conditions were not significantly dissimilar from related Cu-treated ones (LAZ CTRL vs LAZ Cu, and 6ext CTRL vs 6ext Cu). After the 7th day, and until the end of the experiment, significant differences were clearly observed between Cu-treated and non-Cu-treated communities for 41p and LAZ. Conversely, pairwise comparisons showed no significant difference between 6ext CTRL and 6ext Cu, whatever the sampling time.

413 *Representative taxa responsible for structural changes between control and Cu-*
414 *treated conditions*

415 SIMPER analyses revealed that 12 representative families significantly differed initial
416 seawaters, but also control and Cu-treated conditions throughout experiment
417 duration (Figure 5 and Supplementary Table 8). *SAR11 Surface 1* and *Surface 2*
418 *clades*, *Hyphomonadaceae*, and *Family I (Cyanobacteria)* were higher in 41p than in
419 LAZ and 6ext, while *Flavobacteriaceae* and *Rhodobacteraceae* were more abundant
420 in LAZ and 6ext than in 41p (Supplementary Table 8). Finally, major differences
421 between LAZ CTRL and LAZ Cu were explained by the variation of relative
422 abundances of *Hyphomonadaceae*, *Rhodobiaceae*, *Cryomorphaceae* and
423 *Alteromonadaceae*.

424 At a lower taxonomic rank, among the 176 genera recorded, 34 significantly
425 differed control and Cu-treated conditions at the end of the experiment (TF) (Figure 3
426 and Supplementary Table 9). In addition, the genera *Salinihabitans* and *Rhodobium*
427 discriminating 41p CTRL from 41p Cu, decreased in LAZ Cu compared to LAZ CTRL
428 while *Jannaschia*, unknown *Cryomorphaceae*, and *Hirschia* increased in LAZ Cu
429 compared to LAZ CTRL. *Ruegeria* decreased in 6ext Cu compared to 6ext CTRL
430 while *Formosa*, OCS116 clade, and *Methylothera* increased in 6ext Cu compare to
431 6ext CTRL. Finally, representative genera responsible for structural changes
432 between control conditions and Cu-treated conditions for 41p and LAZ were mainly
433 abundant genera while for 6ext, representative genera were mainly rare genera (<
434 1% of all sequences across all samples).

435

Discussion

Copper exposure induced changes in prokaryotic community

In both mesocosm experiments, temporal variations in prokaryotic community structure were observed regardless of metal additions. However, metal addition led to stronger changes of community structure than the experimental setup or the experiment time. Copper exposure killed more than 99% of the ultraphytoplanktonic community. The selective pressure of metals on phytoplankton and more precisely on ultraphytoplankton is now well documented (Echeveste, Agustí and Tovar-Sánchez 2012; Lafabrie *et al.* 2013; Coclet *et al.* 2018). In parallel to the phytoplankton mortality, we observed an increase in heterotrophic prokaryotes' abundance and a decrease in their α -diversity if they originated from the lower and intermediately contaminated sites. This abundance increases thus seemed to relate the development of a limited number of opportunistic *r*-strategists that efficiently use the organic resources released by the phytoplanktonic decay (Odum, 1969; Pianka, 1970). This behavior has already been observed with the highly reactive marine microbial community (Buchan *et al.* 2014; El-Swaiss *et al.* 2015). Taken altogether, these observations highlight the strong interdependence between prokaryotic and phytoplanktonic communities, which could partly explain Cu-responses of microbial communities through indirect trophic relationships.

When considering the widely reported bacterial resistance to metals (Nies 1999; Baker-Austin *et al.* 2006), the response of the prokaryotic community to Cu addition might also be due to the persistence or proliferation of resistant taxa, which can be higher than the extinction rate of sensitive taxa (Wang *et al.* 2015; Qian *et al.* 2017).

Significant differences in community structure between the control and the Cu-added conditions were found at all sampling time. Particularly, *Hirschia* (*Hyphomonadaceae*), *NS4* and *NS5* marine groups, *Formosa* and *Tenacibaculum* (*Flavobacteriaceae*) increased in Cu-treated conditions for both mesocosm experiments, indicating that they could be more resistant or take advantage from copper addition and phytoplankton decay (Pinhassi *et al.* 2004). *Hyphomonadaceae* were known to be abundant in plastic-attached biofilm communities (Zettler, Mincer and Amaral-zettler 2013; Bryant *et al.* 2016; De Tender *et al.* 2017; Oberbeckmann *et al.* 2018), especially in disturbed environments (Briand *et al.*, 2017; Coclet, *et al.*, submitted). However, the results in this paper demonstrate for the first time the significant presence of this group in the prokaryotic community of metal contaminated seawater. *Flavobacteriaceae* was also a major component of the microbial community in strongly disturbed and metal-contaminated environments (Allen *et al.* 2013; Wang *et al.* 2015; Zouch *et al.* 2018; Coclet *et al.* 2019). Finally, the previous observation that *Tenacibaculum* and *Formosa* genera are positively selected in Cu-amended 6ext water as abundant and representative taxa from the most contaminated site (6ext) during a one-month-long survey, is in agreement with the idea of a similar selective pressure *in situ* as that observed in our previous field study experiment (Coclet *et al.* 2019). Conversely, members of *Rhodobacteraceae*, *Family I* (*Cyanobacteria*, mainly *Synechococcus*), *Alteromonadaceae*, and *SAR11 Surface 1 clade* families have been proposed to be sensitive *in situ* to high trace metal levels (Cassier-Chauvat and Chauvat 2014; Coclet *et al.* 2018, 2019; Corcoll *et al.* 2019) or characteristic of marine zones away from anthropogenic influence (Gilbert *et al.* 2009, 2012; Fuhrman, Cram and Needham 2015).

Thus, our study reveals the sensitivity of a part of the prokaryotic community to Cu contamination representative from a harbor, by a combination of direct (i.e. toxicity) or indirect (i.e. related to phytoplankton decay) processes, as suggested recently in a field study (Coclet *et al.* 2019).

Limited effects of Pb contamination on prokaryotic community

In our study, the effect of Pb exposure on prokaryotic communities was far less pronounced than the effect of Cu exposure. Pb addition using nitrate salts appeared to have limited impact on prokaryotic diversity and structure, with no detectable effect on ultraphytoplankton. Moreover, it did not prove to have a synergistic interaction with Cu. The weak toxicity of Pb on prokaryotic communities could be explained by its resistance and detoxification mechanisms, which are very efficient in microbial organisms (Trevors, 1989; Nies, 1999; Nayar *et al.*, 2004; Naik and Dubey, 2013). Additionally, the dichotomy between Cu and Pb effects could be due to the difference in their concentrations added in our mesocosms as well as their bioavailability. Indeed, to mimic the contamination levels observed in Toulon Bay, Pb concentrations were 20-fold lower than Cu concentrations, thus reducing the dose at which the cells were exposed. No data is available in the literature concerning tolerance threshold of marine bacterial communities against Pb contamination. Pb dissolved concentration was constant throughout the experiment, indicating no Pb adsorption on mesocosm materials or on microbial cells. Thus, the surface reactive behavior of Pb might not have induced a reduced bioavailability by reducing its dissolved concentration. Therefore, our study does not allow us to explain precisely this difference of toxicity between Cu and Pb, but the non-essential nature of Pb and potential different ways

to enter the cells than Cu represent hypotheses that should be studied in future research (Han *et al.* 2014).

The relative abundance of *Glaciecola* taxon significantly increased while *Erythrobacter* became undetectable when only Pb was added. *Glaciecola* has been described to be an opportunistic member of prokaryotic communities, its abundance increasing rapidly following an increase in DOC concentration (Sipler *et al.* 2017; von Scheibner, Sommer and Jürgens 2017). Since DOC concentrations in our experiment were similar in all conditions, the ability of *Glaciecola* to take advantage compared to other taxa in a Pb contaminated sea water remains unexplained. The genus *Erythrobacter* was found to be important taxa in surface-attached communities (Zettler, Mincer and Amaral-zettler 2013; Dussud *et al.* 2018; Oberbeckmann *et al.* 2018), and notably in contaminated ecosystems (Catão *et al. submitted*; Briand *et al.*, 2017; Pollet *et al.*, 2018; Coclet, *et al.*, submitted). Thus, the strong decrease of *Erythrobacter* in the water after Pb addition could be explained by the ability of this genus to form biofilms in the mesocosms. Indeed, biofilms are one of the most widely distributed and successful modes of life in water environment (Stoodley *et al.* 2002; Flemming *et al.* 2016) and a strategy that microorganisms might use to protect them from toxic metals (Harrison, Ceri and Turner 2007).

TME exposure history determines Cu effects on prokaryotic communities

Trace metal contamination gradients in Toulon bay have been already described by punctual mappings (Jean *et al.* 2012; Dang *et al.* 2015b; Coclet *et al.* 2018) or during short (Coclet *et al.* 2019) and long-term surveys (Layglon *et al.*, under revision). These studies demonstrated that TME contamination gradients persists at a pluri-

annual scale in Toulon Bay. This led us to assume that prokaryotic communities sampled along this contamination gradient could present different metals exposure histories. Cu addition in both low (41p) and in intermediately (LAZ) contaminated seawaters induced important changes in prokaryotic community structure, while Cu addition in seawater originating from a chronically contaminated site (6ext) only weakly impacted the prokaryotic community structure. Based on the PICT theory (Gustavson *et al.* 1999; Blanck 2002), this observation validates our assumption and demonstrates the adaptive potential of prokaryotic communities to withstand elevated Cu concentrations. This adaptation must have developed rapidly since the water residence time in Toulon Bay has been estimated to a few days (Dufresne, Duffa and Rey 2014), even if this estimation is probably not representative to what happens in very enclosed docks such as the one where we sampled contaminated water (6ext). This adaptive potential could be the result of the presence of a higher proportion of tolerant species in the most contaminated seawater, representative of a strong and chronic metal-exposure history of these communities (Carman, Fleeger and Pomarico 2000). This hypothesis is in agreement with the very small variation of structure after Cu addition in the community sampled from the most contaminated site. Indeed, for this community, we mainly observed changes in the relative abundance of rare taxa.

We hypothesize that mechanisms of toxicity and resistance to Cu co-occur in prokaryotic communities from low and intermediately contaminated sites. The observed dissimilarity in prokaryotic structures between these communities after Cu exposure could be the result of differences in initial community composition in 41p and LAZ seawaters. To the best of our knowledge, our study provides the first

556 evidence that TME exposure history may play a critical role in marine prokaryotic
557 community response to re-current metal contamination.

558

559 **Conclusion**

560 From our mesocosm experiences, we can conclude that harbor-relevant Cu
561 exposure affects abundance, diversity and structure of prokaryotic communities from
562 both low and intermediately TME contaminated seawaters. The effect of Cu addition
563 is visible through both the disappearance of sensitive taxa and the selection of
564 adapted ones. We also demonstrated that TME exposure history has a strong
565 influence on the response of prokaryotic communities to newer-current
566 contamination, as prokaryotic communities from contaminated site were less
567 impacted by Cu addition. Conversely, we showed limited impacts of Pb exposure
568 compared to Cu contamination, probably due to the low toxic potential against
569 prokaryotes and the efficiency of its resistance and detoxification mechanisms.
570 Finally, our observations suggest both direct (toxic) and indirect (trophic) effects of
571 TME contaminations, highlighting the need for studies of the complexity of both biotic
572 and abiotic interactions.

573

574 **Acknowledgements**

575 This study was supported by IMPRECI-M² research program (funded by EC2CO). C.
576 Coclet PhD was funded by the Provence-Alpes-Côte d'Azur region and supported by
577 TPM. The authors wish to thank the French Navy and the LASEM for diving and
578 sampling assistance, C. Martino for building assistance of mesocosm structure
579 (Toulon University, France) for sampling and sample treatment assistance. The
580 authors wish also to thank the Dr. Pedro Lebre to correct language and grammar.

581

582 **Compliance with ethical standards**

583 **Conflict of interest** The authors declare that they have no conflict of interest.

584

585 **References**

586 Allen RC, Tu Y, Nevarez MJ *et al.* The mercury resistance (mer) operon in a marine
587 gliding flavobacterium, *Tenacibaculum discolor* 9A5. *FEMS Microbiol Ecol*
588 2013;**83**:135–148.

589 Baker-Austin C, Wright MS, Stepanauskas R *et al.* Co-selection of antibiotic and
590 metal resistance. *Trends Microbiol* 2006;**14**:176–82.

591 Bell TH, Yergeau E, Maynard C *et al.* Predictable bacterial composition and
592 hydrocarbon degradation in Arctic soils following diesel and nutrient disturbance.
593 *Isme* 2013:1–11.

594 Bissett A, Richardson AE, Baker G *et al.* Life history determines biogeographical
595 patterns of soil bacterial communities over multiple spatial scales. *Mol Ecol*

596 2010;**19**:4315–27.

597 Blanck H. A Critical Review of Procedures and Approaches Used for Assessing
598 Pollution-Induced Community Tolerance (PICT) in Biotic Communities. *Hum Ecol*
599 *Risk Assess* 2002;**8**:1003–34.

600 Blanck H, Wangberg SA, Molander S. Pollution-Induced Community Tolerance - A
601 new ecotoxicological tool. *Functional Testing of Aquatic Biota for Estimating*
602 *Hazards of Chemicals*. Cairns, J. Philadelphia, 1988, 219–30.

603 Briand J-F, Barani A, Garnier C *et al*. Spatio-Temporal Variations of Marine Biofilm
604 Communities Colonizing Artificial Substrata Including Antifouling Coatings in
605 Contrasted French Coastal Environments. *Environ Microbiol* 2017;**74**:585–98.

606 Bryant JA, Clemente TM, Viviani DA *et al*. Diversity and Activity of Communities
607 Inhabiting Plastic Debris in the North Pacific Gyre. *mSystems* 2016;**1**:e00024-16.

608 Buchan A, LeClerc GR, Gulvik CA *et al*. Master recyclers: features and functions of
609 bacteria associated with phytoplankton blooms. *Nat Rev Microbiol* 2014;**12**:686–
610 98.

611 Cabrol L, Quéméneur M, Misson B. Inhibitory effects of sodium azide on microbial
612 growth in experimental resuspension of marine sediment. *J Microbiol Methods*
613 2017;**133**:62–5.

614 Callahan BJ, McMurdie PJ, Rosen MJ *et al*. DADA2 : High-resolution sample
615 inference from Illumina amplicon data. *Nat Methods* 2016;**13**:581.

616 Callender E. *Heavy Metals in the Environment – Historical Trends*. 2nd ed. Published

617 by Elsevier Inc., 2003.

618 Caporaso JG, Kuczynski J, Stombaugh J *et al.* QIIME allows analysis of high-
619 throughput community sequencing data. *Nat Methods* 2010;**7**:335–6.

620 Carman KR, Fleeger JW, Pomarico SM. Does historical exposure to hydrocarbon
621 contamination alter the response of benthic communities to diesel
622 contamination ? *Mar Environ Res* 2000;**49**:255–78.

623 Casas S, Gonzalez J-L, Andral B *et al.* Relation between metal concentration in
624 water and metal content of marine mussels (*Mytilus galloprovincialis*): impact of
625 physiology. *Environ Toxicol Chem* 2008;**27**:1543–52.

626 Cassier-Chauvat C, Chauvat F. Responses to Oxidative and Heavy Metal Stresses in
627 Cyanobacteria: Recent Advances. *Int J Mol Sci* 2014;**16**:871–86.

628 Catão EC, Pollet T, Misson B *et al.* Shear stress as a major driver of marine biofilm
629 communities in the NW Mediterranean Sea. *Front Microbiol* 2019;**10**, DOI:
630 10.3389/fmicb.2019.01768.

631 Cindrić A-M, Garnier C, Oursel B *et al.* Evidencing the natural and anthropogenic
632 processes controlling trace metals dynamic in a highly stratified estuary : The
633 Krka River estuary. *Mar Pollut Bull* 2015;**94**:199–216.

634 Coclet C, Garnier C, D'onofrio S *et al.* Evidence for microbial biofilm community
635 structuring by sediment closeness rather than multichemical contamination or
636 bacterioplankton composition. *Environ Microbiol.*

637 Coclet C, Garnier C, Delpy F *et al.* Metals contamination as a toxic and structuring

638 factor for ultraphytoplankton in a multicontaminated Mediterranean coastal area.
639 2017, DOI: 10.1016/j.pocean.2017.06.006.

640 Coclet C, Garnier C, Delpy F *et al.* Trace metal contamination as a toxic and
641 structuring factor impacting ultraphytoplankton communities in a
642 multicontaminated Mediterranean coastal area. *Prog Oceanogr* 2018;**163**:196–
643 213.

644 Coclet C, Garnier C, Durrieu G *et al.* Changes in bacterioplankton communities
645 resulting from direct and indirect interactions with trace metal gradients in an
646 urbanized marine coastal area. *Front Microbiol* 2019;**10**:250.

647 Corcoll N, Yang J, Backhaus T *et al.* Copper affects composition and functions of
648 microbial communities in marine biofilms at environmentally relevant
649 concentrations. *Front Microbiol* 2019;**9**:1–15.

650 Cossa D, Buscail R, Puig P *et al.* Origin and accumulation of trace elements in
651 sediments of the northwestern Mediterranean margin. *Chem Geol* 2014;**380**:61–
652 73.

653 Dang DH, Lenoble V, Durrieu G *et al.* Sedimentary dynamics of coastal organic
654 matter: An assessment of the porewater size/reactivity model by spectroscopic
655 techniques. *Estuar Coast Shelf Sci* 2014;**151**:100–11.

656 Dang DH, Lenoble V, Durrieu G *et al.* Seasonal variations of coastal sedimentary
657 trace metals cycling: Insight on the effect of manganese and iron
658 (oxy)hydroxides, sulphide and organic matter. *Mar Pollut Bull* 2015a;**92**:113–24.

659 Dang DH, Schäfer J, Brach-Papa C *et al.* Evidencing the Impact of Coastal
 660 Contaminated Sediments on Mussels Through Pb Stable Isotopes Composition.
 661 *Environ Sci Technol* 2015b;**49**:11438–48.

662 Dufresne C, Duffa C, Rey V. Wind-forced circulation model and water exchanges
 663 through the channel in the Bay of Toulon. *Ocean Dyn* 2014;**64**:209–24.

664 Dussud C, Meistertzheim AL, Conan P *et al.* Evidence of niche partitioning among
 665 bacteria living on plastics, organic particles and surrounding seawaters. *Environ*
 666 *Pollut* 2018;**236**:807–16.

667 Echeveste P, Agustí S, Tovar-Sánchez A. Toxic thresholds of cadmium and lead to
 668 oceanic phytoplankton: Cell size and ocean basin-dependent effects. *Environ*
 669 *Toxicol Chem* 2012;**31**:1887–94.

670 El-Swaiss H, Dunn KA, Bielawski JP *et al.* Seasonal assemblages and short-lived
 671 blooms in coastal north-west Atlantic Ocean bacterioplankton. *Environ Microbiol*
 672 2015;**17**:3642–61.

673 Flemming HC, Wingender J, Szewzyk U *et al.* Biofilms: An emergent form of bacterial
 674 life. *Nat Rev Microbiol* 2016;**14**:563–75.

675 Fuhrman JA, Cram JA, Needham DM. Marine microbial community dynamics and
 676 their ecological interpretation. *ISME J* 2015;**13**:133.

677 Ghiglione JF, Conan P, Pujo-Pay M. Diversity of total and active free-living vs.
 678 particle-attached bacteria in the euphotic zone of the NW Mediterranean Sea.
 679 *FEMS Microbiol Lett* 2009;**299**:9–21.

680 Gilbert JA, Field D, Swift P *et al.* The seasonal structure of microbial communities in
681 the Western English Channel. *Environ Microbiol* 2009;**11**:3132–9.

682 Gilbert JA, Steele JA, Caporaso JG *et al.* Defining seasonal marine microbial
683 community dynamics. *ISME J* 2012;**6**:298–308.

684 Goni-Urriza M, Moussard H, Lafabrie C *et al.* Consequences of contamination on the
685 interactions between phytoplankton and bacterioplankton. *Chemosphere*
686 2018;**195**:212–22.

687 Gustavson K, Petersen S, Pedersen B *et al.* Pollution-induced community tolerance
688 (PICT) in coastal phytoplankton communities exposure to copper. *Hydrobiologia*
689 1999;**416**:125–38.

690 Halpern SB, Walbridge S, Sleko AK *et al.* A Global Map of Human Impact on Marine
691 Ecosystems. *Science (80-)* 2008;**319**:948–53.

692 Han S, Zhang Y, Masunaga S *et al.* Relating metal bioavailability to risk assessment
693 for aquatic species: Daliao River watershed, China. *Environ Pollut*
694 2014;**189**:215–22.

695 Harrison JJ, Ceri H, Turner RJ. Multimetal resistance and tolerance in microbial
696 biofilms. *Nat Rev Microbiol* 2007;**5**:928–38.

697 Hughes-Martiny JB, Bohannan BJM, Brown JH *et al.* Microbial biogeography : putting
698 microorganisms on the map. *Nat Rev Microbiol* 2006;**4**:102–12.

699 Jean N, Dumont E, Durrieu G *et al.* Protein expression from zooplankton
700 communities in a metal contaminated NW mediterranean coastal ecosystem.

701   *Mar Environ Res* 2012;**80**:12–26.

702   Lafabrie C, Garrido M, Leboulanger C *et al.* Impact of contaminated-sediment
703   resuspension on phytoplankton in the Biguglia lagoon (Corsica, Mediterranean
704   sea). *Estuar Coast Shelf Sci* 2013;**130**:70–80.

705   Layglon N, Misson B, Durrieu G *et al.* Impact of sediment dredging on seawater
706   quality: a long-term monitoring study of trace metals and microbial community in
707   Toulon Bay (France). *Sci Total Environ.*

708   Levin LA, Boesch DF, Covich A *et al.* The function of marine critical transition zones
709   and the importance of sediment biodiversity. *Ecosystems* 2001;**4**:430–51.

710   Mackey KRM, Buck KN, Casey JR *et al.* Phytoplankton responses to atmospheric
711   metal deposition in the coastal and open-ocean Sargasso Sea. *Front Microbiol*
712   2012;**3**:1–15.

713   Morley NH, Burton JD, Tankere SPC *et al.* Distribution and behaviour of some
714   dissolved trace metals in the western Mediterranean Sea. *Deep Res II*
715   1997;**44**:675–91.

716   Naik MM, Dubey SK. Lead resistant bacteria: Lead resistance mechanisms, their
717   applications in lead bioremediation and biomonitoring. *Ecotoxicol Environ Saf*
718   2013;**98**:1–7.

719   Nayar S, Goh BPL, Chou LM. Environmental impact of heavy metals from dredged
720   and resuspended sediments on phytoplankton and bacteria assessed in in situ
721   mesocosms. *Ecotoxicol Environ Saf* 2004;**59**:349–69.

- 722 Nies DH. Microbial heavy-metal resistance. *Appl Microbiol Biotechnol* 1999;**51**:730–
723 50.
- 724 Oberbeckmann S, Kreikemeyer B, Labrenz M *et al.* Environmental factors support
725 the formation of specific bacterial assemblages on microplastics. *Front Microbiol*
726 2018;**8**:1–12.
- 727 Oksanen J, Blanchet FG, Friendly M *et al.* vegan: Community Ecology Package.
728 Available online at: [https:// CRAN.R-project.org/package=vegan](https://CRAN.R-project.org/package=vegan). 2016.
- 729 Omanović D, Kwokal Ž, Goodwin A *et al.* Trace metal detection in Šibenik Bay,
730 Croatia: Cadmium, lead and copper with anodic stripping voltammetry and
731 manganese via sonoelectrochemistry. A case study. *J Iran Chem Soc*
732 2006;**3**:128–39.
- 733 Oursel B, Garnier C, Durrieu G *et al.* Dynamics and fates of trace metals chronically
734 input in a Mediterranean coastal zone impacted by a large urban area. *Mar*
735 *Pollut Bull* 2013;**69**:137–49.
- 736 Parada AE, Needham DM, Fuhrman JA. Every base matters : assessing small
737 subunit rRNA primers for marine microbiomes with mock communities, time
738 series and global field samples. *Environ Microbiol* 2016;**18**:1403–14.
- 739 Pinhassi J, Sala MM, Havskum H *et al.* Changes in bacterioplankton composition
740 under different phytoplankton regimens. *Appl Environ Microbiol* 2004;**70**:6753–
741 66.
- 742 Pollet T, Berdjeb L, Garnier C *et al.* Prokaryotic community successions and

743 interactions in marine biofilms: the key role of Flavobacteriia. *FEMS Microbiol*
744 *Ecol* 2018;**94**:fiy083.

745 Pringault O, Lafabrie C, Avezac M *et al.* Consequences of contaminant mixture on
746 the dynamics and functional diversity of bacterioplankton in a southwestern
747 Mediterranean coastal ecosystem. *Chemosphere* 2016;**144**:1060–73.

748 Pruesse E, Quast C, Knittel K *et al.* SILVA: A comprehensive online resource for
749 quality checked and aligned ribosomal RNA sequence data compatible with
750 ARB. *Nucleic Acids Res* 2007;**35**:7188–96.

751 Qian J, Ding Q, Guo A *et al.* Alteration in successional trajectories of
752 bacterioplankton communities in response to co-exposure of cadmium and
753 phenanthrene in coastal water microcosms. *Environ Pollut* 2017;**221**:480–90.

754 Quast C, Pruesse E, Yilmaz P *et al.* The SILVA ribosomal RNA gene database
755 project: Improved data processing and web-based tools. *Nucleic Acids Res*
756 2013;**41**:590–6.

757 R Core Team. R: A Language and Environment for Statistical Computing. Vienna: R
758 Foundation for Statistical Computing. 2017.

759 von Scheibner M, Sommer U, Jürgens K. Tight Coupling of *Glaciecola* spp . and
760 Diatoms during Cold-Water Phytoplankton Spring Blooms. *Front Microbiol*
761 2017;**8**:1–11.

762 Serra A, Corcoll N, Guasch H. Copper accumulation and toxicity in fluvial periphyton:
763 The influence of exposure history. *Chemosphere* 2009;**74**:633–41.

- 764 Sheeba VA, Abdulaziz A, Gireeshkumar TR *et al.* Role of heavy metals in structuring
765 the microbial community associated with particulate matter in a tropical estuary.
766 *Environ Pollut* 2017;**231**:589–600.
- 767 Sipler RE, Kellogg CTE, Connelly TL *et al.* Microbial Community Response to
768 Terrestrially Derived Dissolved Organic Matter in the Coastal Arctic. *Front*
769 *Microbiol* 2017;**8**:1–19.
- 770 Sjöstedt J, Langenheder S, Kritzberg E *et al.* Repeated disturbances affect functional
771 but not compositional resistance and resilience in an aquatic bacterioplankton
772 community. *Environ Microbiol Rep* 2018;**10**:493–500.
- 773 Soldo D, Hari R, Sigg L *et al.* Tolerance of *Oocystis nephrocytioides* to copper:
774 Intracellular distribution and extracellular complexation of copper. *Aquat Toxicol*
775 2005;**71**:307–17.
- 776 Stoodley P, Sauer K, Davies DG *et al.* Biofilms as Complex Differentiated
777 Communities. *Annu Rev Microbiol* 2002;**56**:187–209.
- 778 De Tender AC, Devriese IL, Haegeman A *et al.* Temporal dynamics of bacterial and
779 fungal colonization on plastic debris in the North Sea. *Environ Sci Technol*
780 2017;**51**:7350–60.
- 781 Tessier E, Garnier C, Mullot JU *et al.* Study of the spatial and historical distribution of
782 sediment inorganic contamination in the Toulon bay (France). *Mar Pollut Bull*
783 2011;**62**:2075–86.
- 784 The Mermex Group, Durrieu de Madron X, Guieu C *et al.* Marine ecosystems '

785 responses to climatic and anthropogenic forcings in the Mediterranean. *Prog*
786 *Oceanogr* 2011;**91**:97–166.

787 Tlili A. Signification écologique de la tolérance acquise des communautés
788 microbiennes des biofilms de rivières à une contamination d'origine anthropique.
789 2013.

790 Trevors JT. The role of microbial metal resistance and detoxification mechanisms in
791 environmental bioassay research. *Hydrobiologia* 1989;**188–189**:143–7.

792 Turner A. Marine pollution from antifouling paint particles. *Mar Pollut Bull*
793 2010;**60**:159–71.

794 Twining BS, Rauschenberg S, Morton PL *et al.* Metal contents of phytoplankton and
795 labile particulate material in the North Atlantic Ocean. *Prog Oceanogr*
796 2015;**137**:261–83.

797 Wang K, Zhang D, Xiong J *et al.* Response of Bacterioplankton Communities to
798 Cadmium Exposure in Coastal Water Microcosms with High Temporal
799 Variability. *Appl Environ Microbiol* 2015;**81**:231–40.

800 Ward CS, Pan JF, Colman BP *et al.* Conserved Microbial Toxicity Responses for
801 Acute and Chronic Silver Nanoparticle Treatments in Wetland Mesocosms.
802 *Environ Sci Technol* 2019;**53**:3268–76.

803 Xu Y, Sun Q, Yi L *et al.* The source of natural and anthropogenic heavy metals in the
804 sediments of the Minjiang River Estuary (SE China): Implications for historical
805 pollution. *Sci Total Environ* 2014;**493**:729–36.

806 Zettler ER, Mincer TJ, Amaral-zettler LA. Life in the “plastisphere”: microbial
807 communities on plastic marine debris. *Environ Sci Technol* 2013;**47**:7137-7146.

808 Zouch H, Cabrol L, Chifflet S *et al.* Effect of Acidic Industrial Effluent Release on
809 Microbial Diversity and Trace Metal Dynamics During Resuspension of Coastal
810 Sediment. *Front Microbiol* 2018;**9**, DOI: 10.3389/fmicb.2018.03103.

811