Blue LED Irradiation of Iodonium Ylides Gives Diradical Intermediates for Efficient Metal-free Cyclopropanation with Alkenes

Tristan Chidley,^[a] Islam Jameel,^[a] Shafa Rizwan,^[a] Philippe A. Peixoto,^[b] Laurent Pouységu,^[b] Stéphane Quideau,^{*[b]} W. Scott Hopkins^{*[a]} and Graham K. Murphy^{*[a]}

Abstract: A facile and highly chemoselective synthesis of doubly activated cyclopropanes is reported where mixtures of alkenes and β -dicarbonyl-derived iodonium ylides are irradiated with light from blue LEDs. This metal-free synthesis gives cyclopropanes in yields up to 96%, is operative with cyclic and acyclic ylides, and proceeds with a variety of electronically-diverse alkenes. Computational analysis explains the high selectivity observed, which derives from exclusive HOMO to LUMO excitation, instead of free carbene generation. The procedure is operationally simple, uses no photocatalyst, and provides access in one step to important building blocks for complex molecule synthesis.

Recently, the use of hypervalent iodine(III) (HVI) reagents in organic chemistry has undergone tremendous growth, owing to the inherent advantages of mild reaction conditions and broad reactivity, and of recyclable, environmentally-benign by-products.^[1] Cutting edge strategies towards reactions that are catalytic in iodoarene have been developed, including catalytic asymmetric variants.^[2] In addition, recent efforts to merge visible light-mediated photochemistry with HVI-based reactions has revealed new and exciting potential for organic synthesis.^[3] However, compared with the major achievements realized for photoredox catalysis with transition metal complexes,^[4] the intersection of HVI-mediated reactions and visible light photoredox catalysis or photoactivation is still in its infancy.

Our interest in HVI-mediated reactions stems from the movement towards safer and more sustainable organic synthesis practices. By developing new processes in which iodonium ylides serve as diazonium ylide surrogates,^[5] we exploit the increased and often complementary reactivity that they display, which originates from their hypervalent bonding. Iodonium ylides are synthetically versatile,^[6] readily undergoing ionic reactions,^[7] and serving as precursors in C–H insertion,^[8] cycloaddition,^[9] cyclopropanation^[10] or tandem reactions thereof,^[11] as both racemic and asymmetric processes.^[10f,12] The reliance on the use of β -dicarbonyl or similarly β -distabilized species to generate iodonium ylides render these unique carbene precursors ideally functionalized for the synthesis of important synthetic motifs, such as doubly activated cyclopropanes.^[13,14]

Iodonium ylides can be converted to singlet carbenes under irradiation using a mercury (Hg) lamp, and the groups of Hadjiarapoglou,^[9c] Matveeva,^[7c,15] and Spyridis^[9f,16] have reported examples of cyclopropanation and related cycloaddition events. These reactions thus proceed via free carbenes, which are formed upon cleavage of the ylide C–I bond, but the heat generated by the lamp is often an unavoidable source of technical complications and chemical selectivity issues. For example, Hadjiarapoglou showed that irradiating a iodonium ylide **1a** with a Hg lamp leads exclusively to a dihydrofuran in 96% yield, in part due to the heat generated during the reaction, which is sufficient to isomerize any of cyclopropane **3a** formed (Scheme 1a). In this regard, light-emitting diodes (LEDs) constitute new tools for testing the reactivity of iodonium ylides, as light irradiation in the lower-energy 400–500 nm (violet-blue) range can be easily achieved without significant heating, hence enabling the practical, convenient and selective generation of reactive intermediates under much milder conditions. Recently, Davies and Jurberg reported the photochemical decomposition of aryldiazoacetates with blue LEDs in the presence of styrene,^[17,18] however, these cyclopropanation reactions were only successful with donor/acceptor diazo derivatives, as both monostabilized ethyl diazoacetate and distabilized dimethyl 2-diazomalonate derivatives did not absorb in the blue light region. Though they often exhibit similar chemical reactivity, iodonium ylides are zwitterionic and structurally distinct from diazo compounds, because their ylene resonance forms contribute little to their C–I bonding.^[19] Thus, we envisioned that irradiating distabilized iodonium ylides with lower-energy visible light could provide an entry into metal-free carbenoid-type reactivity. Herein, UV-Visible spectroscopy, computational analysis and experimentation were used to elucidate how distabilized cyclic and acyclic iodonium ylides could react with alkenes under blue LED irradiation to provide cyclopropanes (Scheme 1b).

Scheme 1. Light-mediated reactions of iodonium ylides.

[a] T. Chidley, I. Jameel, S. Rizwan, Prof. Dr. W.S. Hopkins and Prof. Dr. G.K. Murphy
Department of Chemistry, University of Waterloo, Waterloo, Ontario, Canada, N2L3G1
E-mail: graham.murphy@uwaterloo.ca

[b] Dr. Philippe A. Peixoto, Prof. Dr. L. Pouységu, Prof. Dr. S. Quideau
Univ. Bordeaux, ISM (CNRS-UMR 5255), 351 cours de la Libération, Talence F-33405 Cedex, France

We began this study by recording the UV-Vis absorption spectra of the dimedone-derived iodonium ylide **1a** at 0.1 M and 0.03 mM concentrations (Figure 1a and 1b, also see Figure SI-1). When overlaid with the emission spectrum of the RGB LEDs, it is clear that excitation of **1a** can only occur in the blue region (see inset). We then tested a series of related iodonium ylides possessing substituents on the phenyl ring of the iodoarene motif, but we found little-to-no variation in their UV-Vis absorption spectra at 0.1 M with iodoarenes equipped with *p*-tolyl (**1b**), *p*-Cl (**1c**), *p*-CN (**1d**), 2,6-dimethoxy-4-methyl (**1g**) and even with the 2-naphthyl (**1h**) as compared to that of the parent iodonium ylide **1a** (see Figure SI-10). Changing the β -dicarbonyl region of the ylide resulted in more significant changes in their absorption spectra. Progressing from the dimedone-(**1a**), to the ketoester-(**1i**), to the diester-(**1j**) derived ylides showed an increasing blue shift (see Figure SI-11), and we anticipated that the decreasing overlap with the blue LED emission spectrum would result in increasingly longer reaction rates for these ylides.

To explore the electronic spectroscopy from a fundamental standpoint, we conducted a computational study of selected iodonium ylides at the time-dependent (TD) DFT level of theory.^[20] First, a search for optimal molecular geometry was performed at the B3LYP/6-311+G(d,p) level of theory.^[21] Upon identifying the global minimum geometry for each ylide, the optimal ground state structure was employed for TD-DFT calculations of the ten lowest-energy excited electronic states. An example of the output from these calculations is provided in Figure 1c, which plots the predicted UV-Vis spectrum for **1a** as relative absorption cross-section versus wavelength. In this, and all other cases examined, we found that the blue LED irradiation drives HOMO to LUMO excitation. For **1a**, the HOMO to LUMO band system is predicted to be the most intense of those calculated (332 nm), and it is well separated from those of higher energies (i.e., shorter wavelengths). While the relative intensity and absorption maximum of the HOMO to LUMO excitation did vary somewhat across the series of iodonium ylides (see Figures SI-1 to SI-6), the nature of the electronic excitation was essentially the same for each iodonium ylide. Specifically, for **1a**, the HOMO of the ylide is predominantly associated with electron density at the carbonyl oxygens and the ylide carbon (Figure 1d). Excitation to the LUMO promotes one electron from the HOMO to an orbital that is predominantly associated with the iodine centre. This interpretation is supported by natural population analysis,^[22] which indicates that 0.26e of electron density is transferred to iodine upon irradiation of the ylide (average determined from ylides **1a,b,g-j**; see Table SI-1). Therefore, unlike related diazo compounds, β -dicarbonyl-derived iodonium ylides are expected to absorb blue light and to generate 1,2-diradical species, such as **A** (with unpaired electron density predominantly on the iodine atom and the ylidic carbon), instead of undergoing heterolytic cleavage to produce free carbenes.

To test the chemical reactivity of iodonium ylides under blue LED (λ_{max} 461 nm) irradiation, we constructed a multi-slot reactor (see Figure SI-15) in which the reaction vial is placed directly above the LED, thus minimizing and standardizing the distance between the two. We found that irradiating ylide **1a** in the presence of 1 equiv of styrene for 2.5 h in CDCl₃ gives cyclopropane **3a** in 71% NMR yield, without formation of the free carbene-derived dimer product **2a** (Table 1, entry 1; see also Table SI-2). Using 2 equiv of styrene led to **3a** in 96% yield, and

Figure 1. a) UV-Vis spectrum of **1a** at 0.1 M, overlaid with emission spectra of RGB LEDs and their λ_{max} values. b) UV-Vis spectrum of **1a** at 0.03 mM. c) Calculated UV-Vis spectrum of **1a**; the dominant 332 nm absorption corresponds to a HOMO to LUMO transition. d) Calculated HOMO of **1a** (bottom), showing electron density predominantly on the β -dicarbonyl motif, and calculated LUMO of **1a** (top), showing predominantly orbital density on the iodine atom. Upon blue LED irradiation, the LUMO becomes populated with a single electron, generating the diradical species **A** depicted in b).

very little improvement was realized when using 4 equiv of styrene (entries 2,3). We found CHCl₃, CH₂Cl₂ or CH₃CN to be also effective solvents, giving **3a** in similar yields within 2 h; however, when the polar protic solvent MeOH was used, neither **2a**, **3a** or O-H insertion products was observed (entries 4-7). Since **1a** was much more soluble in CH₃CN than in the other solvents, it was chosen as the optimal reaction solvent. Using CH₃CN, **3a** was produced and isolated in 95% yield (Table 1, entry 6). As expected (see Figure 1a), irradiating the reaction mixture with either red or green LEDs for 24 h resulted in no ylide consumption, although subsequent irradiation with blue LEDs afforded **3a** in 63% and 79% yields, respectively (entries 8,9). Irradiating **1a** with blue LEDs for 72 h in the absence of

styrene led to complete ylide decomposition (entry 10), again without formation of the free carbene-derived dimer product **2a**.^[8c] Attempting the reaction under ambient light for 24 h, or in the dark for 72 h, resulted in quantitative recovery of **1a** (entries 11,12). Therefore, irradiating **1a** with blue light leads to a mild and highly efficient metal-free cyclopropanation that, unlike related processes employing UV lamps, is entirely chemoselective for the formation of cyclopropane **3a** without any evidence of free carbene intermediates or of isomerisation to the dihydrofuran product (see Scheme 1a).

We then evaluated the impact of the iodoarene substitution pattern on the rate and efficacy of the cyclopropanation reaction by subjecting the dimedone-derived iodonium ylides **1a-1h** to the optimized reaction conditions (Scheme 2). In each case, the ylide was consumed within 2.5 h, affording **3a** in yields ranging from 59 to 95%. It is worth noting that no correlation was observed between reaction rates or yields and either the electron-rich or electron-poor character of the substrates. For example, **3a** was observed in 87% and 89% yields when using either the iodonium ylide **1f** (*p*-OMe) or **1e** (*p*-CF₃). These results are consistent with

Table 1. Optimization of the cyclopropanation reaction.

Entry	Styrene (equiv)	Solvent	Time (h)	Yield ^[a] 2a	Yield ^[a] 3a
1	1.0	CDCl ₃	2.5	—	71
2	2.0	CDCl ₃	1.5	—	96
3	4.0	CDCl ₃	1.5	—	98
4	2.0	CHCl ₃	2	—	95
5	2.0	CH ₂ Cl ₂	2	—	92
6	2.0	CH ₃ CN	1.5	—	98 (95) ^[b]
7	2.0	CH ₃ OH	8	—	—
8 ^[c]	2.0	CH ₃ CN	24	—	0%, ^[d] 63% ^[e]
9 ^[f]	2.0	CH ₃ CN	24	—	0%, ^[d] 79% ^[e]
10	—	CH ₃ CN	72	—	—
11 ^[g]	2.0	CH ₃ CN	24	—	— ^d
12 ^[h]	2.0	CH ₃ CN	72	—	— ^d

[a] ¹H NMR yields using HMDSO as internal standard. [b] Isolated yield.

[c] Irradiation with red LED. [d] Quantitative observation of **1a** by ¹H NMR yield.

[e] ¹H NMR yield after subsequent blue LED irradiation. [f] Irradiation with green LED.

[g] Reaction performed in ambient light. [h] Reaction performed in the dark.

the minimal deviations observed in the UV-Vis absorption spectra of these compounds compared with that of **1a** (see Figure SI-10).

We next submitted the ketoester (**1i**) and diester (**1j**) analogues of **1a** to the same optimized reaction conditions. A significant decrease in reaction rate was observed, with **1i** being consumed in 4 h and **1j** only after 24 h, though the yields

remained excellent. We were pleased to observe that these ylides did not suffer from decomposition upon prolonged irradiation. Interestingly, both the experimental and predicted UV-Vis spectra of **1a**, **1i** and **1j** showed a blue shift upon increasing ester function incorporation (See Figure SI-11). To probe this effect, we conducted another experiment in which ylide **1j** was subjected to the optimal reaction conditions, but irradiating instead with violet LEDs (λ_{max} 397 nm, see Figure SI-12). In this case, ylide **1j** was consumed within 1 h, and the expected cyclopropane product **3d** was observed in 67% ¹H NMR yield, along with 33% of the Meldrum's acid dimer **2b** (see Scheme 2). Thus, the chemoselectivity observed using blue light irradiation derives from exclusive photoexcitation into a diradical intermediate, whose generation is not possible when irradiating with either violet LEDs or Hg lamps.

Scheme 2. Testing the various iodonium ylide derivatives. ¹H NMR yields using HMDSO as the internal standard. [a] Isomerization to **4** occurred during ¹H NMR analysis.

The scope of this cyclopropanation reaction was then evaluated with the aim of demonstrating the viability of our iodonium ylides with various alkenes (Scheme 3).^[23] Ylide **1a** could be efficiently coupled with other styrene derivatives in 1.5 h, and only a modest decrease in yield was observed with the electron-deficient *m*-NO₂ styrene derivative (see **3b**). The Meldrum's acid-derived ylide **1j** was coupled with other additional styrene derivatives bearing either electron-donating or electron-withdrawing groups. The expected cyclopropane products **3d-i** were obtained in yields ranging from 72 to 96%, with the fastest reaction rates occurring with the most electron-rich alkene. Two other styrenes featuring a biphenyl motif were also viable, leading to **3j** and **3k** in 68% and 75% yields after 24 h of irradiation. Additional substitution on the alkene was also tolerated, with both *trans*- and *cis*- β -methylstyrene reacting to give exclusively the *trans*-substituted cyclopropane **3l** in 85% and 81% yield, respectively.^[24,25] Ylide **1j** was then reacted with a variety of non-styrenyl alkenes to varying degrees of success. Cyclopentene and cyclohexene led to cyclopropanes **3m** and **3n** in 28% and 35% yields, while allylbenzene and 1-dodecene respectively gave **3o** in 34% yield and **3p** in 56% yield. In each case, full conversion of **1j** was observed, but we were unable to account for the missing mass balance. Ylide **1i**, which is derived

from barbituric acid, was also effectively coupled with styrene derivatives, affording **3q-s** in 80–84% yields after slightly longer reaction times than those observed for ylide **1j**. Acyclic iodonium ylides were then investigated, and the methyl acetoacetate-derived ylide **1m** was coupled with styrene in 19 h to give **3t** in 82% yield. The dimethyl malonate-derived ylide **1n** was coupled with styrene in 17 h or with *N*-phenylmaleimide in 7 h to furnish **3u** and **3v** in 64 and 46% yields. We observed this ylide **1m** to undergo significant decomposition upon standing at room temperature, which could be overcome by using 20 equiv of styrene, hence affording **3u** in 81% yield over 3 h. In the aim of consuming the ylide at a faster rate, we repeated the experiment using violet LED (λ_{max} 397 nm) irradiation, but **3u** was observed in only 25% yield after 1 h, along with 27% of the corresponding free carbene dimer. We also attempted this reaction using the *o*-iodoanisole-derived iodonium ylide of methyl acetoacetate (**1o**),^[26] which is fully soluble in the reaction mixture, but no improvement was achieved compared with running the reaction as a slurry (64% yield, 22 h). The ensemble of the results of this experimental work, even if not exhaustive, clearly shows that this cyclopropanation reaction

Scheme 3. Mediated cyclopropane synthesis. [a] Using *trans*- β -methylstyrene. [b] Using *cis*- β -methylstyrene. [c] 20 equiv of styrene. [d] Violet LED used. Also 27% of malonate dimer by ¹H NMR. [e] Using soluble ylide **1o**.

between β -dicarbonyl iodonium ylides and alkenes under blue LED irradiation constitutes an efficient and general transformation, which can be performed with either cyclic or acyclic ylides, and with either styrenyl or non-styrenyl alkenes as reaction partners. Reaction rates were generally faster with

increasingly electron-rich alkenes, and reaction yields lower using non-styrenyl alkenes.

Further improvements of these cyclopropanation reactions of iodonium ylides were possible through their in situ synthesis from activated methylene compounds and PhI(OAc)₂.^[9h,10e,11b,27] The rationale behind this possibility is that unstable ylides, which usually decompose upon standing at room temperature, can thus be generated and directly reacted with their alkene partners. To test the feasibility of this experimental option in the context of our photo-initiated process, dimedone (**5**, X = CH₂) or Meldrum's acid (**6**, X = O) were combined base-free with 1 equiv of PhI(OAc)₂ and two equivalents of styrene, then submitted to blue LED irradiation (Scheme 4). Under these conditions, **3a** was generated in 86% yield over 1.5 h, and **3d** in 62% yield over 20 h. Finally, the methyl *p*-nitrophenylacetate derivative **7** was also tested to check if ylides that would be unstable to isolation and purification would also work. The reaction failed in the absence of base, presumably due to the decreased acidity of the methylene protons relative to that of methylene protons of a β -dicarbonyl scaffold. Nevertheless, when 4 equiv of NaH was added, the expected cyclopropane **3w** was obtained in 42% yield, as a 1.7:1 mixture of diastereomers.^[28]

Scheme 4. Testing the tandem in situ ylide formation / cyclopropanation reaction.

Our working hypothesis for the mechanism of this reaction derives from the information gleaned through our computational and experimental studies. As irradiation of an iodonium ylide with blue LEDs does not lead to homodimerization or O–H insertion products with MeOH^[17] (see Table 1), it is improbable that free carbenes are operative in this reaction. Furthermore, we observed an increase of reaction rates when using more electron-rich alkenes, and decreased rates with ylides expressing blue-shifted UV-Vis spectra. We propose that photoexcitation of ylide **1a** leads to reversible formation of the diradical intermediate **A** that may encounter the alkene to forge the diradical **B** (Scheme 5). The higher rates and yields observed with electron-rich alkenes could be attributed to increased stabilization of any benzylic radical character generated in the process. Also, bond rotation within **B** could account for the isomerization required for the formation of **3l** from *cis*- β -methylstyrene (see Schemes 3 and 5). Ring closure would give iodocyclobutane **C**, and reductive elimination of iodobenzene would furnish the cyclopropane end-product **3a**. A drawback of this proposal is that it would require the excited state **A** to be sufficiently long-living to encounter the alkene. Alternatively, given that natural population analysis and adaptive natural partitioning^[19a] both concluded that iodonium ylides hold a full positive charge on the iodine atom in the ground state (eg **1a**), the reaction could instead be initiated by the alkene acting

as a Lewis base toward the iodonium, giving complex **D**.^[10a] If this were to absorb a photon, it could also serve as a precursor of the diradical **B**, which again would need to undergo bond reorganization, relaxation to the ground state, and reductive elimination to generate cyclopropane **3a**.

Scheme 5. Proposed reaction mechanism.

In conclusion, irradiating mixtures of iodonium ylides and alkenes with blue LEDs leads to a mild, metal-free and highly efficient synthesis of cyclopropanes. Computational analysis suggests that photoexcitation leads to a diradical species instead of free carbenes, which accounts for the excellent chemoselectivity observed. Styrenes and non-styrenyl alkenes were both viable, as were both cyclic and acyclic iodonium ylides, showing the potential generality of this new process. Furthermore, a one-pot reaction was developed starting from activated methylene ylide precursors. It is anticipated that this study will open the door to new opportunities for developing blue light-initiated reactions of hypervalent iodine reagents, possibly including catalytic and/or asymmetric variants.

Experimental Section

Into a dry 1 dram (4.0 mL) vial was added iodonium ylide (0.1 mmol, 1 equiv) and CH₃CN (1.0 mL, 0.1 M), and this loaded into photo-reactor #2 and stirred at room temperature for 2 min, giving either a clear solution or a slurry. Styrene (0.023 mL, 0.2 mmol, 2 equiv) was added in one portion and after stirring for 1 min, the blue LED was turned on. The reaction mixture was stirred at room temperature until the reaction was observed to be complete by TLC or NMR analysis. The mixture was then evaporated to dryness and purified by column chromatography, eluting with mixtures of ethyl acetate and hexanes.

Acknowledgements

We would like to acknowledge the Natural Sciences and Engineering Research Council (NSERC) of Canada, the University of Waterloo and the Initiative of Excellence (IdEx) of the University of Bordeaux (Bordeaux-Waterloo joint grant program for emerging cooperation) for funding. This research was enabled in part by support provided by Compute Canada (www.computeCanada.ca). We also thank Dr. Mónica Barra and Dr. Sonny Lee for helpful discussions and instrument access.

Keywords: cyclopropanation • computational analysis • hypervalent iodine • iodonium ylides • LED photochemistry

- [1] a) V. Zhdankin, *Hypervalent Iodine Chemistry: Preparation, Structure, and Synthetic Applications of Polyvalent Iodine Compounds*, John Wiley & Sons, **2013**; b) T. Wirth, *Hypervalent Iodine Chemistry*, Springer-Verlag, Berlin, **2016**; c) A. Yoshimura; V. V. Zhdankin *Chem. Rev.* **2016**, *116*, 3328-3435; d) PATAI'S Chemistry of Functional Groups, I. Marek, B. Olofsson, Z. Rappoport (Ed.), Wiley, New York 2018.
- [2] a) R. D. Richardson; T. Wirth *Angew. Chem., Int. Ed.* **2006**, *45*, 4402-4404; *Angew. Chem.* **2006**, *45*, 4402-4404; b) T. Dohi; Y. Kita *Chem. Commun.* **2009**, 2073-2085; c) F. V. Singh; T. Wirth *Chem. Asian. J.* **2014**, *9*, 950-971; d) A. Flores; E. Cots; J. Bergès; K. Muñoz *Adv. Synth. Catal.* **2019**, *361*, 2-25; e) M. Ochial; K. Miyamoto *Eur. J. Org. Chem.* **2008**, 4229-4239; f) R. Francke *Current Opinion in Electrochemistry* **2019**, *15*, 83-88; g) A. Claraz; G. Masson *Org. Biomol. Chem.* **2018**, *16*, 5386-5402; h) S. Quideau; G. Lyvinec; M. Marguerit; K. Bathany; A. Ozanne-Beaudenon; T. Buffeteau; D. Cavagnat; A. Chénéde *Angew. Chem., Int. Ed.* **2009**, *48*, 4605-4609; *Angew. Chem.* **2009**, *121*, 4675-4679.
- [3] For relevant reviews see a) K. Jia; Y. Chen, *Photochemistry of hypervalent iodine compounds in PATAI'S Chemistry of Functional Groups*, 2018; b) L. Wang; J. Liu *Eur. J. Org. Chem.* **2016**, 1813-1824; For recent examples see c) J.-J. Dai; W.-M. Zhang; Y.-J. Shu; Y.-Y. Sun; J. Xu; Y.-S. Feng; H.-J. Xu *Chem. Commun.* **2016**, 52, 6793-6796; d) H. Huang; G. Zhang; L. Gong; S. Zhang; Y. Chen *J. Am. Chem. Soc.* **2014**, *136*, 2280-2283; e) H. Huang; K. Jia; Y. Chen *Angew. Chem., Int. Ed.* **2015**, *54*, 1881-1884; *Angew. Chem.* **2015**, *127*, 1901-1904; f) G.-X. Li; C. A. Morales-Rivera; Y. Wang; F. Gao; G. He; P. Liu; G. Chen *Chem. Sci.* **2016**, *7*, 6407-6412; g) P. Xu; A. Abdulkader; K. Hu; Y. Cheng; C. Zhu *Chem. Commun.* **2014**, 50, 2308-2310.
- [4] For relevant reviews see a) C. K. Prier; D. A. Rankic; D. W. C. MacMillan *Chem. Rev.* **2013**, *113*, 5322-5363; b) C.-S. Wang; P. H. Dixneuf; J.-F. Soulé *Chem. Rev.* **2018**, *118*, 7532-7585; c) R. C. McAtee; E. J. McClain; C. R. J. Stephenson *Trends in Chemistry* **2019**, *1*, 111-125; d) M. Reckenthäler; A. G. Griesbeck *Adv. Synth. Catal.* **2013**, *355*, 2727-2744; e) N. A. Romero; D. A. Nicewicz *Chem. Rev.* **2016**, *116*, 10075-10166.
- [5] a) J. Tao; T. N. Tuck; G. K. Murphy *Synthesis* **2016**, 48, 772-782; b) J. Tao; C. D. Estrada; G. K. Murphy *Chem. Commun.* **2017**, 53, 9004-9007; c) T. Chidley; G. K. Murphy *Org. Biomol. Chem.* **2018**, *16*, 8486-8490; d) P. E. Ho; J. Tao; G. K. Murphy *Eur. J. Org. Chem.* **2013**, 6540-6544; e) B. A. Laevens; J. Tao; G. K. Murphy *J. Org. Chem.* **2017**, *82*, 11903-11908.
- [6] a) M. S. Yusubov; A. Yoshimura; V. V. Zhdankin *Arkivoc* **2016**, *1*, 342-374; b) S. Chelli; K. Troshin; P. Mayer; S. Lakhdar; A. R. Ofial; H. Mayr *J. Am. Chem. Soc.* **2016**, *138*, 10304-10313.
- [7] a) A. Antos; Y. Elemes; A. Michaelides; J. A. Nyxas; S. Skoulika; L. P. Hadjirapoglou *J. Org. Chem.* **2012**, *77*, 10949-10954; b) E. D. Matveeva; T. A. Podrugina; Y. K. Grishin; A. S. Pavlova; N. S. Zefirov *Russ. J. Org. Chem.* **2007**, *43*, 201-206; c) E. D. Matveeva; T. A. Podrugina; A. S. Pavlova; A. V. Mironov; A. A. Borisenko; R. Gleiter; N. S. Zefirov *J. Org. Chem.* **2009**, *74*, 9428-9432; d) V. V. Zhdankin; O. Maydanovych; J. Herschbach; J. Bruno; E. D. Matveeva; N. S. Zefirov *J. Org. Chem.* **2003**, *68*, 1018-1023; e) M. Ochial; Y. Kitagawa *Tetrahedron Lett.* **1998**, *39*, 5569-5570; f) M. Ochial; Y. Kitagawa; S. Yamamoto *J. Am. Chem. Soc.* **1997**, *119*, 11598-11604.
- [8] a) C. Batsila; E. P. Gogonas; G. Kostakis; L. P. Hadjirapoglou *Org. Lett.* **2003**, *5*, 1511-1514; b) W. Adam; E. P. Gogonas; L. P. Hadjirapoglou *Tetrahedron* **2003**, *59*, 7929-7934; c) M. B. Camacho; A. E. Clark; T. A. Liebrecht; J. P. DeLuca *J. Am. Chem. Soc.* **2000**, *122*, 5210-5211; d) R. M. Moriarty; E. J. May; O. Prakash *Tetrahedron Lett.* **1997**, *38*, 4333-4336.
- [9] a) Y. R. Lee; S. H. Yoon *Synth. Commun.* **2006**, *36*, 1941-1951; b) C. J. Zhu; A. Yoshimura; P. Solntsev; L. Ji; Y. Y. Wei; V. N. Nemykin; V. V. Zhdankin *Chem. Commun.* **2012**, 48, 10108-10110; c) E. P. Gogonas; L. P. Hadjirapoglou *Tetrahedron Lett.* **2000**, *41*, 9299-9303; d) A. Asouti; L. P. Hadjirapoglou *Tetrahedron Lett.* **1998**, *39*, 9073-9076; e) P. Muller; D. Fernandez *Helv. Chim. Acta* **1995**, *78*, 947-958; f) S. Spyroudis; P. Tarantili *J. Org. Chem.* **1993**, *58*, 4885-4889; g) L. Hadjirapoglou; K. Schank *Tetrahedron Lett.* **1989**, *30*, 6673-6676; h) D. Kalpogiannaki; C.-I. Martini; A. Nikopoulou; J. A. Nyxas; V. Pantazi; L. P. Hadjirapoglou *Tetrahedron* **2013**, *69*, 1566-1575; i) W. Adam; E. P. Gogonas; L. P. Hadjirapoglou *J. Org. Chem.* **2003**, *68*, 9155-9158; j) S. Kalogiannis; S. Spyroudis *J. Org. Chem.* **1990**, *55*, 5041-5044.
- [10] a) A. E. Bosnidou; D. Kalpogiannaki; S. Karanestora; J. A. Nixas; L. P. Hadjirapoglou *J. Org. Chem.* **2015**, *80*, 1279-1283; b) S. R. Goudreau; D. Marcoux; A. B. Charette *J. Org. Chem.* **2009**, *74*, 470-473; c) R. M. Moriarty; O. Prakash; R. K. Vaid; L. Zhao *J. Am. Chem. Soc.* **1989**, *111*, 6443-6444; d) R. M. Moriarty; S. Tyagi; M. Kinch *Tetrahedron* **2010**, *66*, 5801-5810; e) R. P. Wurz; A. B. Charette *Org. Lett.* **2005**, *7*, 2313-2316; f) P. Müller; C. Boléa *Helv. Chim. Acta* **2001**, *84*, 1093-1111; g) R. M. Moriarty; J. Kim; L. Guo *Tetrahedron Lett.* **1993**, *34*, 4129-4132; h) P. Müller; Y. Allenbach; E. Robert *Tetrahedron: Asymmetry* **2003**, *14*, 779-785; i) H. T. Borge; T. Hansen *Synlett* **2007**, 55-58.

- [11] a) S. Mo; X. Li; J. Xu *J. Org. Chem.* **2014**, *79*, 9186-9195; b) G. K. Murphy; F. G. West *Org. Lett.* **2006**, *8*, 4359-4361.
- [12] For relevant examples see a) P. Muller; Y. F. Allenbach; S. Chappellet; A. Ghanem *Synthesis* **2006**, 1689-1696; b) B. Moreau; A. B. Charette *J. Am. Chem. Soc.* **2005**, *127*, 18014-18015; c) B. Moreau; D. Alberico; V. N. G. Lindsay; A. B. Charette *Tetrahedron* **2012**, *68*, 3487-3496; d) B. Xu; U. K. Tambar *Angew. Chem., Int. Ed.* **2017**, *56*, 9868-9871; *Angew. Chem.* **2017**, *129*, 10000-10003.
- [13] For reviews on the chemistry of doubly activated cyclopropanes see a) H. U. Reissig; R. Zimmer *Chem. Rev.* **2003**, *103*, 1151-1196; b) M. Yu; B. L. Pagenkopf *Tetrahedron* **2005**, *61*, 321-347; c) M. Rubin; M. Rubina; V. Gevorgyan *Chem. Rev.* **2007**, *107*, 3117-3179; d) C. A. Carson; M. A. Kerr *Chem. Soc. Rev.* **2009**, *38*, 3051-3060; e) H. Nambu; M. Fukumoto; W. Hirota; T. Yakura *Org. Lett.* **2014**, *16*, 4012-4015; f) M. A. Cavitt; L. H. Phun; S. France *Chem. Soc. Rev.* **2014**, *43*, 804-818; g) C.-H. Lin; B. Plietker *Isr. J. Chem.* **2016**, *56*, 409-416; h) H. Nambu; Y. Onuki; N. Ono; T. Yakura *Adv. Synth. Catal.* **2018**, *360*, 2938-2944.
- [14] a) R. Ballini; A. Palmieri; D. Fiorini *Arkivoc* **2007**, 172-194; b) R. P. Wurz; A. B. Charette *Org. Lett.* **2003**, *5*, 2327-2329; c) O. Lifchits; A. B. Charette *Org. Lett.* **2008**, *10*, 2809-2812.
- [15] a) E. D. Matveeva; T. A. Podrugina; A. S. Pavlova; A. V. Mironov; R. Gleiter; N. S. Zefirov *Eur. J. Org. Chem.* **2009**, 2323-2327; b) E. D. Matveeva; T. A. Podrugina; M. A. Taranova; A. A. Borisenko; A. V. Mironov; R. Gleiter; N. S. Zefirov *J. Org. Chem.* **2011**, *76*, 566-572.
- [16] a) M. Papadopolou; S. Spyroudis; A. Varvoglis *J. Org. Chem.* **1985**, *50*, 1509-1511; b) E. Malamidoli-Xenikaki; S. Spyroudis *Synlett* **2008**, 2725-2740.
- [17] I. D. Jurberg; H. M. L. Davies *Chem. Sci.* **2018**, *9*, 5112-5118.
- [18] For further work on blue LED-mediated reactions of phenyldiazoacetate derivatives see a) R. Hommelsheim; Y. Guo; Z. Yang; C. Empel; R. M. Koenigs *Angew. Chem., Int. Ed.* **2019**, *58*, 1203-1207; *Angew. Chem.* **2019**, *131*, 1216-1220; b) S. Jana; R. M. Koenigs *Asian J. Org. Chem.* **2019**, *8*, 683-686; c) Z. Yang; Y. Guo; R. M. Koenigs *Chem. Eur. J.* **2019**, *25*, 6703-6706; d) F. He; R. M. Koenigs *Chem. Commun.* **2019**, 55, 4881-4884; e) J. Yang; J. Wang; H. Huang; G. Qin; Y. Jiang; T. Xiao *Org. Lett.* **2019**, *21*, 2654-2657. For a recent example of LED-mediated generation of carbenes from pyridotriazoles see f) Z. Zhang; D. Yadagiri; V. Gevorgyan *Chem. Sci.* **2019**, *10*, 1039/C9SC02448D.
- [19] a) A. S. Ivanov; I. A. Popov; A. I. Boldyrev; V. V. Zhdankin *Angew. Chem., Int. Ed.* **2014**, *53*, 9617-9621; *Angew. Chem.* **2014**, *126*, 9771-9775.
- [20] a) G. S. Sinclair; R. Tran; J. Tao; W. S. Hopkins; G. K. Murphy *Eur. J. Org. Chem.* **2016**, 4603-4606; b) Z. Zhao; K. C. Y. Ma; C. Y. Legault; G. K. Murphy *Chem. Eur. J.* **2019**, DOI: 10.1002/chem.201804786.
- [21] a) W. J. Hehre; R. Ditchfield; J. A. Pople *J. Chem. Phys.* **1972**, *56*, 2257-2261; b) P. C. Hariharan; J. A. Pople *Theor. Chim. Acta* **1973**, *28*, 213-222.
- [22] a) A. E. Reed; R. B. Weinstock; F. Weinhold *J. Chem. Phys.* **1985**, *83*, 735-746; b) E. D. Glendening; C. R. Landis; F. Weinhold *J. Comput. Chem.* **2013**, *34*, 1429-1437.
- [23] The products of ketoester-derived ylide **1i** underwent rapid conversion to the corresponding dihydrofuran upon purification, so no additional cyclopropanation examples are provided. Cyclopropanations attempted with disulfonyl iodonium ylide **1p** were unsuccessful, as this was inert under the reaction conditions.
- [24] Alkene photoisomerization was not responsible for the stereochemical outcome observed with **3i**. Irradiating *cis*- β -methylstyrene with blue light for 24 h failed to produce any quantity of the *trans*-isomer observable by ¹H NMR.
- [25] There are conflicting literature reports as to whether the characterization data recorded for **3i** corresponds to *syn*- or *anti*-cyclopropane isomer. For its identification as the *syn*-isomer see Y.-R. Lee; J.-H. Choi *Bull. Korean Chem. Soc.* **2006**, *27*, 503-507. For its identification as the *anti*-isomer see Ref. 10h. The recorded crystal structure unambiguously shows compound **3i** to be the *anti*-isomer.
- [26] a) V. V. Zhdankin; J. D. Protasiewicz *Coord. Chem. Rev.* **2014**, *275*, 54-62; b) C. J. Zhu; A. Yoshimura; L. Ji; Y. Y. Wei; V. N. Nemykin; V. V. Zhdankin *Org. Lett.* **2012**, *14*, 3170-3173.
- [27] a) J. Vaitla; K. H. Hopmann; A. Bayer *Org. Lett.* **2017**, *19*, 6688-6691; b) H. T. Bonge; T. Hansen *Tetrahedron Lett.* **2008**, *49*, 57-61.
- [28] a) To rule out a radical mechanism involving photoinitiated decomposition of PhI(OAc)₂ by blue light, it was irradiated for 4 h, after which quantitative ¹H NMR showed it to be 94% unchanged. For an example of radical reactions of iodonium ylides see a) M. Hartmann; Y. Li; C. Mück-Lichtenfeld; A. Studer *Chem. Eur. J.* **2016**, *22*, 3485-3490; For a recent example of visible light-initiated reactions of PhI(OAc)₂ see b) C. F. Meyer; S. M. Hell; A. Misale; A. A. Trabanco; V. Gouverneur *Angew. Chem., Int. Ed.* **2019**, *58*, 8829-8833; *Angew. Chem.* **2019**, *131*, 8921-8925.