



The pneumococcal alternative sigma factor σ X mediates competence shut-off at the cell pole

Calum Johnston, Anne-Lise Soulet, Matthieu Berge, Marc Prudhomme, David de Lemos, Patrice Polard

► To cite this version:

Calum Johnston, Anne-Lise Soulet, Matthieu Berge, Marc Prudhomme, David de Lemos, et al.. The pneumococcal alternative sigma factor σ X mediates competence shut-off at the cell pole. 2020. hal-03031117

HAL Id: hal-03031117

<https://hal.science/hal-03031117>

Preprint submitted on 30 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The pneumococcal alternative sigma factor σ^X mediates competence shut-off at the cell pole

Calum JOHNSTON^{1,2}, Anne-Lise SOULET^{1,2}, Matthieu BERGE^{1,2,3}, Marc PRUDHOMME^{1,2}, David DE LEMOS^{1,2}, Patrice POLARD^{1,2}.

1. Laboratoire de Microbiologie et Génétique Moléculaires, UMR5100, Centre de Biologie Intégrative (CBI), Centre Nationale de la Recherche Scientifique (CNRS), Toulouse, France.

2. Université Paul Sabatier (Toulouse III), Toulouse, France.

3. Dept. Microbiology and Molecular Medicine, Institute of Genetics & Genomics in Geneva (iGE3), Faculty of Medicine, University of Geneva, Geneva, Switzerland.

Keywords

Competence regulation, competence shut-off, alternative sigma factor, σ^X , DprA.

Summary

Bacterial competence for genetic transformation is a well-known species-specific differentiation program driving genome plasticity, antibiotic resistance and virulence in many pathogens. How competence regulation is spatiotemporally integrated in the cell is ill-defined. Here, we unraveled the localization dynamics of the key regulators that master the two intertwined transcription waves controlling competence in *Streptococcus pneumoniae*. The first wave relies on a stress-inducible phosphorelay system, made up of the ComD and ComE proteins, and the second is directed by an alternative sigma factor, σ^x , which includes in its regulon the DprA protein that turns off competence through interaction with phosphorylated ComE. Remarkably, we found that ComD, σ^x and DprA stably co-localize at a single cell pole over the competence period. Furthermore, we provide evidence that σ^x is necessary and sufficient to mediate DprA polar accumulation next to ComD. Finally, we show that through this protein targeting function, σ^x is actively involved in the timely shut-off of the competence cycle, hence preserving cell fitness. Altogether, this study unveils an unprecedented role for a bacterial transcription σ factor in spatially coordinating the negative feedback loop of its own genetic circuit.

46 Introduction

47 In bacteria, sigma (σ) factors are essential transcription effectors that direct the RNA polymerase
48 to and activate RNA synthesis at specific genes promoters. All bacterial species encode a single, highly
49 conserved σ factor that drives the expression of house-keeping genes essential for vegetative growth
50 and cell homeostasis. In addition, many bacteria encode a variable set of alternative σ factors that
51 control specific regulons, providing appropriate properties to the cells in response to various stimuli.
52 These alternative σ factors play pivotal roles in the multifaceted lifestyles of bacteria. They trigger
53 specific developmental programs, such as sporulation or biofilm formation, as well as adapted
54 responses to multiple types of stress and virulence in some pathogenic species (Kazmierczak et al.,
55 2005). How these alternative σ factors are activated in the cell has been extensively studied, revealing
56 multiple mechanisms underlying their finely tuned regulation (Österberg et al., 2011a). However, how
57 these mechanisms are orchestrated spatiotemporally within the cell remains poorly understood.

58 The human pathogen *Streptococcus pneumoniae* (the pneumococcus) possesses a unique
59 alternative σ factor σ^x (Lee and Morrison, 1999). It is key to the regulatory circuit controlling the
60 transient differentiation state of competence. Pneumococcal competence is induced in response to
61 multiple types of stresses, such as antibiotic exposure (Prudhomme et al., 2006; Slager et al., 2014).
62 This induction modifies the transcriptional expression of up to 17% of genes (Aprianto et al., 2018a;
63 Dagkessamanskaia et al., 2004a; Peterson et al., 2004; Slager et al., 2019). Competence is a key feature
64 in the lifestyle of pneumococci as it promotes natural transformation, a horizontal gene transfer
65 process widespread in bacteria that facilitates adaptation by acquisition of new genetic traits (Johnston
66 et al., 2014). In addition, pneumococcal competence development provides the cells with the ability
67 to attack non-competent cells, a scavenging property defined as fratricide (Claverys and Håvarstein,
68 2007), is involved in biofilm formation (Aggarwal et al., 2018; Vidal et al., 2013) and virulence (Johnston
69 et al., 2018; Lin et al., 2016; Lin and Lau, 2019; Zhu et al., 2015).

70 Pneumococcal competence induction is primarily regulated by a positive feedback loop
71 involving the genes encoded by the *comAB* and *comCDE* operons (Figure 1A). The *comC* gene codes for

a peptide pheromone coordinating competence development within the growing cell population. This peptide, accordingly named CSP (Competence Stimulating Peptide), is secreted by the dedicated ComAB transporter (Hui et al., 1995). After export, it promotes autophosphorylation of the membrane-bound two-component system (TCS) histidine kinase (HK) ComD, which in turn phosphorylates its cognate intracellular response regulator (RR) ComE (Figure 1A). Phosphorylated ComE (ComE~P) specifically induces the expression of 25 genes, which include the *comAB* and *comCDE* operons, generating a positive feedback loop that controls competence development. Conversely, unphosphorylated ComE acts as repressor of its own regulon, the expression of which is thus modulated by the ComE/ComE~P ratio (Martin et al., 2013). The ComE regulon includes two identical genes encoding σ^X , named *comX1* and *comX2* (Lee and Morrison, 1999). The σ^X regulon comprises ~60 genes, with ~20 involved in natural transformation (Claverys et al., 2006; Peterson et al., 2004), 5 in fratricide (Claverys and Håvarstein, 2007) but the majority having undefined roles. The reason why the σ^X -encoding gene is duplicated is unknown, the inactivation of one of them having no impact on transformation (Lee and Morrison, 1999). To fully activate transcription, σ^X needs to be assisted by ComW, another protein whose production is controlled by ComE~P (Luo et al., 2004). ComW is proposed to help σ^X association with the RNA polymerase at promoter sequences presenting the consensual 8 bp *cin* box motif (Peterson et al., 2004; Sung and Morrison, 2005). Altogether, ComE~P and σ^X trigger two successive waves of competence (*com*) gene transcription, commonly referred to as early and late, respectively. Importantly, competence shut-off is mediated by the late *com* protein DprA (Mirouze et al., 2013; Weng et al., 2013), which directly interacts with ComE~P to turn-off ComE~P-dependent transcription (Mirouze et al., 2013). In addition to defining the negative feedback loop of the pneumococcal competence regulatory circuit, DprA also plays a crucial, conserved role in transformation by mediating RecA polymerization onto transforming ssDNA to facilitate homologous recombination (Mortier-Barrière et al., 2007; Cheruel et al., 2012) (Figure 1A). Over 8,000 molecules of DprA are produced per competent cell (Mirouze et al., 2013). Although only ~300-600 molecules were required for optimal transformation, full expression of *dprA* was required for optimal

competence shut-off (Johnston et al., 2018). Uncontrolled competence induction in cells lacking DprA results in a large *in vitro* growth defect, and high cellular levels of DprA thus maintain the fitness of the competent population and of resulting transformants (Johnston et al., 2018). In addition, inactivation of *dprA* was shown to be highly detrimental for development of pneumococcal infection, dependent on the ability of cells to develop competence (Lin and Lau, 2019; Zhu et al., 2015). Together, these studies showed that the DprA-mediated shut-off of pneumococcal competence is key for pneumococcal cell fitness.

A hallmark of pneumococcal competence is its tight temporal window, which lasts less than 30 minutes in actively dividing cells (Alloing et al., 1998; Håvarstein et al., 1995). How this regulation is coordinated within the cell remains unknown. Here, we studied the choreography of pneumococcal competence induction and shut-off at the single cell level by tracking the spatiotemporal localization of the main effectors of these processes, DprA, σ^X , ComW, ComD, ComE and exogenous CSP. Remarkably, DprA, σ^X , ComD, CSP and to some extent ComE were found to colocalize at a single cell pole during competence. This study revealed that the entire pneumococcal competence cycle occurs at cell pole, from its induction triggered by ComD, ComE and CSP to its shut-off mediated by DprA and assisted by σ^X . In this regulatory mechanism, σ^X is found to exert an unprecedented role for a σ factor. In addition to directing the transcription of the *dprA* gene, σ^X associates with and anchors DprA at the same cellular pole where ComD and CSP are located, allowing this repressor to interact with newly activated ComE~P and promoting timely extinction of the whole transcriptional regulatory circuit of competence.

Results

DprA displays a polar localization in competent cells, which correlates with competence shut-off.

To investigate the localization of DprA during competence in live cells, we used a fluorescent fusion protein DprA-GFP, produced from the native *dprA* locus (Figure S1A). DprA-GFP was synthesized

during competence and remained stable up to 90 minutes after induction (Figure 1B), similarly to wildtype DprA (Mirouze et al., 2013), without degradation (Figure S1B). A strain possessing DprA-GFP was almost fully functional in transformation (Figure S1C), but partially altered in competence shut-off (Figure S1D). In pneumococcal cells induced with exogenous CSP, the peak of competence induction occurs 15-20 minutes after induction. 15 minutes after competence induction, DprA-GFP showed a diffuse cytoplasmic localization, punctuated by discrete foci of varying intensity (Figure 1C). The distribution and localization of DprA-GFP foci was analyzed by MicrobeJ (Ducret et al., 2016), with results presented as focus density maps ordered by cell length. Spots represent the localization of a DprA-GFP focus on a representative half pneumococcal cell, while spot colour represents density of foci at a particular cellular location. DprA-GFP foci were present in 70% of cells, predominantly at a single cell pole (Figure 1DE). To ascertain whether the polar localization of DprA-GFP was due to the GFP tag or represented functional localization, we carried out immunofluorescence microscopy with competent cells possessing wildtype DprA using α -DprA antibodies. Results showed that native DprA exhibited a similar accumulation pattern as the DprA-GFP foci upon competence induction (Figure 1F).

Next, to explore the relation between these foci and the dual role of DprA in transformation and competence regulation, we investigated focus formation in cells possessing a previously published mutation in DprA impairing its dimerization (DprA^{AR}) (Quevillon-Cheruel et al., 2012). This mutant strongly affected both transformation and competence shut-off (Mirouze et al., 2013; Quevillon-Cheruel et al., 2012). Results showed that 15 minutes after competence induction, DprA^{AR}-GFP did not form foci, despite being produced at wildtype levels (Figure S1E). DprA thus accumulates at the cell pole during competence, dependent on its ability to dimerize. To explore whether the polar foci of DprA were involved in its role in transformation, or competence shut-off, we began by investigating how a *dprA*^{QNQ} mutation, specifically abrogating the interaction between DprA and RecA and thus affecting transformation (Quevillon-Cheruel et al., 2012), affected the localization of DprA-GFP. 15 minutes after competence induction, the DprA^{QNQ}-GFP mutant formed polar foci at wildtype levels (Figure S1F). In addition, the inactivation of *comEC*, encoding for an essential protein of the

transmembrane DNA entry pore (Pestova and Morrison, 1998), or *recA*, encoding the recombinase with which DprA interacts during transformation (Mortier-Barrière et al., 2007), did not alter the frequency or localization of DprA-GFP foci (Figure S1GH). Altogether, these results suggested that the polar foci of DprA-GFP were not related to the DNA entry or recombination steps of transformation but could be linked to competence shut-off.

We recently reported that optimal competence shut-off relies on the maximal cellular concentration of DprA (~8000 molecules), but this level could be reduced by 10 fold while still maintaining wildtype frequency of transformation (Johnston et al., 2018). This conclusion was obtained by expressing *dprA* under the control of the IPTG-inducible P_{lac} promoter (CEP_{lac} -*dprA*), which enables the modulation of the cellular concentration of DprA by varying IPTG concentration in the growth medium. Here, we reproduced these experiments with the DprA-GFP fusion, to test whether its concentration correlates with the formation of polar foci in competent cells. The expression, transformation and competence profiles of a *dprA*⁻ mutant strain harbouring the ectopic CEP_{lac} -*dprA*-*gfp* construct in varying concentrations of IPTG (Figure S2ABC) were equivalent to those reported previously for CEP_{lac} -*dprA* (Johnston et al., 2018). Notably, a steady decrease in DprA-GFP foci was observed as IPTG was reduced (Figure 2AB). When comparing the cellular localization of DprA-GFP foci, a sharp reduction in the proportion of polar foci was observed as IPTG was reduced, with most of the remaining foci observed at midcell and appearing weaker in intensity (Figure 2AC). This shift correlated with a progressive loss of competence shut-off (Figure S2C), presenting a strong link between the presence of polar DprA-GFP foci and the shut-off of pneumococcal competence. Altogether, these results strongly support the notion that the polar foci of DprA-GFP represent the subcellular site where DprA mediates competence shut-off.

Finally, to further explore the temporal dynamics of these DprA-GFP foci, their distribution within competent cells was analyzed over the competence period and beyond. The results are presented in Figure 3A as focus density maps ordered by cell length. The number of cells with foci, as

well as their intensity, was found to increase gradually to reach a maximum of 74% at 30 minutes after competence induction (Figure 3A), with the majority of cells possessing a single focus that persisted long after induction (Figure 3B). Notably, the DprA-GFP foci localization pattern rapidly evolved from a central position to a single cell pole (Figure 3C). DprA-GFP foci were not observed in a particular cell type, with found in small, large or constricted cells throughout competence (Figure 3C). Finally, tracking DprA-GFP foci formed in the cells after 10 minutes of competence induction by time-lapse microscopy showed that once generated, they remained static over 20 minutes (Figure S11J, Movie 1). In conclusion, DprA-GFP forms discrete and static polar foci during competence, with most cells possessing a single focus. This polar localization of DprA correlates with its regulatory role in competent shut-off.

The polar localization of DprA-GFP requires induction of the late com genes

Transcriptional expression of *dprA* is only detected during competence (Aprianto et al., 2018a; Dagkessamanskaia et al., 2004a; Peterson et al., 2004). To explore whether a competence-specific factor was required for the formation of polar DprA-GFP foci during competence, *dprA-gfp* was ectopically expressed from a promoter inducible by the BIP peptide in *dprA*⁻ cells. This BIP-derived induction mimics rapid, strong induction by CSP during competence (Johnston et al., 2016). These cells were found to produce stable DprA-GFP after exposure to BIP, and upon addition of CSP to the growth medium, to transform at wild-type levels and to partially shut-off competence (Figure S3). However, BIP-induced production of DprA-GFP in the absence of CSP resulted in the formation of weak, barely detectable polar foci in only 10% of non-competent cells (Figure 4AB). In comparison, 47% of cells producing DprA-GFP during competence formed bright polar foci (Figure 4AB). This stark increase in DprA-GFP foci showed that a competence-specific factor was crucial for their formation and anchoring at the cell pole. To explore whether the competence-specific factor needed for the polar localization of DprA-GFP in competent cells was part of the early or late *com* regulons, we generated two

constructs allowing us to artificially control DprA-GFP expression and either one or the other of these two connected regulons (Figure 4CD and Supplementary methods). Observation of DprA-GFP in conditions where only late *com* genes were induced revealed the presence of polar foci at wildtype levels (Figure 4EF). Conversely, no foci were observed when DprA-GFP was ectopically expressed with only the early *com* genes (Figure 4E), showing that late *com* regulon expression was required for the polar accumulation of DprA-GFP.

The polar localization of DprA-GFP depends on the alternative sigma factor σ^X

The late *com* regulon is comprised of 62 genes, organized in 18 operons (Claverys et al., 2006; Dagkessamanskaia et al., 2004b; Peterson et al., 2004). To identify the hypothetical late *com* gene product needed for DprA localization at the cell pole, the *cin* boxes that define the late *com* promoters were individually mutated, generating a panel of 18 mutant strains, each lacking the ability to induce a specific late *com* operon. The inactivation was validated by comparing transformation efficiency in three strains, where *cin* box inactivation mirrored gene knockout levels (Table S1, and Supplementary methods). Visualization of the red fluorescent fusion DprA-mKate2 showed that in all 18 mutants, DprA-mKate2 formed foci at levels and localization comparable to wildtype (Table S2). This result contrasted with our previous result (Figure 4E), which suggested that expression of the late *com* regulon was required for formation of polar DprA-GFP foci, causing us to revisit our interpretation of Figure 4C-F. In fact, to express only the early *com* regulon, we inactivated *comX1*, *comX2* and *comW* (Figure 4C), so this strain produced only early *com* proteins, except σ^X and ComW, and lacked DprA-GFP foci (Figure 4E). Conversely, to express only the late *com* regulon, we ectopically expressed *comX* and *comW* (Figure 4D), so this strain produced the late *com* regulon but also the early *com* proteins σ^X and ComW and displayed polar DprA-GFP foci at wildtype levels (Figure 4EF). This led us to consider that the only proteins whose presence correlated directly with the presence of polar DprA-GFP foci were thus σ^X and ComW.

To first investigate whether ComW played a role in the formation of polar DprA-GFP foci, the *comW* gene was inactivated in a strain possessing a *rpoD*^{A171V} mutation, enabling σ^X -RNA polymerase interaction and resulting in late *com* regulon expression in the absence of ComW (Tovpeko et al., 2016). DprA-GFP expressed from the native locus still formed polar foci in this strain at levels comparable to the wildtype strain (Figure S4A). ComW was thus not required for the formation of DprA-GFP foci. In light of this, the only remaining candidate whose presence in competent cells correlated directly with formation of polar DprA-GFP foci was σ^X . Thus, to determine if σ^X alone was necessary and sufficient to localize DprA-GFP to the cell poles, both *comX* and *dprA-gfp* or *dprA-gfp* alone were expressed in non-competent *rpoD*^{wt} cells. Western blot analysis using α -SsbB antibodies indicated that the late *com* regulon was weakly induced when σ^X was ectopically produced in the absence of *comW* (Figure S4B). Cells producing DprA-GFP alone showed polar DprA-GFP foci in 10% of cells (Figure 5A). In contrast, DprA-GFP foci were formed in 37% of cells when σ^X was also produced in non-competent cells (Figure 5A). Importantly, induction of competence in both of these strains resulted in similar foci numbers (Figure S4D). Altogether, this result suggested that σ^X alone was sufficient to stimulate polar foci of DprA-GFP, highlighting an unexpected role for this early competence σ factor only known to act in concert with ComW to induce late competence gene transcription.

σ^X mediates the localization of DprA at the cell pole of competent cells

To investigate how σ^X could be involved in the polar localization of DprA-GFP, we explored how it localized in competent cells. To this end, we generated a *comX1-gfp* construct at the native *comX1* locus combined with a wildtype *comX2* gene (Figure S5 and Supplementary results). Remarkably, σ^X -GFP formed bright polar foci 15 minutes after competence induction (Figure 5A), reminiscent of those formed by DprA-GFP (Figure 1C). A time-course experiment after competence induction showed that σ^X -GFP localized to the cell pole as soon as 4 minutes after CSP addition (Figure 5B), when DprA-GFP forms weak foci at midcell (Figure 3A). In contrast, a ComW-GFP fusion protein displayed a diffuse

cytoplasmic localization in the majority of competent cells, with only 7% of cells possessing weak foci at the cell poles (Figure S5A and Supplementary results). Thus, σ^X -GFP localizes to the cell pole without its partner in transcriptional activation ComW, despite the fact that σ^X is an alternative sigma factor directing RNA polymerase to specific promoters on the chromosome.

Analysis of σ^X -GFP foci distribution showed that they are detected in up to 51% of cells 10 minutes after competence induction, with most cells possessing a single focus (Figure 5BC). The number of cells with foci decreased steadily after this point (Figure 5BCD), contrasting with polar DprA-GFP which remained stable over 60 minutes after induction. Importantly, σ^X -GFP continued to form polar foci in a strain lacking *dprA* (Figure 5E), showing that σ^X does not depend on DprA for its localization. Together, these results strongly supported the notion that σ^X promotes the targeting and assembly of DprA-GFP foci at the cell pole. To further explore this hypothesis, we co-expressed DprA-mTurquoise and σ^X -YFP fluorescent fusions in the same cells and found that 86% of DprA-mTurquoise foci colocalized with σ^X -YFP foci (Figure 5F). DprA and σ^X are thus present at the same pole of the cell at the same time and the polar accumulation of DprA molecules in competent cells depends on σ^X . These results suggested that σ^X could interact with DprA to anchor it to the pole of competent cells. The potential interaction between σ^X and DprA was tested in live competent pneumococcal cells in pull-down experiments. To achieve this, we used GFP-TRAP magnetic beads (Chromotek) to purify σ^X -GFP from competent cells expressing either wildtype DprA or, as a control, the DprA^{AR} mutant that did not accumulate at the cell pole (Figure S1). Results showed that wildtype DprA co-purified with σ^X -GFP from competent cells extracts, but DprA^{AR} did not, revealing that σ^X and DprA interact in live competent pneumococci (Figure 5H). Taken together, these findings reveal that σ^X is responsible for the accumulation of DprA at the cell pole during competence.

Pneumococcal competence induction occurs at the cell pole

We have shown that DprA localization at a single cell pole correlates with the shut-off of competence (Figure 2). Since DprA interacts directly with ComE~P to mediate competence shut-off (Mirouze et al., 2013), this raised the question of the subcellular localization of ComD and ComE, which define a TCS controlling competence regulation. To first explore the localization of ComE in competent cells, a strain was generated expressing a functional *comE-gfp* fluorescent fusion at the native *comE* locus (Figure S6 and Supplementary results). In competent cells, ComE-GFP formed patches around the periphery of the cell, often at the cell pole (Figure 6AB and S7A). A time-lapse experiment showed that these patches were dynamic, navigating around the cell membrane over time (Movie 2). Next, to explore the sub-cellular localization of ComD, we generated a strain expressing *gfp-comD* at the native *comD* locus (Figure S6AC and Supplementary results). The resulting GFP-ComD fusion displayed partial functionality in competence induction and transformation (Figure S6DE). In contrast to ComE-GFP, GFP-ComD formed distinct polar foci of varying intensity in 57% of cells (Figure 6AB), a localization pattern reminiscent of those observed with DprA-GFP and σ^X -GFP (Figures 1 and 5, respectively). Since GFP-ComD was not fully functional, we also analyzed the localization of a synthetic fluorescent exogenous CSP peptide (CSP-HF, Figure S6F) in parallel, to track its interaction with ComD at the cell surface. This fluorescent peptide was found to accumulate at a single cell pole in the majority of competent, wildtype cells (Figure 6AB). In addition, this accumulation was dependent on the presence of ComD (Figure 6A), showing that the polar accumulation of the partially functional GFP-ComD fusion represented a functional subcellular localization during competence. In addition, most cells with ComD-GFP foci possessed a single focus (Figure 6C), which persisted after the shut-off of competence (Figure S7B). Altogether, these findings revealed that activation of the positive feedback loop of competence triggered by CSP interaction with ComD to phosphorylate ComE occurs at the cell pole. Since a single pole is mainly targeted by ComD, this also raised the question of its co-localization with ComX and DprA.

DprA colocalizes with ComD and CSP during competence

To explore the hypothesis that DprA accumulates at the same pole as ComD to mediate competence shut-off, we performed colocalization analyses of DprA with ComE, ComD or CSP respectively, in the same competent cells. Although ComE-YFP foci were dynamic and their localization difficult to analyze, 20% of DprA-mTurquoise foci were nonetheless found to colocalize with ComE-YFP at the cell pole, showing that these proteins can be found at the same pole in the same cells (Figure 6D). In contrast, 76% of DprA-mTurquoise foci colocalized with YFP-ComD (Figure 6E), showing that these foci form at the same pole in the majority of cells. In addition, 73% of DprA-GFP foci colocalized with CSP-HF (Figure 6F). Taken together, these results showed that DprA colocalizes strongly with ComD, generally at one cell pole during pneumococcal competence, further suggesting that the localization of DprA to this cellular location, mediated by σ^X , facilitates pneumococcal competence shut-off. The localization of DprA at the cell pole where ComD appears to interact with CSP means that DprA is present at the time and place that neophosphorylated ComE~P is produced, allowing DprA to interact with the activated regulator at the cell pole and prevent it from accessing its genomic targets, facilitating shut-off.

Two copies of comX are required for optimal competence shut-off

A distinct hallmark of the single pneumococcal alternative sigma factor σ^X of *S. pneumoniae* is that it is produced from two distinct and strictly identical genes, at two distinct loci in the pneumococcal genome, known as *comX1* and *comX2* (Lee and Morrison, 1999). However, inactivation of either of these genes had no effect on the efficiency of transformation as previously reported (Lee and Morrison, 1999) (Figure 7A), suggesting that the expression of either *comX1* or *comX2* is sufficient to induce the late *com* regulon to a level ensuring optimal transformation. Having established that σ^X plays a second key role in competence shut-off, we explored whether this role required both *comX* genes. Inactivation of either *comX* gene not only slightly reduced the peak of late *com* gene expression

but also markedly delayed the rate of competence shut-off (Figure 7B), revealing that reducing the cellular level of σ^X impacts competence shut-off efficiency, presumably because less σ^X is present to promote accumulation of DprA at the cell poles. In addition, this unregulated competence shut-off of single *comX* mutants is accompanied by a reduced growth rate of the cell population, consistent with an alteration of cell fitness linked to altered competence shut-off (Johnston et al., 2018). In conclusion, this finding suggests that pneumococci possess two copies of *comX* to optimize DprA-mediated competence shut-off and maintain the fitness of competent cells.

Pre-competence expression of DprA and σ^X antagonizes competence induction

This study has uncovered a new functional role of pneumococcal σ^X in facilitating DprA-mediated inactivation of ComE-P at the cell pole. To obtain further proof that polar DprA plays a role in competence shut-off, we reasoned that if σ^X localizes DprA to the cell pole to allow it to interact with neophosphorylated ComE~P, then early ectopic expression of DprA and σ^X should antagonize competence induction by interfering with early *com* regulon induction. To test this hypothesis, we expressed DprA alone or both DprA and σ^X prior to CSP addition to the growth medium (Figure 7C), referred hereafter as a pre-competence expression. Pre-competence production of DprA alone did not affect competence induction as shown by monitoring luciferase controlled by an early *com* promoter (Figure 7D). In these conditions, minimal amounts of DprA accumulated at the cell pole (Figure 5A). However, pre-competence expression of both DprA and σ^X resulted in a significantly slower induction of competence (Figure 7E). This suggested that σ^X -mediated pre-localization of DprA to the cell poles in pre-competent cells, as shown in Figure 4G, markedly antagonized CSP induction of competence by allowing DprA to interact with and inactivate neophosphorylated ComE~P. Inactivation of *comW* in this strain to minimize late *com* gene expression did not alter the profiles (Figure 7F), showing that the observed effect was directly attributable to the role of σ^X in localizing DprA to the cell poles. Altogether, these findings further prove that in live cells the targeting of pneumococcal DprA to the cell pole,

promoted by σ^X anchored at this cellular location, mediates the timely antagonization of the competence induction signal.

Discussion

The cell pole defines a competence regulation hub in S. pneumoniae

We report here a spatiotemporal analysis of competence regulation in live pneumococcal cells. We found that the positive regulators ComD, ComE and σ^X , which control the early and late competence expression waves, and the negative regulator DprA colocalize during competence at one cell pole to temporally coordinate its development and shut-off. We have shown that the initial stages of competence induction, relying on the CSP-induced phosphorelay between ComD and ComE, occur at the cell pole (Figure 7G). Most importantly, this study uncovered an unexpected second role for σ^X in competence regulation. In addition to controlling the transcription of the late *com* gene *dprA*, σ^X also mediates the accumulation of DprA molecules mostly at the same pole as ComD, a mechanism that facilitates competence shut-off. σ^X thus controls both induction of the late *com* regulon and the shut-off of the early *com* regulon, including thereby its own expression and by consequence that of its regulon. We found that the σ^X -directed polar localization of DprA in the vicinity of ComD correlates with its role in competence shut-off. We propose that this targeting favours DprA interaction between DprA and neosynthesized ComE~P to promote efficient shut-off (Figure 7H). Importantly, we previously reported that DprA-mediated competence shut-off is crucial to the fitness of competent cells (Johnston et al., 2018; Mirouze et al., 2013), and we revealed here that this vital role of DprA in actively limiting the competence window is orchestrated at the cell pole, which defines a coordination hub of competence regulation. Unexpectedly, DprA is targeted to this hub by σ^X , describing an unprecedented role for an alternative sigma factor.

The pneumococcal polar competence regulation hub is focused around the histidine kinase ComD

An important finding is the discrete accumulation of the pneumococcal HK ComD at the cell pole during competence (Figure 6AB). The localization of HKs has not been extensively studied, but few HKs of two-component signaling systems (TCS) have been found to accumulate at the cell pole. One well-documented example is the *Escherichia coli* HK CheA, which is involved in chemotaxis. CheA localizes along with its cognate RR CheY to a single cell pole, forming a chemotactic cluster with a variety of chemoreceptors. CheY stimulates the production of flagellae at the opposite cell pole, driving chemotaxis (Baker et al., 2006). Another example is the HK PilS in *Pseudomonas aeruginosa*, which also accumulates at the cell pole, and along with its cognate cytoplasmic RR PilR, regulates expression of polar pili (Boyd, 2000). Although ComD tethering at one cell pole is similar to CheA and PilS, major differences exist between these three TCS system. Firstly, regarding how their RRs localize, ComE presents a different localization pattern than CheY and PilR, assembling into patches close to the inner side of the cell membrane and focusing dynamically on the cell pole (Figure 6A and Movie 2). ComE phosphorylation by ComD promotes its dimerization and switch into a transcriptional activator (Martin et al., 2013). ComE~P dimers should then leave the cell pole to interact with genomic targets and turn on early *com* gene expression, producing σ^x and ComW, which then cooperate to turn on late *com* gene expression. Secondly, the ComDE TCS induces its own expression, generating a positive feedback loop (Claverys et al., 2006; Martin et al., 2013), and leading to the strong, rapid induction of competence (Dagkessamanskaia et al., 2004b; Peterson et al., 2004). As found here, this leads to the accumulation of the regulatory proteins at the cell pole hub. In addition, the ComDE TCS controls a complex, multifaceted genetic program involving the altered expression of 17% of the genome (Aprianto et al., 2018b) and is present at a unique cell pole, while the actors of transformation are present at midcell (Bergé et al., 2013). Finally, it is possible that the polar localization of the competence regulatory hub is dictated by the very nature of the competence regulation mechanism itself. This localization could be linked to the transient nature of pneumococcal competence. A spatiotemporally controlled localization for this process would provide a mechanism to allow initial induction followed by repression, thus

preventing toxicity. It is also possible that the purpose of polar accumulation of σ^x is to limit the circulating levels of σ^x . Polar accumulation of σ^x may thus protect the cell from a potentially dangerous hyper-competent state at two levels, firstly by promoting polar DprA accumulation to facilitate shut-off, and secondly by sequestering σ^x itself to prevent over-induction of the late *com* regulon. Although the factor localizing σ^x to the cell pole is unknown, it could be considered an anti-sigma factor in this light. Anti-sigma factors can control alternative sigma factors by interacting directly with them to sequester them to prevent activity until a specific signal is received (Österberg et al., 2011b). Although ComD goes to the same pole as σ^x , it is not the anchor directing σ^x to the pole, since DprA (and thus σ^x) still localizes to the cell pole in the absence of early *com* genes (Figure 4EF).

Two copies of comX ensure optimal fitness of competent cells

This study has uncovered a second role for σ^x in the shut-off of pneumococcal competence, besides its transcriptional role with ComW, which is to associate with the RNA polymerase and direct the expression of the late *com* regulon. This second role is independent of ComW. It promotes accumulation of DprA at the cell pole to facilitate competence shut-off. Pneumococci possess two identical copies of *comX* at distinct locations within the genome, called *comX1* and *comX2* (Lee and Morrison, 1999). It has remained unclear why two copies exist, since inactivation of a single copy of *comX* does not affect transformation efficiency (Figure 7A) (Lee and Morrison, 1999). The finding of a second role for σ^x in the shut-off of competence suggests a different reason for this duplication, since both copies of *comX* are required for optimal competence shut-off (Figure 7B). We thus propose that two copies of *comX* are maintained within the genome to optimize the shut-off of competence. Since unregulated competence is toxic for the cell (Johnston et al., 2018), two copies of *comX* provide a fail-safe in case of loss or inactivation of one *comX* gene by transformation or spontaneous mutation, allowing the cell to nonetheless exit the competent state. In light of the second role of σ^x uncovered here, two copies of *comX* provide a simple yet elegant means for *S. pneumoniae* and close relatives to

ensure they are always equipped to survive competence, allowing cells to reap the potential benefits of competence without the fitness cost associated with unregulated competence.

Polar accumulation of DprA and ComD and heterogeneity in a post-competence cell population

The σ^X -mediated accumulation of DprA in foci at the cell pole depends on a high concentration of DprA molecules in the cell, resulting from the σ^X -driven transcription of the *dprA* gene (Figure 2). This dual feature implies that polar DprA foci are not formed immediately during competence, in line with our findings (Figure 3). We suggest that this short delay provides the opportunity for ComD to phosphorylate ComE and induce competence, before DprA arrives at the cell pole to facilitate shut-off. In addition, the foci formation is dependent on the ability of DprA to dimerize, which alters its both its role in transformation via interaction with RecA and its role in competence shut-off via interaction with ComE (Mirouze et al., 2013; Quevillon-Cheruel et al., 2012). However, DprA polar foci formation is independent of the presence of the transformation pore protein ComEC or the recombinase RecA, suggesting that the observed foci are not linked to the conserved role of DprA in transformation (Figure S1). In addition, the loss of polar foci when reducing cellular levels of DprA-GFP correlates with the loss of competence shut-off, strongly supporting the proposal that DprA accumulation at the cell pole underpins its negative feedback role in competence regulation (Figure 2 and S2).

In a competent population, the presence and intensity of DprA-GFP foci varies from cell to cell (Figure 3). The same heterogeneity is observed with GFP-ComD foci (Fig. S7). Indeed, at the peak of CSP-induced competence, a quarter of cells do not present detectable DprA-GFP or GFP-ComD foci and, in the other cells, the foci exhibit different level of brightness (Figures 3 and S7). In addition, foci are observed in all cell types, meaning that their formation is not determined by a particular stage of the cell-cycle. This highlights a heterogeneity in the pneumococcal competent cell population. Furthermore, DprA-GFP and GFP-ComD foci persist in many cells at least 30 minutes after the shut-off of competence (Figure 3 and S7). It has been shown previously that post-competent cells are unable

to respond to CSP for a period of time after they shut-off competence, a phenomenon known as 'blind to CSP' (Chen and Morrison, 1987; Fox and Hotchkiss, 1957). We suggest that the heterogeneity of polar DprA and ComD accumulation could play a role in this phenomenon. Polar DprA accumulation in a majority of cells could prevent cells from responding to CSP by immediately antagonizing neosynthesized ComE~P, while cells lacking ComD foci may not respond optimally to the competence signal. Our findings explain the suggestion made previously based on a mathematical model simulating competence regulation that a high amount of DprA played a role in this phenomenon (Weyder et al., 2018). This notion is supported by the fact that co-expression of DprA and σ^x prior to CSP addition to the cell culture antagonized competence development (Figure 7C-F). Furthermore, since not all post-competent cells possess detectable polar foci of DprA-GFP or GFP-ComD, we suggest that whether sufficient DprA or ComD has accumulated at the pole of a particular cell should govern whether this cell can respond to an external CSP signal and is thus receptive to a second wave of competence. This produces a mixture of competent and non-competent cells, which may maximize the potential survival of a pneumococcal population.

Concluding remarks

In this study, we have shown that the entire pneumococcal competence regulatory cycle occurs at a single cell pole. This generates an asymmetry at the poles of a competent cell, which can be transmitted to future generations and impact the ability to respond to subsequent competence signals. In addition, we have uncovered a key second role for the competence-dedicated alternative sigma factor σ^x that actively localizes DprA to the polar competence regulatory hub to facilitate competence shut-off. This regulatory mechanism, involving two proteins with other conserved roles in competence and transformation respectively, is pivotal to optimal competence shut-off and maintains the fitness of competent cells. This finding represents the first example of an alternative sigma factor

playing a central role in the extinction of the signal on which its own production depends and broadens our knowledge of the regulatory roles played by bacterial alternative sigma factors.

Materials and Methods

Bacterial strains, transformation and competence

The pneumococcal strains, primers and plasmids used in this study can be found in [Table S3](#). Standard procedures for transformation and growth media were used (Martin et al., 2000). In this study, cells were rendered unable to spontaneously develop competence either by deletion of the *comC* gene (*comC0*) (Dagkessamanskaia et al., 2004a) or by replacing the *comC* gene which encodes CSP1 with an allelic variant encoding CSP2 (Pozzi et al., 1996), since ComD1 is unable to respond to CSP2 (Johnsborg et al., 2006; Weyder et al., 2018). Both of these alterations render cells unable to produce CSP. Unless described, pre-competent cultures were prepared by growing cells to an OD₅₅₀ of 0.1 in C+Y medium (pH 7) before 10-fold concentration and storage at –80°C as 100 µL aliquots. Antibiotic concentrations (µg mL⁻¹) used for the selection of *S. pneumoniae* transformants were: chloramphenicol (Cm), 4.5; erythromycin, 0.05; kanamycin (Kan), 250; spectinomycin (Spc), 100; streptomycin (Sm), 200; trimethoprim (Trim), 20. For the monitoring of growth and *luc* expression, precultures were gently thawed and aliquots were inoculated (1 in 100) in luciferin-containing (Prudhomme and Claverys, 2007) C+Y medium and distributed (300 µl per well) into a 96-well white microplate with clear bottom. Transformation was carried out as previously described (Martin et al., 2000). 100 µL aliquots of pre-competent cells were resuspended in 900 µL fresh C+Y medium with 100 ng mL⁻¹ CSP and appropriate IPTG concentrations and incubated at 37°C for 10 min. Transforming DNA was then added to a 100 µL aliquot of this culture, followed by incubation at 30°C for 20 min. Cells were then diluted and plated on 10 mL CAT agar with 5% horse blood and appropriate concentrations of IPTG before incubation at 37°C for 2 h. A second 10 mL layer of CAT agar with appropriate antibiotic was added to plates to select transformants, and plates without antibiotic were used as comparison

to calculate transformation efficiency where appropriate. Plates were incubated overnight at 37°C. To compare transformation efficiencies, transforming DNA was either R304 (Mortier-Barrière et al., 1998, p.) genomic DNA or a 3,434 bp PCR fragment amplified with primer pair MB117-MB120 as noted (Marie et al., 2017), both possessing an *rpsL41* point mutation conferring streptomycin resistance. To track competence profiles, a previously described protocol was used (Prudhomme and Claverys, 2007). Relative luminescence unit (RLU) and OD values were recorded throughout incubation at 37°C in a Varioskan luminometer (ThermoFisher). The *comC-luc* and *ssbB-luc* reporter genes were transferred from R825 or R895 as previously described (Bergé et al., 2002; Chastanet et al., 2001). *CEP_{lac}-dprA-gfp* strains were grown in varying concentrations of IPTG from the beginning of growth, as previously described (Johnston et al., 2018). Detailed information regarding the construction of new plasmids and strains can be found in the [Supplementary Information](#).

Fluorescence microscopy and image analysis

Pneumococcal precultures grown in C+Y medium at 37°C to an OD₅₅₀ of 0.1 were induced with either CSP (100 ng mL⁻¹) or BIP (250 ng mL⁻¹) peptide. At indicated times post induction, 1 mL samples were collected, cooled down by addition of 500 mL cold medium, pelleted (3 min, 3,000 g) and resuspended in 1 mL C+Y medium. 2 µL of this suspension were spotted on a microscope slide containing a slab of 1.2% C+Y agarose as previously described (Bergé et al., 2013) before imaging. Unless stated, images were visualized 15 minutes after competence induction, at the peak of competence gene expression. To generate movies, images were taken of the same fields of vision at varying time points during incubation at 37°C. Images were captured and processed using the Nis-Elements AR software (Nikon). Images were analyzed using MicrobeJ, a plug-in of ImageJ (Ducret et al., 2016). Data was analyzed in R and unless stated, represented as focus density maps plotted on the longitudinal axis of half cells ordered by cell length. Each spot represents the localization of an individual focus, and spot colour represents focus density at a specific location on the half cell. Cells

with >0 foci shown for each time point. In cells possessing >1 foci, foci were represented adjacently on cells of the same length.

Western blots

To compare the expression profiles of competence proteins after competence induction, time course Western blots were carried out. Cells were diluted 100-fold in 10 mL C+Y medium pH 7 and grown to OD 0.1. Where appropriate, cells were induced with CSP (100 ng mL⁻¹) or BIP (250 ng mL⁻¹) peptide. At indicated time points, OD₅₅₀ measurements were taken and 500 µL of culture was recovered. Samples were centrifuged (3 min, 3,000 g) and pellets were resuspended in 40 µL of TE 1x supplemented with 0.01% DOC and 0.02% SDS. Samples were then incubated for 10 minutes at 37°C before addition of 40 µL 2x sample buffer with 10% β-mercaptoethanol, followed by incubation at 85°C for 10 minutes. Samples were then normalized compared to the initial OD₅₅₀ reading, and loaded onto SDS-PAGE gels (BIORAD). Samples were migrated for 30 min at 200V, and transferred onto nitrocellulose membrane using a Transblot Turbo (BIORAD). Membranes were blocked for 1h at room temperature in 1x TBS with 0.1% Tween20 and 10% milk, before two washes in 1x TBS with 0.1% Tween20 and probing with primary antibodies (1/10,000 as noted) in 1x TBS with 0.1% Tween20 and 5% milk overnight at 4°C. After a further four washes in 1x TBS with 0.1% Tween20, membranes were probed with anti-rabbit secondary antibody (1/10,000) for 1h 30 min, followed by another four washes in 1x TBS with 0.1% Tween20. Membranes were activated using Clarity Max ECL (BIORAD) and visualized in a ChemiDoc Touch (BIORAD).

Co-Immunoprecipitation

Co-immunoprecipitation was done using magnetic GFP-Trap beads as per manufacturer's instructions (Chromotek). Briefly, cells were inoculated 1/100 in 25 mL of C+Y medium pH 7 and grown to OD₅₅₀ 0.1. Competence was induced by addition of 100 ng mL⁻¹ CSP, and cells were incubated for 10 min at 37°C. Cultures were mixed with 25 mL cold buffer A (10 mM Tris pH 7.5, 150 mM NaCl) and

centrifuged for 15 min at 5,000 g. Pellets were washed twice with 10 mL cold buffer A, and stored at -80°C until use. After defrosting, pellets were resuspended in 1 mL buffer B (10 mM Tris pH 7.5, 150 mM NaCl, 0.2mM EDTA, 0.1% TritonX100, 1 M DTT) and incubated for 10 minutes on ice, followed by 10 min at 37°C, and a further 10 min on ice. Samples were then sonicated (2 x 30s with 10s pause) and centrifuged for 30 min at 4°C and 16,000 g. After normalizing the protein concentrations in the samples, 20 µg mL⁻¹ RNase A and 50 µg mL⁻¹ DNase I were added and samples were tumbled end over end at 4°C for 30 min. 75 µl of GFP-Trap beads were added to the samples, which were then tumbled end over end at 4°C for 2 h 30 min. GFP-Trap beads were purified by magnetism and washed twice in 500 µL ice cold dilution buffer (10 mM Tris pH 7.5, 150 mM NaCl, 0.5mM EDTA), before being resuspended in 2x sample buffer + 10% β-mercaptoethanol and incubated at 95°C for 10 minutes. Samples were then run on SDS-PAGE gel and Western blots carried out as described above.

Pre-competence expression of DprA and σ^X.

Cells (R4500, CEPlac-dprA, dprA::spc, CEPIIR- R4509, R4511,) were grown to OD₄₉₂ 0.2 in 2 mL C+Y medium (pH7.6) with 50 µM IPTG. After centrifugation for 5 minutes at 5,000 rpm, cells were resuspended in 1 mL C+Y medium and stored at -80°C in 100 µL aliquots until required. Aliquots were resuspended in 900 µL fresh C+Y medium (pH7.6) and diluted 1/10 in a 96-ell plate in C+Y medium (pH 7.6) with luciferin and 50 µM IPTG to induce DprA expression. After 25 minutes, BIP (250ng µL⁻¹) was added where noted to induce σ^X. 20 minutes later, CSP (100ng µL⁻¹) was added to induce competence for 20 minutes. Luminometric and photometric reading were taken every 2 minutes during this time to report induction of *comC-luc*.

Acknowledgements

We thank Nathalie Campo and Mathieu Bergé for critical reading of the manuscript, and the rest of the Polard lab for helpful discussions. We thank Jérôme Rech for help creating Movies. We

thank Jan-Willem Veening for kind donation of the pMK111 plasmid. This work was funded by the Agence Nationale de la Recherche (Grants ANR-10-BLAN-1331 and ANR-13-BSV8-0022).

Author Contributions

Conceptualization, C. J., M. P., P. P.; Methodology, C. J., P. P.; Investigation, C. J., A. L. S., M. B., M. P., D. D. L.; Writing – Original draft, C. J.; Writing – Review and editing, C. J., P. P.; Funding acquisition, P. P.

References

- Aggarwal, S.D., Eutsey, R., West-Roberts, J., Domenech, A., Xu, W., Abdullah, I.T., Mitchell, A.P., Veening, J.-W., Yesilkaya, H., Hiller, N.L., 2018. Function of BriC peptide in the pneumococcal competence and virulence portfolio. *PLoS Pathog.* 14, e1007328. <https://doi.org/10.1371/journal.ppat.1007328>
- Akerley, B.J., Rubin, E.J., Camilli, A., Lampe, D.J., Robertson, H.M., Mekalanos, J.J., 1998. Systematic identification of essential genes by in vitro mariner mutagenesis. *Proc. Natl. Acad. Sci. U. S. A.* 95, 8927–8932. <https://doi.org/10.1073/pnas.95.15.8927>
- Alloing, G., Martin, B., Granadel, C., Claverys, J.P., 1998. Development of competence in *Streptococcus pneumoniae*: pheromone autoinduction and control of quorum sensing by the oligopeptide permease. *Mol. Microbiol.* 29, 75–83. <https://doi.org/10.1046/j.1365-2958.1998.00904.x>
- Aprianto, R., Slager, J., Holsappel, S., Veening, J.-W., 2018a. High-resolution analysis of the pneumococcal transcriptome under a wide range of infection-relevant conditions. *Nucleic Acids Res.* 46, 9990–10006. <https://doi.org/10.1093/nar/gky750>
- Aprianto, R., Slager, J., Holsappel, S., Veening, J.-W., 2018b. High-resolution analysis of the pneumococcal transcriptome under a wide range of infection-relevant conditions. *Nucleic Acids Res.* 46, 9990–10006. <https://doi.org/10.1093/nar/gky750>
- Baker, M.D., Wolanin, P.M., Stock, J.B., 2006. Signal transduction in bacterial chemotaxis. *BioEssays News Rev. Mol. Cell. Dev. Biol.* 28, 9–22. <https://doi.org/10.1002/bies.20343>
- Bergé, M., Moscoso, M., Prudhomme, M., Martin, B., Claverys, J.-P., 2002. Uptake of transforming DNA in Gram-positive bacteria: a view from *Streptococcus pneumoniae*. *Mol. Microbiol.* 45, 411–421.
- Bergé, M.J., Kamgoué, A., Martin, B., Polard, P., Campo, N., Claverys, J.-P., 2013. Midcell recruitment of the DNA uptake and virulence nuclease, EndA, for pneumococcal transformation. *PLoS Pathog.* 9, e1003596. <https://doi.org/10.1371/journal.ppat.1003596>
- Bergé, M.J., Mercy, C., Mortier-Barrière, I., VanNieuwenhze, M.S., Brun, Y.V., Grangeasse, C., Polard, P., Campo, N., 2017. A programmed cell division delay preserves genome integrity during natural genetic transformation in *Streptococcus pneumoniae*. *Nat. Commun.* 8. <https://doi.org/10.1038/s41467-017-01716-9>
- Boyd, J.M., 2000. Localization of the histidine kinase PilS to the poles of *Pseudomonas aeruginosa* and identification of a localization domain. *Mol. Microbiol.* 36, 153–162. <https://doi.org/10.1046/j.1365-2958.2000.01836.x>

- Campbell, E.A., Choi, S.Y., Masure, H.R., 1998. A competence regulon in *Streptococcus pneumoniae* revealed by genomic analysis. *Mol. Microbiol.* 27, 929–939. <https://doi.org/10.1046/j.1365-2958.1998.00737.x>
- Caymaris, S., Bootsma, H.J., Martin, B., Hermans, P.W.M., Prudhomme, M., Claverys, J.-P., 2010. The global nutritional regulator CodY is an essential protein in the human pathogen *Streptococcus pneumoniae*. *Mol. Microbiol.* 78, 344–360.
- Chastanet, A., Prudhomme, M., Claverys, J.P., Msadek, T., 2001. Regulation of *Streptococcus pneumoniae* *clp* genes and their role in competence development and stress survival. *J. Bacteriol.* 183, 7295–7307. <https://doi.org/10.1128/JB.183.24.7295-7307.2001>
- Chen, J.D., Morrison, D.A., 1987. Modulation of competence for genetic transformation in *Streptococcus pneumoniae*. *J. Gen. Microbiol.* 133, 1959–1967. <https://doi.org/10.1099/00221287-133-7-1959>
- Claverys, J.-P., Håvarstein, L.S., 2007. Cannibalism and fratricide: mechanisms and raisons d’être. *Nat. Rev. Microbiol.* 5, 219–229. <https://doi.org/10.1038/nrmicro1613>
- Claverys, J.-P., Prudhomme, M., Martin, B., 2006. Induction of competence regulons as a general response to stress in gram-positive bacteria. *Annu. Rev. Microbiol.* 60, 451–475. <https://doi.org/10.1146/annurev.micro.60.080805.142139>
- Dagkessamanskaia, A., Moscoso, M., Hénard, V., Guiral, S., Overweg, K., Reuter, M., Martin, B., Wells, J., Claverys, J.-P., 2004a. Interconnection of competence, stress and CiaR regulons in *Streptococcus pneumoniae*: competence triggers stationary phase autolysis of *ciaR* mutant cells. *Mol. Microbiol.* 51, 1071–1086.
- Dagkessamanskaia, A., Moscoso, M., Hénard, V., Guiral, S., Overweg, K., Reuter, M., Martin, B., Wells, J., Claverys, J.-P., 2004b. Interconnection of competence, stress and CiaR regulons in *Streptococcus pneumoniae*: competence triggers stationary phase autolysis of *ciaR* mutant cells. *Mol. Microbiol.* 51, 1071–1086.
- Ducret, A., Quardokus, E.M., Brun, Y.V., 2016. MicrobeJ, a tool for high throughput bacterial cell detection and quantitative analysis. *Nat. Microbiol.* 1, 16077. <https://doi.org/10.1038/nmicrobiol.2016.77>
- Fox, M.S., Hotchkiss, R.D., 1957. Initiation of bacterial transformation. *Nature* 179, 1322–1325. <https://doi.org/10.1038/1791322a0>
- Guiral, S., Mitchell, T.J., Martin, B., Claverys, J.-P., 2005. Competence-programmed predation of noncompetent cells in the human pathogen *Streptococcus pneumoniae*: genetic requirements. *Proc. Natl. Acad. Sci. U. S. A.* 102, 8710–8715. <https://doi.org/10.1073/pnas.0500879102>
- Håvarstein, L.S., Coomaraswamy, G., Morrison, D.A., 1995. An unmodified heptadecapeptide pheromone induces competence for genetic transformation in *Streptococcus pneumoniae*. *Proc. Natl. Acad. Sci. U. S. A.* 92, 11140–11144. <https://doi.org/10.1073/pnas.92.24.11140>
- Hui, F.M., Zhou, L., Morrison, D.A., 1995. Competence for genetic transformation in *Streptococcus pneumoniae*: organization of a regulatory locus with homology to two lactococcal A secretion genes. *Gene* 153, 25–31. [https://doi.org/10.1016/0378-1119\(94\)00841-f](https://doi.org/10.1016/0378-1119(94)00841-f)
- Johnsborg, O., Kristiansen, P.E., Blomqvist, T., Håvarstein, L.S., 2006. A hydrophobic patch in the competence-stimulating Peptide, a pneumococcal competence pheromone, is essential for specificity and biological activity. *J. Bacteriol.* 188, 1744–1749. <https://doi.org/10.1128/JB.188.5.1744-1749.2006>
- Johnston, C., Hauser, C., Hermans, P.W.M., Martin, B., Polard, P., Bootsma, H.J., Claverys, J.-P., 2016. Fine-tuning of choline metabolism is important for pneumococcal colonization. *Mol. Microbiol.* 100, 972–988. <https://doi.org/10.1111/mmi.13360>
- Johnston, C., Martin, B., Fichant, G., Polard, P., Claverys, J.-P., 2014. Bacterial transformation: distribution, shared mechanisms and divergent control. *Nat. Rev. Microbiol.* 12, 181–196. <https://doi.org/10.1038/nrmicro3199>

- Johnston, C., Mortier-Barrière, I., Khemici, V., Polard, P., 2018. Fine-tuning cellular levels of DprA ensures transformant fitness in the human pathogen *Streptococcus pneumoniae*. *Mol. Microbiol.* 109, 663–675. <https://doi.org/10.1111/mmi.14068>
- Kazmierczak, M.J., Wiedmann, M., Boor, K.J., 2005. Alternative sigma factors and their roles in bacterial virulence. *Microbiol. Mol. Biol. Rev.* MMBR 69, 527–543. <https://doi.org/10.1128/MMBR.69.4.527-543.2005>
- Kloosterman, T.G., Hendriksen, W.T., Bijlsma, J.J.E., Bootsma, H.J., van Hijum, S.A.F.T., Kok, J., Hermans, P.W.M., Kuipers, O.P., 2006. Regulation of glutamine and glutamate metabolism by GlnR and GlnA in *Streptococcus pneumoniae*. *J. Biol. Chem.* 281, 25097–25109. <https://doi.org/10.1074/jbc.M601661200>
- Lee, M.S., Morrison, D.A., 1999. Identification of a new regulator in *Streptococcus pneumoniae* linking quorum sensing to competence for genetic transformation. *J. Bacteriol.* 181, 5004–5016.
- Lin, J., Lau, G.W., 2019. DprA-dependent exit from the competent state regulates multifaceted *Streptococcus pneumoniae* virulence. *Infect. Immun.* <https://doi.org/10.1128/IAI.00349-19>
- Lin, J., Zhu, L., Lau, G.W., 2016. Disentangling competence for genetic transformation and virulence in *Streptococcus pneumoniae*. *Curr. Genet.* 62, 97–103. <https://doi.org/10.1007/s00294-015-0520-z>
- Luo, P., Li, H., Morrison, D.A., 2004. Identification of ComW as a new component in the regulation of genetic transformation in *Streptococcus pneumoniae*. *Mol. Microbiol.* 54, 172–183. <https://doi.org/10.1111/j.1365-2958.2004.04254.x>
- Marie, L., Rapisarda, C., Morales, V., Bergé, M., Perry, T., Soulet, A.-L., Gruget, C., Remaut, H., Fronzes, R., Polard, P., 2017. Bacterial RadA is a DnaB-type helicase interacting with RecA to promote bidirectional D-loop extension. *Nat. Commun.* 8, 15638. <https://doi.org/10.1038/ncomms15638>
- Martin, B., García, P., Castanié, M.P., Claverys, J.P., 1995. The *recA* gene of *Streptococcus pneumoniae* is part of a competence-induced operon and controls lysogenic induction. *Mol. Microbiol.* 15, 367–379.
- Martin, B., Prats, H., Claverys, J.P., 1985. Cloning of the *hexA* mismatch-repair gene of *Streptococcus pneumoniae* and identification of the product. *Gene* 34, 293–303. [https://doi.org/10.1016/0378-1119\(85\)90138-6](https://doi.org/10.1016/0378-1119(85)90138-6)
- Martin, B., Prudhomme, M., Alloing, G., Granadel, C., Claverys, J.P., 2000. Cross-regulation of competence pheromone production and export in the early control of transformation in *Streptococcus pneumoniae*. *Mol. Microbiol.* 38, 867–878.
- Martin, B., Soulet, A.-L., Mirouze, N., Prudhomme, M., Mortier-Barrière, I., Granadel, C., Noirot-Gros, M.-F., Noirot, P., Polard, P., Claverys, J.-P., 2013. ComE/ComE~P interplay dictates activation or extinction status of pneumococcal X-state (competence). *Mol. Microbiol.* 87, 394–411. <https://doi.org/10.1111/mmi.12104>
- Mirouze, N., Bergé, M.A., Soulet, A.-L., Mortier-Barrière, I., Quentin, Y., Fichant, G., Granadel, C., Noirot-Gros, M.-F., Noirot, P., Polard, P., Martin, B., Claverys, J.-P., 2013. Direct involvement of DprA, the transformation-dedicated RecA loader, in the shut-off of pneumococcal competence. *Proc. Natl. Acad. Sci. U. S. A.* 110, E1035–1044. <https://doi.org/10.1073/pnas.1219868110>
- Mortier-Barrière, I., Campo, N., Bergé, M.A., Prudhomme, M., Polard, P., 2019. Natural Genetic Transformation: A Direct Route to Easy Insertion of Chimeric Genes into the Pneumococcal Chromosome. *Methods Mol. Biol.* Clifton NJ 1968, 63–78. https://doi.org/10.1007/978-1-4939-9199-0_6
- Mortier-Barrière, I., de Saizieu, A., Claverys, J.P., Martin, B., 1998. Competence-specific induction of *recA* is required for full recombination proficiency during transformation in *Streptococcus pneumoniae*. *Mol. Microbiol.* 27, 159–170.
- Mortier-Barrière, I., Velten, M., Dupaigne, P., Mirouze, N., Piétrement, O., McGovern, S., Fichant, G., Martin, B., Noirot, P., Le Cam, E., Polard, P., Claverys, J.-P., 2007. A key presynaptic role in

transformation for a widespread bacterial protein: DprA conveys incoming ssDNA to RecA. *Cell* 130, 824–836. <https://doi.org/10.1016/j.cell.2007.07.038>

Österberg, S., del Peso-Santos, T., Shingler, V., 2011a. Regulation of alternative sigma factor use. *Annu. Rev. Microbiol.* 65, 37–55. <https://doi.org/10.1146/annurev.micro.112408.134219>

Österberg, S., del Peso-Santos, T., Shingler, V., 2011b. Regulation of alternative sigma factor use. *Annu. Rev. Microbiol.* 65, 37–55. <https://doi.org/10.1146/annurev.micro.112408.134219>

Pestova, E.V., Morrison, D.A., 1998. Isolation and characterization of three *Streptococcus pneumoniae* transformation-specific loci by use of a lacZ reporter insertion vector. *J. Bacteriol.* 180, 2701–2710.

Peterson, S.N., Sung, C.K., Cline, R., Desai, B.V., Snesrud, E.C., Luo, P., Walling, J., Li, H., Mintz, M., Tsegaye, G., Burr, P.C., Do, Y., Ahn, S., Gilbert, J., Fleischmann, R.D., Morrison, D.A., 2004. Identification of competence pheromone responsive genes in *Streptococcus pneumoniae* by use of DNA microarrays. *Mol. Microbiol.* 51, 1051–1070.

Pozzi, G., Masala, L., Iannelli, F., Manganelli, R., Havarstein, L.S., Piccoli, L., Simon, D., Morrison, D.A., 1996. Competence for genetic transformation in encapsulated strains of *Streptococcus pneumoniae*: two allelic variants of the peptide pheromone. *J. Bacteriol.* 178, 6087–6090. <https://doi.org/10.1128/jb.178.20.6087-6090.1996>

Prudhomme, M., Attaiech, L., Sanchez, G., Martin, B., Claverys, J.-P., 2006. Antibiotic stress induces genetic transformability in the human pathogen *Streptococcus pneumoniae*. *Science* 313, 89–92. <https://doi.org/10.1126/science.1127912>

Prudhomme, M., Claverys, J.-P., 2007. There will be a light: the use of luc transcriptional fusions in living pneumococcal cells., in: *The Molecular Biology of Streptococci*. R. Hakenbeck, and G.S. Chatwal, pp. 519–524.

Quevillon-Cheruel, S., Campo, N., Mirouze, N., Mortier-Barrière, I., Brooks, M.A., Boudes, M., Durand, D., Soulet, A.-L., Lisboa, J., Noirot, P., Martin, B., van Tilbeurgh, H., Noirot-Gros, M.-F., Claverys, J.-P., Polard, P., 2012. Structure-function analysis of pneumococcal DprA protein reveals that dimerization is crucial for loading RecA recombinase onto DNA during transformation. *Proc. Natl. Acad. Sci. U. S. A.* 109, E2466–2475. <https://doi.org/10.1073/pnas.1205638109>

Salles, C., Créancier, L., Claverys, J.P., Méjean, V., 1992. The high level streptomycin resistance gene from *Streptococcus pneumoniae* is a homologue of the ribosomal protein S12 gene from *Escherichia coli*. *Nucleic Acids Res.* 20, 6103. <https://doi.org/10.1093/nar/20.22.6103>

Sanchez-Puelles, J.M., Ronda, C., Garcia, J.L., Garcia, P., Lopez, R., Garcia, E., 1986. Searching for autolysin functions. Characterization of a pneumococcal mutant deleted in the *lytA* gene. *Eur. J. Biochem.* 158, 289–293. <https://doi.org/10.1111/j.1432-1033.1986.tb09749.x>

Slager, J., Aprianto, R., Veening, J.-W., 2019. Refining the Pneumococcal Competence Regulon by RNA Sequencing. *J. Bacteriol.* 201. <https://doi.org/10.1128/JB.00780-18>

Slager, J., Kjos, M., Attaiech, L., Veening, J.-W., 2014. Antibiotic-induced replication stress triggers bacterial competence by increasing gene dosage near the origin. *Cell* 157, 395–406. <https://doi.org/10.1016/j.cell.2014.01.068>

Sung, C.K., Li, H., Claverys, J.P., Morrison, D.A., 2001. An rpsL cassette, janus, for gene replacement through negative selection in *Streptococcus pneumoniae*. *Appl. Environ. Microbiol.* 67, 5190–5196. <https://doi.org/10.1128/AEM.67.11.5190-5196.2001>

Sung, C.K., Morrison, D.A., 2005. Two distinct functions of ComW in stabilization and activation of the alternative sigma factor ComX in *Streptococcus pneumoniae*. *J. Bacteriol.* 187, 3052–3061. <https://doi.org/10.1128/JB.187.9.3052-3061.2005>

Tovpeko, Y., Bai, J., Morrison, D.A., 2016. Competence for Genetic Transformation in *Streptococcus pneumoniae*: Mutations in σ A Bypass the ComW Requirement for Late Gene Expression. *J. Bacteriol.* 198, 2370–2378. <https://doi.org/10.1128/JB.00354-16>

van Raaphorst, R., Kjos, M., Veening, J.-W., 2017. Chromosome segregation drives division site selection in *Streptococcus pneumoniae*. *Proc. Natl. Acad. Sci. U. S. A.* 114, E5959–E5968. <https://doi.org/10.1073/pnas.1620608114>

- Vidal, J.E., Howery, K.E., Ludewick, H.P., Nava, P., Klugman, K.P., 2013. Quorum-sensing systems LuxS/autoinducer 2 and Com regulate *Streptococcus pneumoniae* biofilms in a bioreactor with living cultures of human respiratory cells. *Infect. Immun.* 81, 1341–1353.
<https://doi.org/10.1128/IAI.01096-12>
- Weng, L., Piotrowski, A., Morrison, D.A., 2013. Exit from competence for genetic transformation in *Streptococcus pneumoniae* is regulated at multiple levels. *PLoS One* 8, e64197.
<https://doi.org/10.1371/journal.pone.0064197>
- Weyder, M., Prudhomme, M., Bergé, M., Polard, P., Fichant, G., 2018. Dynamic Modeling of *Streptococcus pneumoniae* Competence Provides Regulatory Mechanistic Insights Into Its Tight Temporal Regulation. *Front. Microbiol.* 9, 1637.
<https://doi.org/10.3389/fmicb.2018.01637>
- Zhu, L., Lin, J., Kuang, Z., Vidal, J.E., Lau, G.W., 2015. Deletion analysis of *Streptococcus pneumoniae* late competence genes distinguishes virulence determinants that are dependent or independent of competence induction. *Mol. Microbiol.* 97, 151–165.
<https://doi.org/10.1111/mmi.13016>

Figure Legends

Figure 1: DprA: Localization and roles in competence and transformation.

(A) (1) Pre-CSP is exported and matured by the ComAB transporter, and then phosphorylates the histidine kinase ComD. (2) ComD transphosphorylates ComE, which then stimulates the expression of 17 early *com* genes, including two copies of *comX*. (3) These encode an alternative sigma factor σ^x , which controls late *com* genes including *dprA*. (4) DprA dimers load RecA onto ssDNA to mediate transformation and interact with ComE~P to shut-off competence. (5) Transforming DNA is internalized in single strand form and is protected from degradation by DprA and RecA (6), which then mediate transformation (7). Orange arrows, early *com* promoters; purple arrows, late *com* promoters.

(B) Western blot tracking cellular levels of DprA-GFP after competence induction in strain R3728. α -DprA antibody used. (C) Sample fluorescence microscopy images of R3728 strain producing DprA-GFP 15 minutes after competence induction. Scale bars, 1 μ m. (D) Schematic representation of focus density maps with half cells represented as vertical lines in ascending size order and localization of foci represented along the length axis of each half cell. Black half-cells represent those presented, and grey those not presented. (E) DprA-GFP accumulates at the cell poles during competence. 1290 cells and 1128 foci analyzed (F) Sample immunofluorescence microscopy images of a strain producing wildtype

DprA (R1502; wildtype) and a strain lacking DprA (R2018; *dprA*⁻) fixed 15 minutes after competence induction. Scale bars, 1 μ m.

Figure 2: Reducing cellular DprA-GFP levels results in loss of polar accumulation and competence shut-off

(A) Focus density maps of DprA-GFP foci at different cellular levels during competence. *CEP_{lac}-dprA-gfp* from strain R4262. Cellular DprA-GFP levels were controlled by growing cells in a gradient of IPTG. 1.5 μ M IPTG, 11267 cells and 791 foci analyzed; 3 μ M IPTG, 10623 cells and 748 foci analyzed; 6 μ M IPTG, 10603 cells and 743 foci analyzed; 12.5 μ M IPTG, 6985 cells and 1010 foci analyzed; 25 μ M IPTG, 2945 cells and 1345 foci analyzed; 50 μ M IPTG, 3678 cells and 1964 foci analyzed. Sample microscopy images of strain R4262 in varying IPTG concentrations. Scale bars, 1 μ m. (B) Reducing cellular levels of DprA-GFP reduces the number of cells with foci. Error bars represent triplicate repeats. (C) Reducing cellular levels of DprA-GFP results specifically in loss of polar foci. Error bars represent triplicate repeats.

Figure 3: Analysis of the cellular localization of DprA-GFP

(A) DprA-GFP foci persist at the cell pole after competence shut-off Data at different time points after CSP addition represented as in [Figure 1E](#). 4 minutes, 8336 cells and 1383 foci analyzed; 6 minutes, 2871 cells and 1674 foci analyzed; 10 minutes, 2614 cells and 1502 foci analyzed; 15 minutes, 1290 cells and 1128 foci analyzed; 20 minutes, 2110 cells and 1900 foci analyzed; 30 minutes, 789 cells and 831 foci analyzed; 60 minutes, 842 cells and 839 foci analyzed. Sample microscopy images of strain R3728 at varying times after competence induction. Scale bars, 1 μ m. (B) Most competent cells possess a single polar focus of DprA-GFP. Error bars represent triplicate repeats. (C) Most cells possess polar DprA-GFP foci. Along the length of a cell of arbitrary length 1, polar foci are found between positions 0-0.15 and 0.85-1, midcell foci are found between 0.35 and 0.65, and anything in between is localized as betwixt. Error bars represent triplicate repeats.

Figure 4: Polar accumulation of DprA-GFP appears to depend on late *com* regulon expression

(A) Sample fluorescence microscopy images of strain R4060 producing DprA-GFP 15 minutes after induction with BIP or BIP and CSP. Scale bars, 1 μ m. (B) Competence induction is required for optimal accumulation of DprA-GFP at the cell poles. Focus density maps as in Figure 1E. BIP+, 7845 cells and 790 foci analyzed; BIP+ CSP+, 2707 cells and 1478 foci analyzed. (C) Genetic context strain R4107 expressing *dprA-gfp* and only the late *com* regulon. P_{BIP} and P_X as in panel A, P_E represents early *com* promoter controlled by ComE. Light blue circle, ComW; light green oval, RNA polymerase; purple hexagon, σ^X . (D) Genetic context of strain R4140 expressing *CEP_R-dprA-gfp* and only the early *com* regulon. P_{BIP} as in panel A, P_E as in panel D. (E) Sample fluorescence microscopy images of strains producing DprA-GFP with only late (R4107) or only early (R4140) *com* operons 15 minutes after competence induction. Scale bars, 2 μ m. (F) Induction of the late *com* regulon is required for accumulation of DprA-GFP at the cell poles. Focus density maps as in Figure 1E. 1988 cells and 1824 foci analyzed. (G) Focus density maps produced as in Figure 1E from images where DprA-GFP was produced outside of competence in presence or absence of σ^X . DprA-GFP alone, 7845 cells and 790 foci analyzed; DprA-GFP + σ^X , 3545 cells and 1355 foci analyzed. Strains used: DprA-GFP alone, R4060; DprA-GFP and σ^X , R4489.

Figure 5: σ^X -GFP interacts directly with DprA at the cell pole during competence

(A) Sample fluorescence microscopy images of strain R4451 producing σ^X -GFP from *comX1* and wildtype σ^X from *comX2* 15 minutes after competence induction. Scale bars, 1 μ m. (B) σ^X -GFP accumulates at the cell poles during competence. Focus density maps presented as in Figure 1E. 4 minutes, 7544 cells and 489 foci analyzed; 6 minutes, 5442 cells and 1711 foci analyzed; 10 minutes, 4358 cells and 2691 foci analyzed; 15 minutes, 3746 cells and 2144 foci analyzed; 20 minutes, 4211 cells and 1754 foci analyzed; 30 minutes, 4695 cells and 1920 foci analyzed; 60 minutes, 5713 cells and 1016 foci analyzed. (C) Most cells have a single σ^X -GFP focus. Data from the time-course experiment presented in panel B showing the number of foci per cell at each time point. Error bars represent

triplicate repeats. (D) DprA-GFP foci persist in cells longer than σ^X -GFP foci. Comparison of cells with foci at different timepoints from timecourse experiments. DprA-GFP from **Figure 3A**, σ^X -GFP from panel B. Error bars represent triplicate repeats. (E) Accumulation of σ^X at the cell poles does not depend on DprA. Sample microscopy images of a *comX1-gfp, dprA⁻* strain (R4469). Focus density maps generated from cells visualized 15 minutes after competence induction presented as in **Figure 1E**. 1104 cells and 638 foci analyzed. (F) σ^X and DprA colocalize at the cell pole. Colocalization of σ^X -YFP and DprA-mTurquoise in R4473 cells visualized 15 minutes after competence induction. 7460 cells and 3504 DprA-mTurquoise foci analyzed. Scale bars, 1 μ m. (G) DprA is copurified with σ^X -GFP while DprA^{AR} is not. Western blot of pull-down experiment carried out on strains producing σ^X -GFP and either DprA (R4451) or DprA^{AR} (R4514) 10 minutes after competence induction. WCE, whole cell extract; FT1, flow through; E, eluate.

Figure 6: The main actors of competence induction and shut-off colocalize at the cell pole

(A) Sample fluorescence microscopy images of cells producing ComE-GFP (R4010), GFP-ComD (R3914) or wildtype cells (R1501) and *comD⁻* cells (R1745) exposed to CSP-HF. Scale bars, 1 μ m. GFP-ComD, 1933 cells and 1397 foci; CSP-HF, 2105 cells and 2121 foci. (B) Focus density maps of cells producing GFP-ComD (R3914) and wildtype (R1501) cells exposed to CSP-HF 15 minutes after competence induction. Data presented as in **Figure 1E**. GFP-ComD, 1933 cells and 1397 foci analyzed; CSP-HF, 2105 cells and 2121 foci analyzed. (C) Number of foci present in cells possessing foci of different fluorescent fusions 15 minutes after competence induction. Data taken from panel B except for DprA-GFP, taken from **Figure 3**. (D) DprA-mTurquoise and ComE-YFP colocalization in competent R4176 cells visualized by fluorescence microscopy 15 minutes after competence induction. 3143 cells and 2126 DprA-mTurquoise foci analyzed. Scale bars, 1 μ m. (E) DprA-mTurquoise and YFP-ComD colocalization in competent R4111 cells visualized by fluorescence microscopy 15 minutes after competence induction. 2857 cells and 1335 DprA-mTurquoise foci analyzed. Scale bars, 1 μ m. (F) DprA-GFP and CSP-HF

colocalization in competent R4062 cells visualized by fluorescence microscopy 15 minutes after competence induction. 7588 cells and 3663 DprA-mTurquoise foci analyzed. Scale bars, 1 μ m.

Figure 7: Pre-competence expression of DprA and σ^x antagonizes competence induction

(A) Inactivation of *comX1* or *comX2* does not impact transformation efficiency. Error bars represent triplicate repeats. (B) Inactivation of *comX1* or *comX2* delays the shut-off of competence. Data is plotted as Relative light units, corrected by optical density (RLU/OD) against OD. Error bars represent triplicate repeats. (C) Visual representation of experiment exploring the impact of pre-competence expression of DprA and σ^x on induction. (D) BIP induction of cells lacking *CEPII_R-comX* does not alter competence induction. Cells possessing *comC-luc* and *CEP_{lac}-dprA* (R4511) were treated as described in Panel C. Error bars represent triplicate repeats. (E) Production of σ^x and DprA prior to competence induction antagonizes competence. Cells possessing *comC-luc* and *CEP_{lac}-dprA* and *CEPII_R-comX* (R4500) were treated as in panel C. Error bars represent triplicate repeats. (F) Inactivation of *comW* does not alter the antagonization of competence induction mediated by early DprA and σ^x production. Cells possessing *comC-luc*, *CEP_{lac}-dprA*, *CEPII_R-comX* and *Δ comW::trim* (R4509) were treated as in panel C. Error bars represent triplicate repeats. (G) Polar competence induction in *Streptococcus pneumoniae*. (1) Extracellular CSP interacts with ComD at the cell poles, prompting ComD autophosphorylation. (2) Patches of ComE navigate around the cell membrane. (3) Polar ComD phosphorylates ComE. (4) Active ComE~P dimers leave the cell poles to interact with genomic targets, inducing the early *com* regulon and (5) launching an autocatalytic feedback loop. (6) Among the early *com* genes, *comX1*, *comX2* and *comW* produce σ^x and its activator ComW, which induce the late *com* regulon, including DprA. Orange arrows, early *com* promoters; purple arrows, late *com* promoters. C, *comC*; D, *comD*; E, *comE*; X1/2, *comX1/comX2*; W, *comW*; RNAP, RNA polymerase. (H) Polar shut-off of pneumococcal competence. (1) σ^x interacts directly with DprA, promoting accumulation of DprA at the cell pole, generating a polar DprA 'cloud' in competent cells, near ComD (2). (3) Polar DprA interacts directly with neosynthesized ComE~P, preventing the regulator from accessing its genomic targets. (4)

904 Unphosphorylated ComE interacts with early com promoters, acting as a repressor to prevent
905 induction (Martin et al., 2013), promoting extinction of the competence signal, resulting in
906 competence shut-off (5).

907

908

909

910

911

912

913

914

915

916

917

918

919

920

921

922

923

924

925

926

927

928

929




Figure 1




Figure 2






Figure 3




Figure 4




Figure 5




Figure 6




Figure 7