**Differential impact of top-down and bottom-up forces in structuring
freshwater bacterial communities**

Pradeep Ram AS^{1*}, Keshri J^{1,2} and Sime-Ngando T¹

*¹Laboratoire Microorganismes: Génome et Environnement, UMR CNRS 6023, Université
Clermont-Auvergne, 63178 Aubière Cedex, France*

*²Present address: College of Veterinary Medicine, Western University of Health Sciences,
Pomona, CA-91766, USA*

Running title: Bottom-up and top-down effects on bacterial community

Accepted in FEMS Microbiology Ecology

*Correspondence: A.S. Pradeep Ram,

Laboratoire Microorganismes : Génome et Environnement,

UMR CNRS 6023,

Université Clermont-Auvergne,

1 Impasse Amélie Murat, BP 80026,

63178 Aubière Cedex,

FRANCE

E-mail : pradeep_ram.angia_sriram@uca.fr

Phone : +33 4 73 40 74 63

Fax : +33 4 73 40 76 70

ABSTRACT

Limited data exists on the simultaneous impact of bottom-up (nutrients) and top-down (viruses and heterotrophic nanoflagellates) forces in shaping freshwater bacterial communities. In our laboratory microcosms, nutrient additions (organic and inorganic) and viral reduction approach led to the proliferation of high nucleic acid (HNA) bacterial subpopulation without an increase in phage abundance. High viral-mediated bacterial lysis in the presence of nanoflagellates yielded high proportion of low nucleic acid bacterial subpopulation. 16S rRNA gene sequence analysis indicated that members of class Proteobacteria and Bacteroidetes evoked differential responses to nutrients and mortality forces thereby resulting in differences ($p < 0.001$) in bacterial community composition and diversity, as observed from ANOSIM and UniFrac analysis. Bacterial species richness (Chao) and diversity (Shannon) index was significantly higher ($p < 0.001$) in the presence of both the top-down factors and viruses alone, whereas lower host diversity was observed under nutrient relaxation of growth limiting substrates due to the explosive growth of opportunistic HNA bacterial subpopulation. Our results are in agreement with the theoretical model of 'Killing the Winner' where the availability of growth limiting substrates can act as a stimulating factor for host community composition while top-down forces can operate in the control of host diversity.

Keywords: bacteria, viruses, nutrients, bacterial community, 16S rRNA gene sequencing, freshwater microcosms

INTRODUCTION

Top-down control of bacterial populations especially by viruses has received attention in recent years since the finding of its high numbers in a wide variety of aquatic systems (Weitz and Wilhelm 2012; Sime-Ngando 2014). Together with bacterivorous nanoflagellates, viruses are now considered to be a major ecological determinant in regulating the structure and function of heterotrophic microbial communities (Suttle 2007; Sandaa *et al.* 2009). The recognition of viruses as dynamic components of planktonic communities through their destruction up to 60% of daily produced bacterial biomass in freshwater systems has challenged the conventional view that bacterial communities were largely thought to be impacted by bottom up factors. Among the interacting forces structuring bacterial communities, most attention has been on bottom-up effects (physical and chemical parameters), with lesser focus given to bacterial interaction with other segments of a microbial community such as viral or protistan activity (Bouvy *et al.* 2011; Pradeep Ram *et al.* 2016).

Within the microbial part of aquatic systems, bottom-up (nutrient supply) and top-down (protistan grazing and viral lysis) processes are known to influence and control bacterial community composition and diversity in time and space. Viruses and protists can directly impact bacterial communities either through their host specific lysis and size selective grazing respectively or indirectly through the alteration of organic pools by mortality processes (Liu *et al.* 2015). Previous studies carried out in freshwater systems have revealed that under nutrient limiting conditions, synergistic interactions between grazing and viral lysis tend to generate growth resources for fueling bacterial activity (Simek *et al.* 2001; Pradeep Ram and Sime-Ngando 2008), however their net influence on the taxonomical resolution of bacterial communities is less clearly understood. Detail examination of how the trophic interactions such as the supply of resources and mortality factors that lead to alteration of bacterial community composition and their diversity have been less studied. In seawater microcosms, relatively few moderate changes in bacterial community amidst of high viral abundance and their apparent high turnover made it difficult to distinguish statistically significant effects of viruses on bacterial community composition (Schwalbach *et al.* 2004). In perialpine lakes, multivariate ordination analyses suggested that bottom-up factors (nutrients and temperature) was the driving force behind the bacterial community shifts in spite of prevailing high viral abundances (Berdjeb *et al.* 2011). Considering the complexity and the importance of interactions between bottom-up and top-down factors, we used a reconstitution approach

using incubation of bacteria with viruses and virus-free treatments together with nutrient manipulation, and both the sources of mortality in parallel to simultaneously examine the effect of both top-down and bottom-up control on bacterial community.

In this study we extend the line of investigation to a natural freshwater system (Lake Pavin, France) to investigate the variable effect and impact of bottom-up (substrate) and top-down (viruses and heterotrophic nanoflagellates) on the bacterial community. Lake Pavin is a high altitude, meromictic freshwater lake where its late season (autumn) is characterized by lowest annual concentration of nutrient (carbon and inorganic) and chlorophyll pigments, both of which are known to have direct influence on bacterial physiology (inferred from the nucleic acid content, high versus low) and henceforth on viral life styles, namely lytic and lysogeny (Palesse *et al.* 2014). Therefore the strategies adopted by the viruses, whether to kill or not to kill can have variable impacts on certain members of bacterial community (Bouvier *et al.* 2007; Pradeep Ram and Sime-Ngando 2008; Chen *et al.* 2019). Several bacterial groups are known to have specific preferences for nutrient uptake especially during the nutrient depleted season, henceforth it can be expected that different members of the group with varying physiology and activity which tend to occupy different niches could signal changes in host-virus interactions (Øvreås *et al.* 2003; Liu *et al.* 2015).

Here we hypothesize that the relative abundances of low and high nucleic acid content bacterial subpopulation possessing differing physiological state and metabolism can strongly be influenced by the presence or type of mortality factors and nutrient supply, which eventually can evoke significant patterns and variations in bacterial community composition and diversity estimates. We also investigated the extent to which we can link bacterial diversity estimates and community composition to top-down and nutrient availability respectively. In this study, high throughput sequencing of 16S rRNA genes using Illumina MiSeq platform was the method of preference for capturing the effect of top-down and bottom-up forcing's on host taxonomical changes in bacterial community because they not only facilitates a rapid and high resolution of host community but also provides information on bacteria-virus interactions at the community level.

MATERIALS AND METHODS

Study site, sampling and experimental design

Lake water samples for the experiment was collected from oligo-mesotrophic Lake Pavin (see Palesse *et al.* [2008] for detailed characteristics) at euphotic depth (5m) using Van Dorn bottle

at the deepest central point of the lake (45°29'N ; 2°56'E) on 16 October 2015 (autumn season). The water samples collected in triplicates (i.e. three replicate hauls) were filled in clean insulated carboys and transported to laboratory well protected from heat and light radiations. Upon arrival, the samples were immediately processed for microcosm set up.

Preparation of water samples for the experimental setup is shown in **Fig. 1**. The experiment involved three fractions, 1). “grazer fraction” (containing bacteria, viruses and heterotrophic nanoflagellates, BVF) via filtration through a succession of nylon fibre sieves (150, 63 and 25 μm porosities), 2). “grazer free fraction” (containing bacteria and viruses, BV) via filtration through 1 μm pore-size filter (cellulose acetate, Sartorius) and 3). “viral reduced fraction” (containing bacteria with depleted viral abundance, B) obtained from washing and flushing out the viruses in 10 fold concentrated samples (by ultrafiltration, Model DC 10LA, Amicon®) with ultrafiltrate (virus free water) according to Weinbeuer and Suttle (1996). For obtaining bacteria and viruses twice that of *in situ* abundances (BV2x and B2x), the concentrated samples were diluted accordingly with ultrafiltrate to obtain the desired concentration. Both grazer free (BV) and viral reduced fractions (B), which received organic and inorganic nutrients, individually and in combinations, including organic carbon (glucose, 100 μM C), nitrate (NaNO_3 , 25 μM N) and phosphate (NaH_2PO_4 , 5 μM P) in three different treatments were designated as BV-C, BV-NP, BV-CNP (containing bacteria and viruses at *in situ* abundances) and B-C, B-NP, B-CNP (represents viral reduced fraction) respectively. The experimental samples were equally distributed in polycarbonate bottles (5L) and incubated in the dark at *in situ* temperature ($14.7 \pm 1^\circ\text{C}$). The samples were taken at every 12 hour interval from time zero (T0) to 108 hours (T108) for bacterial and viral abundances, and at 24 hour interval from T0 to T96 for viral lytic infection and community composition. Prior to use, all glass wares, tubings and filtration devices coming in contact with the experimental samples were acid washed (10% HCl) and thoroughly rinsed with de-ionized water. The experimental microcosms were set up within 4 hours from the time of sampling.

Our reconstitution approach involving filtration and dilution procedures were successful in obtaining desired concentration of bacterial and viral abundances among different microcosms. Viral reduction treatment was chosen to observe the response and development of bacterial communities under reduced viral top-down pressure. The followed procedure was effective in reducing viral abundance by about two fold when compared to its *in situ* concentration.

Flow cytometry

Single-celled bacteria and viruses were counted using flow cytometer (FACS Calibur, Becton Dickinson, Franklin Lake, NJ, USA). The optimized flow cytometry protocol (different sample dilutions, threshold levels, flow rates etc.) that was previously validated for the enumeration of bacteria and viruses from Lake Pavin was adopted (Palesse *et al.* 2014). Samples (2 ml each) were fixed with paraformaldehyde (0.5% final concentration) for 30 minutes in the dark at 4°C. Following fixation, the samples were diluted in 0.02 µm filtered TE buffer (10 mM Tris-HCl and 1 mM EDTA, pH 8), and stained with SYBR Green I (10,000 fold dilution of commercial stock, Molecular Probes, Oregon, USA). The mixture was incubated for 5 min at ambient temperature, heated for 10 min at 80°C in the dark, and cooled for 5 min at room temperature prior to analysis. The samples were then passed through flow cytometer equipped with an air-cooled laser providing 15 mW at 488 nm with the standard filter set-up (Brussaard *et al.* 2010). Bacteria and viruses were discriminated based on side light scatter versus green fluorescence (530 nm wave-length, fluorescence channel 1 of the instrument) plot. Flow cytometry list modes were analyzed using CellQuest Pro software (BD Biosciences, San Jose, CA, USA; version 4.0). TE buffer blanks were routinely examined to control for contamination of the equipment and reagents.

Flow cytometry allowed the identification and quantification of two major bacterial fractions: low nucleic acid content (LNA cells) and high nucleic acid content (HNA cells) based on the differences in the individual cell fluorescence (related to nucleic acid content) and in the side and forward light scattering signal on the cytogram (Gasol *et al.* 1999; Bouvier *et al.* 2007).

Heterotrophic nanoflagellate abundance

Samples for the measurements of heterotrophic nanoflagellate (HNF) abundance were fixed immediately after sampling with 1% glutaraldehyde (final concentration). Primulin-stained HNF collected on 0.8-µm polycarbonate black filters (25 mm diameter) were counted under UV excitation under a LEICA epifluorescent microscope as described by Caron (1983). A total of 200-400 nanoflagellates from each slide were counted along several transects (SD < 10%). All solutions were filter-sterilized, and a blank was routinely examined to control for contamination of equipment and reagents.

Viral lytic infection

Viral lytic infection was inferred from the frequency of visibly infected cells (FVIC) according to Pradeep Ram and Sime-Ngando (2008). Bacterial cells contained in 8 ml of

glutaraldehyde fixed water samples (final concentration 1%) were collected on copper electron microscope grids (400-mesh, carbon-coated Formvar film) by ultracentrifugation (Beckman coulter SW 40 Ti Swing-Out-Rotor at 70 000 x g for 20 minutes at 4°C). Each grid was stained at room temperature (ca. 20°C) for 30 s with Uranyl acetate (2 % w/w, pH = 4), rinsed twice with 0.02 µm-filtered distilled water, and dried on a filter paper. Grids were examined using a JEOL 1200Ex transmission electron microscope (TEM) operated at 80kV at a magnification of 20,000 to 60,000x to distinguish between bacterial cells with and without intracellular viruses. A bacterium was considered infected when at least five viruses, identified by their shape and size, were clearly visible inside the host cell. At least 400-600 bacterial cells were inspected per grid to determine FVIC. Because mature phages are visible only late in the infection cycle, FVIC counts were converted to the frequency of infected cells (FIC) using the equation $FIC = 9.524 FVIC - 3.256$ (Weinbauer *et al.* 2002).

DNA Extraction

Bacteria from the experimental water samples (150-200 ml) were collected onto 0.2µm white polycarbonate membrane filters (Sartorius, Germany) by applying low vacuum and stored for a week at -20°C. The total DNA was then extracted from the filters by using Fast DNA spin kit for soil (MP Biomedicals, Germany), which included two consecutive bead-beating steps, first, for 40 s, followed by a second bead beating 60 s to disrupt additional, previously unlysed cells. Remaining steps were followed as per manufacturer's protocol. Yield and quality of the extracted DNA samples were checked on 0.8% agarose gel, and the extracted genomic DNA was quantified, checked for purity at A260/280 nm by NanoDrop spectrophotometer (Thermo Scientific, USA.). The DNA was stored at -20 °C until further processing.

16S rRNA gene amplification and sequencing

PCR amplification was carried out to amplify V3–V4 conserved regions of bacterial 16S rRNA gene sequences in triplicate using the 16S rRNA gene specific primers (forward primer 5'-CTTTCCTACACGACGCTCTTCCGATCTACGGRAGGCAGCAG-3' and reverse 5'-GGAGTTCAGACGTGTGCTCTTCCGATCTTACCAGGGTATCTAATCCT-3') (Liu *et al.* 2007). These primers include the Illumina adapter overhang nucleotide sequences (underlined segment) as well as V3-V4-specific sequences producing an amplicon of 460 bp in length. The PCR library preparation was carried out using KAPA Hi Fi Hot Start Ready-mix PCR Kit (KAPA BIOSYSTEMS® U.S.A.). Briefly, each 25 µL of PCR reaction contains 10 ng/µL (1 µL) of genomic DNA template, 12.5 µL 2X Mastermix KAPA Hi Fidelity DNA polymerase

(1 U), 10 μ M each primer and nuclease free water. PCR reactions were carried out with initial denaturation step at 95 °C for 3 min followed by 24 cycles of 98 °C for 30 s, 65 °C for 30 s, and 72 °C for 30 s and ended with an extension step at 72 °C for 5 min. The PCR products were confirmed by 1.5% agarose gel electrophoresis. Amplicons from respective samples were pooled together in order to decrease the PCR biasness led by stochastic fluctuations in amplification efficiencies and sequenced (paired-end at a length of 300 nucleotides in each direction) on the Illumina MiSeq platform at the INRA, Toulouse (France). Pooling of PCR amplicons has not been significantly shown to influence Illumina based sequencing data and such procedure has been previously done in samples obtained from freshwater systems (Pascualt *et al.* 2014). Sequence data were received as .fastq files and has been archived in NCBI Sequence Read Archive under the accession number PRJNA479735.

Sequence data analysis

Sequence processing was performed using mothur software, version 1.39.1 (Schloss *et al.* 2009). The paired-end MiSeq Illumina reads from the samples were aligned and converted to contigs yielding 2,007,708 reads. Sequence analysis of each combined single fasta file was processed using the mothur MiSeq SOP accessed on the 10 December 2016 (Kozich *et al.* 2013). Sequences having ambiguous characters, homopolymers longer than 8 bp were removed. High-quality sequences were aligned against the mothur version of SILVA bacterial reference sequences (Pruesse *et al.* 2007) pre clustered, chimeras and singletons removed and OTUs were defined at 97% identity. Sequence processing was followed as described in Keshri *et al.* (2017). Alpha and Beta diversity was calculated on rarified data. Non-metric multidimensional scaling (NMDS) was used to visualize the similarity in bacterial communities between samples. Similarity among bacterial communities among different experimental treatments was also assessed using theayc distance matrix (Yue and Clayton 2005).

Physico chemical parameters

Water temperature and dissolved oxygen profiles were determined *in situ* using an YSI Pro ODO^{TM} probe (Yellow Springs, Ohio, USA). Phytoplankton concentration was estimated *in situ* as μ g equivalent Chlorophyll-*a* L⁻¹ (Chl) using a submersible fluorescence photometer (Fluoroprobe; BBE-Moldaenke, Kiel, Germany). Total organic carbon and total nitrogen concentrations were determined by high temperature catalytic oxidation method (680°C) using a TOC analyzer (Shimadzu TOC-V CPN, Japan).

Statistical analysis

The response of bacterial and viral parameters to different nutrient (inorganic and organic) additions was tested by analysis of variance (ANOVA), with a significant response at $p < 0.05$. Potential relationships among various measured variables were tested by Pearson's correlation analysis. All statistical analyses were performed using Minitab's software for Windows (Release 17, Minitab). Differences in bacterial community composition among the experimental treatments were assessed using UniFrac weighted distance metric (Lozupone *et al.* 2006). The randomization/permutation procedure analysis of similarities (ANOSIM) was used to evaluate the significant differences between groups. The ANOSIM statistic R displays the degree of separation between groups and is calculated by the difference of the between-group and within-group mean rank similarities. The value of $R = 1$ denotes complete separation whereas $R = 0$ suggests no separation (Clarke and Gorley 2001).

RESULTS

Initial conditions

The experiment was carried out during the clear water phase where the water temperature ($14 \pm 0.2^\circ\text{C}$), dissolved oxygen ($8.2 \pm 0.4 \text{ mg l}^{-1}$), organic carbon ($1.8 \pm 0.3 \text{ mg l}^{-1}$), total nitrogen ($0.2 \pm 0.02 \text{ mg l}^{-1}$) and chlorophyll *a* ($1.7 \pm 0.3 \text{ mg l}^{-1}$) concentrations were typical of the studied season. Phosphate concentration was below the detection limit. At the start of the experiment (T0), the mean viral and bacterial abundance were $1.8 \pm 0.2 \times 10^7 \text{ ml}^{-1}$ and $0.9 \pm 0.1 \times 10^6 \text{ cells ml}^{-1}$ respectively, with virus to bacteria ratio of 20. Low nucleic acid bacterial cells (LNA) were 8 times higher than high nucleic acid bacterial cells (HNA). Actinobacteria (69.1 %), Proteobacteria (17.3 %), and Bacteroidetes (10.5 %) together accounted for 97% of the total bacterial phyla at start of the experiment. HNF abundance estimated the start of the experiment was $1.7 \pm 0.2 \times 10^3 \text{ cells ml}^{-1}$.

Bacterial and viral abundance

The results presented here correspond to mean value (triplicates) of different incubation times (T0 to T108). Addition of nutrients (carbon+nitrogen+phosphorous) evoked a strong and significant ($F_{1,26}=29.4$, $p < 0.0001$) increase in bacterial abundance from T48 and T12 in BV-CNP and B-CNP compared to T0 (**Fig. 2A&B**). The maxima in bacterial abundance that was observed at T108 ($12.6 \pm 0.5 \times 10^6 \text{ cells ml}^{-1}$) and T72 ($15.0 \pm 0.8 \times 10^6 \text{ cells ml}^{-1}$) in BV-

CNP and B-CNP respectively was one order of magnitude higher than the abundance at the start of the experiment. In both the cases the increase in bacteria was never accompanied by proportional increase in viral abundances, thereby resulting in significantly lower ($F_{1,24}=27.5$, $p < 0.0001$) virus to bacteria ratio (range = 1.1-2.2) compared to microcosms without nutrient additions (range = 11.1-23.1). Except in CNP treatments, viruses followed host abundance in the grazer and grazer free fraction resulting in a significant correlation ($r = 0.88$, $p < 0.001$) between the two. Generally, in the viral reduced microcosms continuous or moderate decline in viral abundance was observed from start to end of the experiment (**Fig. 2B**)

Addition of nutrient limiting substrates and viral reductions contributed to the variability of HNA and LNA populations with incubation time. In the microcosms without nutrient additions (BVF, BV and BV2X), LNA was always found to dominate HNA cells (LNA:HNA = 1.4 to 13.9), whereas in microcosms with carbon supplements the reverse was observed (HNA:LNA = 1.0 to 15.2) (**Fig. 3**). In all viral reduced microcosms, HNA was the dominant subpopulation (**Fig. 3**).

Phage infection

Viral lysis was more evident in the grazer treatment where the highest percentage of viral infection (30.3%) coincided with the maxima of HNF abundance at T48 (**Supplementary material, Fig. S1**). Overall, viral lysis in the grazer and grazer free fraction varied by a factor of 10 and was significantly ($F_{1,18}=30.1$, $p < 0.0001$) higher compared viral reduced treatments (**Fig. 4A&B**). High viral lysis did not bring about significant variability in virus to bacteria ratio suggesting a constant production and loss. The significant increase in host population following nutrient addition in grazer free fraction was never accompanied by increase in lytic infection.

16S rRNA gene sequencing analysis

A total of 1,168,347 effective sequences after quality and chimera removal were analyzed further with over 15,000 sequences in each library. Clustering at 97% similarity resulted in a total of 9,183 OTUs. All of the rarefaction curves reached saturation indicating that sequencing depth was considerable enough to cover the whole bacterial diversity (**Supplementary material, Fig. S2**).

Bacterial community diversity and composition in different microcosms

Incubations of the resident bacterial assemblage with the presence of viruses and grazers (BVF), viruses alone (BV) and under viral reduction (B) yielded distinct signatures in bacterial diversity and community composition. The variation in bacterial richness (Chao = 328-383) and diversity (Shannon = 3.58-3.75) was higher in the presence of both the loss factors throughout the experiment. Overall, the bacterial diversity under reduced viral concentrations was significantly lower ($p < 0.001$) compared to microcosms with *in situ* viral abundances (**Supplementary material, Table S1**). Addition of glucose and inorganic nutrients in BV and B microcosms promoted shifts in bacterial community dominants as a function of carbon source preference, resulting in lower bacterial diversity.

In microcosms, the distribution and relative abundance of different bacterial groups which was visible at the genus level. Out of 16 different phyla along with unclassified bacterial members with varying abundance, 3 phyla namely Actinobacteria, Proteobacteria and Bacteroidetes, approximately accounted for 99% of the total sequences with varying abundances. The members of Actinobacteria (represented by genera *clade hgcI*, *CL500-29 marine group*, *Candidatus Planktophila*) which were the dominant group at the start of the experiment decreased linearly with time. Proteobacterial members such as Alphaproteobacteria, Betaproteobacteria, Gammaproteobacteria and Bacteroidetes were highly dynamic and showed variable response with progress in incubation time in different microcosms.

Within the class Betaproteobacteria, *Limnohabitans* increased 3 to 5 fold in BV-CNP treatment, whereas in the viral reduced fractions a notable increase in the relative abundance of genera *Delftia*, *Duganella* and *Pelomonas* was observed (**Fig. 5**). Similarly the increase of *Caulobacter* belonging to class Alphaproteobacteria by 6.7 fold compared to start of the experiment was evident in B-NP microcosms. The strong increase of Gammaproteobacteria by 53 and 70.3 fold in the BV-CNP and B-CNP treatment respectively eventually coincided with high percentage increase of HNA cells. This response of Gammaproteobacteria to carbon addition was represented by members belonging to genera *Pseudomonas* and *Acinetobacter* in viral reduced and *Pseudomonas* only in microcosm containing *in situ* abundances. Among the class Bacteroidetes, the genera *Flavobacterium* and *Sediminibacterium* which showed ten and two-fold increase respectively in the presence of both the loss factors only. Overall the sequences from all the microcosms were classified into 29 classes, 69 orders, 135 families and 281 genera.

Two dimensional non-metric multidimensional scaling accounted for bacterial community differences (stress = 0.08, $r^2 = 0.97$) (**Fig. 6**). The variations towards first axis of

non-metric multidimensional scaling is mostly contributed by OTU2 (*Pseudomonas*) OTU43 (env.OPS_17_ge from family Sphingobacteriales), OTU122 (*Polaromonas*) and OTU100 (*Rhodococcus*). However, OTU10 (*Sediminibacterium*), OTU66 (Microbacteriaceae), OTU5 (*Pseudomonas*) and OTU3 (*Pseudomonas*) contributed most to the variations of second axis. Based on the distribution of OTUs, ANOSIM and weighted UniFrac analyses indicated significant differences ($p < 0.001$) in bacterial community composition in grazer, grazer free and viral reduced microcosms (**Supplementary material Table S2A&B**). The clustering of samples from microcosms with ambient and reduced viral abundance is clearly visible except in the case of CNP supplemented samples. The dendrogram based on thetacy calculators confirmed the hierarchical clustering of bacterial community composition in samples in two clades of the tree and raw samples branched closely with samples from ambient viral abundances microcosms (**Supplementary material Fig. S3**).

DISCUSSION

This study provides evidence on the differential impact of top-down (viruses and grazers) and bottom-up (nutrients) forces on bacterial community structure (diversity and composition) which caused shifts in bacterial subpopulation with differing physiological state and metabolism. Two dimensional non-metric multidimensional scaling accounted for structural differences in bacterial community, whereas ANOSIM and UniFrac analysis highlighted for significant differences in bacterial communities in experimental microcosms.

Nutrient impact on bacterial physiology

The response of natural bacterial communities to nutrients additions can be linked closely to the state of the ecosystem (Øvreås *et al.* 2003). During the studied period of the year, multiple nutrient limitation of bacterial community was evident with large proliferation of high nucleic acid content bacterial subpopulation in experimental microcosms supplemented with organic and inorganic nutrients (Fig. 2). However, the inability of bacterial cells to divide but increase in their biovolume in microcosms with addition of carbon source alone is suggestive of strong co-limitation of phosphate source (Carlsson and Caron 2001). Phosphate is known to enhance cell division, whereas their strong limitation could result in the storage of organic carbon in cells without dividing and thus maintain higher carbon metabolism (del Giorgio and Cole 1998). The dominance of LNA at the start of the experiment and subsequent increase of HNA cells following the addition of carbon source

suggest that these two groups may differentiate according to the prevailing nutrient environment and thus play different ecological roles. The explosive growth of HNA cells with nutrient additions is consistent with the reports that an enrichment of copiotrophic bacterial phylotypes (such as those comprise the HNA subpopulation) coincides with bacterial productivity and biomass maxima (Nelson *et al.* 2014; Mojica *et al.* 2019). Therefore, nutrient conditions which are bound to vary at different time periods especially in natural freshwater system could serve as a potential influencing factor in altering the bacterial physiological state with high and low concentrations favoring HNA and LNA cells respectively (Palesse *et al.* 2014; Liu *et al.* 2015).

The expression of HNA cells under nutrient relaxation could be a part of growth strategy where certain bacteria such as *Vibrio* are capable of surviving extended periods of carbon starvation and grow rapidly at high substrate concentrations (Øvreås *et al.* 2003). Our results support and suggest the fact that HNA cells could be fast growing, diverse and be more adapted to changing environments, using their repertoire of transport and regulation genes (Philippot *et al.* 2010). Such versatility and fast growing nature of HNA cells which possess more flexible genes allow them to exploit pulses of nutrients (Mojica *et al.* 2019) and occupy more ecological niches (Schattenhofer *et al.* 2011). Cytometric signatures have revealed that both HNA and LNA cells may represent physiologically and ecologically distinct bacterial populations that have components which are intrinsic to themselves and this distribution may in fact be linked to more general adaptive processes within the bacteria (Bouvier *et al.* 2007; Mojica *et al.* 2019). This bimodal distribution could probably play a key role in the functioning of bacterioplankton communities.

Linking top-down and bottom-up effects to bacterial community dynamics

Top-down forces and nutrients had variable impact on bacterial community. High viral lysis in the presence of top-down forces in non-amended microcosms containing microbial populations at *in situ* abundances and low viral impact on bacterial population with nutrient additions brought about changes in bacterial community composition and diversity. The explosive increase of fast-growing opportunistic bacteria (i.e. HNA cells) in nutrient spiked microcosms was never accompanied by the increase in phage abundance and infection, thus differing with the reports from marine systems where the limiting nutrient was found to constrain bacterial growth and consequently phage-mediated host cell lysis (Williamson and Paul 2004). Low viral impact on bacterial communities could likely be due to tradeoff which can exists between bacterial traits of antiviral defense (reduction and masking of receptors or

viral adsorption) and resource competition, where organic enrichment may lower resource competition and allow bacterial communities to pay the costs involved in increasing antiviral defense (Thingstad *et al.* 2005). Moreover extracellular proteases produced by rapidly growing bacteria can largely play a role in the hydrolytic destruction or modification of viral capsids thus alleviating viral lytic pressure leading to uncoupling between bacterial and viral production (Motegi and Nagata, 2009). In microbial model systems, the magnitude of trade-off between host and phages are largely influenced by environmental factors especially resources (nutrients) where its limitation can result in the coexistence of both susceptible and resistant hosts in natural communities (Friman and Buckling 2013; Record *et al.* 2016). Such a scenario of viral defense trade-offs through the expression of host defense mechanisms such as CRISPR-Cas (Barrangou *et al.* 2007) and restriction enzymes (Labrie *et al.* 2010) can have predictable effects on their community structure, properties, dynamics and also on their response to environmental changes (Stearns 1989; Jessup and Bohannan 2008). Host defense strategies which tend to contribute to the shape of trade-off curve in a predator-prey system can be either concave (cheap defense) or convex (costly defense) depending on the presence of ancestral and coevolved predators respectively (Huang *et al.* 2017). The presence of such dynamical trade-off between host and phage, which vary with the environment can eventually affect their ecological interactions and subsequently drive evolutionary changes (Jessup and Bohannan 2008).

The application of next generation sequencing to sequence 16S rRNA genes facilitated in obtaining greater information on taxonomic resolution of bacterial community and their diversity pattern in microcosms. Bacterial diversity was highest in the presence of both the top-down factors indicating that they had tighter control on bacterial communities (Zhang *et al.* 2007). Contrastingly, the addition of carbon source resulted in lower bacterial diversity estimates due to the proliferation of certain communities which have the ability to rapidly consume available resources (Sandaa *et al.* 2009; Topper *et al.* 2012). It has been previously demonstrated that both nutrient and loss factors tend to operate simultaneously in freshwater systems where the former can evoke strong changes in the physiology and metabolic activity of certain members of bacterial population which can eventually make those susceptible communities selectively vulnerable to the later, thus impacting their diversity (Storesund *et al.* 2015; Pradeep Ram *et al.* 2016). The observed differences in bacterial communities in response to top-down and bottom-up factors in different treatments with the progress of experiment was highlighted by ANOSIM and weighted UniFrac analyses which is based on the distribution of OTUs where significant difference between the bacterial groups are

reflected in the differences in their community composition. This was duly supported by two dimensional non-metric multidimensional scaling based on thetacy dissimilarity scores to account for community differences.

The members of class Gamma-, Beta- and Alphaproteobacteria which evoked contrasting and differential responses to nutrient additions and predators are known to belong to HNA subpopulation (Schattenhofer *et al.* 2011; Mojica *et al.* 2019), while lineages from Actinobacteria were selected against in all the treatments. The low detection of Gammaproteobacteria under high viral infection rates in the presence top-down factors could be explained by their potential vulnerability to loss factors, whereas their sharp increase in with nutrient supplements is suggestive of their opportunistic lifestyle and adaptability to grow under high-nutrient concentrations (Øvreås *et al.* 2003; Pinhassi and Berman 2003), much quicker than the viruses could infect them. It was interesting to note that even within *Gammaproteobacteria*, different genera such as *Pseudomonas* and *Alteromonas* had variable responses to nutrient additions and viral reductions respectively.

Although it is very well known that Betaproteobacteria are more substrate responsive, their significant presence of genus *Limnohabitans* amid both the top-down factors are justified by their ability to efficiently utilize regenerated nutrients from grazing and viral lysis under nutrient limiting conditions (Simek *et al.* 2001; Pradeep Ram and Sime-Ngando 2008). The high expression of minor genera *Pelomonas*, *Delftia*, and *Duganella* (Betaproteobacteria) and genus *Caulobacter* (Alphaproteobacteria) under reduced viral conditions are suggestive of their vulnerability to viral infection wherein the scenario of host characteristics could prevail over density (Bouvier and del Giorgio 2007). The increased representation of *Sphingobacteria* and *Flavobacteria* (Bacteroidetes) in the presence of both the top down factors are suggestive of its ability to change their morphology (filamentous and floc formation) to escape grazing and at the same time efficiently use regenerated nutrients (Hornak *et al.* 2005).

The generalizations from a single microcosm experiment should be done with care, even when the experiment was carried out with a defined set of controls, parallels and factorial design. The initial bacterial community composition that was determined at the start of the experiment in the grazer fraction was also applied to other fractions (grazer free and viral reduced) assuming that the initial differences between treatments may overall be small. However such assumption could perhaps be associated with certain amount of bias in our experimental microcosms. Despite of methodological constraints and pit falls associated with our chosen experimental approach, we were able to demonstrate the variable influence of top-down factors especially viral lysis and nutrients on bacterial diversity and community

composition. The expression of HNA over LNA subpopulation under nutrient enriched and viral depletion was reflected in the changes in bacterial community composition at the genus level. Our study is in agreement within the conceptual idea of simple model version of ‘Killing the Winner (KtW)’ theory, which postulates that bacterial community diversity is a feature that is essentially top-down controlled by viruses, while community composition is bottom-up controlled by competition for limiting nutrients (Töpper *et al.* 2012; Storesund *et al.* 2015). Such a model where the two separate control mechanisms for composition and diversity could be envisaged in our study and therefore can be extended for freshwater bacterial communities. Therefore accordingly, our findings indicate that the competition for growth limiting nutrients can act as a structuring factor for the host community composition which occupies available niches, while top-down factors control the identity and abundance of bacterial hosts within each niche.

ACKNOWLEDGEMENTS

JK was supported by a postdoctoral fellowship from the Université Clermont Auvergne (formerly Université Blaise Pascal), Clermont Ferrand (France). We thank J. Colombet and F. Perriere for their technical assistance in flow cytometry and nutrient analysis respectively. We appreciate two anonymous reviewers and editor for their time, effort and valuable contributions towards this manuscript.

CONFLICT OF INTEREST: None

REFERENCES

- Barrangou R, Fremaux C, Deveau *et al.* Crispr provides acquired resistance against viruses in prokaryotes. *Science* 2007;**315**:1709-12.
- Berdjeb L, Ghiglione J-F, Jacquet S. Bottom-up versus top-down control of hypo- and epilimnion free-living bacterial community structures in two neighboring freshwater lakes. *Appl Environ Microbiol* 2011;**77**:3591–9.
- Bouvier T, del Giorgio PA, Gasol JM. A comparative study of cytometric characteristics of high and low nucleic acid bacterioplankton cells from different aquatic systems. *Environ Microbiol* 2007;**9**:2050–66.
- Bouvier T, del Giorgio PA. Key role of selective viral-induced mortality in determining marine bacterial community composition. *Environ Microbiol* 2007;**9**:287–97.

- 530 Bouvy M, Bettarel Y, Bouvier C *et al.* Trophic interactions between viruses, bacteria and
 531 nanoflagellates under various nutrient conditions and simulated climate change.
 532 *Environ Microbiol* 2011;**13**:1842–57.
- 533 Brussaard C, Payet JP, Winter C *et al.* Quantification of aquatic viruses by flow cytometry.
 534 In: Wilhelm SW, Weinbauer MG, Suttle C (eds). *Manual of Aquatic Viral Ecology*:
 535 ASLO, Chapter 11, 2010, 102–9.
- 536 Carlsson P, Caron DA. Seasonal variation of phosphorous limitation of bacterial growth in a
 537 small lake. *Limnol Oceanogr* 2001;**46**:108-20.
- 538 Caron DA. Techniques for enumeration of heterotrophic and phototrophic nanoplankton using
 539 epifluorescent microscopy, and comparison with other procedures. *Appl Environ*
 540 *Microbiol* 1983;**46**:491-98.
- 541 Chen X, Ma R, Yang Y *et al.* Viral regulation on bacterial community impacted by lysis-
 542 lysogeny switch: A microcosm experiment in eutrophic coastal waters. *Front*
 543 *Microbiol* 2019;**10**:1763.
- 544 Clarke KR, Gorley RN. *PRIMER v5: User manual/tutorial*. PRIMER-E, Plymouth:UK,
 545 2001, 91.
- 546 del Giorgio PA, Cole JJ. Bacterial growth efficiency in natural aquatic ecosystems. *Annu Rev*
 547 *Ecol Syst* 1998;**29**:503–41.
- 548 Friman V-P, Buckling A. Effects of predation on real-time host-parasite coevolutionary
 549 dynamics. *Ecol Lett* 2013;**16**:39-46.
- 550 Gasol JM, Zweifel UL, Peters F *et al.* Significance of size and nucleic acid content
 551 heterogeneity as measured by flow cytometry in natural planktonic bacteria. *Appl*
 552 *Environ Microbiol* 1999;**65**:4475–83.
- 553 Hornak K, Masin M, Jezbera J *et al.* Effects of decreased resource availability, protozoan
 554 grazing and viral impact on the structure of bacterioplankton assemblage in a canyon-
 555 shaped reservoir. *FEMS Microbiol Ecol* 2005;**52**:315–27.
- 556 Huang W, Traulsen A, Werner B *et al.* Dynamical trade-offs arise from antagonistic
 557 coevolution and decrease intraspecific diversity. *Nat Commun* 2017;**8**:2059.
- 558 Jessup CM, Bohannan BJM. The shape of ecological trade-off varies with environment. *Ecol*
 559 *Lett* 2008;**11**:947-59.
- 560 Keshri J, Pradeep Ram AS, Colombet J *et al.* Differential impact of lytic viruses on the
 561 taxonomical resolution of freshwater bacterioplankton community structure. *Water*
 562 *Res* 2017;**124**:129–38.

- 563 Kozich JJ, Westcott SL, Baxter NT *et al.* Development of a dual-index sequencing strategy
564 and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina
565 sequencing platform. *Appl Environ Microbiol* 2013;**79**:5112–20.
- 566 Labrie SJ, Samson JE, Moineau S. Bacteriophage resistance mechanisms. *Nat Rev Microbiol*
567 2010;**8**:317–27.
- 568 Liu H, Yuan X, Xu J *et al.* Effects of viruses on bacterial functions under contrasting
569 nutritional conditions for four species of bacteria isolated from Hong Kong waters. *Sci*
570 *Rep* 2015;**5**:14217.
- 571 Liu L, Yang J, Yu Z *et al.* The biogeography of abundant and rare bacterioplankton in the
572 lakes and reservoirs of China. *ISME J* 2015;**9**:2068–77.
- 573 Liu Z, Lozupone C, Hamady M *et al.* Short pyrosequencing reads suffice for accurate
574 microbial community analysis. *Nucleic Acids Res* 2007;**35**:e120.
- 575 Lozupone C, Hamady M, Knight R. UniFrac—an online tool for comparing microbial
576 community diversity in a phylogenetic context. *BMC Bioinformatics* 2006;**7**:371.
- 577 Mojica KDA, Carlson CA, Behrenfeld MJ. Regulation of low and high nucleic acid
578 fluorescent heterotrophic prokaryote subpopulations and links to viral-induced
579 mortality within natural prokaryote-virus communities. *Microb Ecol* 2019,
580 DOI:10.1007/s00248-019-01393-9.
- 581 Motegi C, Nagata T. Addition of monomeric and polymeric organic substrates alleviates viral
582 lytic pressure on bacterial communities in coastal seawaters. *Aquat Microb Ecol*
583 2009;**57**:343–50.
- 584 Nelson CE, Carlson CA, Ewart CS *et al.* Community differentiation and population
585 enrichment of Sargasso Sea bacterioplankton in the euphotic zone of a mesoscale
586 mode-water eddy. *Environ Microbiol* 2014;**16**:871–87.
- 587 Øvreås L, Bourne D, Sandaa RA *et al.* Response of bacterial and viral communities to
588 nutrient manipulations in seawater microcosms. *Aquat Microb Ecol* 2003;**31**:109–21.
- 589 Palesse S, Colombet J, Pradeep Ram AS *et al.* Linking host prokaryotic physiology to viral
590 lifestyle dynamics in a temperate freshwater lake (Lake Pavin, France). *Microb Ecol*
591 2014;**68**:740–50.
- 592 Pascault N, Roux S, Artigas J *et al.* A high-throughput sequencing ecotoxicology study of
593 freshwater bacterial communities and their responses to tebuconazole. *FEMS Microb*
594 *Ecol* 2014;**90**:563–74.
- 595 Philippot L, Andersson SGE, Battin TJ *et al.* The ecological coherence of high bacterial
596 taxonomic ranks. *Nat Rev Microbiol* 2010;**8**:523–9.

- 597 Pinhassi J, Berman T. Differential growth response of colony-forming α - and γ -proteobacteria
 598 in dilution culture and nutrient addition experiments from Lake Kinneret (Israel), the
 599 Eastern Mediterranean Sea, and the Gulf of Eilat. *Appl Environ Microbiol*
 600 2003;**69**:199–211.
- 601 Pradeep Ram AS, Chaibi-Slouma S, Keshri J *et al.* Functional responses of bacterioplankton
 602 diversity and metabolism to experimental bottom-up and top-down forcings. *Microb*
 603 *Ecol* 2016;**72**:347–58.
- 604 Pradeep Ram AS, Sime-Ngando T. Functional responses of prokaryotes and viruses to grazer
 605 effects and nutrient additions in freshwater microcosms. *ISME J* 2008;**2**:498–509.
- 606 Pruesse E, Quast C, Knittel K *et al.* SILVA, a comprehensive online resource for quality
 607 checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic*
 608 *Acids Res* 2007;**35**:7188–96.
- 609 Record NR, Talmy D, Våge S. Quantifying tradeoffs for marine viruses. *Front Microbiol*
 610 2016;**3**:251.
- 611 Sandaa R-A, Gomez-Consarnau L, Pinhassi J *et al.* Viral control of bacterial biodiversity—
 612 evidence from a nutrient enriched marine mesocosm experiment. *Environ Microbiol*
 613 2009;**11**:2585–97.
- 614 Schattenhofer M, Wulf J, Kostadinov I *et al.* Phylogenetic characterization of picoplanktonic
 615 populations with high and low nucleic acid content in the North Atlantic Ocean. *Syst*
 616 *Appl Microbiol* 2011;**34**:470–5.
- 617 Schloss PD, Westcott SL, Ryabin T *et al.* Introducing mothur: open-source, platform-
 618 independent, community-supported software for describing and comparing microbial
 619 communities. *Appl Environ Microbiol* 2009;**75**:7537–41.
- 620 Schwalbach M, Hewson I, Fuhrman J. Viral effects on bacterial community composition in
 621 marine plankton microcosms. *Aquat Microb Ecol* 2004;**34**:117–27.
- 622 Simek K, Pernthaler J, Weinbauer MG *et al.* Changes in bacterial community composition,
 623 dynamics and viral mortality rates associated with enhanced flagellate grazing in a
 624 meso-eutrophic reservoir. *Appl Environ Microbiol* 2001;**67**:2723–33.
- 625 Sime-Ngando T. Environmental bacteriophages: viruses of microbes in aquatic systems.
 626 *Front Microbiol* 2014;**5**:1–14.
- 627 Stearns SC. Trade-offs in life-history evolution. *Funct Ecol* 1989;**3**:259–68.
- 628 Storesund JE, Erga SR, Ray JL *et al.* Top-down and bottom-up control on bacterial diversity
 629 in a western Norwegian deep-silled fjord. *FEMS Microb Ecol* 2015;**91**:7.

- 630 Suttle CA. Marine viruses—major players in the global ecosystem. *Nat Rev Microbiol* 2007;**5**:
631 801–12.
- 632 Thingstad TF, Øvreås L, Egge JK *et al.* Use of non-limiting substrates to increase size; a
633 generic strategy to simultaneously optimize uptake and minimize predation in pelagic
634 osmotrophs? *Ecol Lett* 2005;**8**:675–82.
- 635 Töpper B, Thingstad TF, Sandaa RA. Effects of differences in organic supply on bacterial
636 diversity subject to viral lysis. *FEMS Microb Ecol* 2012;**83**:202–13.
- 637 Weinbauer MG, Hornak K, Jezbera J *et al.* Synergistic and antagonistic effects of viral lysis
638 and protistan grazing on bacterial biomass, production and diversity. *Environ*
639 *Microbiol* 2007;**9**:777–88.
- 640 Weinbauer MG, Suttle CA. Potential significance of lysogeny to bacteriophage production
641 and bacterial mortality in coastal waters of the Gulf of Mexico. *Appl Environ*
642 *Microbiol* 1996;**62**:4374–80.
- 643 Weinbauer MG, Winter C, Höfle MG. Reconsidering transmission electron microscopy based
644 estimates of viral infection of bacterioplankton using conversion factors derived from
645 natural communities. *Aquat Microb Ecol* 2002;**27**:103–10.
- 646 Weitz JS, Wilhelm SW. Ocean viruses and their effects on microbial communities and
647 biogeochemical cycles. *F1000 Biol Rep* 2012;**4**:17.
- 648 Williamson SJ, Paul JH. Nutrient stimulation of lytic phage production in bacterial
649 populations of the Gulf of Mexico. *Aquat Microb Ecol* 2004;**36**:9–17.
- 650 Yue JC, Clayton MK. A similarity measure based on species proportions. *Commun Stat –*
651 *Theor M* 2005;**34**:2123–31.
- 652 Zhang R, Weinbauer MG, Qian PY. Viruses and flagellate sustain apparent richness and
653 reduce biomass accumulation of bacterioplankton in coastal marine waters. *Environ*
654 *Microbiol* 2007;**9**:3008–18.