Wave and Tide Controls on Rip Current Activity and Drowning Incidents in Southwest France

Bruno Castelle, Tim Scott, Rob Brander, R Jak Mccarroll, Eric Tellier, Elias de Korte, Léana Tackuy, Arthur Robinet, Bruno Simonnet, Louis-Rachid Salmi

► To cite this version:

Bruno Castelle, Tim Scott, Rob Brander, R Jak Mccarroll, Eric Tellier, et al.. Wave and Tide Controls on Rip Current Activity and Drowning Incidents in Southwest France. *Journal of Coastal Research*, 2020, 95, 10.2112/si95-150.1 . hal-03044408

HAL Id: hal-03044408

<https://hal.science/hal-03044408>

Submitted on 14 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Wave and Tide Controls on Rip Current Activity and Drowning Incidents in Southwest France

Bruno Castelle^{*,†}, Tim Scott[‡], Rob Brander[§], R. Jak McCarroll[‡], Eric Tellier^{†,‡,§}, Elias de Korte^{§§}, Léana Tackuy[†], Arthur Robinet^{†,††}, Bruno Simonnet^{‡,§}, and Louis-Rachid Salmi^{†,‡,§}

[†]CNRS, Université de Bordeaux
UMR EPOC 6554,
OASU
Pessac, France

[‡]Coastal Processes
Research Group
School of Biological and
Marine Sciences
University of Plymouth
Plymouth, UK

[§]School of Biological,
Earth and
Environmental
Sciences
University of New South
Wales
Sydney, Australia

^{††}INSERM, Université
de Bordeaux
U1219 ISPED
Bordeaux population
health research
Bordeaux, France

www.cerf-jcr.org

^{‡‡}CHU de Bordeaux
Pôle Urgences adultes,
SAMU-SMUR
Pôle de Santé Publique,
SIM/UMES
Bordeaux, France

^{§§}IMAR, Department of
Physical Geography,
Faculty of Geosciences,
Utrecht University
Utrecht, Netherlands

^{†††}BRGM, French
Geological Survey
Pessac, France

www.JCRonline.org

ABSTRACT

Castelle, B.; Scott, T.; Brander, R.W.; McCarroll, R.J.; Tellier, E.; de Korte, E.; Tackuy, L.; Robinet, A.; Simonnet, B. and Salmi, L.-R., 2020. Wave and tide controls on rip current activity and drowning incidents in Southwest France. In: Malvárez, G. and Navas, F. (eds.), *Global Coastal Issues of 2020. Journal of Coastal Research*, Special Issue No. 95, pp. 769–774. Coconut Creek (Florida), ISSN 0749-0208.

The meso-macrotidal sandy surf beaches of southwest France are a popular destination visited by millions of tourists each summer, who potentially expose themselves to surf zone hazards and particularly to rip currents flowing through the inner-bar rip channels. 281 non-fatal and fatal drowning incidents, most of them caused by rip currents, recorded by lifeguards during the summer 2007, 2009 and 2015 were combined with measured and hindcast wave and tide data. Results show that drownings occur disproportionately near neap low tide, under shore-normal incident waves and average to above-average height and period, with drowning incidents tending to occur in clusters with particular days of mass incidents. An XBeach model is implemented on measured rip-channelled bathymetries to address the influence of offshore wave conditions and tidal elevation on rip flow dynamics and resulting hazard. Simulations show that rip flow increases with increasing wave height, increasing wave period and increasingly shore-normal incidence, which is consistent with the increased number of drowning incidents for such conditions. Although more incidents also tend to occur on warm sunny days with light winds, presumably driving more exposure to the rip current hazard, this highlights the dominance of the physical hazard on the life risk along this stretch of coast.

ADDITIONAL INDEX WORDS: *Rip current, drowning incident, environmental controls, beach safety.*

INTRODUCTION

It is well established that rip currents are the leading cause of both fatal and non-fatal drowning incidents on surf beaches worldwide. Rips are strong, narrow seaward flowing currents often originating close to the shoreline (Castelle *et al.*, 2016a; Dalrymple *et al.*, 2011; MacMahan *et al.*, 2006) which, although potentially forming through a variety of driving mechanisms (Castelle *et al.*, 2016a), are essentially driven by the action of breaking waves. One of the most common rip types flows through channels incised in nearshore sandbars (e.g., Houser *et al.*, 2013, MacMahan *et al.*, 2006). Channel rip activity, and resulting physical hazard, is

primarily controlled by offshore wave conditions (e.g., Brander, 1999; Bruneau *et al.*, 2011) and tide elevation (e.g., Austin *et al.*, 2014). Surf zone morphology is also critical to rip activity, with deeper rip channels resulting in more intense rips (e.g., McCarroll *et al.*, 2018, Moulton *et al.*, 2017).

The number of drowning incidents, or the overall drowning risk at a given coast, not only depends on the physical hazard (primarily rip currents), but also on beachgoer exposure. Warm sunny days with low winds typically result in increased beach attendance and beachgoer exposure to hazards and, in turn, more life risk. Understanding the environmental controls on life risk related to rip currents along the coast typically requires accurate drowning incident records combined with detailed wave, tide and weather data at the time of the incidents (Scott *et al.*, 2014).

The southwest Gironde-Landes coast of France hosts 230 km of meso-macrotidal sandy surf beaches (Figure 1a) that are visited by

DOI: 10.2112/SI95-150.1 received 31 March 2019; accepted in revision 13 February 2020.

*Corresponding author: bruno.castelle@u-bordeaux.fr

©Coastal Education and Research Foundation, Inc. 2020

Figure 1. (a) Location map of the Gironde and Landes coasts in southwest France. Yellow stars indicate tide gauges, and the red and pink polygons show the location of the WavewatchIII model output grid point and weather station, respectively. (b) Aerial photograph of Truc Vert beach taken at low tide showing the presence of a classic rip current system along this coast (credit V. Marieu).

millions of tourists each summer. Beaches are relatively straight (Figure 1b) and are exposed to high-energy ocean waves coming from the WNW direction (Castelle *et al.*, 2017). Beaches are double-barred and rip-dominated (*e.g.*, Bruneau *et al.*, 2009) with rip current causing approximately 80% of drowning incidents (Castelle *et al.*, 2018). Here we examine extensive environmental data to address the primary environmental controls on the full drowning risks along this stretch of coast, with the overarching goal to improve future beach safety management and education of beach users in southwest France.

METHODS

Drowning Data

A comprehensive surf-zone injury (SZI) dataset, that comprised injury report forms filed for every incident, documented a total of 2523 SZIs over 186 sample days during the summers of 2007, 2009 and 2015 (Castelle *et al.*, 2018). Amongst all the data analyzed in Castelle *et al.* (2018), here we address 281 drowning incidents (Figure 2a,b) using: date and time of the incident; beach location; activity (*e.g.*, wading, surfing, bodyboarding); cause of injury (*e.g.*, rip current, shore-break waves); and injury type (*e.g.*, drowning, spine injury). The drowning stage was also provided according to a 4-stage classification widely used in France, from the milder to the most severe: (1) exhaustion, but no sign of aspiration of water; (2) moderate respiratory impairment, anxiety; (3) altered consciousness, severe respiratory impairment or acute pulmonary edema, tachycardia or hypotension; and (4) coma, respiratory or cardiac arrest. 11.4% of the drowning incidents were severe (stage 3 and 4, Figure 2b), with most occurring outside of the supervised bathing zone (Figure 2d).

Environmental Data

Each incident was associated with: a significant wave height H_s , mean and peak wave periods T_{02} and T_p , respectively, and wave angle θ from a 6-hourly wave hindcast (Bouidière *et al.*, 2013) at grid point 1.3232°W, 44.7374°N (Figure 1a) in approximately 30-m depth. A tidal component analysis of a 3-month time series of continuous, storm-free, tide gauge data (Figure 1a) was used to generate tide elevation time series.

Figure 2. (a) Surf zone incidents related to injury type. (b) Drowning incidents related to drowning stage and location of the incident related to (c) milder drowning (stage 1 & 2) and (d) severe drowning (stage 3 & 4) with respect to supervised bathing zone.

Figure 3. Time series during the 2009 summer of: (a) number of daily fatal and non-fatal drowning incidents; (b) significant wave height H_s ; (c) peak (T_p , black line) and mean (T_{02} , blue line) wave periods; (d) wave direction θ ; (e) water level η with respect to mean sea level. In (b,c,d) the summer mean is indicated by the horizontal dashed line

Figure 4. Environmental controls on drowning incidents: normalized frequency distributions F_n during the summers of 2007, 2009 and 2015 (light grey region), referred to as 'average' background distribution, of (a,b) significant wave height H_s ; (c,d) mean wave period T_{02} ; (e,f) wave direction θ ; (g,h) water level η . In all panels the dark grey bars show the difference between the surfing related and the 'average' background distributions and the vertical dashed lines in (a-f) show the background T_{op} and bottom panels are for milder (stage 1 & 2) and severe (stage 3 & 4) drowning incidents, respectively.

Numerical Modelling

An XBeach model (Roelvink *et al.*, 2009) was implemented on a 10 m x 10 m regular grid to address the influence of offshore wave conditions and tidal elevation on rip flow dynamics and resulting hazard. The model, which is depth-averaged and wave-group resolving, has been demonstrated to be suitable for this purpose even when default settings were used (Castelle *et al.*, 2016b). The model was based on detailed topo-bathymetry collected during an intensive field experiment performed at Biscarosse on June 13-17 2007 (Bruneau *et al.*, 2009, 2011), which is broadly representative of the coastal morphology. For more detail on the model implementation at Biscarosse the reader is referred to Castelle *et al.* (2016b, 2019).

RESULTS

Figure 3 shows time series of wave/tide conditions and daily drowning incidents during the 2009 summer. Drowning incidents tend to occur in clusters. Overall, there appears to be no single environmental factor controlling the occurrence of drowning incidents (also considering weather factors, not shown). Instead, it is possible that certain combinations of environmental controls are conducive to causing drowning incidents. To better understand these combinations promoting risky conditions, the approach of Scott *et al.* (2014) was utilized whereby the environmental conditions during which incidents occurred were compared with the 'average' background conditions of the summer seasons. The average frequency distribution of these parameters was compared with those computed from the environmental parameters associated with each recorded rip incident. Differences between the distributions therefore provides an indication of environmental

conditions that may be driving drowning incidents (Scott *et al.*, 2014).

Figure 4 shows that, for both milder and severe drowning incidents, larger waves (Figure 4a,b) with longer period (Figure 4c,d) and more shore-normal incidence ($\sim 280^\circ$, Figure 4e,f) increase the likelihood of drownings. Near-neap-low tide levels and, to a lesser extent, extremely-high-tide levels are also over-represented. Interestingly, these trends appear more pronounced for the severe drowning incidents (lower panels in Figure 4), although the reasonably small number of such incidents ($n = 32$) prevents from drawing conclusions. Additional analysis (not shown) also indicates that more drowning incidents occur on warm, sunny days with light winds (Castelle *et al.*, 2019), presumably as a result of increased beach attendance and increased beachgoer exposure to physical hazard.

Figure 5 shows time series of drowning incidents, wave and tide conditions of the four days when the largest number drowning incidents in the entire dataset were reported. These four days were associated with average or above average wave height and period (Figure 5i-l) and near shore-normal wave incidence (not shown), but not for any particular tide range or low tide timing within the day. However, drowning incidents clearly occurred preferably for lower tide levels, consistent with Figure 4g,h, with some milder drowning incidents occurring for extremely high water level occurring in the afternoon (Figure 5d), likely when beach attendance is maximized with beachgoers exposing themselves to high-tide swash rips (Castelle *et al.*, 2019). Severe drowning incidents can also occur in clusters before or after the supervised hours (11AM-7PM) around low tide levels even during the night (Figure 5d). These rare mass drowning events

Figure 5. Time series on the four days with the largest number of drowning incidents in the entire dataset showing: (a-d) distribution drowning incidents with drowning stage colored; (e-h) tide elevation η ; (i-l) significant wave height H_s and peak wave period T_p (e) with the horizontal dashed lines indicating the summer means.

Figure 6. Wave-driven currents with (a) breaking wave patterns (white areas) and (b) flow intensity coloured at Biscarrosse beach for the reference simulation with $H_s = 1.5$ m, $T_p = 10$ s, $\theta = 280^\circ$, $\eta = -1$ m. 20-min averaged rip flow velocity against (c) H_s , (d) T_p and (e) η .

are typically associated with risky behavior of an individual or group of individuals, for example when four drowning incidents (including one stage 3 and one stage 4) concomitantly occurred at Biscarrosse late at night on July 27, 2007.

To further explore the influence of wave and tide conditions on rip current activity in southwest France, we modelled nearshore hydrodynamics using morphology surveyed in mid-June 2007 at Biscarrosse, varying H_s from 0.25 m to 3 m (0.25-m interval), T_p from 4 s to 20 s (2-s interval), θ from 280° to 320° (5° interval) and η from -2.2 m to 2.2 m (0.2-m interval). The reference simulation was chosen for $H_s = 1.5$ m, $T_p = 10$ s, $\theta = 280^\circ$ and $\eta = -1$ m, which are the conditions corresponding to a disproportionate amount of drowning incidents observed in our dataset (Figure 4). The reference simulation (Figure 6a,b) shows intense rip currents through the four rip channels, although these rip currents contrast in terms of shape, width and intensity. Figure 6c-e shows that rip flow intensity averaged over the four rip current systems largely increases for increasing wave height (Figure 6c) and increasing wave period (Figure 6d) although rip flow speed tends to stabilize for $H_s > 1.8$ m and $T_p > 15$ s. More intense rips also occur at lower tide levels, peaking around mean low water neap (Figure 6e), in line with maximized incident data (Figure 4g,h). Rip flow speed is, however, largely variable alongshore (see the large error bars in Figure 6c-e) owing to the large variability in rip channel morphology. These results indicate that wave and tide conditions for which rip flow speed is maximized are approximately the same as those for which more drowning incidents are reported.

CONCLUSIONS

On the high-energy surf beaches of southwest France, summer drowning incidents occur disproportionately near neap low tide, under shore-normally incident waves and average to above-average wave height and period. This is consistent with numerical modelling results that show flow intensity maximized for shore-normal incidence, around neap low tide levels and for increased

wave height and period. This highlights the dominance of the physical hazard on the life risk along this stretch of coast and the potential for the development of drowning risk predictors along this coast.

ACKNOWLEDGMENTS

BC was funded by project SONO (ANR-17-CE01-0014) from the Agence Nationale de la Recherche (ANR). Truc Vert beach is labelled by the Service National d'Observation (SNO) Dynalit (<https://www.dynalit.fr>). Weather station and tide gauge data were provided by the Météo France Radome network and the SHOM, respectively.

LITERATURE CITED

- Austin, M.J.; Masselink, G.; Scott, T.M., and Russell, P.E., 2014. Water level controls on macro-tidal rip currents. *Continental Shelf Research*, 75, 28-40.
- Boudière, E.; Maisondieu, C.; Arduin, F.; Accensi, M.; Pineau-Guillou, L., and Lepesqueur, J., 2013. A suitable metocean hindcast database for the design of marine energy converters. *International Journal of Marine Energy*, 3-4, 40-52.
- Brander, R.W., 1999. Field observations on the morphodynamic evolution of a low-energy rip current system. *Marine Geology*, 157, 199-217.
- Bruneau, N.; Bonneton, P.; Castelle, B., and Pedreros, R., 2011. Modeling rip current circulations and vorticity in a high-energy meso-environment. *Journal of Geophysical Research – Oceans*, 116, C07026.
- Bruneau, N.; Castelle, B.; Bonneton, P.; Pedreros, R.; Almar, R.; Bonneton, N.; Bretel, P.; Parisot, J.-P., and Sénéchal, N., 2009. Field observations of an evolving rip current on a meso-macrotidal well-developed inner bar and rip morphology. *Continental Shelf Research*, 29, 1650-1662.

- Castelle, B.; Scott, T.; Brander, R.W., and McCarroll, R.J., 2016a. Rip current types, circulation and hazard. *Earth-Science Reviews*, 163, 1-21.
- Castelle, B.; McCarroll, R.J.; Brander, R.W.; Scott, T., and Dubarbier, B., 2016b. Modelling the alongshore variability of optimum rip current escape strategies on a multiple rip-channelled beach. *Natural Hazards*, 81, 664-686.
- Castelle, B.; Bujan, S.; Ferreira, S., and Dodet, G., 2017. Foredune morphological changes and beach recovery from the extreme 2013/2014 winter at a high-energy sandy coast. *Marine Geology*, 385, 41-55, 2017.
- Castelle, B.; Scott, T.; Brander, R.W.; McCarroll, R.J.; Robinet, R.; Tellier, E.; de Korte, E.; Simonnet, B., and Salmi, L.-R., 2019. Environmental controls on surf zone injuries on high-energy beaches. *Natural Hazards and Earth System Sciences*, 19, 2183-2205.
- Dalrymple, R.A.; MacMahan, J.H.; Reniers, A.J.H.M., and Nelko, V., 2011. Rip currents. *Annual Reviews in Fluid Mechanics*, 43, 551-581.
- Houser, C.; Arnott, R.; Ulzhofer, S., and Barrett, G., 2013. Nearshore circulation over transverse bar and rip morphology with oblique wave forcing. *Earth Surface Processes and Landforms*, 38, 1269-1279.
- MacMahan, J.H.; Thornton E.B., and Reniers, A.J.H.M., 2006. Rip current review. *Coastal Engineering*, 53(2), 191-208.
- McCarroll, R.J.; Brander, R.W.; Scott, T., and Castelle, B., 2018. Bathymetric controls on rotational surfzone currents. *Journal of Geophysical Research - Earth Surface*, 123, doi: 10.1029/2017JF004491.
- Moulton, M.; Elgar, S.; Raubenheimer, B.; Warner, J.C., and Kumar, N., 2017. Rip currents and alongshore flows in single channels dredged in the surf zone. *Journal of Geophysical Research - Oceans*, 122, 3799-3816, doi: 10.1002/2016JC012222.
- Roelvink, J.A.; Reniers, A.J.H.M.; van Dongeren, A.; de Vries, J. V.; McCall, R., and Lescinski, J., 2009. Modelling storm impacts on beaches, dunes and barrier islands. *Coastal Engineering*, 56, 1133-1152.
- Scott, T.M.; Masselink, G.; Austin, M.J., and Russell, P., 2014. Controls on macrotidal rip current circulation and hazard. *Geomorphology*, 214, 198-215.