Do Equids Live longer than Grazing Bovids?

Morgane Tidière, Patrick Duncan, Jean-François Lemaître Lemaître, Jean-Michel Gaillard, Laurie Bingaman Lackey, Dennis Müller, Marcus Clauss

► To cite this version:

Morgane Tidière, Patrick Duncan, Jean-François Lemaître Lemaître, Jean-Michel Gaillard, Laurie Bingaman Lackey, et al.. Do Equids Live longer than Grazing Bovids?. Journal of Mammalian Evolution, 2020, 27 (4), pp.809-816. 10.1007/s10914-019-09483-8 . hal-03060300

HAL Id: hal-03060300

<https://hal.science/hal-03060300>

Submitted on 15 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Do equids live longer than grazing bovids?

2 Morgane Tidière^{1*}, Patrick Duncan², Jean-François Lemaître¹, Jean-Michel Gaillard¹, Laurie
3 Bingaman Lackey³, Dennis W.H. Müller⁴ and Marcus Clauss⁵

4 ¹Université de Lyon, F-69000, Lyon; Université Lyon 1; CNRS, UMR5558, Laboratoire de
5 Biométrie et Biologie Evolutive, F-69622, Villeurbanne, France.

6 ²Centre d'Etudes Biologiques de Chizé, Station d'Ecologie de Chizé-La Rochelle, UMR 7173
7 CNRS, 79360 Villiers-en-Bois, France.

8 ³World Association of Zoos and Aquariums (WAZA), Carrer de Roger de Llúria, 2, 2-2, Bar-
9 celona, Spain.

10 ⁴Zoological Garden Halle (Saale), Fasanenstr. 5a, 06114 Halle (Saale), Germany.

11 ⁵Clinic for Zoo Animals, Exotic Pets and Wildlife, Vetsuisse Faculty, University of Zurich,
12 Winterthurerstr. 260, 8057 Zurich, Switzerland.

13 *Corresponding authors: mtidiere@gmail.com

14 **ABSTRACT**

15 A large part of the diversity of longevity and actuarial senescence (i.e. the progressive decline
16 of survival probabilities with age) across vertebrates can be related to body size, phylogeny
17 and the species' position on the slow-fast continuum of life histories. However, differences in
18 mortality patterns between ecologically similar species, such as bovids and equids, remain
19 poorly understood. Equids are commonly understood to outlive bovid species relative to their
20 body mass, despite very similar feeding niches. Comparing survival patterns of 13 bovid and
21 10 equid sub-species, our findings confirm that equids outlive bovid species, with a higher
22 adult survival rate and a delayed onset of senescence for equids, but no difference of rate of
23 actuarial senescence. These differences are associated with a slower generation time and
24 longer inter-birth interval, due to a longer gestation period, for equids compared to bovids.
25 Finally, our results suggest that all biological times (i.e. all life history traits expressed in time
26 units) have evolved synchronously in bovids, whereas in equids gestation time and inter-birth
27 interval either were never in synchrony with, or have slowed down relative to other biological
28 times. Our findings suggest the existence of different selection pressures, or different
29 constraints, on specific time-related traits between these two mammalian orders.

30 **Keywords:** actuarial senescence; biological times; demography; pace of life; slow-fast con-
31 tinuum

32 **Running head:** Equids vs bovids' actuarial senescence

33 **Word count:** 5734 words

34 INTRODUCTION

35 Actuarial senescence, defined as the progressive decrease of survival with increasing age, is a
36 ubiquitous process across the tree of life (Nussey et al. 2013; Jones et al. 2014). Striking dif-
37 ferences in actuarial senescence and lifespan have been found in related species, such as the
38 naked mole rat (*Heterocephalus glaber*) which lives nine times longer than similar-sized ro-
39 dents (Buffenstein 2008; Ruby et al. 2018). Such differences across phylogenetically related
40 species or across species sharing the same lifestyle remain poorly understood and are current-
41 ly attracting much interest (Gorbunova et al. 2014; Healy et al. 2014). However, little work
42 has been done, so far, to understand variation in senescence between distantly related species
43 that have converged into closely similar niches.

44 Here we compare patterns of actuarial senescence (hereafter ‘senescence’) between
45 taxa from two orders of mammals, equids (Perissodactyl) and grazing bovids (Artiodactyl),
46 which are large herbivores whose evolution separated 55 million years ago (Janis 1976).
47 Many of the grazing species, among the equids and bovids, have evolved in parallel in similar
48 environments since the Miocene (Janis 1976). These two groups of herbivores display evolu-
49 tionary convergence in key morphological characters, such as elongated limbs and hyp-
50 sodonty (i.e. the height of the molar teeth), and occupy similar feeding niches despite having
51 different digestive systems (Janis 1976). Moreover, it is commonly assumed that horses live
52 longer than cattle. For instance, in the AnAge database (Tacutu et al. 2012), the maximum
53 lifespan recorded for the seven species of extant equids ranges from 30.1 years for kiang
54 (*Equus kiang*) to 57 years for domestic horse (*E. caballus*), while for grazing bovids the max-
55 imum longevity recorded is 36.1 years for the lowland anoa (*Bubalus depressicornis*).

56 Among bovids, grazers live longer and senesce later and more slowly than do brows-
57 ers, both in the wild (Gaillard et al. 2008) and in captive populations (Lemaître et al. 2013;
58 Tidière et al. 2015). This difference is even more marked in captivity, probably because the

food provided in zoos is more suitable for grazers than for browsers (Müller et al. 2010, 2011; Lemaître et al. 2013; Tidière et al. 2015). The extant equids are typically grazers (Schulz and Kaiser 2013), while bovids comprise species ranging over a continuum of dietary niches from grazers to browsers (Hofmann 1989; Gordon and Prins 2008); therefore, we only used data from grazers among the bovids to perform a valid comparison of longevity and senescence patterns across these groups. The use of captive populations allows patterns of longevity and senescence of the two orders to be studied independently of environmental variation, which is likely to be a key driver of mortality patterns (Gaillard et al. 1993). In captivity, animals are protected against starvation and predation, and competition for resources is minimal or absent. Using a dataset of 15 species (comprising 23 species or subspecies; 13 bovids and 10 equids) living in captivity, we found strong support for longer longevity in equids than in bovids after taking into account allometric relationships. This difference is associated with higher adult survival and delayed onsets of senescence for equids, but there is no difference between the two orders in the rate of actuarial senescence.

A major axis of variation in longevity and senescence across species is included in the slow-fast continuum of the life histories (Stearns 1983). Indeed, variation in life history traits among species is structured along a continuum of biological times that contrasts species with short longevity, early reproduction, high annual fecundity, and early onset and/or fast rates of senescence (at the continuum's fast end), with species with long longevity, delayed reproduction, low annual fecundity, and late onset and/or slow rates of senescence (at its slow end, e.g. Gaillard et al. 1989; Ricklefs 1998, 2010; Jones et al. 2008; Bronikowski and Vleck 2010). We, therefore, assessed whether observed differences of longevity between bovids and equids simply correspond to a different ranking over the slow-fast continuum as measured by the generation time (Gaillard et al. 2005). Two plausible, but different, demographic scenarios were tested: (1) all biological times or (2) only certain biological times are slower in equids

than in bovids, leading to a difference in the pace of life (as represented by longer generation times in equids). Causes of the differences could be selection pressures for a slower pace of life in equids, or for a faster pace of life in bovids, with potentially different selection pressures and constraints acting on the different components of generation time. Evolutionary processes behind the differences in extant equids and bovids are discussed.

METHODS

Datasets

The life tables of the zoo populations were compiled from the Species360 database, which contains data since 1817 from more than 1,000 member institutions in over 90 countries worldwide. Information on sex, dates of birth and death is available for each individual. Assuming that the total number of individuals living in zoos for a given species corresponds to a worldwide meta-population, we computed population parameters for species living in captivity. We constructed life tables available for females of equid and grazing bovid species for which the exact dates of birth and death are known. We considered all dead and living females but, to avoid any overestimation of the species' longevity due to some individuals never being recorded as dead, we removed individuals still alive in January 2014 with an age more than 125% older than the maximum age at death reported in AnAge, as longevity can be longer in zoos today than 20 years ago due to the improvement of captive conditions (e.g. Wich et al. 2009 for orang-utan, *Pongo* sp.). To ensure that the inclusion of all cohorts did not bias the results, the same analyses were done with cohorts for which all individuals are dead, defined as cohorts for which all females were already dead in January 2014. For some species, wild-type and domestic populations are kept in zoos (e.g. *Bos javanicus*). For these species, only the wild-type taxa were retained in the analyses. We obtained a final dataset of 23 taxa including 13 grazing bovids and 10 equids (Fig. S1).

Survival and actuarial senescence metrics

For each species, five different, and complementary, metrics were used to assess senescence patterns: maximum age at death, longevity, average adult survival rate, age at the onset of senescence and rate of senescence. The maximum age at death was determined as the maximum age at death observed in the database (Table S1). For the four other metrics, we built the capture history of each female and obtained age-specific survival estimates using the CMR (capture-mark-recapture) software E-SURGE (Choquet et al. 2009) (see appendix S1 for more details). From these age-specific survival estimates, we calculated the proportion of individuals alive at each age (l_x , with $l_0=1$) and longevity was measured as the age at which 90% of individuals from the initial cohort were dead (age x at which $l_x=0.1$) (Table S1). This avoids spurious estimates due to different sample sizes or exceptionally long lives of a few individuals (Moorad et al. 2012).

To assess actuarial senescence, we fitted different age-dependent models from birth onwards: constant survival, two age-classes, full age-dependence, and several Gompertz models with different onsets of actuarial senescence (see appendix S1 for more details). To select the best survival model, we used a model selection procedure based on the Akaike Information Criterion (AIC, Burnham and Anderson 2002). For each species, we retained the model with the lowest AIC (Table S2). For 10 of the 13 bovids and 9 of the 10 equids, a Gompertz model provided a better fit than either the constant or the two-age class model, and was more parsimonious than the full age-dependent model (Table S2). The age at which survival begins to decline, corresponding to the age at the onset of senescence (Péron et al. 2010), varied among taxa (Table S1). In addition, from the selected Gompertz model, we measured both adult survival rate as the mean annual survival estimated between 1 year of age and the onset of senescence (Table S1) and rate of senescence as the exponential rate of mortality increase with age (Table S1). On a logit scale, age variation in survival was,

therefore, constrained to be linear from the age at the onset of senescence onwards (Gaillard et al. 2004). For four species (*Connochaetes taurinus albojubatus*, *Kobus kob leche*, *Syncerus caffer* and *Equus quagga boehmi*), the full age-dependent model had a lower AIC than any Gompertz model. For these four taxa, only the maximal age at death and longevity were analyzed.

Mean values for the traits of adult females in each taxon

Mean body mass (BMA, in kg) for adult females was extracted from the literature (Table S3) to correct for allometric constraints. When available, body mass measures from wild populations were preferred.

The pace of life of a given taxon was measured as the generation time using the method described in Gaillard et al. (2005) (see appendix S1 for more details). As life history traits with a currency of time, we considered gestation length (GL, in days), age at first parturition (AFP, in days), and the inter-birth interval (IBI, in days) to test the potential difference of life history strategy between equids and bovids (Table S3). Values for GL, AFP, and IBI were extracted from the literature (Table S3).

Analyses

We first tested the differences in survival and actuarial senescence metrics between bovids and equids, accounting for body mass, because large species live longer and senesce later and more slowly than small species (e.g. Jones et al. 2008; Ricklefs 2010; Lemaître et al. 2014; Tidière et al. 2015). Dependent variables included maximum age at death, longevity, adult survival rate, onset of senescence and rate of senescence. The full models included body mass and order (as factor) as independent variables.

In a second step, we tested whether the differences in survival and actuarial senescence metrics between bovids and equids parallel a difference in life history strategy. For

each trait corresponding to a biological time included in the study (i.e. generation time, gestation length, inter-birth interval and age at first parturition), the full models included body mass and order (as factor) as independent variables.

In a third step, we ran a normed Principal Component Analysis (PCA, package ‘*ade4*’, (Dray and Dufour 2007) for bovids (N=5 taxa) and equids (N=7 taxa) separately, with generation time, longevity, age at first parturition, gestation length and inter-birth interval as biological times. Under scenario 1, all biological times in both equids and bovids should contribute equally to the overall time scale (i.e. equi-correlation sensu Morrison 1967), which corresponds to the slow-fast continuum (Gaillard et al. 2016). In contrast, under scenario 2, equi-correlation should not be observed, meaning that the overall time scale is driven more strongly by some specific biological times and less by others.

Analyses were not corrected for phylogenetic relatedness beyond the inclusion of order as a factor because the number of species was rather small and did not allow estimating reliably the phylogenetic inertia. All traits were log-transformed to improve the normality of the distributions, except order (entered as a factor) and adult survival rate that was logit-transformed. Analyses were performed using the linear model (lm) in R version 2.14.0 (R Development Core Team 2016) and parameter estimates are given $\pm$ SE and/or 95% confidence intervals.

The datasets generated during and/or analysed during the current study are available in the supplementary data. However, the raw longevity dataset used to compute the survival and senescence metrics are available upon request and with permission of Species360.

RESULTS

Differences in survival and actuarial senescence parameters

As expected, all survival metrics (i.e. maximum age at death, longevity and adult survival

rate) increased with adult body mass (Tables 1 and S4, Fig. 1). Moreover, for a given body mass, equids had a maximum age at death 6.9 years (95%CI [3.9; 10.0], Table 1, Fig. 1A) longer and a longevity 5.5 years (95%CI [2.4; 8.6], Table 1, Fig. 1B) longer than bovids. Equids also had a higher adult survival (+ 0.018, 95%CI [0.002; 0.035], Table 1, Fig. 1C) than bovids. Order was the only factor included in the selected model to explain variation observed in the onset of actuarial senescence (Table S4): equids began to senesce 5.89 years (95%CI [2.33; 9.44], Table 1) later than bovids, irrespective of body mass (Fig. S2). Finally, observed variation in the rate of senescence was explained only by the age at the onset of senescence (Table S4), with the rate of senescence increasing when the onset was late (Table 1, Fig. S3). However, the model also including body mass was competitive (Δ AIC = -1.31, Table S4) that for a given onset of senescence, the rate of senescence tended to decrease with increasing body mass (β = -0.342 ± 0.196). There was no detectable difference between bovids and equids in the rates of senescence in any of the models.

Using only extinct cohorts in the dataset results were similar to those obtained with all individuals (Tables S5-S7): the order factor was included in the selected models for all survival metrics, which indicates that equids outlived bovids and had higher adult survival. For senescence metrics, the selected models did not include body mass or the order factor. However, for the onset of senescence, the AIC of the model with order as a factor was only 0.51 AIC units lower than the selected model with equids senescing 4.21 ± 1.57 years later than bovids. Finally, the rate of senescence was only influenced by the onset of senescence.

Differences in life history traits

As expected, irrespective of the order, the four biological times were positively correlated to adult body mass (Table 2 and S8, Fig. S4). Bovid had shorter generation times (by 2.31 years, Fig. S4A), shorter gestation lengths (by 89.2 days, Fig. S4B) and shorter inter-birth

intervals (by 157.3 days, Fig. S4C) than equids. On the other hand, no detectable difference of age at first parturition was found between equids and bovids (Fig. S4D, Tables 2 and S8).

Evolutionary pathways

In absence of species-specific selection pressure on a particular trait, generation time is expected to be isometrically related to other biological times such as longevity, age at first reproduction, inter-birth interval and gestation length. The generation time thus constitutes an integrative measure of the time scale at which the life cycle of a given species is completed (Gaillard et al. 2005; Staerk et al. 2019). Here, we assessed whether equids displayed longer generation times either, as a part of a generally slower pace of life compared to bovids, in which case all biological times should be isometrically associated, or whether the two groups differed in only one particular biological time such as gestation length, which would lead to longer generation times in equids than bovids. We found that all five of the biological times contributed almost equally to the bovids' pace of life, although the gestation length was slightly less tightly associated with other time-related traits (Table S9, Fig. 2A). Moreover, the generation time was equivalent to the PC1 (Fig. 2A), which means that generation time provides an overall measure of the pace of life in bovids. On the other hand, the picture was markedly different for equids. Age at first parturition, instead of generation time (Fig. 2B), was the closest to PC1 and was independent of inter-birth interval and gestation length (Table S9). Longevity was negatively associated with the overall time scale (Table S9, Fig. 2B). These findings demonstrate that, while bovids closely match the biological time syndrome hypothesis, the variation among biological times was uncoupled in equids. This suggests the existence of different selection pressures, or different constraints, on specific time-related traits between the families.

DISCUSSION

Our findings confirm that, for a given body mass, equids live longer than bovids, whatever the metric used (maximum age at death or longevity). This difference in longevity between bovids and equids is associated with both higher survival and a delay in the onset of actuarial senescence in equids, but there is no difference in the rates of actuarial senescence. This is consistent with several recent studies highlighting that the onset of senescence has to be considered when analyzing senescence, and is more responsive than the rate of senescence (Péron et al. 2010; Hammers et al. 2013; Gamelon et al. 2014; Tidière et al. 2015), although the latter is still preferentially used in comparative analyses (e.g. Jones et al. 2008; Ricklefs 2010). Moreover, the analyses of covariation among biological times in bovid and equid taxa indicate that a synchronization of all biological times evolved in bovids, whereas biological times were uncoupled in equids. This could be the result of a specific selection for longer gestation lengths in equids, a physiological constraint that prevents the evolution of shorter gestation lengths in equids, a stronger selection for a faster pace of life in bovids, or a combination of any of these factors.

Tooth wear is an essential process in mammalian ageing (Skogland 1988; King et al. 2005), and there is evidence that tooth height is related to longevity within species (Veiberg et al. 2007). The difference in longevity we reported may be explained by a difference in a key morphological adaptation of grazers, hypsodonty. Comparison of estimates of hypsodonty index for six bovid and seven equid taxa (see Table S3) showed that equids have hypsodonty about 50% higher than bovids (difference = 2.49; $t = 4.69$; $df = 10.678$; $p\text{-value} < 0.001$, Fig. S5). Dental anatomy could, therefore, provide a mechanical explanation for the greater longevity of equids relative to similar-sized bovids, through a longer functional duration of the chewing tooth-row. It has been suggested that wear rates should be less in ruminants, because much of the chewing occurs during rumination, on material that has already been washed of grit in the forestomach (Hatt et al. 2019). However, the limited information

available has not yet demonstrated different rates of tooth wear between equids and bovids (Damuth and Janis 2014). More data like that measured in red deer (*Cervus elaphus*) by Péres-Barberia and Gordon (1998) are needed on tooth wear in free-ranging equids and grazing bovids to conclude firmly whether the greater hypsodonty in equids plays a key role in explaining their greater longevity.

The longer generation time of equids demonstrates that they have a slower pace of life than similar-sized grazing bovids. This difference of pace of life is partly due to a difference of inter-birth interval, itself caused in part by the longer gestation length of equids, and partly to their higher adult survival, which also contributes to their slow pace of life. Two different concepts exist for the occurrence of a slow pace of life. It could be the result of specific selection pressures that facilitate its evolution as a specific adaptation to resource-limited niches. Alternatively, species characterized by a slower pace of life might be displaced by species with a faster pace of life except in resource-poor niches (Vermeij 1994). There are differences in susceptibility to resource constraints between bovids and equids: the abundance of wildebeest (i.e. a grazing bovid) is much more affected by food resources than zebras (a similar-sized equid) (Grange and Duncan 2006). **Whether longer gestation periods evolved in equids from a common ancestor of the two groups, perhaps a Condylarth (Rose 1996), which had shorter gestation periods, or whether a generally slower pace of life is a characteristic of the equid lineage, cannot be determined from our study, since the common ancestor's life history is of course unknown.**

Most bovid species have an inter-birth interval of less than one year (mean of 334.00 days, 95%CI [286.54;381.46]), which allows them to produce one litter per year. On the other hand, equids display mean inter-birth intervals of about 1 year and 4 months (491.29 days, 95%CI [440.34;542.23]). The bovid life history strategy thus allows breeding seasonally to match plant phenology, which is clearly an advantage in many grassland systems. Decreasing

the gestation length may have been part of the bovid evolution towards seasonal breeding (Zerbe et al. 2012), and the shorter generation times may have allowed bovids to increase their demographic rates and abundances. This may give them an advantage in resource rich environments where predation pressure is higher than in poor environments (Melis et al. 2013). Indeed, the faster pace of life allows higher population growth rates, leading to a better demographic response to predation pressure (Grange et al. 2004). A shorter generation time allows faster evolution and this may also have facilitated their species diversity. For instance, the shorter gestation length in bovids compared to giraffids has been invoked to explain the replacement of a diverse giraffid community with a diverse bovid one over evolutionary time (Clauss and Rössner 2014).

To conclude, equids live longer than grazing bovids of similar size. This is because equids have higher survival, and their actuarial senescence begins at a later age. Equids also have greater hypsodonty, which could be a key morphological adaptation allowing greater longevity in equids, though more data are needed on wear rates in the wild to reach a firm conclusion. Grazing bovids have a faster pace of life, with shorter gestation periods. The shorter gestation periods of bovids may be associated with accelerated rates of intrauterine growth (Baur 1977), although the morphological and physiological mechanisms involved remain to be discovered. This scenario considers equids to be constrained by slower intrauterine growth and a slower pace of life, which makes them mainly competitive in environments with limited resources. Alternatively, a longer gestation length for equids may be a specifically evolved adaptation to increased survival in these environments. The fast pace of life in bovids may be a key to the rapid evolution of the bovids since the Miocene; today they dominate communities of grazing animals, in species diversity and in abundance.

ACKNOWLEDGMENTS

We are very grateful to the many people who generously provided data on live weights

and on the animals' life histories so that we could carry out the analyses with confidence, in particular Kristin Brabender (Hortobagy National Park, Hungary), Frédéric Joly (Takh, France), Jim Kao (Taipei Zoo), Klaus Mueller-Schilling (Hannover Zoo), Peter Novellie (Nelson Mandela Metropolitan University, South Africa), Kate Schoenecker (US Geological Survey, USA), Juan Soy (Al Wabra Wildlife Preservation, Qatar), Beatrice Steck (Basel Zoo, Switzerland), Tim Thier and Martha Fischer (St. Louis Zoo, USA), and Tim Woodfine, Tanya Langenhorst and Danielle Free (Marwell Wildlife, UK) who provided data on liveweights, and/or on the animals' life histories; and Patricia Moehlman and Sarah King (IUCN Equid Specialist Group), Ulrike Rademacher (Stuttgart Zoo, Germany), Belinda Low Mackey (Grevy's Zebra Trust, USA) and Laurent Tatin (*Conservatoire des Espaces Naturels*, Provence-Alpes-Côte d'Azur, France) who kindly linked us to sources of knowledge. We are grateful to one anonymous referee for helpful comments on a previous draft.

REFERENCES

- Baur R (1977) Morphometry of the Placental Exchange Area. Springer
- Bronikowski AM, Vleck D (2010) Metabolism, body size and life span: a case study in evolutionarily divergent populations of the garter snake (*Thamnophis elegans*). Integr Comp Biol 50:880–887
- Buffenstein R (2008) Negligible senescence in the longest living rodent, the naked mole-rat: insights from a successfully aging species. J Comp Physiol B 178:439–445
- Burnham KP, Anderson DR (2002) Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach. Springer, New York
- Choquet R, Rouan L, Pradel R (2009) Program E-SURGE: a software application for fitting multievent models. Environ Ecol Stat 3:8–39
- Clauss M, Rössner GE (2014) Old world ruminant morphophysiology, life history, and fossil record: exploring key innovations of a diversification sequence. Ann Zool Fenn

329 51:80–94

330 Damuth J, Janis CM (2014) A comparison of observed molar wear rates in extant herbivorous
 331 mammals. *Ann Zool Fenn* 51:188–200

332 Dray S, Dufour AB (2007) The ade4 package: implementing the duality diagram for
 333 ecologists. *J Stat Softw* 22:1–20

334 Gaillard J-M, Delorme D, Boutin J-M, Van Laere G, Boisaubert B, Pradel R (1993) Roe deer
 335 survival patterns: a comparative analysis of contrasting populations. *J Anim Ecol*
 336 62:778–791

337 Gaillard J-M, Duncan P, van Wieren SE, Loison A, Klein F, Maillard D (2008) Managing
 338 large herbivores in theory and practice: is the game the same for browsing and grazing
 339 species. In: Gordon IJ, Prins HHT (eds) *The Ecology of Browsing and Grazing*.
 340 Springer, Berlin, Heidelberg, pp 293–307

341 Gaillard J-M, Lemaître J-F, Berger V, Bonenfant C, Devillard S, Douhard M, Gamelon M,
 342 Plard F, Lebreton J-D (2016) Life history axes of variation. In: Kliman RM (ed) *The*
 343 *Encyclopedia of Evolutionary Biology*. Academic Press, pp 312–323

344 Gaillard J-M, Pontier D, Allaine D, Lebreton J-D, Trouvilliez J, Clobert J (1989) An analysis
 345 of demographic tactics in birds and mammals. *Oikos* 56:59–76

346 Gaillard J-M, Yoccoz NG, Lebreton J-D, Bonenfant C, Devillard S, Loison A, Pontier D,
 347 Allaine D (2005) Generation time: a reliable metric to measure life-history variation
 348 among mammalian populations. *Am Nat* 166:119–123

349 Gaillard J-M, Viallefont A, Loison A, Festa-Bianchet M (2004) Assessing senescence patterns
 350 in populations of large mammals. *Anim Biodivers Conserv* 27:47–58

351 Gamelon M, Focardi S, Gaillard J-M, Gimenez O, Bonenfant C, Franzetti B, Choquet R,
 352 Ronchi F, Baubet E, Lemaître J-F (2014) Do age-specific survival patterns of wild
 353 boar fit current evolutionary theories of senescence? *Evolution* 68:3636–3643

354 Gorbunova V, Seluanov A, Zhang Z, Gladyshev VN, Vijg J (2014) Comparative genetics of
355 longevity and cancer: insights from long-lived rodents. *Nat Rev Genet* 15:531–540

356 Gordon IJ, Prins HHT (2008) *The Ecology of Browsing and Grazing*. Ecological Studies.
357 Springer-Verlag, Berlin Heidelberg

358 Grange S, Duncan P (2006) Bottom-up and top-down processes in African ungulate
359 communities: resources and predation acting on the relative abundance of zebra and
360 grazing bovids. *Ecography* 29:899–907

361 Grange S, Duncan P, Gaillard J-M, Sinclair ARE, Gogan PJP, Packer C, Hofer H, East M
362 (2004) What limits the Serengeti zebra population? *Oecologia* 140:523–532

363 Hammers M, Richardson DS, Burke T, Komdeur J (2013) The impact of reproductive
364 investment and early-life environmental conditions on senescence: support for the
365 disposable soma hypothesis. *J Evol Biol* 26:1999–2007

366 Hatt J-M, Codron D, Müller DWH, Ackermans NL, Martin LF, Kircher PR, Hummal J,
367 Clauss M (2019) The rumen washes off abrasives before heavy-duty chewing in
368 ruminants. *Mammal Biol* 97:104-111

369 Healy K, Guillerme T, Finlay S, Kane A, Kelly SBA, McClean D, Kelly DJ, Donohue I,
370 Jackson AL, Cooper N (2014) Ecology and mode-of-life explain lifespan variation in
371 birds and mammals. *Proc R Soc Lond B Biol Sci* 281:20140298

372 Hofmann RR (1989) Evolutionary steps of ecophysiological adaptation and diversification of
373 ruminants: a comparative view of their digestive system. *Oecologia* 78:443–457

374 Janis C (1976) The evolutionary strategy of the Equidae and the origins of rumen and cecal
375 digestion. *Evolution* 30:757–774

376 Jones OR, Gaillard J-M, Tuljapurkar S, Alho JS, Armitage KB, Becker PH, Bize P, Brommer
377 J, Charmantier A, Charpentier M, Clutton-Brock T, Dobson FS, Festa-Bianchet M,
378 Gustafsson L, Jensen H, Jones CG, Lillandt B-G, McCleery R, Merilä J, Neuhaus P,

379 Nicoll MAC, Norris K, Oli MK, Pemberton J, Pietiäinen H, Ringsby TH, Roulin A,
 380 Saether B-E, Setchell JM, Sheldon BC, Thompson PM, Weimerskirch H, Jean
 381 Wickings E, Coulson T (2008) Senescence rates are determined by ranking on the fast-
 382 slow life-history continuum. *Ecol Lett* 11:664–673

383 Jones OR, Scheuerlein A, Salguero-Gómez R, Camarda CG, Schaible R, Casper BB,
 384 Dahlgren JP, Ehrlén J, García MB, Menges ES, Quintana-Ascencio PF, Caswell H,
 385 Baudisch A, Vaupel JW (2014) Diversity of ageing across the tree of life. *Nature*
 386 505:169–173

387 King SJ, Arrigo-Nelson SJ, Pochron ST, Semprebon GM, Godfrey LR, Wright PC, Jernvall J
 388 (2005) Dental senescence in a long-lived primate links infant survival to rainfall. *Proc*
 389 *Natl Acad Sci* 102:16579–16583

390 Lemaître J-F, Gaillard J-M, Bingaman Lackey L, Clauss M, Müller DWH (2013) Comparing
 391 free-ranging and captive populations reveals intra-specific variation in aging rates in
 392 large herbivores. *Exp Gerontol* 48:162–167

393 Lemaître J-F, Müller DWH, Clauss M (2014) A test of the metabolic theory of ecology with
 394 two longevity data sets reveals no common cause of scaling in biological times:
 395 longevity scaling in mammals. *Mammal Rev* 44:204–214

396 Melis C, Nilsen EB, Panzacchi M, Linnell JDC, Odden J (2013) Roe deer face competing
 397 risks between predators along a gradient in abundance. *Ecosphere* 4:art111

398 Moorad JA, Promislow DEL, Flesness N, Miller RA (2012) A comparative assessment of
 399 univariate longevity measures using zoological animal records. *Aging Cell* 11:940–
 400 948

401 Morrison DF (1967) *Multivariate Statistical Methods*. McGraw-Hill

402 Müller DWH, Bingaman Lackey L, Streich WJ, Fickel J, Hatt J-M, Clauss M (2011) Mating
 403 system, feeding type and ex situ conservation effort determine life expectancy in

404 captive ruminants. *Proc R Soc B Biol Sci* 278:2076–2080

405 Müller DWH, Bingaman Lackey L, Streich WJ, Hatt J-M, Clauss M (2010) Relevance of
 406 management and feeding regimens on life expectancy in captive deer. *Am J Vet Res*
 407 71:275–280

408 Nussey DH, Froy H, Lemaître J-F, Gaillard J-M, Austad SN (2013) Senescence in natural
 409 populations of animals: widespread evidence and its implications for bio-gerontology.
 410 *Ageing Res Rev* 12:214–225

411 Pérez-Barberia FJ, Gordon IJ (1998) The influence of molar occlusal surface area on the
 412 assimilation efficiency, chewing behaviour and diet selection of red deer. *J Zool*
 413 245:307-316

414 Péron G, Gimenez O, Charmantier A, Gaillard J-M, Crochet P-A (2010) Age at the onset of
 415 senescence in birds and mammals is predicted by early-life performance. *Proc R Soc*
 416 *B Biol Sci* 277:2849–2856

417 R Development Core Team (2016) A language and environment for statistical computing. R
 418 Foundation for Statistical Computing, Vienna

419 Ricklefs RE (1998) Evolutionary theories of aging: confirmation of a fundamental prediction,
 420 with implications for the genetic basis and evolution of life span. *Am Nat* 152:24–44

421 Ricklefs RE (2010) Life-history connections to rates of aging in terrestrial vertebrates. *Proc*
 422 *Natl Acad Sci* 107:10314–10319

423 Rose KD (1996) On the origin of the order Artiodactyla. *Proc Natl Acad Sci* 93:1705-1709

424 Ruby JG, Smith M, Buffenstein R (2018) Naked mole-rat mortality rates defy Gompertzian
 425 laws by not increasing with age. *eLife* 7

426 Schulz E, Kaiser TM (2013) Historical distribution, habitat requirements and feeding ecology
 427 of the genus *Equus* (Perissodactyla). *Mammal Rev* 43:111–123

428 Skogland T (1988) Tooth wear by food limitation and its life history consequences in wild

429 reindeer. *Oikos* 51:238–242

430 Staerk J, Conde DA, Ronget V, Lemaître J-F, Gaillard J-M, Colchero F (2019) Performance of
 431 generation time approxiamtions for extinction risk assessments. *J App Ecol* 56:1436–
 432 1446

433 Stearns SC (1983) The influence of size and phylogeny on patterns of covariation among life-
 434 history traits in the mammals. *Oikos* 41:173–187

435 Tacutu R, Craig T, Budovsky A, Wuttke D, Lehmann G, Taranukha D, Costa J, Fraifeld VE,
 436 de Magalhães JP (2012) Human Ageing Genomic Resources: integrated databases and
 437 tools for the biology and genetics of ageing. *Nucleic Acids Res* 41:D1027–D1033

438 Tidière M, Gaillard J-M, Müller DWH, Bingaman Lackey L, Gimenez O, Clauss M, Lemaître
 439 J-F (2015) Does sexual selection shape sex differences in longevity and senescence
 440 patterns across vertebrates? A review and new insights from captive ruminants.
 441 *Evolution* 69:3123–3140

442 Veiberg V, Mysterud A, Gaillard J-M, Delorme D, Laere GV, Klein F (2007) Bigger teeth for
 443 longer life? Longevity and molar height in two roe deer populations. *Biol Lett* 3:268–
 444 270

445 Vermeij GJ (1994) The evolutionary interaction among species: selection, escalation, and
 446 coevolution. *Annu Rev Ecol Syst* 25:219–236

447 Wich SA, Shumaker RW, Perkins L, de Vries H (2009) Captive and wild orangutan (*Pongo*
 448 sp.) survivorship: a comparison and the influence of management. *Am J Primatol*
 449 71:680–686

450 Zerbe P, Clauss M, Codron D, Bingaman Lackey L, Rensch E, Streich JW, Hatt J-M, Müller
 451 DWH (2012) Reproductive seasonality in captive wild ruminants: implications for
 452 biogeographical adaptation, photoperiodic control, and life history. *Biol Rev* 87:965–
 453 990

454 **Table 1.** Parameter estimates from the linear models of the relationship between survival (i.e.
455 maximum age at death, longevity, and adult survival rate) or actuarial senescence (i.e. onset
456 and rate of senescence) metrics and the mean female adult body mass among 23 taxa of large
457 herbivores belonging to two orders of mammals.

		β	95% CI	t value	Adjusted-R ²
Maximum age at death N= 23 taxa	Intercept	2.496	2.033; 2.960	10.56	0.61
	Order (<i>Equidae</i>)	0.225	0.133; 0.318	4.79	
	Adult body mass	0.142	0.059; 0.225	3.34	
Longevity N= 23 taxa	Intercept	1.787	1.234; 2.341	6.33	0.63
	Order (<i>Equidae</i>)	0.225	0.115; 0.335	4.01	
	Adult body mass	0.228	0.129; 0.328	4.50	
Adult survival rate N= 19 taxa	Intercept	1.050	-0.224; 2.324	1.62	0.47
	Order (<i>Equidae</i>)	0.385	0.153; 0.617	3.25	
	Adult body mass	0.321	0.097; 0.544	2.81	
Onset of senescence N= 19 taxa	Intercept	2.056	1.845; 2.267	19.08	0.34
	Order (<i>Equidae</i>)	0.498	0.191; 0.804	3.18	
Rate of senescence N= 19 taxa	Intercept	-0.926	-2.123; 0.271	-1.52	0.31
	Onset	0.791	0.276; 1.305	3.01	

458

459 **Table 2.** Parameter estimates from the linear models of the relationship between biological
460 times and the mean female adult body mass among 23 taxa of large herbivores belonging to
461 two orders of mammals.

		β	95% CI	t value	Adjusted-R ²
Generation time N= 12 taxa	Intercept	0.107	-0.668; 0.881	0.27	0.78
	Order (<i>Equidae</i>)	0.179	0.002; 0.356	1.98	
	Adult body mass	0.310	0.156; 0.464	3.95	
Gestation length N= 15 taxa	Intercept	4.844	4.542; 5.146	31.42	0.89
	Order (<i>Equidae</i>)	0.243	0.179; 0.307	7.44	
	Adult body mass	0.143	0.087; 0.200	4.99	
Inter-birth interval N= 12 taxa	Intercept	5.050	4.260; 5.840	12.53	0.72
	Order (<i>Equidae</i>)	0.295	0.115; 0.476	3.21	
	Adult body mass	0.150	-0.006; 0.307	1.88	
Age at first parturition N= 14 taxa	Intercept	5.582	4.670; 6.494	12.00	0.39
	Adult body mass	0.260	0.094; 0.426	3.07	

462

463 **Figure legends**

464 **Figure 1.** Relationship between maximum age at death (A), longevity (B), or adult survival
465 (C) and taxon-specific body mass of adult females for 13 bovids (open dots) and 10 equids
466 (full dots). For a similar body mass, equids live longer and have a higher adult survival than
467 bovids.

468 **Figure 2.** Normed Principal Component Analyses performed on 5 biological times for 5 bovid
469 and 7 equid taxa separately. All biological times are positively correlated among bovids,
470 whereas for equids there is an uncoupling of biological times.