Trophic conditions influence widespread distribution of Aster-Like nanoparticles within aquatic environments

Maxime Fuster, Hermine H. Billard, Marie Mandart, Johannes Steiger,
Telesphore Sime-Ngando, Jonathan Colombet

► To cite this version:

Maxime Fuster, Hermine H. Billard, Marie Mandart, Johannes Steiger, Telesphore Sime-Ngando, et al.. Trophic conditions influence widespread distribution of Aster-Like nanoparticles within aquatic environments. Microbial ecology, 2020, 80 (4), pp.741-745. 10.1007/s00248-020-01541-6 . hal-03084202

HAL Id: hal-03084202

<https://hal.science/hal-03084202>

Submitted on 20 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution| 4.0 International License

Title

Trophic conditions influence widespread distribution of Aster Like Nanoparticles within aquatic environments

Authors

Maxime Fuster¹, Hermine Billard¹, Marie Mandart¹, Johannes Steiger²,
Télesphore Sime-Ngando¹ and Jonathan Colombet^{1*}

Affiliations

¹Université Clermont Auvergne, CNRS, Laboratoire Microorganismes : Genome, Environnement (LMGE), F-63000 Clermont-Ferrand, France

² Université Clermont Auvergne, CNRS, GEOLAB, F-63000 Clermont-Ferrand, France.

* Address correspondence to: jonathan.colombet@uca.fr

Abstract

Aster Like Nanoparticles (ALNs) are newly described femto-entities. Their ecology (*e.g.* geographic distribution, spatial dynamic, preferences, forcing factors) is still unknown. Here we report that these entities, which have largely been ignored until now, can develop or maintain themselves in most aquatic environments in the Loire River catchment, France. We observed a significant influence of the trophic state on ALN ecological distributions. A positive relationship between prokaryotic abundance and ALN ($r^2 = 0.72$, $p < 0.01$) has been identified, but its exact nature remains to be clarified. Combined with their ubiquitous distribution and high abundances (up to 7.9×10^6 ALNs.mL⁻¹) recorded in our samples, this probably makes ALNs an overlooked functional component in aquatic ecosystems.

The discovery of the importance of extracellular vesicles [1], bacteria CPR (*Candidate Phyla Radiation*), archaea DPANN (*Diapherotrites*, *Parvarchaeota*, *Aenigmarchaeota*, *Nanoarchaeota*, *Nanohaloarchaea*) [2,3] and ‘biomimetic mineralo-organic particles’ [4] permitted to reconsider the diversity and ecological role of the femtoplankton, hitherto confined to the sole viruses, some <0.2µm filterable prokaryotes [5]. The recent discovery of “*Aster Like Nanoparticles*” (ALNs) in pelagic aquatic environments showed that the femtoplankton still hosts many unknown and undervalued entities [6]. In a previous, unprecedented study on these entities, the authors have shown that ALNs are amorphous star-shaped entities suspected to be organic in nature (composed mainly of carbon, oxygen, calcium and nitrogen with traces of potassium). Pleomorphic, these entities present three main morphotypes, with 4, 11 and 20 arms, and a reduced biovolume, lower than of the smallest known prokaryote. These original characteristics, combined with their sensitivity to biocidal treatments and their ability to grow in the absence of cells, raise questions about their exact nature and origin [6]. Their high seasonal abundances (up to $9.0 \pm 0.5 \times 10^7$ entities·mL⁻¹), in the range of virus-like particles, may exceed those of prokaryotes by up to about one order of magnitude [6]. Added to observations of phenotypic close contact with prokaryotes [6], such numerical abundance levels probably make ALNs a significant actor in the functioning of aquatic systems. For example, its calcium composition [6] could have a significant impact on calcium homeostasis of aquatic environments. Without yet knowing their full biological nature, we hypothesized that their distribution and dynamics in a wide range of aquatic environments will open a window on the understanding of their ecological significance.

Except for first findings obtained from lake samples [6], we still lack information on the ecological distribution and spatio-temporal dynamics of ALNs, in relation to the potential environmental forcing factors.

Results acquired in this second study demonstrate that ALNs are overlooked femto-entities, widely distributed and that their dynamics are controlled by trophic status, in particular by the biological environment. Identification and quantification of ALNs and of their morphological variants are described in Colombet and collaborators [6]. Twenty-five study sites distributed over the Loire River catchment (France) were selected to represent different aquatic systems (rivers, canal, marshes, lakes, mudflat, and coastal ocean) as well as longitudinal gradient from the source to the estuary (Figure 1a). Sampling sites were classified into three trophic states according to all abiotic and biotic parameters measured *in-situ*. Further details on the methodology used are provided in the supplementary text and data.

ALNs have the ability to colonize different aquatic environments: lakes, rivers, marshes, coastal oceanic waters, except source areas and altitude lakes (Figure 1). They were found in environments with contrasting physico-chemical (*e.g.* conductivity, oxygen, nutrients) and biological (*e.g.* virus, prokaryotes, algae) characteristics (Supplementary Table S1). This defines them as tolerant and ubiquitous entities capable of maintaining themselves in the majority of continental and coastal aquatic environments. The abundances of ALNs recorded ($1.1 \times 10^7 \pm 0.17 \times 10^7$ ALNs.mL⁻¹) indicate preferential development in plain lakes, marshes, Loire estuary and in lesser extent within free-flowing river channels (Figure 1b-c). Detection of ALNs has also been mentioned in the tropical estuarine system of Ha Long Bay (Vietnam) and in the Saloum River in Senegal, thus broadening the habitat preferences and potential distribution of these entities [6].

Fig 1. Sampling site locations (1-25) and distribution of ALNs in different aquatic environments along the watershed of Loire River (France). **a.** Site locations are plotted on the map of the Loire watershed. Each site or river continuum is characterized by ALN concentration range. **b.** Distribution of the ALN concentrations and their different forms along the river continuum. **c.** Comparisons of the ALN concentrations and their different forms between the different aquatic environments sampled. Statistical differences between aquatic environments (test U, *** = $p < 0.01$) are provided in the inserted table (S= Source area, I= Intermediate Loire, D= Downstream Loire, A= Altitude lakes, P=Plain lakes, M=Marshes, NS=Not significant). m= mean.

87 However, sampling of these studies was sporadic, and these conclusions must be weighted by
88 the high seasonal or even daily fluctuations of ALNs recorded previously [6]. The sampling
89 period, *i.e.* spring, was based on the results obtained from the unique study on seasonal
90 monitoring of ALNs in a eutrophic lake [6]. We recognize the need of additional data on the
91 seasonal variability of ALNs in different aquatic environments in order to optimize sampling
92 periods.

93
94 The presence of ALNs in the coastal ocean samples raises questions about their potential for
95 development in the marine environment and their possible transfer from continental waters to
96 the oceans. We suggest that the gradual and significant increase of ALN abundance
97 (undetectable to 8.8×10^5 ALNs.mL⁻¹) from the source to the estuary (Figure 1b) can be
98 explained through the ALN downstream transfer along the river continuum, as observed for
99 other entities or pollutants [7-9]. The occurrence of 11-budding-arm forms in the downstream
100 area suggests that ALNs could be transported from the upstream catchment to more favorable
101 conditions for their development downstream. The occurrence of the different forms of ALN
102 indicates dominance of 20-arm forms (69 % of total ALNs) in marshes, while 11-arm forms
103 dominate in lakes and rivers (38 % and 47 % of total ALNs, respectively; Figure 1c).
104 Environmental conditions probably select and/or favor the development of one type in respect
105 to another, through a process that remains unexplained and which could either reflect the
106 gradual development from one form to another or the occurrence and development of rather
107 independent forms. In order to better understand spatial distributions of ALNs, future
108 investigations will have to demonstrate if a form of resistance distinct from forms of
109 propagation able to develop over time (*i.e.* assuming distinct growth stages of the same
110 organism) [6], exist.

112 These observations also point to the hypothesis that ALNs could be present in all aquatic
113 environments but are selected by environmental factors (Baas Becking hypothesis) [10]. Such
114 an assumption is supported by the capacity of ALNs to develop in unrelated biomes, from
115 temperate regions separated by hundreds of kilometers as in our study, to tropical regions [6].
116 According to the parameters recorded, we reported a significant influence of the trophic state
117 on ecological distributions of ALNs(Fig. 2a) with a significant increase from undetectable
118 ALNs.mL⁻¹ in the environments classified in the first trophic state, to 2.2×10^6 ALNs.mL⁻¹ in
119 those classified in the third trophic state [11]. In this study, the trophic state was estimated
120 according to the Trophic State Index, a reference method for the classification of aquatic
121 environments [12].

122
123 Therefore, we suggest that ALNs, in addition to certain microbial components such as
124 phytoplankton communities [13], may serve as a putative indicator of trophic conditions of
125 aquatic environments. Whatever the aquatic system considered, it seems that biological features
126 prevail over physico-chemical ones in the regulation of the distribution and abundances of
127 ALNs (Fig. 2b). The major correlative factor identified is the abundance of prokaryotes (r^2
128 =0.72, $p < 0.01$; Fig. 2b). A significant statistical correlation and the observation of direct
129 physical contact between prokaryotes and ALNs [6], suggest an interaction between these two
130 components. Previous study showed that ALNs can grow without a potential host [6], however,
131 the presence of prokaryotes could promote their development through mechanisms that remain
132 unknown [6]. Therefore, future investigations are needed to elucidate the nature of this mutual
133 relationship which, perhaps, could be related to either bottom-up resource-driven or to top-
134 down host-symbiont interactions.

b

	Prokaryotes	Viruses	Microphytoplankton	Picocyanobacteria	Picophytoplankton	T°C	pH	O ₂	Conductivity	P	N	C
ALNs	0.72	0.55	0.53	0.66	0.5	0.47	0.46	-0.3	0.45	0.31	0.19	0.36
Prokaryotes		0.71	0.68	0.67	0.69	0.44	0.5	-0.18	0.45	0.32	0.14	0.61
Viruses			0.38	0.46	0.37	0.36	0.42	-0.02	0.28	0.12	-0.12	0.59
Microphytoplankton				0.5	0.57	0.1	0.16	-0.16	0.15	0.32	0.23	0.56
Picocyanobacteria					0.6	0.6	0.46	-0.12	0.6	0.15	0.3	0.23
Picophytoplankton						0.46	0.35	-0.19	0.36	0.2	0.15	0.44
T°C							0.51	-0.41	0.79	-0.09	0.13	0.08
pH								0.33	0.63	0.27	0.13	0.17
O ₂									-0.1	0.1	-0.08	-0.13
Conductivity										-0.04	0.17	0.09
P											0.38	0.4
N												-0.09

Spearman correlation coefficient > 0.60 / < -0.60
 p-value < 0.01

Fig 2. Relationships between ALNs and physical-chemical and biological environment. **a.** Comparison of the average (test U, $p < 0.01$) of ALNs according to the low (state 1 = sites 1, 2, 3, 4), medium (state 2 = sites 5, 6, 8, 11, 14, 15, 16, 18, 19, 20, 24, 25) and high (state 3 = sites 7, 9, 10, 12, 13, 17, 21, 22, 23) trophic levels of the different aquatic environments. **b.** Correlation analysis (Spearman test, $p < 0.01$) between the different physical chemical and biological parameters of the different sampling points. T°C = temperature, O₂=dissolved oxygen, P= total organic phosphorus, N= total organic nitrogen, C= total organic carbon.

Our findings reinforced the hypothesis of the biological origin of ALNs [6] and the hypothesis that ALNs are ubiquitous and widespread within continental freshwaters and coastal brackish waters. The estimation of the ecological importance of femtoplanktonic entities depends closely on their abundance and functions. This is demonstrated for viruses [14,15], extracellular vesicles, ultramicro-prokaryotes [2,3], and suspected for non-biological entities such as 'biomimetic mineralo-organic particles' [4]. The average abundances of 9.9×10^5 and 7.9×10^6 we recorded respectively in plain lakes and marshes area (see fig.1), are in the range of those of prokaryotes or viruses. The strong empirical relationship with prokaryotes and the regulation by their environment implies that ALNs are additional and so far, ignored actors in aquatic ecosystems. Furthermore, to fully understand the ecological significance of ALNs in plankton dynamics, it is crucial to validate their biological nature.

Fundings

MF was supported by a PhD fellowship from the CPER 2015-2020 SYMBIOSE challenge program (French Ministry of Research, UCA, CNRS, INRA, Auvergne-Rhône-Alpes Region, FEDER). This study is a contribution to the “C NO LIMIT” project funded by the Interdisciplinary Mission of the French National Center of Scientific Research (CNRS) Program X-life, 2018 edition. This research was also financed by the French government IDEX-ISITE initiative 16-IDEX-0001 (CAP 20-25).

Acknowledgments

This study was supported by the CYSTEM platform UCA-PARTNER (University of Clermont Auvergne UCA), Clermont-Ferrand, France) and the Laboratory Microorganisms: Genome and Environment (LMGE, UMR6023 CNRS-UCA, Clermont-Ferrand, France).

Conflict of Interest

The authors declare no conflict of interest.

Authors contribution

MF, HB, MM, JC designed the study and analyzed data. All authors wrote the manuscript, contributed critically and gave final approval for publication.

References

- [1] Soler N, Krupovic M, Marguet E, Forterre P (2015) Membrane vesicles in natural environments: a major challenge in viral ecology. *The ISME Journal*. 9(4):793–796. doi:10.1038/ismej.2014.184
- [2] Castelle C.J, Banfield J.F (2018) Major New Microbial Groups Expand Diversity and Alter our Understanding of the Tree of Life. *Cell*. 172:1181–1197. <https://doi.org/10.1016/j.cell.2018.02.016>
- [3] Ghuneim L.-A.J, Jones D.L, Golyshin P.N, Golyshina O.V. (2018) Nano-Sized and Filterable Bacteria and Archaea: Biodiversity and Function. *Front Microbiol*. 9. doi:10.3389/fmicb.2018.01971
- [4] Wu C.-Y, Martel J, Wong T-Y, Young D, Liu C-C, Lin C-W, Young J.D (2016). Formation and characteristics of biomimetic mineralo-organic particles in natural surface water. *Scientific Reports*. 6(1). doi:10.1038/srep28817
- [5] Sieburth J.M, Smetacek V, Lenz J (1978) Pelagic ecosystem structure: Heterotrophic compartments of the plankton and their relationship to plankton size fractions. *Limnology and Oceanography*. 23(6):1256–1263. doi:10.4319/lo.1978.23.6.1256
- [6] Colombet J, Billard H, Viguès B, Balor S, Boulé C, Geay L, Benzerara K, Menguy N, Ilango G, Fuster M, et al (2019) Discovery of High Abundances of Aster-Like Nanoparticles in Pelagic Environments: Characterization and Dynamics. *Frontiers in Microbiology*. 10. doi:10.3389/fmicb.2019.02376

211 [7] Harvey E, Gounand I, Little C.J, Fronhofer E.A, Altermatt F (2017) Upstream trophic structure modulates
212 downstream community dynamics via resource subsidies. *Ecology and Evolution*.7(15):5724–5731.
213 doi:10.1002/ece3.3144

214 [8] Dann L.M, Smith R.J, Jeffries T.C, McKerral J.C, Fairweather P.G, Oliver R.L, Mitchell J.G (2017) Persistence,
215 loss and appearance of bacteria upstream and downstream of a river system. *Marine and Freshwater Research*.
216 68(5):851. doi:10.1071/mf16010

217 [9] Cho K.H, Pachepsky Y.A, Oliver D.M, Muirhead R.W, Park Y, Quilliam R.S, Shelton, D.R. (2016) Modeling fate
218 and transport of fecally-derived microorganisms at the watershed scale: State of the science and future
219 opportunities. *Water Research*. 100:38–56. doi:10.1016/j.watres.2016.04.064

220 [10] Baas Becking, L.G.M., and Nicolai, E. (1934). On the ecology of a Sphagnum Bog. *Blumea - Biodivers. Evol.*
221 *Biogeogr. Plants* 1, 10–45.

222 [11] Carlson R.E., J. Simpson (1996) A Coordinator's Guide to Volunteer Lake Monitoring Methods. North American
223 Lake Management Society. 96 pp.

224 [12] Carlson, R. E. (1977) A trophic state index for lakes. *Limnol. Oceanogr.* 22:361-369.
225

226 [13] Martinet J, Guédant P, Descloux S (2016) Phytoplankton community and trophic status assessment of a newly
227 impounded sub-tropical reservoir: case study of the Nam Theun 2 Reservoir (Lao PDR, Southeast Asia).
228 *Hydroécologie Appliquée*. 19:173–195. <https://doi.org/10.1051/hydro/2015006>

229 [14] Sime-Ngando T, Colombet J (2009) Virus and prophages in aquatic ecosystems. *Can. J. Microbiol.* 55:95–109.

230 [15] Brussaard C.P.D (2004) Viral Control of Phytoplankton Populations—a Review. *J Eukaryot Microbiol.* 51:125–
231 138

232