**Characterization of a novel type III alcohol dehydrogenase from *Thermococcus*
barophilus Ch5**

Likui Zhang^{1,2#}, Donghao Jiang², Yuting Li², Leilei Wu², Qing Liu², Kunming Dong²
and Philippe Oger^{3#}

¹Guangling College, Yangzhou University, China

²College of Environmental Science and Engineering, Marine Science & Technology
Institute, Yangzhou University, China

³Univ Lyon, INSA de Lyon, CNRS UMR 5240, Villeurbanne, France

[#]Corresponding author: Dr. Likui Zhang

Tel: +86-514-89795882

Fax: +86-514-87357891

E-mail address: lkzhang@yzu.edu.cn

Corresponding author: Prof. Philippe Oger

E-mail address: philippe.oger@insa-lyon.fr

Abstract

The genome of the hyperthermophilic and piezophilic euryarchaeon *Thermococcus barophilus* Ch5 encodes three putative alcohol dehydrogenases (Tba ADHs). Herein, we characterized Tba ADH₅₄₇ biochemically and probed its mechanism by mutational studies. Our data demonstrate that Tba ADH₅₄₇ can oxidize ethanol and reduce acetaldehyde at high temperature with the same optimal temperature (75°C) and exhibit similar thermostability for oxidation and reduction reactions. However, Tba ADH₅₄₇ has different optimal pH for oxidation and reduction: 8.5 for oxidation and 7.0 for reduction. Tba ADH₅₄₇ is dependent on a divalent ion for its oxidation activity, among which Mn²⁺ is optimal. However, Tba ADH₅₄₇ displays about 20% reduction activity without a divalent ion, and the maximal activity with Fe²⁺. Furthermore, Tba ADH₅₄₇ showcases a strong substrate preference for 1-butanol and 1-hexanol over ethanol and other alcohols. Similarly, Tba ADH₅₄₇ prefers butylaldehyde to acetaldehyde as its reduction substrate. Mutational studies showed that the mutations of residues D195, H199, H262 and H234 to Ala result in the significant activity loss of Tba ADH₅₄₇, suggesting that residues D195, H199, H262 and H234 are responsible for catalysis. Overall, Tba ADH₅₄₇ is a thermoactive ADH with novel biochemical characteristics, thereby allowing this enzyme to be a potential biocatalyst.

Keywords: Hyperthermophilic Archaea; Alcohol dehydrogenase; Biochemical characteristics

1 Introduction

Alcohol dehydrogenases (ADHs), which are members of the oxidoreductase family, are able to catalyze the cofactor-dependent oxidation of ethanol and other alcohols to produce the corresponding aldehydes and ketones [1]. Thus, ADHs play important roles in the detoxification and metabolism of primary and secondary alcohols. Based on their cofactor requirements, ADHs are classified into four types: nicotinamide adenine dinucleotide (NAD) dependent ADH; pyrroloquinoline quinone (PQQ) -dependent ADH; heme or cofactor F420 -dependent ADH; flavine adenine dinucleotide -(FAD) dependent ADH [2]. In addition, NAD-dependent ADHs are further subdivided into three types depending on the alcohols metabolized: short-chain ADHs exemplified by the drosophila ADH [3] (type I), medium-chain or zinc ADHs (type II or Zn-ADH) exemplified by the horse liver ADH [4], and long-chain or iron ADHs (type III or Fe-ADH) exemplified by the *Zymomonas* ADH2 [5, 6]. These three families display low similarity and arose independently throughout evolution, suggesting that these ADHs harbor different structures and mechanisms of reaction.

Several type III ADHs have been biochemically characterized since a type III ADH was first reported from *Zymomonas mobilis*, including *Saccharomyces cerevisiae* ADH4 [7], *Escherichia coli* L-1,2-propanediol oxidoreductase [8], *Clostridium acetobutylicum* ADH1 [9], *C. acetobutylicum* butanol dehydrogenases [10], and *Bacillus methanolicus* methanol dehydrogenase [11], and a human iron-activated ADH [12]. Crystal structures of the type III ADHs from bacteria and

1 archaea show that they possess a very highly conserved catalytic cleft where Fe^{2+}
2
3 ion is coordinated with three His residues and one Asp residue [13-16]. Currently,
4
5 several type III ADHs from hyperthermophilic *Thermococcus* have been reported since
6
7 the first type III ADH was characterized from *Thermococcus litoralis* [17], including
8
9 *Thermococcus* strain ES-1 [18-19], *Thermococcus* strain AN1 [20], *Thermococcus*
10
11 *hydrothermalis* [21], and *Thermococcus thio还原ens* [15] and *Pyrococcus furiosus*
12
13 [22]. All these reported type III ADHs from HA have one common characteristic:
14
15 high thermostability, which is distinct from other ADHs from mesophiles. Although
16
17 several ADHs from HA have been found, thermostable ADHs are still in demand due
18
19 to unique characteristics, demonstrating a variety of substrate specificities and
20
21 physiological functions.
22
23
24
25
26
27
28
29
30

31 *Thermococcus barophilus* Ch5, which was isolated from a deep-sea
32
33 hydrothermal field of the Mid-Atlantic Ridge (Logachev field chimney, 3,020 m
34
35 depth), is a thermo-piezophilic euryarchaeon, growing optimally at 88°C under
36
37 40 MPa of hydrostatic pressure [23]. The genome of *T. barophilus* Ch5 encodes 2,679
38
39 proteins, two thirds of which have no associated functions [24]. Furthermore, several
40
41 thermostable enzymes from *T. barophilus* Ch5 have been characterized [25-29] and
42
43 have been shown to have distinct characteristics from other homologues found in
44
45 other extremophilic archaeal models. Thus, *T. barophilus* Ch5 is an important model
46
47 for studying structural and function relationship of thermostable proteins or enzymes
48
49 present in hyperthermophilic archaea as well as a great untapped source of novel extreme
50
51 thermophilic enzymes for industrial processes.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Three putative alcohol dehydrogenases (Tba ADHs) are encoded in the genome of *T. barophilus* Ch5, including one short-chain ADH (GenBank: ALM74123) and two putative type III ADHs (GenBank: ALM74514 and ALM74601). Interestingly, these two putative Fe-ADHs from *T. barophilus* Ch5 have only 26% similarity in amino acid compositions, suggesting that they might harbor distinct characteristics. Currently, no ADHs from *T. barophilus* Ch5 has been biochemically characterized. In this work, we report the cloning, expression, biochemical characterization and mutational studies of the recombinant Tba ADH₅₄₇ encoded by gene TBCH5v1_0547 (Genbank: ALM74514) and is conserved in all *Thermococcales* species, suggesting that this ADH might be a putative type III alcohol dehydrogenase with novel characterization.

2 Materials and methods

2.1 Materials

The pET-30a(+) vector was purchased from Novagen (Merck, Darmstadt, Germany). Plasmid Extraction Kit, PCR Cycle Pure Kit, and Gel Extraction Kit were purchased from Omega (Guangzhou, China). *Escherichia coli* DH5 α cells and *E. coli* BL21 (DE3) cells were purchased from Transgene (Beijing, China); dNTPs, T4 DNA ligase, *Nde*I, *Xho*I, and Pfu DNA polymerase were purchased from Thermo Scientific (Waltham, MA). Chemicals were purchased from Amresco (WA, USA).

2.2 Phylogenetic analyses

The genes coding for Tba ADH₅₄₇ and Tba ADH₆₄₁ were used to identify their homologues in all *Thermococcales* species which genomes are sequenced. ADH

1 sequences were aligned in Seaview with Muscle [30]. Conserved sites were selected
2
3 with GBlocks. PhyML v3.696 was employed to construct a maximum likelihood
4
5 unrooted phylogenetic tree. Bootstrapping was performed with 1000 replicates. The
6
7 sequence accession numbers for each individual protein is labeled in the tree. The tree
8
9 was inferred with PhyML. The scale bar represents the average number of
10
11 substitutions per site. Numbers at branches represent bootstrap values (1000 replicates,
12
13 values > 50% are shown).
14
15
16
17
18
19

20 2.3 Sequence alignment

21
22 The target ADH sequences from archaea, bacteria and eukaryotes were retrieved
23
24 from NCBI. The sequences were aligned by DNAMAN software version 6. The
25
26 alignment figure was output with a graphic file.
27
28
29
30

31 2.4 Cloning of the gene encoding *Tba ADH₅₄₇*

32
33 We amplified the TBCH5v1_0547 gene encoding a putative type III alcohol
34
35 dehydrogenase using the *T. barophilus* Ch5 genomic DNA as a template and Pfu
36
37 DNA polymerase in the presence of the forward primer and the reverse primer. The
38
39 sequences of these two primers are listed in Table 1. The amplified PCR product and
40
41 the pET-30a(+) vector were digested by *NdeI* and *XhoI*, and the cleaved DNA
42
43 fragments were ligated by T4 DNA ligase to yield the recombinant plasmid
44
45 pET30a(+)-*Tba ADH₅₄₇*, and transformed into *E. coli* DH5 α cells. After sequence
46
47 verification, the recombinant plasmid pET30a(+)-*Tba ADH₅₄₇* was transformed into *E.*
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
coli BL21 (DE3) cells for expression of the *Tba ADH₅₄₇* protein harboring a 6 x
His-tag in its C-terminus.

2.5 Overexpression and purification of Tba ADH₅₄₇

For the Tba ADH₅₄₇ protein expression, the expression strain *E. coli* BL21 (DE3) harboring the pET30a(+)-Tba ADH₅₄₇ plasmid was cultured at 37°C in LB medium containing 10 µg/mL kanamycin until the OD₆₀₀ reached 0.6. Then, the expression of the Tba ADH₅₄₇ gene was induced by addition of isopropyl thiogalactoside (IPTG, 0.1 mM). Expression was performed for 12 hat room temperature.

Cells were harvested by centrifugation (5,000 g) at room temperature. The collected pellet was resuspended in Ni column buffer A containing 20 mM Tris-HCl (pH 8.0), 1 mM dithiothreitol (DTT), 500 mM NaCl, 50 mM imidazole and 10% glycerol. Cells were disrupted by ultrasonication at 4°C. Cell debris was removed by centrifugation (16,000 g) at 4°C. The collected supernatant was heated at 70°C for 20 min to inactivate the *E. coli* proteins. After centrifugation at 16,000 g and 4°C, the resulting supernatant was loaded onto a HisTrap FF column (GE Healthcare, Uppsala, Sweden). The Tba ADH₅₄₇ protein was eluted with NCGTM Chromatography System (Bio-Rad, Hercules, CA, USA) by a linear gradient of 50–500 mM imidazole with a Ni column buffer B containing 20 mM Tris-HCl (pH 8.0), 1 mM DTT, 500 mM NaCl, 500 mM imidazole and 10% glycerol. The collected fractions containing the His-tagged Tba ADH₅₄₇ protein were analyzed by migration on a 10% SDS-PAGE. The purified Tba ADH₅₄₇ protein fractions were combined and dialyzed against a storage buffer containing 50 mM Tris-HCl (pH 8.0), 1 mM DTT, 50% glycerol and 50 mM NaCl, and stored at -80°C. The Tba ADH₅₄₇ protein concentration was determined by measuring the absorbance at 280 nm. The theoretical molar extinction

coefficient of Tba ADH₅₄₇ protein is predicted to be 33,350 M⁻¹ cm⁻¹.

2.6 Construction, overexpression, and purification of the Tba ADH₅₄₇ mutants

Following the manufacturer's instructions, the Tba ADH₅₄₇ mutants were constructed by Site-Directed Mutagenesis Kit (Transgene, China) using the wild-type Tba ADH₅₄₇ gene as a template. The sequences of the mutagenic primers are listed in Table 1. The D195A, H199A, H262A, H266A, and H274A mutant plasmids were verified by sequencing, and the mutant proteins were overexpressed, purified and quantified as described for the wild-type protein.

2.7 Enzymatic assays of Tba ADH₅₄₇

The standard Tba ADH₅₄₇ enzymatic assay was performed in a total volume of 3 ml containing 100 mM Tris-HCl (pH 8.5), 0.67 mM NAD⁺, 0.1 mM MgCl₂, 0.46 M ethanol and 500 nM Tba ADH₅₄₇ at 75°C for 30 min and was terminated by placing samples on ice. Alternatively, the standard reaction for the reductive reaction in a total volume of 3 ml including 100 mM Tris-HCl (pH 8.5), 0.3 mM NADH, 0.46 M acetaldehyde, 0.1 mM FeSO₄, and 500 nM Tba ADH₅₄₇ was conducted at 75°C for 30 min and were terminated by placing samples on ice. Tba ADH₅₄₇ activity was determined spectrophotometrically at 340 nm by measuring the amount of NADH produced or reduced. The extinction coefficient (ϵ_{340}) was 6,220 M⁻¹ cm⁻¹. All reactions were performed in triplicate and the average values were reported. One unit of enzyme was defined as the amount of enzyme producing or reducing 1 μ mol of NADH per min.

The optimal reaction temperature of the Tba ADH₅₄₇ activity was investigated in

the standard conditions but varying temperature in a 40°C to 100°C range for oxidation and in a 40°C to 80°C for reduction. The optimal pH of the Tba ADH₅₄₇ activity was determined at 75°C varying the pH of the reactions between 6.5 and 10 for oxidation and between 6.0 and 8.0 for reduction. The varied pHs were adjusted with five different buffers (all at 20 mM concentrations): sodium phosphate-NaOH (pH 6.0, pH 6.5 and pH 7.0), Tris-HCl (pH 7.5, pH 8.0, and pH 8.5 at the desired temperature), and Gly-NaOH (pH 9.0, pH 9.5 and pH 10.0). The requirement for a divalent ion for the activity of Tba ADH₅₄₇ was assayed in presence of 10 mM EDTA or 0.1 mM Fe²⁺, Mg²⁺, Mn²⁺, Co²⁺, Cu²⁺, Ca²⁺, Ni²⁺ or Zn²⁺. The substrate specificity of the Tba ADH₅₄₇ activity was investigated with ethanol, methanol, isopentanol, isopropanol, ethylene glycol, 1-butanol, 1-hexanol or glycerol as the substrates for oxidation, and with acetone, butylaldehyde, caproaldehyde or acetaldehyde as the substrates for reduction.

2.8 Kinetic assays of Tba ADH₅₄₇

The kinetic parameters of Tba ADH₅₄₇ were obtained by performing reactions at different ethanol concentrations (1 mM, 5 mM, 10 mM, 20 mM, 40 mM, 80 mM, 100 mM and 200 mM) for oxidation and at different acetaldehyde concentrations (1 mM, 2.5 mM, 5 mM, 10 mM, 20 mM, 40 mM, 80 mM, and 100 mM) for reduction. The corresponding K_m and V_{max} values were calculated using the linear regression and the values given represented the calculated mean.

2.9 Circular dichroism (CD) measurements

The wild-type (WT), D195A, H199A, H262A, H266A and H274A mutant Tba

ADH₅₄₇ proteins were dialyzed into 50 mM PBS (phosphate-buffered saline) pH7.5 for CD analysis. The CD spectra were recorded at 20°C from 200 nm to 250 nm using a J-810 spectropolarimeter (JASCO, Japan) and a cuvette of path length 0.2 cm. The spectra were generated with 0.2 mg/ml WT and mutant Tba ADH₅₄₇ proteins at a scanning rate of 50 nm/min, and triplicate spectrum readings were collected per sample. The CD spectral data were reported as mean residue ellipticity [θ] by using the program KaleidaGraph (Synergy Software).

3 Results

3.1 Tba ADH₅₄₇ is a putative type III alcohol dehydrogenase

Two putative NAD-ADHs of type III (Fe-ADH) are encoded in the genome of *T. barophilus* Ch5: Tba ADH₆₄₁ and Tba ADH₅₄₇. As shown in Fig. 1A, Tba ADH₅₄₇ and Tba ADH₆₄₁ are clustered in different branches. Tba ADH₅₄₇ is conserved in all *Thermococcales* species (Fig. 1B), including *Thermococcus*, *Pyrococcus* and *Palaeococcus* genera. Tba ADH₅₄₇ is the most closely related to ADH from *T. paralvinellae* (Fig. 1B), and this ADH displays a relative closer relation with the ADH homologues from *Palaeococcus* than the ADH homologues from *Pyrococcus* in *Thermococcales*. Surprisingly, Tba ADH₅₄₇ is not conserved in other Archaea (Fig. 1B). It is only distantly related to bacterial homologues (Fig. 1B). Tba ADH₅₄₇ displays 26~30% similarity with Tba ADH₆₄₁ and the homologues from *Thermococcus* strain ES-1, *T. hydrothermalis*, *T. litoralis*, *Thermococcus* strain AN1, *Thermococcus kodakarensis* (ADH2), *Pyrococcus furiosus*, *Z. mobilis* and *S. cerevisiae*. In contrast, Tba ADH₅₄₇ exhibits 71~75% similarity with the homologues

1 from *Thermococcus* strain AN1, *T. thioeducens* and *T. kodakarensis* (ADH1),
2
3 suggesting that Tba ADH₅₄₇ might be more similar in function to *T. zilligii*, *T.*
4
5
6 *thioeducens* and *T. kodakarensis* (ADH1) ADHs. Thus, the low similarity of two type
7
8
9 III ADHs from *T. barophilus* Ch5 suggests that they might have distinct biochemical
10
11 characteristics.
12

13
14 As shown in Fig. 2, Tba ADH₅₄₇ possesses three conserved motifs: motif I
15
16 (GGGS), motif II (DxxxH), and motif III (a conserved motif with three histidine
17
18 residues), which are present in type III ADHs. However, the amino acid variations
19
20 and three residues are missing in the highly conserved motif III in Tba ADH₅₄₇ and
21
22
23 ADHs from *T. thioeducens*, *T. kodakarensis* (ADH1) and *Thermococcus* strain AN1.
24
25
26 Furthermore, thirty four residues are strictly conserved in all thirteen proteins (Fig. 2),
27
28
29 including seven glycine residues, six proline residues, six alanine residues, four
30
31
32 histidine residues, three aspartic acid residues, two lysine residues, two glutamic acid
33
34
35 residues, one threonine, one leucine and one isoleucine. Overall, Tba ADH₅₄₇
36
37
38 possesses the conserved motifs present in type III ADHs, however, displays low
39
40
41 similarity with the homologues exemplified by Tba ADH₆₄₁ ADH.
42
43

44 3.2 Cloning, expression and purification of Tba ADH₅₄₇ protein

45
46

47 In this work, we cloned the Tba ADH₅₄₇ gene encoded by TBCH5v1_0547
48
49 (Genbank: ALM74514) into a pET30a(+) expression vector and overexpressed it in *E.*
50
51
52 *coli* BL21 (DE3) cells. As shown in Fig. 3A, the gene encoding the Tba ADH₅₄₇
53
54
55 protein was successfully induced after addition of IPTG and the expressed protein was
56
57
58 recovered in the supernatant after ultrasonication disruption. After heat treatment at
59
60
61

70°C for 30 min, most of *E. coli* proteins were inactivated and removed after centrifuge, however, the Tba ADH₅₄₇ protein can withstand this heat treatment. Last, the recombinant Tba ADH₅₄₇ protein was purified to near homogeneity after Ni-column affinity purification (Fig. 3A). The purified 6 x His-tagged protein displayed an approximate MW of 42 kDa (Fig. 3A), consistent with the deduced amino acid sequence.

3.3 Effect of temperature on Tba ADH₅₄₇ activity

Since *T. barophilus* Ch5 is a hyperthermophile [23], we expect its proteins or enzymes to be highly thermophilic and thermostable. We first investigated the thermophilicity of Tba ADH₅₄₇ by performing enzyme activity assays at varied temperatures ranging from 40°C to 100°C. As shown in Fig. 3B, Tba ADH₅₄₇ only retained about 10% and 30% relative activity at 40°C and 100°C, respectively. Tba ADH₅₄₇ displayed maximal activity at 75°C, which makes it a thermophilic enzyme.

Next, we determined whether Tba ADH₅₄₇ has reduction activity. Using acetaldehyde as the substrate, we performed the reduction reaction catalyzed by Tba ADH₅₄₇ in the presence of NADH at various temperatures ranging from 40°C to 80°C since NADH is unstable at temperatures higher than 80°C. We found that Tba ADH₅₄₇ has the reduction activity, and also displays maximal activity for acetaldehyde reduction at 75°C (Fig. 3B). Thus, Tba ADH₅₄₇ has the same optimal reaction temperature for both ethanol oxidation and acetaldehyde reduction under our standard reaction conditions.

3.4 Thermostability of Tba ADH₅₄₇

To probe the thermostability of Tba ADH₅₄₇, we incubated this ADH at varied temperatures for different times and then used the heated enzyme to perform the ethanol oxidation and acetaldehyde reduction reactions. We selected the heated Tba ADH₅₄₇ at 90, 95 and 100°C for 30 min for analyzing its reduction activity. As shown in Fig. 4A, the reduction activity of Tba ADH₅₄₇ decreased as the heating temperature increased. However, this ADH still retained 20% reduction activity even after this ADH was heated at 100°C for 30 min.

On the other hand, we employed the heated Tba ADH₅₄₇ at 100°C at varied times to analyze its oxidation reaction. Compared with the control reaction without the heating treatment, the oxidation activity of Tba ADH₅₄₇ was lowered as the heating time extended, (Fig. 4B). However, Tba ADH₅₄₇ still harbored about 20% oxidation activity after heating at 100°C for 30 min, which is consistent with the observation for the reduction activity of the enzyme heated at 100°C for 30 min. Overall, Tba ADH₅₄₇ is a thermophilic and thermostable ADH, suggesting its potential application in ethanol oxidation and acetaldehyde reduction at high temperature.

3.5 Effect of pH on Tba ADH₅₄₇ activity

Next, we investigated the optimal pH for ethanol oxidation and acetaldehyde reduction of Tba ADH₅₄₇. As shown in Fig. 5A, no Tba ADH₅₄₇ activity was observed below pH 6.5 and only a residual activity was detected at pH 10. Tba ADH₅₄₇ can oxidize ethanol in the pH range from 7.0 to 10.0. At pH 8.5, Tba ADH₅₄₇ displayed maximal activity, suggesting that pH 8.5 is the optimal pH for the reduction reaction of this ADH.

1 The effect of pH on reduction reaction of Tba ADH₅₄₇ was also determined using
2
3 varied pH buffers ranging from 6.0 to 8.0 since NADH is unstable at above pH 8.0
4
5 (Fig. 5A). Tba ADH₅₄₇ exhibited maximal reduction activity at pH7.0, suggesting that
6
7 its optimal reaction pH is 7.0, which is lower than that of its oxidation reaction. Thus,
8
9 Tba ADH₅₄₇ displays distinct optimal pH adaptation for oxidation and reduction
10
11 reactions.
12
13
14
15

16 3.6 Effect of divalent ions on Tba ADH₅₄₇ activity

17
18 The metal requirements of archaeal NAD-dependent ADHs are very variable,
19
20 thus, we investigated the metal dependency of Tba ADH₅₄₇ activity for ethanol
21
22 oxidation. In the absence of the addition of a divalent ion or in the presence of 10 mM
23
24 EDTA (the latter should chelate the divalent ions eventually co-purified with the Tba
25
26 ADH₅₄₇ enzyme), Tba ADH₅₄₇ yielded no measurable activity (Fig. 5B). The activity
27
28 in Tba ADH₅₄₇ is restored to some extent upon addition of all the divalent ions tested.
29
30 Maximum Tba ADH₅₄₇ activity was observed in the presence of 0.1 mM Mn²⁺. By
31
32 contrast, the lowest activity of Tba ADH₅₄₇ was found in the presence of 0.1 mM Zn²⁺,
33
34 Cu²⁺, Ca²⁺, Ni²⁺, Co²⁺, Mg²⁺ or Fe²⁺, which stimulated Tba ADH₅₄₇ activity with less
35
36 than 50% of that observed with Mn²⁺. Thus, Tba ADH₅₄₇ is dependent on a divalent
37
38 ion for ethanol oxidation, which is sharply contrasted to the currently reported ADHs.
39
40
41
42
43
44
45
46
47
48
49

50 Compared with the oxidation reaction, Tba ADH₅₄₇ displays about 30%
51
52 reduction activity in the absence of divalent ion or in the presence of EDTA (Fig. 5B).
53
54 Surprisingly, this ADH exhibits maximal reduction activity in the presence of Fe²⁺,
55
56 suggesting that Fe²⁺ is the optimal divalent ion for its reduction activity, which is
57
58
59
60
61
62
63
64
65

sharply contrasted with the observation that Mn^{2+} is the optimal ion for the oxidation activity of this ADH. Compared with the control reactions without divalent ion or without EDTA, Ca^{2+} slightly inhibited the reduction activity whereas Cu^{2+} , Mg^{2+} , or Ni^{2+} stimulated the reduction activity of this ADH with varied degrees. On the other hand, Zn^{2+} or Mn^{2+} enables this ADH to retain similar reduction activity to the control reactions. Thus, Tba ADH₅₄₇ displays distinct divalent ion adaptation for acetaldehyde reduction.

3.7 Substrate specificity of Tba ADH₅₄₇

We determined the substrate specificity of Tba ADH₅₄₇ under the optimal conditions (75°C, pH 8.5 and 0.1 mM Mn^{2+}) for oxidation. Quite strikingly, we found that the recombinant Tba ADH₅₄₇ exhibited the highest activity in the presence of 1-butanol or 1-hexanol and not in the presence of ethanol, suggesting that 1-butanol and 1-hexanol are more suitable substrates for this enzyme than is ethanol (Fig. 6A). Only a few other primary alcohols were possible substrates for Tba ADH₅₄₇, including ethylene glycol (40% activity), isopentanol (25% activity), isopropanol (10% activity) and glycerol (5% activity). In contrast, we observed that Tba ADH₅₄₇ is fully inactive on methanol. Thus, Tba ADH₅₄₇ displays a preference for substrate in the order from high to low: 1-butanol > 1-hexanol > ethanol > ethylene glycol > isopentanol > isopropanol > glycerol. Combining all these observations, we can conclude that the optimal reaction conditions for Tba ADH₅₄₇ activity with 1-butanol as a substrate are: 75°C, pH 8.5 and 0.1 mM Mn^{2+} . Under the optimal reaction conditions, we measured the specific activity of Tba ADH₅₄₇ to be 931 U/mg for 1-butanol oxidation.

Next, we determined the substrate specificity of reduction reaction of Tba ADH₅₄₇ using acetone, acetaldehyde, caproaldehyde and butylaldehyde as the substrates. Similar to the oxidation reaction with 1-butanol as a preference substrate, Tba ADH₅₄₇ displayed higher reduction activity in the presence of butylaldehyde than in the presence of acetaldehyde (Fig. 6B), suggesting that butylaldehyde is more suitable substrate than acetaldehyde for reduction reaction of this ADH. Compared with the acetaldehyde reduction reaction, Tba ADH₅₄₇ retained significantly lowered reduction activity for acetone and caproaldehyde reduction, suggesting that acetone and caproaldehyde are also possible substrates for reduction activity of this ADH. Thus, the optimal reaction conditions for Tba ADH₅₄₇ reduction activity with butylaldehyde as a substrate are: 75°C, pH 7.0 and 0.1 mM Fe²⁺. The specific activity of Tba ADH₅₄₇ was measured to be 3924 U/mg for butylaldehyde reduction, suggesting that this ADH has a higher activity for aldehyde reduction than alcohol oxidation.

3.8 K_m and V_{max} for Tba ADH₅₄₇

We determined the kinetic parameters of Tba ADH₅₄₇ using the ethanol at varied concentrations ranging from 1 mM to 200 mM as the substrates under the optimal oxidation conditions as described above. The K_m and V_{max} values were calculated to be 92 mM and 8×10^{-3} mM/min from the fitted Lineweaver-Burk plot (Fig. 7A), respectively. Considering that 500 nM Tba ADH₅₄₇ was employed in oxidation reaction, the K_{cat} was further measured to be 0.27 s⁻¹.

Furthermore, we examined the kinetic parameters of Tba ADH₅₄₇ using the

1 acetaldehyde at varied concentrations ranging from 1 mM to 80 mM as the substrates
2
3 under the optimal reaction condition (75°C, pH 7.0 and 0.1 mM Fe²⁺). As shown in
4
5 Fig. 7B, the K_m and V_{max} values of Tba ADH₅₄₇ for acetaldehyde reduction were
6
7 estimated to be 34.5 mM and 0.266 mM/min from the fitted Lineweaver-Burk plot,
8
9 respectively. The corresponding K_{cat} for acetaldehyde reduction was further calculated
10
11 to be 69 s⁻¹, which is about 256-fold higher than that for ethanol oxidation. By
12
13 contrast, the K_m value for acetaldehyde reduction of Tba ADH₅₄₇ is about 2.7-fold
14
15 lower than that for ethanol oxidation. Thus, Tba ADH₅₄₇ displays higher catalytic
16
17 efficiency for acetaldehyde reduction than for ethanol oxidation.
18
19
20
21
22
23
24

25 *3.9 Mutational studies of Tba ADH₅₄₇*

26
27 As shown in Fig. 2A, Tba ADH₅₄₇ harbors three conserved motifs with several
28
29 conserved amino acid residues present other ADHs. To reveal key amino acid residues
30
31 for catalysis of Tba ADH₅₄₇, we constructed the D195A, H199A, H262A, H266A, and
32
33 H274A mutants. Following the wild-type protein expression and purification, we
34
35 purified five Tba ADH₅₄₇ mutant proteins (Fig. 8A).
36
37
38
39
40
41

42 To investigate whether the D195A, H199A, H262A, H266A, and H274A mutants
43
44 cause the overall structure change of Tba ADH₅₄₇, we performed the CD analysis of
45
46 the WT and mutant enzymes. The D195A, H199A, H262A, H266A, and H274A
47
48 mutants displayed different conformational change from the wild-type Tba ADH₅₄₇
49
50 with varied degrees (Fig. 8B). Specifically, the D195A, H266A and H274A mutants
51
52 exhibited similar overall structure (Fig. 8B), which are distinct from the WT enzyme.
53
54 Furthermore, the H199A substitution caused maximal change of overall structure of
55
56
57
58
59
60
61
62
63
64
65

1 the WT enzyme, followed by three substitutions (D195A, H266A and H274A) and the
2
3 H262A substitution (Fig. 8B). Thus, our observations demonstrate that the D195A,
4
5 H199A, H262A, H266A, and H274A substitutions disrupt the overall structure of the
6
7
8
9 WT enzyme.

10
11 Next, we investigated the ethanol oxidation and acetaldehyde reduction activities
12
13 of these five mutants. Compared with the wild-type protein, the D195A mutant
14
15 displayed no reduction activity and the slight weak oxidation activity (Fig. 8C),
16
17 suggesting that D195 is a key residue for catalysis of this ADH. Similarly, the H262A
18
19 mutant harbored the weak oxidation activity (2.9%) and reduction activity (6.4%),
20
21 thereby indicating that H262 is also a key residue for catalysis. However, the H266A
22
23 mutant had the higher reduction activity (134%) but the lower oxidation activity (36%)
24
25 than the wild-type protein, suggesting that the mutation of H266 to Ala stimulates its
26
27 reduction activity but suppresses its oxidation activity. Furthermore, the H199A and
28
29 H274A mutants retained 10%~20% oxidation and reduction activity, demonstrating
30
31 that residues H199 and H274 are involved partially in catalysis. Overall, residues
32
33 D195 and H262 in Tba ADH₅₄₇ are essential for catalysis, and residues H199 and
34
35 H274 are also involved partially in catalysis.

4. Discussion

46
47 In this work, we present the biochemical characteristics and mutational studies of
48
49 Tba ADH₅₄₇ which is one of the ADHs of the hyperthermophilic and piezophilic
50
51 euryarchaeon *T. barophilus* Ch5. Tba ADH₅₄₇ belongs to a cluster of type III archaeal
52
53 ADHs (Table 2) since this enzyme harbors three conserved motifs present in
54
55
56
57
58
59
60
61
62
63
64
65

1 iron-containing ADHs. However, Tba ADH₅₄₇ displays low amino acid similarity
2
3 with the reported iron-containing ADHs from hyperthermophiles, including *T.*
4
5
6 *litoralis* [17], *T. hydrothermalis* [21], *Thermococcus* strain ES-1 [18], *P. furiosus* [22]
7
8 and *Thermotoga hypogea* [31], suggesting this ADH might be a novel type III ADH.
9
10 Comparison of biochemical characteristics of iron-containing ADHs from
11
12 hyperthermophiles is summarized in Table 2, displaying distinct properties for
13
14 oxidation or reduction.
15
16
17
18
19

20 As expected, Tba ADH₅₄₇ displays maximal enzyme activity at high
21
22 temperatures since it originates from the hyperthermophilic euryarchaeon *T.*
23
24 *barophilus* Ch5. In comparison to other thermostable type III ADHs, Tba ADH₅₄₇
25
26 displays thermophilicity 5~10°C degrees lower than that reported for ADHs from *T.*
27
28 *hydrothermalis* (80°C) [21], *T. litoralis* (85°C) [17] and *Thermococcus* strain AN1
29
30 (85°C) [20]. Interestingly, the optimal temperatures of ADHs from *Thermococcus*
31
32 strain ES-1, *P. furiosus* and *T. hypogea* exceeded 100°C and could not be clearly
33
34 determined [18, 22, 31]. To our knowledge, we first measured the optimal temperature
35
36 of acetaldehyde reduction, demonstrating the same optimal temperature for ethanol
37
38 oxidation under our assay conditions. Overall, Tba ADH₅₄₇ harbors a relatively low
39
40 optimal temperature for its activity, compared with the reported type III ADHs from
41
42 hyperthermophiles.
43
44
45
46
47
48
49
50
51
52

53 In this work, we revealed that Tba ADH₅₄₇ retains about 40% and 20% activity
54
55 after being heated at 95°C and 100°C for 30 min, respectively. The half-life of *P.*
56
57 *furiosus* ADH was reported to be 7 h at 95°C [22] while the half-life of *T. litoralis*
58
59
60
61
62
63
64
65

ADH was about 0.3 h at 95°C [17]. *Thermococcus* strain ES-1 and *T. hypogea* ADHs retain 50% activity after being heated at 95°C for 4 h and at 90°C for 2 h, respectively [32, 34]. A half-life of *Thermococcus* strain AN1 ADH was measured to be only 16 min at 80°C [20]. Furthermore, *T. hydrothermalis* ADH abolishes 50% activity at 80°C for 0.25 h [21]. Thus, the type III ADHs from hyperthermophiles exhibit strong thermostability, but their degree of thermostability varies with different their host sources.

Maximal activity of Tba ADH₅₄₇ for ethanol oxidation was observed at pH 8.5, which is similar to other type III ADHs from *T. litoralis* and *Thermococcus* strain ES-1 [17-18]. Tba ADH₅₄₇ differs significantly from other extremophilic type III ADHs displaying a lower pH optimum than the type III-ADHs from *T. hydrothermalis*, *P. furiosus* and *T. hypogea* [21-22, 31]. However, maximal activity of Tba ADH₅₄₇ for acetaldehyde reduction was measured at pH 7.0, suggesting that this ADH displays various optimal pH for oxidation and reduction reactions. Similar observations are found in the reported ADHs from *Thermococcus* strain ES-1 [18], *T. hydrothermalis* [21], *P. furiosus* [22], and *T. hypogea* [31].

Although the members of the Fe-ADH family are described as iron-activated alcohol dehydrogenases, these enzymes can be activated by a range of divalent cations, including zinc, nickel, magnesium, copper, cobalt, or manganese (e.g., [32-35], besides iron. To date, divalent cations are needed for a type III ADH activity, among which Fe is essentially ADHs from *Thermococcales*. However, we found that Tba ADH₅₄₇ is the only known example of a type III archaeal ADH that is dependent

on a divalent ion for its oxidation activity. Interestingly, this ADH displays maximal oxidation activity in the presence of added Mn^{2+} . However, Tba ADH₅₄₇ can catalyze acetaldehyde reduction reaction in the absence of a divalent ion and displays maximal activity in the presence of Fe^{2+} for acetaldehyde reduction.

Among the characterized archaeal ADHs capable of oxidizing primary alcohols, all others isolated from *Thermococcales* species exhibit a greater preference for 1-butanol over ethanol as a substrate as observed for Tba ADH₅₄₇. Similar to the ADHs from *Thermococcus* strain AN1 [20], *T. litoralis* [17] and *Thermococcus* strain ES-1 [18], Tba ADH₅₄₇ shows a higher preference for primary alcohols of longer chain such as 1-hexyl alcohols. However, AdhD from *P. furiosus* essentially cannot process ethanol and has a strong preference for 1-butanol [36], and the ADH from *T. litoralis* has a marked preference for aliphatic primary alcohols [17]. In contrast to Tba ADH₅₄₇, *Thermococcus sibiricus* ADH, which belongs to the short-chain ADH family, demonstrates a preference for ethanol over 1-butanol, but is most probably a secondary alcohol dehydrogenase as shown by its strong preference for 2-butanol over 1-butanol [37]. In addition, similar to the ADHs from all the *Thermococcales* species listed above, Tba ADH₅₄₇ cannot utilize methanol as a substrate. In this respect, Tba ADH₅₄₇ and ADHs from the other *Thermococcales* differ from the ADHs found outside the *Thermococcales* order, such as *Picrophilus torridus* ADH [38] or *Thermoplasma acidophilum* ADH [39], which retain some level of activity with methanol.

As shown in Table 2, the K_m values of Tba ADH₅₄₇ for ethanol oxidation and

1 acetaldehyde reduction are higher than those of the reported ADHs from the other
2
3 *Thermococcales* order. Furthermore, the K_{cat} of Tba ADH₅₄₇ for ethanol oxidation is
4
5 lower than that of the reported ADHs from the other *Thermococcales* order. However,
6
7 the K_{cat} of Tba ADH₅₄₇ for acetaldehyde reduction is maximum among the all the
8
9 reported ADHs listed above. Importantly, the K_{cat}/K_m of Tba ADH₅₄₇ for acetaldehyde
10
11 reduction is 670-fold higher than that for ethanol oxidation, suggesting that this ADH
12
13 displays higher catalytic efficiencies for acetaldehyde reduction than ethanol
14
15 oxidation as suggested by Fernandez et al. who proposed that type III ADHs from
16
17 microorganisms contributed to aldehyde detoxication rather than alcohol oxidation
18
19 [40]. Similarly, 13-fold, 49-fold and 3.3-fold higher K_{cat}/K_m values for acetaldehyde
20
21 reduction than ethanol oxidation are observed for the ADHs from *Thermococcus*
22
23 strain ES-1, *P. furiosus* and *T. hypogea* [18, 22, 31], respectively. Thus, Tba ADH₅₄₇
24
25 displays higher K_{cat}/K_m values for acetaldehyde reduction over ethanol oxidation than
26
27 other ADHs from hyperthermophiles listed above. Since *T. barophilus* Ch5 is an
28
29 anaerobic euryarchaeon [23], the catalytic activity of Tba ADH₅₄₇ for ethanol synthesis
30
31 is far greater than that for ethanol oxidation, which might recover the NAD⁺ required
32
33 for another Entner–Doudoroff glycolytic (ED) pathway under anaerobic conditions.
34
35

36
37 Alignment of amino acid sequences from type III ADH demonstrates that they
38
39 possess a conserved motif III with the three highly conserved histidine residues.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

1 His267 could interact with the substrate, and therefore should be involved in catalysis,
2
3 but not in metal-binding [13]. Similar to *E. coli* propanediol oxidoreductase, Tba
4
5 ADH₅₄₇ harbors the corresponding residues H262, H266 and H274 in the conserved
6
7 motif III. Our mutational data show that the H262A, H266A and H274A Tba ADH₅₄₇
8
9 mutants retain 5%, 40%, and 25% oxidation activity, respectively. Thus, these
10
11 observations suggest that a combination of residues H262, H266 and H274 is essential
12
13 for catalysis. Surprisingly, the mutation of H266 to Ala results in 1.4-fold increase in
14
15 reduction activity while the mutations of H262 and H274 to Ala lead to 75% and 90%
16
17 reduction activity loss, respectively, which might be caused by overall conformational
18
19 disruption due to the H262A, H266A and H274A substitutions. To our knowledge, we
20
21 first revealed the function the three highly conserved histidine residues in the motif III
22
23 of type III ADH in acetaldehyde reduction using Tba ADH₅₄₇ as a target,
24
25 demonstrating that residue H262 and H274 are involved in catalyzing acetaldehyde
26
27 reduction reaction.
28
29
30
31
32
33
34
35
36
37
38

39 In addition to the motif II, motif II (DxxxH) with a highly conserved His residue
40
41 and an Asp residue is widely present in type III ADHs. The crystal structure of *T.*
42
43 *thioreducens* ADH shows that the conserved residues Asp193 and His197 interact
44
45 with Fe atom plus residues His260 and His272 located in the conserved motif III [15].
46
47 The conserved His residue and an Asp residue in motif II are observed in the crystal
48
49 structures of other type III ADHs. However, the biochemical function of the
50
51 conserved His residue and an Asp residue in motif II has not been clarified. In this
52
53 work, we constructed the D195A and H199A Tba ADH₅₄₇ mutants and investigated
54
55
56
57
58
59
60
61
62
63
64
65

1 their activity for ethanol oxidation and acetaldehyde reduction. Note that residues
2
3 D195 and H199 are analogous to residues D193 and H197 in *T. thioeducens* ADH.
4
5 Our observations demonstrate that the D195A mutant retains slight weak activity for
6
7 ethanol oxidation, and lacks acetaldehyde reduction activity. Furthermore, the H199A
8
9 mutant harbors the compromised activity. Similar to the H262A, H266A and H274A
10
11 substitutions, the D195A and H199A substitutions disrupt the overall structural
12
13 change, which might cause the activity loss of the wild-type Tba ADH₅₄₇. Thus,
14
15 residues D195 and H199 in Tba ADH₅₄₇ are responsible for catalysis and maintaining
16
17 their structural roles.
18
19
20
21
22
23
24

25 **5. Conclusion**

26
27 Herein, we present biochemical characterization of Tba ADH₅₄₇, displaying same
28
29 optimal temperature (75°C) for oxidation and reduction activities. However, this ADH
30
31 has distinct optimal pH for ethanol oxidation and acetaldehyde reduction.
32
33 Interestingly, Tba ADH₅₄₇ can catalyze ethanol oxidation with a divalent ion, which is
34
35 sharply contrasted to the reported ADHs. Furthermore, Tba ADH₅₄₇ exhibits distinct
36
37 divalent ion preference: Mn²⁺ for ethanol oxidation and Fe²⁺ for acetaldehyde
38
39 reduction. Primary alcohols are the favored substrates for oxidation of Tba ADH₅₄₇,
40
41 with 1-butanol yielding the highest efficiency. Furthermore, butyaldehyde is a more
42
43 suitable substrate for reduction of Tba ADH₅₄₇ than acetaldehyde. We revealed that
44
45 residue D195, H199, H262 and H234 in Tba ADH₅₄₇ are involved in catalysis by
46
47 mutational studies. Overall, our findings suggest that Tba ADH₅₄₇ might have a
48
49 biotechnological potential for the biotransformation reaction due to its unique
50
51 biochemical characteristics and high activity at high temperature.
52
53
54
55
56
57
58
59
60
61

Author Contributions

LZ and PO designed experiments; DJ, YL and LW performed experiments; LZ, KD, QL and PO analyzed data; LZ, KD, QL and PO wrote and revised the paper.

Declaration of Competing Interest

All authors declare that there is no conflict of interests regarding the publication of this paper.

Funding

This work was supported by the Provincial Natural Science Foundation of Jiangsu Province (No. BK20191219 and No. BK20180940), High Level Talent Support Program of Yangzhou University and the Academic Leader of Middle and Young People of Yangzhou University Grant.

Figure legends

Fig. 1. Phylogenetic analysis of Tba ADH₆₄₁ and Tba ADH₅₄₇. A. Phylogenetic tree of Tba ADH₆₄₁ and Tba ADH₅₄₇ among bacteria and archaea. B. Phylogenetic tree of Tba ADH₅₄₇ among bacteria and archaea.

Fig. 2. Sequence alignment of type III ADHs from bacteria, archaea and eukaryotes. Three conserved motifs (motif I: GGGS; motif II: DxxxH; motif III: the three highly conserved histidine residues) are boxed. The consensus residues are colored. The residues for engineering Tba ADH₅₄₇ mutants are marked by an asterisk. Tba: *Thermococcus barophilus* (GenBank: ALM74514.1 for ADH₅₄₇ and ALM74601 for ADH₆₄₁); Tan: *Thermococcus* strain AN1, (GenBank: WP_010477684); Tth: *Thermococcus thioreducens* (GenBank: ASJ12775); Tko: *Thermococcus kodakarensis*

(GenBank: WP_011249959 for ADH1 and WP_083755808 for ADH2); Tpa: *Thermococcus paralvinellae* (GenBank: AHF80371); Thy: *Thermococcus hydrothermalis* (GenBank: CAA74334); Tsp: *Thermococcus* sp. (GenBank: AAB63011); Tli: *Thermococcus litoralis* (GenBank: AAB63011); Pfu: *Pyrococcus furiosus* (GenBank: WP_011011187); Zmo: *Zymomonas mobilis* (GenBank: TWE26216); Sce: *Saccharomyces cerevisiae* (GenBank: GAX68104).

Fig. 3. Tba ADH₅₄₇ can catalyze ethanol oxidation and acetaldehyde reduction reaction at high temperature. A. Overexpression and purification of Tba ADH₅₄₇ protein. M: protein marker. B. The optimal reaction temperature of Tba ADH₅₄₇ activity. The ethanol oxidation reactions of Tba ADH₅₄₇ were performed in the presence of Mn²⁺ at different temperatures at pH 8.5. The acetaldehyde reduction reactions of Tba ADH₅₄₇ were performed in the presence of Fe²⁺ at different temperatures at pH 7.0.

Fig. 4. Thermostability of Tba ADH₅₄₇ activity. A. The thermostability for reduction reaction catalyzed by Tba ADH₅₄₇. The heated Tba ADH₅₄₇ at varied temperatures for 30 min was employed to perform the acetaldehyde reduction reactions in the presence of Fe²⁺ at 75°C at pH 7.0. CK: the reaction with the non-heated enzyme. B. The thermostability for oxidation reaction catalyzed by Tba ADH₅₄₇. The heated Tba ADH₅₄₇ at 100°C for varied times was employed to perform the ethanol oxidation reactions in the presence of Mn²⁺ at 75°C at pH 8.5. CK: the reaction with the non-heated enzyme.

Fig. 5. Effects pH and divalent metal ions on Tba ADH₅₄₇ activity. A. Effect of pH on

Tba ADH₅₄₇ activity. The ethanol oxidation reactions of Tba ADH₅₄₇ were performed in the presence of Mn²⁺ at 75°C at varied pHs. The acetaldehyde reduction reactions of Tba ADH₅₄₇ were performed in the presence of Fe²⁺ at 75°C at varied pHs. CK: the reaction without enzyme. B. Effects divalent metal ions on Tba ADH₅₄₇ activity. The ethanol oxidation reactions of Tba ADH₅₄₇ were performed in the presence of varied divalent metal ions at 75°C at pH 8.5. The acetaldehyde reduction reactions of Tba ADH₅₄₇ were performed in the presence of varied divalent metal ions at 75°C at pH 7.0. CK: the reaction without enzyme. EDTA: the reaction with 10 mM EDTA.

Fig. 6. Substrate specificity of the oxidation activity and the reduction activity of Tba ADH₅₄₇. A. Substrate specificity of the oxidation activity of Tba ADH₅₄₇. The oxidation reactions of Tba ADH₅₄₇ were performed in the presence of Mn²⁺ at 75°C at pH 8.5 using varied alcohols as the substrates. B. Substrate specificity of the reduction activity of Tba ADH₅₄₇. The reduction reactions of Tba ADH₅₄₇ were performed in the presence of Fe²⁺ at 75°C at pH 7.0 using varied aldehydes as the substrates.

Fig. 7. The Lineweaver-Burk plots for ethanol oxidation and acetaldehyde reduction of Tba ADH₅₄₇. (A) The Lineweaver-Burk plot for ethanol oxidation of Tba ADH₅₄₇. The oxidation reactions of Tba ADH₅₄₇ were performed in the presence of Mn²⁺ at 75°C at pH 8.5 using ethanol with varied concentrations as the substrates. The Lineweaver-Burk curve was generated by plotting the reciprocal of each ethanol concentration as a horizontal ordinate and the reciprocal of each reaction rate as a vertical coordinate with a linear regression. (B) The Lineweaver-Burk plot for acetaldehyde reduction of Tba ADH₅₄₇. The reduction reactions of Tba ADH₅₄₇ were

performed in the presence of Fe^{2+} at 75°C at pH 7.0 using acetaldehyde with varied concentrations as the substrates. The Lineweaver-Burk curve was generated by plotting the reciprocal of each acetaldehyde concentration as a horizontal ordinate and the reciprocal of each reaction rate as a vertical coordinate with a linear regression.

Fig. 8. Purification and the activity assays of the Tba ADH₅₄₇ mutants. A. Purification of the WT, D195A, H199A, H262A, H266A and H274A mutant Tba ADH₅₄₇ proteins. WT: wild-type; M: protein marker. B. CD analysis of the WT and mutant Tba ADH₅₄₇ proteins. The mean residue ellipticity was recorded with different color line for the WT and mutant Tba ADH₅₄₇ proteins as indicated by monitoring changes in secondary structure with scanning from 200nm to 250 nm. The CD spectra of the WT, D195A, H199A, H262A, H266A and H274A mutant Tba ADH₅₄₇ proteins are colored with black, red, yellow, green, cyan and blue, respectively. C. The activity assays of the Tba ADH₅₄₇ D195A, H199A, H262A, H266A and H274A mutants. The oxidation reactions of the Tba ADH₅₄₇ mutants were performed in the presence of Mn^{2+} at 75°C at pH 8.5 using ethanol as the substrate. The reduction reactions of the Tba ADH₅₄₇ mutants were performed in the presence of Fe^{2+} at 75°C at pH 7.0 using acetaldehyde as the substrate.

References

1. J.A. Littlechild, J.E. Guy, M.N. Isupov, Hyperthermophilic dehydrogenase enzymes, *Biochem. Soc. Trans.* 32 (2004) 255-258.
2. H. Radianingtyas, P.C. Wright, Alcohol dehydrogenases from thermophilic and hyperthermophilic archaea and bacteria, *FEMS Microbiol. Rev.* 27 (2003) 593-616.
3. D.R. Thatcher, The complete amino acid sequence of three alcohol dehydrogenase alleloenzymes (*Adh*-N11, *Adh*-s and *Adh*-UF) from the fruitfly *Drosophila melanogaster*, *Biochem. J.* 187 (1980) 875-886.
4. H. Eklund, B. Nordström, E. Zeppezauer, G. Soederlund, I. Ohlsson, T. Boiwe, B.-O. Soederberg, O. Tapia, C.-I. Brändén, Å. Åkeson, Three-dimensional structure of horse liver alcohol dehydrogenase at 2.4 Å resolution, *J. Mol. Biol.* 102 (1976) 27-59.
5. D. Neale, R. K. Scopes, J. M. Delly, R. E. H. Wettenhall, The two alcohol dehydrogenases of *Zymomonas mobilis*: purification by differential dye ligand chromatography, molecular characterization and physiological roles, *Eur. J. Biochem.* 154 (1986) 119-124.
6. T. Conway, G. W. Sewell, Y. A. Osman, L. O. Ingram, Cloning and sequencing of the alcohol dehydrogenase II gene from *Zymomonas mobilis*, *J. Bacteriol.* 169 (1987) 2591-2597.

- 1 7. V. M. Williamson, C. E. Paquin, Homology of *Saccharomyces cerevisiae* ADH4
2
3 to an iron-activated alcohol dehydrogenase from *Zymomonas mobilis*, Mol. Gen.
4
5 Genet. 209 (1987) 374-381.
6
7
- 8 8. T. Conway, L. O. Ingram, Similarity of *Escherichia coli* propanediol
9
10 oxidoreductase (fucO product) and an unusual alcohol dehydrogenase from
11
12 *Zymomonas mobilis* and *Saccharomyces cerevisiae*, J. Bacteriol. 171 (1989)
13
14 3754-3759.
15
16
17
- 18 9. J. S. Youngleson, W. A. Jones, D. T. Jones, D. R. Woods, Molecular analysis and
19
20 nucleotide sequence of the *adh1* gene encoding an NADPH-dependent butanol
21
22 dehydrogenase in the gram-positive anaerobe *Clostridium acetobutylicum*, Gene
23
24 78 (1989) 355-364.
25
26
27
- 28 10. K. A. Walter, G. N. Bennett, E. T. Papoutsakis, Molecular characterization of two
29
30 *Clostridium acetobutylicum* ATCC 824 butanol dehydrogenase isozyme genes, J.
31
32 Bacteriol. 174 (1992) 7149-7158.
33
34
35
- 36 11. G. E. De Vries, N. Arfman, P. Terpstra, L. Dijkhuizen. Cloning, expression, and
37
38 sequence analysis of the *Bacillus methanolicus* C1 methanol dehydrogenase gene.
39
40 J. Bacteriol. 174 (1992) 5346-5353.
41
42
43
- 44 12. Y. Deng, Z. Wang, S. Gu, C. Ji, K. Ying, Y. Xie, Y. Mao, Cloning and
45
46 characterization of a novel human alcohol dehydrogenase gene (ADHFe1), DNA
47
48 Seq. 13 (2002) 301-306.
49
50
51
- 52 13. C. Montella, L. Bellsollell, R. Pérez-Luque, J. Badía, L. Baldoma, M. Coll, J.
53
54 Aguilar, Crystal structure of an iron-dependent group III dehydrogenase that
55
56
57
58
59
60
61
62
63
64
65

- interconverts L-lactaldehyde and L-1,2-propanediol in *Escherichia coli*. J Bacteriol. 187 (2005) 4957-4966.
14. J. Moon, H. Lee, S. Park, J. Song, M. Park, H. Park, J. Sun, J. Park, B. Kim, J. Kim, Structures of iron-dependent alcohol dehydrogenase 2 from *Zymomonas mobilis* ZM4 with and without NAD⁺ cofactor, J. Mol. Biol. 407 (2011) 413-424.
15. S.B. Larson, J.A. Jones, A. McPherson, The structure of an iron-containing alcohol dehydrogenase from a hyperthermophilic archaeon in two chemical states, Acta Crystallogr. F Struct. Biol. Commun. 75 (2019) 217-226.
16. R. Schwarzenbacher, F. von Delft, J. M. Canaves, L. S. Brinen, X. Dai, A. Deacon, M. et al, Crystal structure of an iron-containing 1,3-propanediol dehydrogenase (TM0920) from *Thermotoga maritime* at 1.3 Å resolution. Proteins, 54 (2004) 174–177.
17. K. Ma, F.T. Robb, M.W. Adams, Purification and characterization of NADP-specific alcohol dehydrogenase and glutamate dehydrogenase from the hyperthermophilic archaeon *Thermococcus litoralis*, Appl. Environ. Microbiol. 60 (1994) 562-568.
18. K. Ma, H.J. Loessner, J. Heider, M.K. Johnson, M.W.W. Adams, Effects of elemental sulfur on the metabolism of the deep-sea hyperthermophilic archaeon *Thermococcus* strain ES-1: characterization of a sulfur-regulated, non-heme iron alcohol dehydrogenase, J. Bacteriol. 177 (1995) 4748-4756.
19. X. Ying, A.M. Grunden, L. Nie, M.W. Adams, K. Ma, Molecular characterization of the recombinant iron-containing alcohol dehydrogenase from the

- 1 hyperthermophilic Archaeon, *Thermococcus* strain ES1, Extremophiles, 13 (2009)
2
3 299-311.
4
5
6 20. D. Li, K.J. Stevenson, Purification and sequence analysis of a novel
7
8 NADP(H)-dependent type III alcohol dehydrogenase from *Thermococcus* strain
9 AN1, J. Bacteriol. 179 (1997) 4433-4437.
10
11
12
13
14 21. E. Antoine, J.L. Rolland, J.P. Raffin, J. Dietrich, Cloning and over-expression in
15
16 *Escherichia coli* of the gene encoding NADPH group III alcohol dehydrogenase
17 from *Thermococcus hydrothermalis*. Characterization and comparison of the
18 native and the recombinant enzymes, Eur. J. Biochem. 264 (1999) 880-889.
19
20
21
22
23 22. Ma K, Adams MWW (1999) An unusual oxygen-sensitive, iron and
24
25 zinc-containing alcohol dehydrogenase from the hyperthermophilic archaeon
26
27 *Pyrococcus furiosus*. J Bacteriol 181:1163-1170.
28
29
30
31
32
33 23. V.T. Marteinsson, J.L. Birrien, A.L. Reysenbach, M. Vernet, D. Marie, A.
34
35 Gambacorta, P Messner, U.B. Sleytr, D. Prieur, *Thermococcus barophilus* sp.
36 nov., a new barophilic and hyperthermophilic archaeon isolated under high
37
38 hydrostatic pressure from a deep-sea hydrothermal vent, Int.J. Syst. Bacteriol. 49
39
40
41
42
43
44
45 (1999) 351-359.
46
47
48 24. P. Oger, T.G. Sokolova, D.A. Kozhevnikova, E.A. Taranov, P. Vannier, H.S. Lee,
49
50 K.K. Kwon, S.G. Kang, J.H. Lee, E.A. Bonch-Osmolovskaya, A. V. Lebedinsky,
51
52 Complete genome sequence of the hyperthermophilic and piezophilic archaeon
53
54 *Thermococcus barophilus* Ch5, capable of growth at the expense of
55
56
57
58
59
60
61
62
63
64
65 hydrogenogenesis from carbon monoxide and formate, Genome Announc. 4

- (2016).
25. K.M. Kwon, S.G. Kang, T.G. Sokolova, S.S. Cho, Y.J. Kim, C.H. Kim, ST. Kwon, Characterization of a family B DNA polymerase from *Thermococcus barophilus* Ch5 and its application for long and accurate PCR, *Enzyme Microb. Technol.* 86 (2016) 117-126.
26. Y. Wang, L. Zhang, X. Zhu, Y. Li, H. Shi, P. Oger, Z. Yang, Biochemical characterization of a thermostable endonuclease V from the hyperthermophilic euryarchaeon *Thermococcus barophilus* Ch5, *Int. J. Biol. Macromol.* 117 (2018) 17-24.
27. H. Shi, Q. Gan, D. Jiang, Y. Wu, Y. Yin, H. Hou, H. Chen, Y. Xu, L. Miao, Z. Yang, P. Oger, Biochemical characterization and mutational studies of a thermostable uracil DNA glycosylase from the hyperthermophilic euryarchaeon *Thermococcus barophilus* Ch5, *Int. J. Biol. Macromol.* 134 (2019) 846-855.
28. H. Shi, Y. Huang, Q. Gan, M. Rui, H. Chen, C. Tu, Z. Yang, P. Oger, L. Zhang, Biochemical characterization of a thermostable DNA ligase from the hyperthermophilic euryarchaeon *Thermococcus barophilus* Ch5, *Appl. Microbiol. Biotechnol.* 103 (2019) 3795-3806.
29. Q. Gan, M. He, H. Shi, Z. Yang, P. Oger, L. Ran, L. Zhang, Characterization of a Family IV uracil DNA glycosylase from the hyperthermophilic euryarchaeon *Thermococcus barophilus* Ch5, *Int. J. Biol. Macromol.* 146 (2020) 475-481.
30. M. Gouy, S. Guindon, O. Gascuel, SeaView version 4: a multiplatform graphical user interface for sequence alignment and phylogenetic tree building, *Mol. Biol.*

- Evol. 27 (2010) 221-224.
31. X. Ying, Y. Wang, H. Badiei, V. Karanassios, K. Ma, Purification and characterization of an iron-containing alcohol dehydrogenase in extremely thermophilic bacterium *Thermotoga hypogea*, Arch. Microbiol. 187 (2007) 499-510.
32. S. Elleuche, K. Fodor, B. Klippel, H.A. von der, M. Wilmanns, G. Antranikian, Structural and biochemical characterisation of a NAD(+)-dependent alcohol dehydrogenase from *Oenococcus oeni* as a new model molecule for industrial biotechnology applications, Appl. Microbiol. Biotechnol. 97 (2013) 8963-8975.
33. G. Sulzenbacher, K. Alvarez, R.H. Van Den Heuvel, C. Versluis, S. Spinelli, V. Campanacci, et al. Crystal structure of *E. coli* alcohol dehydrogenase YqhD: evidence of a covalently modified NADP coenzyme, J. Mol. Biol. 342 (2004) 489-502.
34. S. Elleuche, K. Fodor, H.A. von der, B. Klippel, M. Wilmanns, G. Antranikian, Group III alcohol dehydrogenase from *Pectobacterium atrosepticum*: insights into enzymatic activity and organization of the metal ion-containing region, Appl. Microbiol. Biotechnol. 98 (2014) 4041-4051.
35. J. Extance, S.J. Crennell, K. Eley, R. Cripps, D.W. Hough, M.J. Danson, Structure of a bifunctional alcohol dehydrogenase involved in bioethanol generation in *Geobacillus thermoglucosidasius*, Acta Crystallogr. D Biol. Crystallogr. 69 (2013) 2104-2115.
36. R. Machielsen, A.R. Uria, S.W. Kengen, J. van der Oost, Production and

- 1 characterization of a thermostable alcohol dehydrogenase that belongs to the
2
3
4 aldo-keto reductase superfamily, Appl. Environ. Microbiol. 72 (2006) 233-238.
5
6 37. T.N. Stekhanova, A.V. Mardanov, E.Y. Bezsudnova, V.M. Gumerov, N.V. Ravin,
7
8 K.G. Skryabin, V.O. Popov, Characterization of a thermostable short-chain
9
10 alcohol dehydrogenase from the hyperthermophilic archaeon *Thermococcus*
11
12 *sibiricus*, Appl. Environ. Microbiol. 76 (2010) 4096-4098.
13
14
15
16
17 38. M. Hess, G. Antranikian, Archaeal alcohol dehydrogenase active at increased
18
19 temperatures and in the presence of organic solvents, Appl. Microbiol. Biotechnol.
20
21
22 77 (2008) 1003-1013.
23
24
25
26 39. E.N. Marino-Marmolejo, A.D. León-Rodríguez, A.P.B. de la Rosa, L. Santos,
27
28 Heterologous expression and characterization of an alcohol dehydrogenase from
29
30 the archaeon *Thermoplasma acidophilum*, Mol. Biotechnol. 42 (2009) 61-67.
31
32
33
34 40. M.R. Fernandez, J.A. Biosca, A. Norin, H. JoÅrnvall, X. PareÂs, Class III alcohol
35
36 dehydrogenase from *Saccharomyces cerevisiae*: Structural and enzymatic features
37
38 differ toward the human/mammalian forms in a manner consistent with functional
39
40 needs in formaldehyde detoxication, FEBS Lett. 370 (1995) 23-26.
41
42
43
44 41. N. Obradors, E. Cabiscol, J. Aguilar, J. Ros, Site-directed mutagenesis studies of
45
46 the metal-binding center of the iron-dependent propanediol oxidoreductase from
47
48 *Escherichia coli*, Eur. J. Biochem. 258 (1998) 207-213.
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1 Sequences of the oligonucleotides used to clone the Tba ADH₅₄₇ gene and construct its mutants

Name	Sequence (5'-3')
Tba ADH ₅₄₇ F	GGGAATTCCATATGCAGTTCTTCAGCTTAAAGAC
Tba ADH ₅₄₇ R	CCGCTCGAGATCATAGAAGGCTCTCTTATAC
D195A F	AAGGAACAGCGGACTGGCTGTTCTGGTGCAC
D195A R	GCCAGTCCGCTGTTCCCTTGCAACTTCTTTTG
H199A F	GACTGGATGTTCTGGTGGCCGGAATTGAAGCT
H199A R	GCCACCAGAACATCCAGTCCGCTGTTCCCTTGC
H262A F	CCCGTTTAGGCTTGTGCGCCAGCTTAAGCCAT
H262A R	GCGCACAAGCCTAAACGGGCATTAAGAAAAGC
H266A F	TGTGCCACAGCTTAAGCGCTAAAGCGGCTTGG
H266A R	GCGCTTAAGCTGTGGCACAAGCCTAAACGGGC
H274A F	CGGCTTGGAATTGCCCCTGCCGGCTTGTTAAAT
H274A R	GCAGGGGCAATCCAAGCCGCTTTATGGCTTAA

The italic bases represent restriction sites.

The substitution bases are underlined.

Table 2 Comparisons of biochemical characteristics of iron-containing ADH members from hydrothermophiles.

Organism		T/°C	pH	K_m	K_{cat}	K_{cat}/K_m	Reference
		optimum	optimum	(mM)	(s ⁻¹)	(s ⁻¹ M ⁻¹)	
<i>T. barophilus</i>	Oxidation ^a	75	8.5	92	0.27	2.9	This work
	Reduction ^b	75	7.0	34.5	69	2.0 x 10 ³	
<i>Thermococcus</i> strain AN1	Oxidation	85	6.8-7.0	ND	ND	ND	[20]
	Reduction	ND	7.0	0.12	ND	ND	
<i>T. litoralis</i>	Oxidation	85	8.8	11	26	2.3 x 10 ³	[17]
	Reduction	ND	ND	0.4	ND	ND	
<i>T. hydrothermalis</i>	Oxidation	80	10.5	ND	ND	ND	[21]
	Reduction	ND	7.5	ND	ND	ND	
<i>Thermococcus</i> strain ES-1	Oxidation	>95	8.8-10.4	8.0	48	6.0 x 10 ³	[18]
	Reduction	ND	ND	0.25	19	7.6 x 10 ⁴	
<i>P. furiosus</i>	Oxidation	>95	9.4-10.2	29.4	19	6.5 x 10 ²	[22]
	Reduction	ND	ND	0.17	5.5	3.2 x 10 ⁴	
<i>T. hypogea</i>	Oxidation	>95	11.0	9.7	14	1.5 x 10 ³	[31]
	Reduction	ND	8.0	3.1	15	5.0 x 10 ³	

a: Ethanol oxidation reaction; b: Acetaldehyde reduction; ND: Not determined

Tba_ADH547	MFFSLKTRIVL..GEGSLSYIKSVAKKHSR.....VLIFSSSKSMRVHGFNLSTNNVVEDANAEEVAITGVPAE..EVEYVE	74
Tan_ADH	MFQLKTRIEE..GEGSLKALSREVRGYER.....VLILSSSSMKRHGFLGEVEDCVREAGAEVFSVTNLPAE..GVETVE	72
Tth_ADH	MFWLKTRIEE..GEGSLSRLSREVKGHER.....VLILASGSMKRHGFLSEAEDYVKEAGAEVFSIAGLPAE..SVEVIE	72
Tko_ADH1	MFWLKTRISE..GEDSLESLEKDEAKDHER.....VLILASGSMKRHGFLSEAEDYVREAGAEVMSISGLPAE..SVEVIE	72
Tba_ADH641	MLWESQIPINQIFEMRCKTTNYFGVGAINKFYDIAKDLKE..RGITKVILVTGRSSYKKCGAWDVVKPALEENGIEYVHYDKIGPN..TVDMDV	91
Tpa_ADH	MLWESQIPINQIFELRCRTIDYFGVGAINKFYDIAKDLKENRGISRVILVTGKSSYKKCGAWDVVKPALEENGIEYVHYDKVGP..TVDMDV	92
Thy_ADH	MVWESHVSINQVFEMRCKTTNYFGLCAIHKFNDIVRELKG..KGVDKVILVTGXSSYKKCGAWDVVRPALEENGVEYVHYDKVGAN..TVDMD	91
Tsp_ADH	MVWESHVPINQVFELRCRTIDYFGLCAIHKFNGFVRELKG..KGVDRVILVTGXSSYKKCGAWDVVRPALEENGVEYVHYDKVGAN..TVDMD	91
Tli_ADH	MLWESRLPINQVFELRCRTIDYFGVGAINKFYDIAKDLKENRDIDKVLVLTGKSSYKKCGAWDVVKPALEKYGYIEYVHYDKVGP..TVDMD	92
Tko_ADH2	MAGQSFDLNFWHPSFFLPFLQNLCDSSNKNFGRKRLPELLER..HGLSG....KALMVTGRSSTKKTGLLYRVQELLKKAGVESIVFDRVIPN..ISTHVD	94
Pfu_ADH	MFEISIIYLP..TEIVFGPGK...LEMLPKLVKK..HGLSG....KALIVTGRSSTKETGVLYRVQELLKQAGVESIVFDKIIPN..ISTHVD	80
Zmo_ADH	MASST..FYIPFVNEMEGGSLEKAIKDLNGSGF....KNALIVSDAFMNKSGVVKQVADLLKAQGINSAVYDGVMPN..TAVTAVL	78
Sce_ADH	MSSVTGFYIPPIISFFGEGALEETADYIKNKDY....KALIVTDPGIAAIGLSGRVQKMLEERGLNVAIYDKTQPN..NIANVT	79
Consensus	an	
Tba_ADH547	SIMPKVR.EFQPDLLVALGGGSGVIIDVAVKAVKFYDAPELKFEFVAFISRFKPKFIPKLKTPLIAIPSTISGAGSEVSAASVLKKGDIKYN..LVSFEIAP	171
Tan_ADH	ELLPKLREEFQPDLLVALGGGSGVIIDAKALKVFYDAPEVKFEEIAFIDRFSPKPFVPLKLTPLIAIPSTISGAGSEVSASSVLKKGVEKYN..IVSLEIAP	170
Tth_ADH	EFLPKVR.EFGPDLLVAMGGGSGVIIDTKALKVFYDAPELNFGEIAFIDRFSPKPFVPLKLTLLIAIPSTISGAGSEVSGASVLKKGVEKYN..IVTPEIAP	169
Tko_ADH1	EFLPKVR.EFGPDLLIAMGGGSGVIIDTKALKVFYDAPELVFEEIAFINRFSRPFVPLKLTPLIAIPSTISGAGSEVSAASVLKKNGIKYN..IVTPEIAP	169
Tba_ADH641	EAAQMGR.EFGAQAVIGIGGGSPIDISAKGVAILLEYTDKDARQLYELK.....FTPKKAKPVIAINTTHGTGTEVDRFAVASILEKEYKPAIAYDCIYF	184
Tpa_ADH	EAAQMGR.EFGAQAVIGIGGGSPIDISAKGVAILLEYTDKTARELYKLQ.....FSRPAKPIIAINTTHGTGTEVDRFAVASILEEYKPAIAYDCIYF	185
Thy_ADH	EAAEMGR.EFGAQAVIGIGGGSPIDISAKGVAILLEYPDKTARDLYEFR.....FTPVKAKPIIAINTTHGTGTEVDRFAVASIPEKEYKPAIAYDCIYF	184
Tsp_ADH	EAAEMGR.EFGAQAVIGIGGGSPIDISAKGVAILLEYPDKTARDLYEFR.....FTPVKAKPVIAINTTHGAGTEVDRFAVATPEKEYKPAIAYDCIYF	184
Tli_ADH	EAKEMGV.EFGAQAVIGIGGGSPIDISAKGVAILLEYPDKTGKDLYEFK.....FKPTKAKPIIAINTTHGTGTEVDRFAVASILEKEYKPAIAYDFIYF	185
Tko_ADH2	EAAELAR.RENVDFVVGGLGGGSAIDISAKAIAMTAKSGGKYWDYVPAVGGG....KKPSGALPIVAIPTTHGTGTEADPYAVITNPETKEKQGIGYDVLFF	189
Pfu_ADH	EGAEIAR.KENVSFVVGGLGGGSAIDISAKAIAMTAAAGGKYWDYVPAVGGG....KKPTGALPIVAIPTTHGTGTEADPYAVITNPETKEKQGIGYDVLFF	175
Zmo_ADH	EGLKILK.DNNSDFVISLGGGSPHICAKAIALVATNGGEVVDYEGIDK.....SKKPALPLMSINTTACTASEMTRFCIITDEVHRVKMAIVDRHVT	170
Sce_ADH	AGLKVLK.EQNSEIVVSIIGGGSAHINAKAIALLATNGGEIGDYEGVNQ.....SKKAALPLFAINTTACTASEMTRFTIISNEKKIKMAIIDNNVT	171
Consensus	gggs d k i t g e III p	
Tba_ADH547	EFAILERLRLPRTMPKEVARNSGILVVLVHGIEAYTTTAATFSDAMAIKAIKLVYKWLPLSVQG..DEKARENVHYAATMAGIAFLNARLGLCHSLSHKAA	269
Tan_ADH	EVAILERLRLPRTMPGEVARNSGILVVLVHGIEAYTTKAANFSDTMAINAIAKTVYFWLPLSVKG..DEEARARVHYAATMAGIAFLNARLGLCHAMSHKAA	268
Tth_ADH	DVAILERLRLPRTMPPEVARNSGILVVLVHGIEAYTTKVASFSDAMAIKAIKTVYFWLPLSVKG..DEEARARVHYAATMAGIAFLNARLGLCHAMSHKAA	267
Tko_ADH1	EVAILERLRLPRTMPREARNSGILVVLVHGIEAYTTKVAGFSDAMAIRAIAKTVFEWLPKSVEG..DPEARARMHYAATMAGIAFLNARLGLAHAMSHKAA	267
Tba_ADH641	LYAIDELALTVKLPADQTRYVTIDALNHVNEAATTKVASEYSILLAQETARLIFKYIPEALNHPDNLQARYYLLYASAIAGISFDNGLLHFTHALEHPLS	284
Tpa_ADH	LYSIDELTLMTKLPADQTRYVTIDALNHVNEAATTKVASEYSILLAKETARLIFKYIPEALIHDPNLQAKYLLYASAIAGISFDNGLLHFTHALEHPLS	285
Thy_ADH	LYAIDELALTTKLPPEQTLVYTIIDALNHVTEAATTKVANEYSILLAKEAARLIFTYIPEALNHPDNLQARYALLYASAIAGISFDNGLLHFTHALEHPLS	284
Tsp_ADH	LYAIDELSLTVKLSPEQTLVYTIIDALNHVTEAATTKLANEYSILLAKEAARLIFTYIPEALKHPDNLQARYALLYASAIAGISFDNGLLHFTHALEHPLS	284
Tli_ADH	LYAIDELALMTKLPANQTRYVTIDALNHVNEAATTKATNEYSILLAQETVRLISKYIPEALAHPENLQARYYLLYASAIAGISFDNGLLHFTHALEHPLS	285
Tko_ADH2	KFSIVDEEVMVTLPCNQTVYTSMAFYHSIEAFLNVRACFYSVDVLDLSMRVVYIPIAYENLQDLARTQLAWASTEAGITETLTGVIANHALEHGLS	289
Pfu_ADH	KFSIVDEELMLTLPKDQTVYTSMAFYHSIEAFLNVRANFYSVDVLDLSMRVVIYIPLAYENLRNLEARTQLAWASTEAGITETVTGVVANHALEHGLS	275
Zmo_ADH	MVSVNDELMLVMGMPKGLTAATGMALTAFAFAYSSTAATEITDACLKAASMIANIKTACDNGKDMFAREAMAYAQFLAGMAFNNASLGYVHAMAHQLG	270
Sce_ADH	AVAVNDELSTMFGLPALTAATGLTALTACIEAYVSTASNEITDACLKGIDLINESTVAAYKDGKDKKRTDMCYEYLAGMAFNNASLGYVHALAHQLG	271
Consensus	dp d h ea p a l a a ag h h	
Tba_ADH547	WIA...EHGLLNAIFLEYVMEFN..MRSEYARKRYAEIARELG...FNTAQELIEVIREFNEMLGVPK..SELVSEEEFLSKLDEMSEK..AYHDPLINF	358
Tan_ADH	WIG...EHGLLNAIFLEYVMEFNA.ERSEYARKRYAEIARELG...FNTAKDLIEVVKELNEMLGVPG..SELVDEEAFVSKVEEMANK..AYRDGLIPF	358
Tth_ADH	WIG...EHGLLNAVFLLEYVMEFNA.SKSDYARRRYAEIARELG...FQTAKDLIEVVKELNEMLGVPK..LGELVDEETFASKVEEMAER..TYHDGLIAF	357
Tko_ADH1	WVG...EHGLLNAIFLEYVMEFNA.ERSEYARKRYAEIARELG...LKSAAELIKAVRELNERLGVPK..SELVDEETFVERVEEMAER..AYRDGLIAF	357
Tba_ADH641	AVKPELEHGLGLAMLLAEVIRKHIYPATAPILAIEYRPLVPEAKG.VPG.EALKVAKAVEKWLFSIGITEKIDTVGFGEDDVKKLAEAMTTPSLDLLLSL	382
Tpa_ADH	AVKPDLEHGLGLAILLEAVIRKHIYPATARILAEYRPLVPEVKG.VPG.EAELVAKKVEEWLFAIGITQKIDTVGFSEDDVKRLTELAMTTPSLNLLLSL	383
Thy_ADH	AVKPDLEHGLGLAMLLAEVIRQIYPATAKILAIEYRPLVPEAKG.VPG.EVELVARRVEEWLFSIGITEKIDTVGFTEGVDVKLTQLAMTTPSLDLLLSM	382
Tsp_ADH	AVKPDFEHGIGLAMLLAEVIRHIYPATAKILAIEYRPLVPEAKG.VPG.EVELVARRVEEWLFSIGITEKIDTVGFSETDVNRLTELAMKTTPSLNLLLSM	382
Tli_ADH	AVKPDLEHGLGLAMLLAEVVKHIYPATAEILAIEYRPIAPEAKG.YPG.EAEFVAKRIEQWLFISIGIKEKIDTVGFGEDDIDKLTKLAMETPGLDGLLAM	383
Tko_ADH2	GFYPEVEHGLGLCILGELYFEYILDHAYERLAIVGREVFGVYES.NDRKAAELAVKKLRDFQSLFGVNKKRELGVKEEDIPKMAETAYR..MMKGVIDV	386
Pfu_ADH	GFYPEVEHGLGLCILGELYFEYILDYAYEKLAIVGREVFGVYEP.NDRKAAELAIKKLRDFQSLFGVNKKRELGVKEEDIPEMARTAYR..MMKPVIEA	372
Zmo_ADH	GYV.NLEHGVCAVALLHVLAYNASVAGRLKDVGVAMGLDIANLGDKEGAETIQAVRDLAASIGIPANITELGAKKEDVPLADHALK....DACALT	365
Sce_ADH	GFY.HLEHGVCAVALLHVLQEANMQCPKAKKRLDEIALHFGAS....QEDPEETIKALHVLNRTMNIPIRNIKELGVKTEDFEILAEHAMH....DACHLT	362
Consensus	phg p 1	
Tba_ADH547	NEVEPSIEEIKELYKRAFYD...	378
Tan_ADH	NEVEPKPEEIRELYLRAFYGR..	379
Tth_ADH	NEVEPKPEEIKELYLKAYRGE..	378
Tko_ADH1	NEVEPKPEEIRELYLKALNAL..	378
Tba_ADH641	AEIPATKETVEAIYRDSLYPLSE	405
Tpa_ADH	AEVEATKETIEAIYRDSLYPLSK	406
Thy_ADH	AEVEATKERIAAIYRDSLYPIGR	405
Tsp_ADH	AEGEATRERIAAIYRDSLYPISR	405
Tli_ADH	AEVEANEDVIRAIYRDSLYPLSE	406
Tko_ADH2	TEGNLSVEDLEEYIRRAY.....	404
Pfu_ADH	TEGDLKVEDLEEYIRRAY.....	390
Zmo_ADH	NRQGDQKEVEELFLSAF.....	383
Sce_ADH	NEVQFTKEQVVAIIKKAYEY...	382
Consensus	p	

A Figure

B

AFigure

[Click here to
access/download;Fig](#)

B

A

B

A

B

A

[Click here to
access/download;Figu](#)

B