



**Revised stratigraphic framework for the lower
Anti-Atlas Supergroup based on U–Pb geochronology of
magmatic and detrital zircons (Zenaga and Bou
Azzer-El Graara inliers, Anti-Atlas Belt, Morocco)**

Abdelhak Ait Lahna, Nasrddine Youbi, Colombo Celso Gaeta Tassinari,
Miguel Angelo Stipp Basei, Richard Ernst, Latifa Chaib, Abdelhafed Barzouk,
João Mata, Andreas Gärtner, Hassan Admou, et al.

► **To cite this version:**

Abdelhak Ait Lahna, Nasrddine Youbi, Colombo Celso Gaeta Tassinari, Miguel Angelo Stipp Basei, Richard Ernst, et al.. Revised stratigraphic framework for the lower Anti-Atlas Supergroup based on U–Pb geochronology of magmatic and detrital zircons (Zenaga and Bou Azzer-El Graara inliers, Anti-Atlas Belt, Morocco). *Journal of African Earth Sciences*, 2020, 171, pp.103946. 10.1016/j.jafrearsci.2020.103946 . hal-03096023

HAL Id: hal-03096023

<https://hal.science/hal-03096023>

Submitted on 6 Jan 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Journal Pre-proof

Revised stratigraphic framework for the lower Anti-Atlas supergroup based on U–Pb geochronology of magmatic and detrital zircons (Zenaga and Bou Azzer-El Graara inliers, Anti-Atlas Belt, Morocco)

Abdelhak Ait Lahna, Nasrddine Youbi, Colombo Celso Gaeta Tassinari, Miguel Angelo Stipp Basei, Richard E. Ernst, Latifa Chaib, Abdelhafed Barzouk, João Mata, Andreas Gärtner, Hassan Admou, Moulay Ahmed Boumehdi, Ulf Söderlund, Mohamed Khalil Bensalah, Jean-Louis Bodinier, Lhou Maacha, Andrey Bekker

PII: S1464-343X(20)30197-7

DOI: <https://doi.org/10.1016/j.jafrearsci.2020.103946>

Reference: AES 103946

To appear in: *Journal of African Earth Sciences*

Received Date: 11 December 2019

Revised Date: 28 June 2020

Accepted Date: 30 June 2020

Please cite this article as: Lahna, A.A., Youbi, N., Gaeta Tassinari, C.C., Stipp Basei, M.A., Ernst, R.E., Chaib, L., Barzouk, A., Mata, Joã., Gärtner, A., Admou, H., Boumehdi, M.A., Söderlund, U., Bensalah, M.K., Bodinier, J.-L., Maacha, L., Bekker, A., Revised stratigraphic framework for the lower Anti-Atlas supergroup based on U–Pb geochronology of magmatic and detrital zircons (Zenaga and Bou Azzer-El Graara inliers, Anti-Atlas Belt, Morocco), *Journal of African Earth Sciences* (2020), doi: <https://doi.org/10.1016/j.jafrearsci.2020.103946>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Ltd.



**Revised stratigraphic framework for the lower Anti-Atlas Supergroup
based on U-Pb geochronology of magmatic and detrital zircons (Zenaga and Bou
Azzer-El Graara inliers, Anti-Atlas Belt, Morocco)**

Abdelhak Ait Lahna^{1*}, Nasrddine Youbi^{1,2}, Colombo Celso Gaeta Tassinari³, Miguel Angelo
Stipp Basei⁴, Richard E. Ernst^{5,6}, Latifa Chaib¹, Abdelhafed Barzouk¹, João Mata², Andreas
Gärtner⁷, Hassan Admou¹, Moulay Ahmed Boumehdi^{1,2}, Ulf Söderlund^{8,9}, Mohamed Khalil
Bensalah^{1,2}, Jean-Louis Bodinier^{10,11}, Lhou Maacha¹², Andrey Bekker^{13,14}

¹Department of Geology, Faculty of Sciences-Semlalia, Cadi Ayyad University, Prince Moulay Abdellah
Boulevard, P.O. Box 2390, Marrakech, Morocco: aitlahna.abdelhak@gmail.com;
youbi@uca.ac.ma; latifa_918@yahoo.fr; barzoukabdelhafed@gmail.com; admou@uca.ac.ma;
boumehdi@uca.ac.ma; bensalah@uca.ac.ma.

²Instituto Dom Luiz, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal:
jmata@fc.ul.pt

³Centro de Pesquisas Geocronológicas (CPGeo), Instituto de Geociências (IG), Universidade de São Paulo-
USP, Caixa Postal 11348, CEP 05422-970, São Paulo, SP, Brazil: ccgtassi@usp.br.

⁴Instituto de Geociências (IG), Departamento de Mineralogia e Geotectônica (GMG), Universidade de São
Paulo-USP, Rua do Lago, 562 Cidade Universitária 05508-080 São Paulo, SP, Brazil:
baseimas@usp.br

⁵Department of Earth Sciences, Carleton University, 1125 Colonel By Drive, Ottawa, Canada K1S 5B6:
Richard.Ernst@carleton.ca

⁶Faculty of Geology and Geography, Tomsk State University, 36 Lenin Ave, Tomsk 634050, Russia:
Richard.Ernst@ernstgeosciences.com

⁷Senckenberg Naturhistorische Sammlungen Dresden, Museum für Mineralogie und Geologie, Sektion
 Geochronologie, GeoPlasma Lab, Königsbrücker Landstraße 159, 01109 Dresden, Germany:
Andreas.Gaertner@senckenberg.de

⁸Department of Geology, Lund University, Sölvegatan 12, SE-223 62 Lund, Sweden:
Ulf.Soderlund@geol.lu.se

⁹The Swedish Museum of Natural History, SE-114 18 Stockholm, Sweden

¹⁰Mohammed VI Polytechnic University, Geology and Sustainable Mining Department, Hay Moulay Rachid,
 Ben Guerir, 43150, Morocco: JeanLouis.Bodinier@um6p.ma

¹¹Géosciences Montpellier, Université de Montpellier 2, CNRS, Cc 60, Place Eugène Bataillon, 34095
 Montpellier Cedex 05, France: bodin@gm.univ-montp2.fr

¹²Managem Group, Twin Center, Tour A, Angle Boulevards Zerkouni & Al Massira Al Khadra, P.O. Box
 5199, Casablanca, Morocco: L.maacha@managemgroup.com

¹³Department of Earth and Planetary Sciences, University of California, Riverside, CA 92521, USA:
andrey.bekker@ucr.edu

¹⁴Department of Geology, University of Johannesburg, Auckland Park 2006, South Africa

*Corresponding author: aitlahna.abdelhak@gmail.com

Revised for Journal of African Earth Sciences

Running Title “Revised stratigraphic framework for the lower Anti-Atlas Supergroup” by Ait Lahna
 et al., 2020.

23959 words

18 figures & 2 tables

10 supplementary data tables and figures

Abstract

U-Pb geochronology of magmatic and detrital zircons (Zenaga and Bou Azzer-El Graara inliers, Anti-Atlas Belt, Morocco) and a reassessment of the published constraints suggest a revised stratigraphic framework for the lower Anti-Atlas Supergroup. Five, major unconformity-bounded lithostratigraphic packages are here distinguished: the two lower units of Paleoproterozoic age are named the Tasserda-Taghatine Group (2030-1706 Ma) and the Oumoula (Mimount) Formation (ca. 1745-1650 Ma); the third unit of Paleoproterozoic to Neoproterozoic age (ca. 1650 to > 883 Ma) is the Tizi n'Taghatine Group; the fourth and fifth units of Neoproterozoic age are the ca. 883 Ma Tachdamt and the ca. 700 Ma Bleida formations. Implications of this revised stratigraphic framework include: 1) the Tasserda-Taghatine Group might be linked to the post-orogenic collapse after the Eburnean Orogeny; 2) the Tizi n'Taghatine Group might be ca. 1.1 Ga in age based on proposed correlation with the Taoudeni Basin succession in Mauritania; 3) the Bleida Formation likely reflects deposition in the foreland basin at the early stage of the Pan-African Orogeny; 4) the Oumoula (Mimount) Formation, Tizi n'Taghatine Group, and Tachdamt Formation potentially record extensional events within the Nuna/Columbia and Rodinia supercontinents; 5) the provenance of the lower Anti-Atlas Supergroup (based on our new detrital zircon dating) is mainly from the West African craton along with possible contributions from other cratons such as Amazonia and the Sahara Metacraton; 6) the flood basalt sequence of the Tachdamt Formation likely belongs to the ca. 885-883 Ma intraplate Iguerda-Taïfast Large Igneous Province (LIP) event defined by previously dated dykes in the Iguerda and Taïfast inliers; and 7) the 1650 Ma Zenaga LIP can be potentially linked with LIP magmatism in Baltica and Laurentia.

Key Words: Morocco; Anti-Atlas Belt; Proterozoic; lower Anti-Atlas Supergroup; U-Pb geochronology

Highlights

► The Taghdout mafic sill of the Ifzwane Suite yielded an age of 1676 ± 37 Ma

- 77 ► The maximum depositional age of the lower Anti-Atlas Supergroup ranges from 2049 Ma
78 (bottom) to 700 Ma (top) and the provenance is mainly from the WAC
- 79 ► A new 5-part lithostratigraphic framework for the Proterozoic lower Anti-Atlas Supergroup of
80 the Anti-Atlas Belt is proposed
- 81 ► The 1650-1640 Ma Zenaga and 885-883 Ma Iguerda-Taïfast LIPs provide important stratigraphic
82 constraints
- 83 ► Correlation between the lower Anti-Atlas Supergroup and the Taoudeni Basin succession in
84 Mauritania
- 85

1. Introduction

Substantial advances in the understanding of the geological evolution of several major Precambrian accretionary orogens of North America and Baltica (Hoffman, 1988; Chacko et al., 2000; Corrigan et al., 2005, 2009; Pehrsson et al., 2013; Bogdanova et al., 2008; 2016) have been made in recent years. However, there is still a fundamental gap in the knowledge of Precambrian orogenic belts in Africa. Much of the uncertainty about the accretionary history of the Precambrian African orogens stems from the limited database of reliable radiometric ages on structurally important units, combined with a lack of detailed studies using modern kinematic analysis in these deformed regions as well as of cutting-edge research using sedimentology and sequence stratigraphy.

One of the crucial Neoproterozoic African orogens is exposed in the Anti-Atlas Belt of Morocco located in the northern part of the West African Craton (WAC, Fig. 1). Extensive outcrops of the Neoproterozoic Pan-African Belt and underlying Paleoproterozoic foreland (Michard et al., 2008; Gasquet et al., 2008; Soulaïmani et al., 2018 and references therein) are well exposed and have been studied for a long time (e.g. Choubert, 1963). Traditionally, the Precambrian of the Anti-Atlas Belt has been divided into three broad lithosomes (e.g. Choubert, 1963): the P I (Archean to Paleoproterozoic), P II (early and middle Neoproterozoic), and P III (late Neoproterozoic) (Fig. 2). This lithostratigraphic subdivision has recently been challenged (Walsh et al., 2002; 2012; Thomas et al., 2002; 2004; Michard et al., 2008; Gasquet et al., 2008; Youbi et al., 2013). In the conventional interpretation, the oldest, Paleoproterozoic rocks (PI) of the Anti-Atlas Belt are subdivided into a series of complexes (e.g. the Zenaga and Kerdous complexes in the Siroua and Kerdous basement inliers, respectively). The rocks of the P II lithosome deposited on this basement have collectively been called the *Anti-Atlas Supergroup*, within which there are four groups of volcano-sedimentary units (e.g. Lkest-Taghdout, Bou Azzer, Iriri, and Saghro groups) and they are conventionally inferred to be Neoproterozoic in age. Various intrusive suites are also recognized (e.g. Ifzwane and Toudma Suites). These volcano-sedimentary units have been related to the earliest passive margin, oceanic, and island-arc settings with inferred ages between ca. 800 and 660-640

Ma. The earliest Pan-African deformation at 660-640 Ma resulted from closure of the ocean basin, SW-directed thrusting, and accretion of the island-arc remnants.

Recent studies provided new structural, geochronological, and geochemical data, however many critical questions about the accretionary history of the Anti-Atlas Belt are still debated (e.g. Soulaïmani et al., 2018 and references therein). Such questions include the age of obducted oceanic crust segments (Siroua and Bou Azzer El Graara ophiolites), relationship between arc terranes to the north with ophiolites and an associated blueschist facies belt to the south, the age and nature of the basement in these terranes, the degree of interaction of Neoproterozoic magmas with the basement, and the timing of accretion of these terranes to the West African Craton. Finally, discrimination between the post-Eburnean sedimentary succession, recently dated to be late Paleoproterozoic in age (Ikenne et al., 2017; Soulaïmani et al., 2019; Youbi et al., 2019; this work), and the Neoproterozoic sedimentary sequence (Leblanc and Moussine-Pouchkine, 1994; Bouougri and Saquaque, 2004; Blein et al., 2014a,b; Soulaïmani et al., 2018; Youbi et al., 2018) remains uncertain.

The lower Anti-Atlas Supergroup of the Anti-Atlas Belt (Lkest-Taghdout Group *sensu* Thomas et al., 2004; Figs. 1D, 2 and 3) is one of the most important Proterozoic lithostratigraphic units on the WAC, yet with poor age constraints. It consists essentially of basalts, quartzites, and stromatolitic/oolitic carbonates, intruded by dolerite dyke swarms and sill complexes (Ifzwane and Toudma Suites). The lower Anti-Atlas Supergroup was previously thought to be ca. 1000-800 Ma in age on the basis of (1) stromatolites characteristic for the Neoproterozoic carbonate successions (Choubert, 1962), and (2) the Rb/Sr age of the contact-metamorphosed host rocks associated with mafic dykes (789 ± 10 Ma) in the Tachdamt Subinlier of the Bou Azzer El Graara Inlier (Clauer, 1976; Cahen et al., 1984).

The present study is based on field observations and new U-Pb ages for igneous zircons from dolerite sill intruding carbonates from the lower Anti-Atlas Supergroup in the Zenaga Inlier and detrital zircon grains from four samples from the lower, middle, and upper parts of the lower Anti-

Atlas Supergroup in the Zenaga and Bou Azzer El Graara inliers. We propose a new lithostratigraphy for the Proterozoic lower Anti-Atlas Supergroup of the Anti-Atlas Belt based on a discussion of our data in combination with the previously published data (Abati et al., 2010; Walsh et al., 2012; Letsch, 2018; Bouougri et al., 2020) and constrain the provenance for the five, major unconformity-bounded packages of the supergroup.

2. Geological background

2.1. The West African Craton

The West African Craton (WAC, Fig. 1) extends across northwestern Africa, and consists of two Archean basement shields, the Reguibat Shield to the north and the Leo-Man Shield to the south. The Archean nuclei (3.5-2.7 Ga in age) of these shields are juxtaposed against an array of Paleoproterozoic domains (2.35-2.00 Ga in age) made up of greenstone belts and regions of extensive tonalite-trondhjemite-granodiorite (TTG) plutons, which are overlain by Meso- to Neoproterozoic and younger sedimentary basins. Smaller basement inliers are composed of Paleoproterozoic rocks: the Kayes and Kedougou-Kenieba inliers to the west, and those of the Anti-Atlas Belt to the north (Ennih and Liégeois, 2008; Abati et al., 2012). The WAC remained largely stable since 2 Ga (Schofield et al., 2006; Jessell et al., 2016). The basement of the WAC has evolved through several major orogenic cycles: the Paleoproterozoic Leonian cycle (several episodes between 3.40 and 3.10 Ga with uncertain chronology, related to continental accretion; e.g. Thiéblemont et al., 2004; Koffi et al., 2020), the Liberian cycle (2.85-2.70 Ga; Key et al., 2008; Koffi et al., 2020), the Paleoproterozoic Eburnean and Birimian cycles (2.15-2.07 Ga and 2.27-1.96 Ga, respectively; Abouchami et al., 1990; Boher et al., 1992; Egal et al., 2002; Schofield et al., 2006; Baratoux et al., 2011; Grenholm et al., 2019; McFarlane et al., 2019) and the late Neoproterozoic Pan-African orogenic event (885-550 Ma; Thomas et al., 2002; 2004; Gasquet et al., 2005; 2008; Walsh et al., 2012; Youbi et al., 2018). The WAC hosts four major intracratonic basins: the large central Taoudeni Basin and the smaller Tindouf and Volta basins, to the north and southwest, respectively, and the Gourma aulacogen to the east (Moussine-Pouchkine and Bertrand-

Sarfati, 1978; Villeneuve, 2005; Deynoux et al., 2006; Lottaroli et al., 2009). These basins comprise mostly Mesoproterozoic-Neoproterozoic to Devonian-Carboniferous sediments. The WAC is surrounded by Pan-African belts, including the Trans-Saharan and Anti-Atlas belts (Ennih and Liégeois, 2008; Liégeois et al., 2013; Toummite et al., 2013; Brahimi et al., 2018).

Early compilations of WAC lithostratigraphy suggested an absence of Mesoproterozoic events or rocks, indicating a tectonically quiescent period between 1.7 and 1.0 Ga (e.g. Ennih and Liégeois, 2008 and references therein). The lack of detrital zircons of that age in the pre-Cryogenian strata is considered characteristic of sediments sourced from the WAC (e.g. Abati et al., 2010; Gärtner et al., 2013; 2017; 2018). However, recent studies in the northern WAC (El Bahat et al., 2013; Kouyaté et al., 2013; Söderlund et al., 2013; Youbi et al., 2013; Ikenne et al., 2017) demonstrated important intraplate magmatic events based on dating of mafic dykes by the U-Pb method on baddeleyite and zircon. Six distinct events are distinguished throughout the inliers: ca. 2040 Ma Tagragra of Tata, 1750 Ma Tagragra of Akka, 1650 Ma Zenaga, 1416–1380 Ma Bas Drâa, and 885-883 Ma Iguerda-Taïfast events, and 612-600 Ma pulse of the Central Iapetus Magmatic Province (CIMP; Walsh et al., 2002; Gasquet et al., 2004; El Bahat et al., 2013; Kouyaté et al., 2013; Söderlund et al., 2013; Youbi et al., 2013; El Bahat et al., 2017; Ikenne et al., 2017; Bouougri et al., 2020; Youbi et al., 2020). Re-Os geochronology of the sedimentary succession of the Atar Group in the Taoudeni Basin of Mauritania indicated depositional ages ranging from 1109 ± 22 to 1105 ± 37 Ma (Rooney et al., 2010). These Mesoproterozoic ages are confirmed by the study of microfossils (Beghin et al., 2017) and chemostratigraphy (Kah et al., 2012). Recently, Baratoux et al. (2019) identified four new magmatic events 1.79-1.76, 1.58, 1.53, and 0.91-0.87 Ga in the southern WAC that add additional evidence for Mesoproterozoic mafic intrusive activity on this craton. Ar-Ar ages of ca. 1.24 Ga for the 1575 and 1520 Ma dyke swarms are considered to record a thermal overprint (Tapsoba et al., 2018).

2.2. The Anti-Atlas Belt

The ENE-WSW trending Anti-Atlas Belt (800 km long and 200 km wide) of Morocco is the most important segment of the Neoproterozoic Pan-African Orogen in the northern part of the WAC (Fig. 1D). The Precambrian basement crops out in several erosional inliers (“boutonniers” in French), containing Late Ediacaran and younger units (Bas Drâa, Ifni, Kerdous, Tagragra of Akka, Tagragra of Tata, Iggherm, Siroua, Zenaga, Bou Azzer, Saghro, and Ougnat; Fig. 1D) and distributed along two major fault systems (South Atlas Fault, SAF and Anti-Atlas Major Fault, AAMF; Fig. 1D).

The Anti-Atlas Belt contains the Paleoproterozoic basement (affected by the Eburnean Orogeny) unconformably overlain by the lower Anti-Atlas Supergroup (= Lkest-Taghdout Group *sensu* Thomas et al., 2004; = Tizi n’Taghatine Group *sensu* Bouougri and Saquaque, 2004), which is in turn unconformably overlain by the Ouarzazate Supergroup related to the Pan-African Orogeny. The two major periods of tectonothermal activity, associated with crustal accretion, have been recognized during the Proterozoic (Fig. 2) in the Anti-Atlas Belt and are described below.

(i) *The Paleoproterozoic Eburnean Orogeny (2.20-2.07 Ga*; Abouchami et al., 1990; Boher et al., 1992; Egal et al., 2002; Schofield et al., 2006; Baratoux et al., 2011; McFarlane et al., 2019). The Paleoproterozoic basement rocks are found exclusively SW of the AAMF (Choubert, 1963), but they may also constitute the basement to the Pan-African Orogen to the NE (Ennih and Liégeois, 2001; Liégeois et al., 2006; Gasquet et al., 2008). The Paleoproterozoic basement is composed of metasedimentary rocks, granites, paragneisses, and migmatites, with U-Pb zircon ages ranging from 2200 to 2030 Ma (e.g. Aït Malek et al., 1998; Walsh et al., 2002; Thomas et al., 2002; Gasquet et al., 2004, 2008; O’Connor, 2010).

(ii) *The Neoproterozoic Pan-African Orogeny (885–550 Ma)*. The Neoproterozoic sequences of the Anti-Atlas Belt are composed of units that were involved in the Pan-African Orogeny and the unconformably overlying post-collisional volcano-sedimentary rocks of the Ouarzazate Supergroup, which are a part of the Central Iapetus Magmatic Province (CIMP; Doblas et al., 2002; Maloof et al., 2005; Gasquet et al., 2005, 2008; Ernst and Youbi, 2017; Tuduri et al., 2018; Youbi et al., 2020). The Neoproterozoic rocks underlying the Ouarzazate Supergroup are subdivided into the

215 lower units (Lkest-Taghdout, Bou-Azzer, and Iri groups), affected by all Pan-African orogenic
 216 events, and the upper units (Saghro and Bou Salda groups), only affected by the latest stage of the
 217 Pan-African Orogeny (e.g. Walsh et al., 2002; Thomas et al., 2004; Gasquet et al., 2005; 2008;
 218 Alvaro et al., 2014).

219 The lower Anti-Atlas Supergroup consists essentially of basalt, quartzite, and stromatolitic/oolitic
 220 carbonate, cut by undated dolerite dyke swarms and sill complexes (Thomas et al., 2004; Bouougri
 221 and Saquaque, 2004). It marks the onset of a period of rifting at the northern portion of the WAC
 222 during the late Neoproterozoic. The rifting culminated with the opening of an oceanic basin on the
 223 northern edge of the WAC. The relics of the oceanic crust (Bou-Azzer Group) are preserved in the
 224 Bou-Azzer and Siroua inliers as highly sheared allochthonous ophiolite complexes (El Hadi et al.,
 225 2010; Blein et al., 2014a and references therein). The ophiolitic assemblage is mainly made up of
 226 ultramafic rocks (serpentinites with few chromite pods), but also includes mafic meta-cumulates,
 227 meta-basaltic sheeted dykes, and, to a smaller extent, pillow lavas (Leblanc, 1975, 1981; Admou
 228 and Juteau, 1998). Their geochemical signatures point to emplacement in a supra-subduction zone
 229 (Bodinier et al., 1984; Naidoo et al., 1991; Ahmed et al., 2005; Hodel et al., 2017; Triantafyllou et
 230 al., 2019; 2020). Dating of plagiogranite intrusions in the Siroua Inlier indicated formation of an
 231 oceanic crust at ca. 760 Ma (U-Pb zircon age; Samson et al., 2004). Given the absence of reliable
 232 radiometric data on the age of the lower Anti-Atlas Supergroup, ca. 760 Ma was considered as the
 233 minimum age of its formation. The rifting might be associated with the late Neoproterozoic
 234 extension within the supercontinent Rodinia (Gasquet et al., 2008). Following the opening of the
 235 ocean basin, a subduction phase was initiated with island-arc development (the Iri Group in the
 236 Siroua Inlier; Thomas et al., 2002; Triantafyllou, 2015; 2016). The cores and overgrowths of zircon
 237 grains from the Iri Group migmatite yielded two U-Pb SHRIMP ages at 743 ± 14 and 663 ± 13
 238 Ma, respectively (Thomas et al., 2002). The older age obtained from the cores is interpreted as the
 239 age of crystallization and emplacement of the migmatite protolith in an island-arc setting, whereas
 240 the younger age is thought to represent the timing of ophiolite obduction and arc accretion to the

northern edge of the WAC in association with the regional metamorphism and the main orogenic event in the Anti-Atlas Belt (Thomas et al., 2002).

The development of the Anti-Atlas Belt oceanic arc complexes occurred during three magmatic flare-ups at ca. 750, 700, and 650 Ma that were interspersed with an early major tectonic episode between 730-700 Ma (Triantafyllou et al., 2018; 2020). A garnet Sm-Nd date of 647.2 ± 1.7 Ma was recently obtained for the Tasriwine ophiolite complex in the Siroua Inlier (Inglis et al., 2016). This age is more than 15 million years younger than the previous age estimate by Thomas et al. (2002). North of the AAMF in the eastern Anti-Atlas Belt (Figs. 1D and 2), the oldest rocks are thick deposits of basalt and turbidite of the Saghro Group, interpreted as either a distal-slope equivalent of the upper part of the lower Anti-Atlas Supergroup, deposited between the WAC to the south and an island arc (Saquaque et al., 1989; Thomas et al., 2002), or slivers of the stretched Eburnean crust (Hindermeyer, 1953; Ouguir et al., 1996; Fekkak et al., 2001; 2002; Massironi et al., 2007; Schiavo et al., 2007; Michard et al., 2017).

Preliminary detrital zircon data reported by Liégeois et al. (2006) and summarized in Gasquet et al. (2008) suggest that the sediments of the Saghro Group in the Kelaa Mgouna section were derived from sources ranging in age from about 880 to 615 Ma. The later post-collisional, extensional stage (605–542 Ma) is characterized by molasse deposition and magmatism of the Ouarzazate Supergroup, which was followed by the foreland development in the Saharan cratonic basin to the south (Taroudant and Tata groups; e.g. Doblas et al., 2002; Maloof et al., 2005; Gasquet et al., 2005, 2008; Hefferan et al., 2014; Karaoui et al., 2014; 2015; Tuduri et al., 2018; Geyer and Landing, 2006; 2020; Youbi et al., 2020).

2.3. The lower Anti-Atlas Supergroup

Since our study is focused on the lower Anti-Atlas Supergroup, we will give here an updated review of its lithology, stratigraphy, and nomenclature. The lower Anti-Atlas Supergroup, known in the Anti-Atlas literature as the Taghdout (Lkest/Tizi n'Taghatine) Group, contains a sedimentary

267 succession deposited along the northern margin of the WAC (e.g. Thomas et al., 2002, 2004;
 268 Bouougri and Saquaque, 2004; Alvaro et al., 2014). The lower part of the lower Anti-Atlas
 269 Supergroup shows a progressive change of sedimentary facies from SW to NE representing a broad
 270 proximal to distal trend: conglomerates, sandstones, and quartz sandstones predominate in the Ifni
 271 and Kerdous inliers, while quartz sandstones, carbonates, shales, turbidites, and basaltic lava flows
 272 (the so called “Série des Calcaires et Quartzites”; see Choubert, 1962) developed in the Igherm,
 273 Siroua, Zenaga, and Bou Azzer El Graara inliers (Bouougri and Saquaque, 2004; Thomas et al.,
 274 2004; Alvaro et al., 2014). The scarcity of structural studies combined with modern sedimentology
 275 and sequence stratigraphy to test correlations between the groups in different inliers has led to a
 276 proliferation of synonymous names in the lithostratigraphic terminology: e.g. the Jbel Lkest Group
 277 in the Kerdous Inlier (Hassenforder, 1987; O’Connor, 2010), the Bleida-Tachdamt or Bleida Group
 278 in the Bou Azzer El Graara Inlier (Leblanc and Moussine-Pouchkine, 1994; Blein et al., 2014a), and
 279 the Taghdout Group (= Tizi n’Taghatine) in the Zenaga Inlier (Thomas et al., 2002, 2004; Bouougri
 280 and Saquaque, 2004). Correlation between these groups (recently combined into the Lkest-
 281 Taghdout Group, equivalent to the lower Anti-Atlas Supergroup; Gasquet et al., 2008; Álvaro et al.,
 282 2014) is both lithostratigraphic and structural. The only lithostratigraphic correlation linking the
 283 Siroua-Zenaga and Bou Azzer El Graara inliers was proposed by Bouougri and Saquaque (2004)
 284 who considered the section of the Tizi n’Taghatine mountain pass as a stratotype for the group
 285 named after this locality. Although their lithostratigraphy requires revision, we will refer to it in this
 286 work. Two reasons motivated this choice: (1) the dated samples of this study are from the same area
 287 that Bouougri and Saquaque (2004) focused on, and (2) the Siroua-Zenaga and Bou Azzer El
 288 Graara inliers are the only areas where modern sedimentology and sequence stratigraphy were
 289 applied (Fig. 1D). However, we will use the name of the lower Anti-Atlas Supergroup (an
 290 equivalent of the Lkest-Taghdout Group) because, although the latter group is widely known in the
 291 literature (Gasquet et al., 2008; Álvaro et al., 2014), it combines genetically unrelated sequences
 292 deposited in superimposed sedimentary basins and in different tectonic settings with long-lasting

hiatuses along unconformities. Five, major unconformity-bounded lithostratigraphic packages are further here distinguished: the lower Tasserda-Taghatine Group, the second Oumoula (Mimount) Formation, the third Tizi n'Taghatine Group, the fourth Tachdamt Formation, and the uppermost Bleida Formation.

For a long time, until 2013, available data on the lower Anti-Atlas Supergroup suggested that the continental passive margin developed during the Tonian between about 1000 Ma (based on stromatolite morphology pointing to a Neoproterozoic age; Choubert, 1963) and ca. 787 Ma (Rb-Sr isochron age of contact metamorphism of the host rocks associated with mafic dykes at Tachdamt; Clauer, 1976; Cahen et al., 1984). However, these age constraints are still debated (Gasquet et al., 2008; Alvaro et al., 2014; Ikenne et al., 2017; Soulaïmani et al., 2018; 2019; Youbi et al., 2019). There are too few reliable geochronologic age constraints to tightly constrain the age of the lower Anti-Atlas Supergroup. It unconformably overlies the Eburnean basement, including the post-tectonic 2032 ± 5 Ma Azguemerzi granodiorite of the Zenaga Inlier, and in turn is unconformably overlain by the subhorizontal volcanic and sedimentary succession of the Late Ediacaran (ca. 606-541 Ma) Ouarzazate Supergroup (Figs. 6A, B). The lower Anti-Atlas Supergroup records a low-grade tectonometamorphic event related to the Pan-African Orogeny dated at 663 ± 13 Ma (Thomas et al., 2002) and 647.2 ± 1.7 Ma (Inglis et al., 2016) in the Siroua Inlier. Consequently, the depositional age of the lower Anti-Atlas Supergroup can be broadly bracketed between 2032 ± 5 Ma and $663 \pm 13 / 647.2 \pm 1.7$ Ma.

The lower Anti-Atlas Supergroup sediments are intruded by abundant dolerite and gabbro dykes and sills, also found as dyke swarms and sill complexes in the Paleoproterozoic basement of the Zenaga (Fig. 4) and several other inliers (e.g. Iggherm, Tagragra of Akka, Tagragra of Tata, Kerdous, and Bas Drâa). These rocks have been collectively called the Ifzwane Suite in the Zenaga and Siroua inliers (Thomas et al., 2004), and the Toudma Suite in the Kerdous Inlier (Thomas et al., 2004; O'Connor, 2010).

Letsch (2018) showed that the quartzites of the Mimount Formation (*sensu* Thomas et al., 2002) and the Oumoula Formation (*sensu* Bouougri and Saquaque, 2004), overlying the basal, carbonate-quartzite-metapelite units near the Town of Taghdout at the northern margin of the Zenaga Inlier (see Thomas et al., 2002), do not contain detrital zircons younger than ca. 1987 Ma (Benziane, 2007; Walsh et al., 2012) and ca. 1809 Ma (Abati et al., 2010), respectively. Neoproterozoic detrital zircon grains only appear in the overlying formations of the Saghro and Bou Salda groups of the Siroua Inlier (Abati et al., 2010). Detrital zircons from the Mimount and Oumoula formations display exclusively Paleoproterozoic ages, suggesting a ca. 1890 Ma maximum depositional age for the upper parts of the formations. In the Ifni Inlier, detrital zircons from quartzites of the upper part of the Lkest Group (the basal group of the lower Anti-Atlas Supergroup) exhibit two main clusters in distribution of SHRIMP $^{207}\text{Pb}/^{206}\text{Pb}$ ages, 2060-1890 and 1810-1640 Ma (Schulte et al., 2016). The maximum depositional age is given by a weighted average $^{207}\text{Pb}/^{206}\text{Pb}$ date of 1744 ± 26 Ma based on four youngest concordant grains.

3. Sampling and Analytical Procedures

3.1. Sampling

In order to investigate the maximum depositional age of the lower Anti-Atlas Supergroup of the Anti-Atlas Belt, we collected five samples from two key inliers within the Anti-Atlas Orogen (Zenaga and Bou Azzer El Graara) for geochronological investigation. The Taghdout sill that intruded to the lower part of the lower Anti-Atlas Supergroup in the Zenaga Inlier was sampled for geochemistry and zircon U-Pb geochronology: sample TA1 was collected at the centre of the sill and herein dated at 1676 ± 37 Ma and sample TA2 is from the margin of the sill. Four metasedimentary samples were collected for detrital zircon U-Pb geochronology: one sample from metasilstones of the Tasserda Formation of the Zenaga Inlier and three samples from the Agoummy, Tachdamt, and Bleida formations of the lower Anti-Atlas Supergroup in the Bou Azzer

Inlier (Figs. 3, 4, and 5). Sample site locations, a brief description of the samples, and their ages are provided in Table 1.

3.2. Analytical procedures

3.2.1. Whole-rock geochemistry

Elemental concentrations of whole rocks were obtained at “Service d'Analyse des Roches et des Minéraux” at the Geochemical and Petrographical Research Center in Nancy (SARM laboratory-CRPG-CNRS, Nancy, France). To avoid contamination, crushing and pulverizing were undertaken using agate mills. Powders were treated with LiBO_2 , dissolved with HNO_3 , and analyzed and calibrated with international geostandards (Basalt BE-N, Basalt BR, Dolerite WS-E, Microgabbro PM-S). Major and trace elements were analyzed using inductively coupled plasma atomic emission spectroscopy (ICP-AES) and Inductively Coupled Mass Spectrometry (ICP-MS), respectively. The analytical procedures are described in detail at <http://helium.crpq.cnrs-nancy.fr/SARM/>.

3.2.2. U-Pb Geochronology

Zircon grains from all studied samples were separated from 3-5 kg of crushed rocks using conventional heavy liquid and magnetic separation techniques (jaw crusher, disk grinder, Wilfley table, Frantz isodynamic magnetic separator, and density separation using bromoform and methylene iodide) at the Centro de Pesquisas Geocronológicas, Universidade de São Paulo (CPGeo-USP), Brazil. About 20 to 100 zircon grains from each sample were mounted in epoxy resin, polished to approximately half of mean grain thickness for further imaging with transmitted light and cathodoluminescence to resolve internal grain texture. After coating with Au, cathodoluminescence (CL) images of zircon grains were obtained using a Quanta 250 FEG scanning electron microscope equipped with Mono CL3 + cathodoluminescence spectroscope (Centaurus) at the Centro de Pesquisas Geocronológicas of the University of São Paulo (CPGeo-USP), Brazil. The conditions used in CL analysis were as follows: 60 μA emission current, 15.0 kV

accelerating voltage, 7 μm beam diameter, 200 μs acquisition time, and a resolution of 1024×884 pixels.

3.2.2.1 Sensitive High Resolution Ion Microprobe (SHRIMP II) analysis

Igneous zircon grains from the dolerite sample TA 1 and detrital zircon grains from the quartzite samples BA21 and BA24 were analyzed by the U-Pb isotopic technique using a SHRIMP II at CPGeo-USP following the analytical procedures of Williams (1998). The crystals were mounted, together with the TEMORA-2 reference zircon (Black et al., 2004), in epoxy and polished to expose the interior of the grains. Correction for common Pb was made based on ^{204}Pb measured, and the typical error for the $^{206}\text{Pb}/^{238}\text{U}$ ratio was less than 2%; uranium abundance and U/Pb ratios were calibrated against the TEMORA-2 reference material, and the ages were calculated using Isoplot® version 3.0 software application (Ludwig, 2003). Errors are reported as 1σ deviations and ages have been calculated at the 95% confidence level.

3.2.2.2 Laser Ablation Inductively Coupled Plasma Mass Spectrometry (LA-ICP-MS) analysis

U-Pb analyses of detrital zircon grains from the samples TG-15-1 and ALBL29 were performed using a Neptune multicollector inductively coupled plasma mass spectrometer (MC-ICP-MS) and an Analyte G2 excimer laser ablation (LA) system at CPGeo-USP, Brazil. The U-Pb analysis was performed in the following order: two blanks, two 612 NIST standard glasses, three external standards, 13 unknown samples, two external standards, and two blanks. An interpolation of the four blank measurement sequences, using the bracketing method with two blanks before and two after the samples, was subtracted from all the seven measured masses (202, 204, 206, 207, 208, 232, and 238). The ^{235}U isotope was not measured, but calculated using the relative abundance $^{238}\text{U}/^{235}\text{U}=137.88$. The ^{204}Hg interference on ^{204}Pb was corrected using ^{202}Hg and assuming $^{204}\text{Hg}/^{202}\text{Hg} = 4.350360$. Using the partially corrected $^{207}\text{Pb}/^{206}\text{Pb}$ ratio as an estimate for the age and the equations of Stacey and Kramers (1975), the $^{206}\text{Pb}/^{204}\text{Pb}$, $^{207}\text{Pb}/^{204}\text{Pb}$, and $^{208}\text{Pb}/^{204}\text{Pb}$ relative abundances were calculated and corrected for common (non-radiogenic) Pb fraction.

Bracketing measurement of unknowns with NIST-612 standard glasses (two measurement sequences before and after) was used to convert the total Pb, Th, and U signals to ppm concentrations.

A final bracketing (with three measurements before and two after unknown) with the GJ-1 reference zircon (Jackson et al., 2004) was used to correct the effect of the instrumental fractionation on four ratios ($^{206}\text{Pb}/^{238}\text{U}$, $^{207}\text{Pb}/^{235}\text{U}$, $^{207}\text{Pb}/^{206}\text{Pb}$, and $^{208}\text{Pb}/^{232}\text{Th}$) before they were used to calculate the respective ages. Errors in all arithmetic operations were algebraically propagated. Least squares fitting was used for the best line fit and all the data are reported at the 1σ level calculated with Isoplot 3.70 (Ludwig, 2008). A U-Pb analysis is concordant when it overlaps with the Concordia within uncertainty. So, it seems to be appropriate to exclude results with a low level of concordance ($^{206}\text{Pb}/^{238}\text{U}$ age / $^{207}\text{Pb}/^{206}\text{Pb}$ age $\times 100$), but very large errors that overlap with the Concordia from interpretation. Thus, an interpretation with respect to the obtained ages was done for all grains within the concordance interval of 90-110 % ($^{206}\text{Pb}/^{238}\text{U}$ age / $^{207}\text{Pb}/^{206}\text{Pb}$ age $\times 100$) which is often used (e.g. Spencer et al., 2016). Discordant analyses were generally interpreted with caution. Analyses with discordance of more than 10% were discarded. For zircons younger than 1300 Ma, $^{206}\text{Pb}/^{238}\text{U}$ ages were preferred, while for older zircons the $^{207}\text{Pb}/^{206}\text{Pb}$ ages were used. The detrital zircon age distribution by probability density was performed using the AgeDisplay application (an EXCEL workbook to evaluate and display univariate geochronological data using binned frequency histograms and probability density distributions) (Sircombe, 2004).

4. Results

4.1. The Great Mafic Sill of the Ifzwane Suite at Taghdout Town

4.1.1. Field observations

The Taghdout sill was collected near Taghdout Town, ca. 9 km NW of Tazenakht City (Table 1 and Figs. 3 and 4), from a 227 m thick, well-preserved, massive sill belonging to the Ifzwane Suite (dolerite dyke swarm and sill complex of the Zenaga Inlier, of likely multiple ages as discussed in

section 2.1). The sill intruded into the lower part of the lower Anti-Atlas Supergroup (Fig. 6C), which mainly consists of carbonate and mixed siliciclastic-carbonate sequence of the Ifrane-n-Taghatine Formation (Bouougri and Saquaque, 2004). The sill can be traced in outcrops for more than one km in length and has thick chilled margins of marble grading away from the intrusion into carbonate host rock (Figs. 3, 4, and 6D). The following new samples were used in this study: TA1 (centre of the sill) and TA2 (border of the sill) from the Taghdout sill of the Zenaga Inlier.

4.1.2. Petrographic features

Fresh samples of the Taghdout sill are dark-grey, but due to widespread hematitic alteration the rock appears reddish. However, the original texture and primary igneous mineralogy are usually well preserved. The texture varies from a coarse-grained doleritic texture in the centre of the sill to a fine-grained, microlitic porphyric texture at the border of the sill (Fig. 7).

The primary igneous minerals consist of clinopyroxene (augite and locally pigeonite), very rare orthopyroxene, plagioclase, and oxides. Apatite and zircon are present as accessory minerals; amphibole and, more rarely, biotite are found around pyroxene crystals or as irregular patches in the groundmass. Although tholeiitic mafic rocks are generally characterized by low alkaline content ($\text{Na}_2\text{O}+\text{K}_2\text{O}$), granophyric quartz/K-feldspar mesostasis was observed in coarse-grained portions of the sill, consistent with the examples previously described in literature (e.g. Fodor, 2009; Block et al., 2015). The secondary mineral assemblage includes albite, chlorite, actinolite, epidote, titanite, calcite, and quartz, which is characteristic for the greenschist facies of metamorphism.

4.1.3. Geochronology: U-Pb dating of the Great Taghdout sill

The zircon grains from the sample TA1 range in length from 120 to 150 μm . Cathodoluminescence (CL) images (Figure S1 of supplementary data) show that most zircons are stubby sub-euhedral to euhedral, with a typical oscillatory zoning within a large uniform central zone, followed by much finer, poorly resolvable oscillatory zoned bands. Some grains lack any zoning (Figure S1 of supplementary data). We analyzed 18 spots on 13 zircon grains (Table S1 of supplementary data).

High Th/U ratios (between 1.18 and 3.38) support a magmatic origin for the zircons (Fig. 8) (cf. Möller et al., 2003; Hoskin and Schaltegger 2003; Linnemann et al., 2007). Most of the analyzed spots with high U and Th concentrations show evidence for significant Pb loss. On the Wetherill diagram, these thirteen analyses are aligned along a discordia that shows two intercepts with the concordia (Fig. 9A): an upper intercept with an age of 1639 ± 34 Ma and a lower intercept with an age of 688 ± 65 Ma. The age of 1639 ± 34 Ma is interpreted to reflect the time of magmatic emplacement of the sill, whereas the 688 ± 65 Ma age is thought to correspond to the time of a metamorphic and /or thermal overprint event. However, these intercept ages have a very high Mean Square Weighted Deviation (MSWD = 6.8), making this age poorly constrained. In contrast, analyses with low Th and U concentrations (Th <1000 ppm, spots 2.1, 7.1, and 13.1) produced nearly concordant $^{207}\text{Pb}/^{206}\text{Pb}$ ages. We therefore calculated the weighted mean average $^{207}\text{Pb}/^{206}\text{Pb}$ age for those spots characterized by low Th and U concentrations, obtaining a mean $^{207}\text{Pb}/^{206}\text{Pb}$ age of 1676 ± 37 Ma ($n = 3$, MSWD = 3.1; Fig. 9B).

4.1.4. Geochemical characteristics

The major and trace element composition of previously studied and new samples (TA1 and TA2) are listed in Table 2. On the total alkali versus silica diagram (Fig. 10A), the studied samples TA1 and TA2 and the ca. 1.6 Ga doleritic dykes and sills previously analyzed by Youbi et al. (2013) plot in the subalkaline basalt field. These additional dolerites are: DZ 36 (dated at 1640 ± 2 Ma and 1641 ± 3 Ma; Kouyaté et al., 2013) and DZ 40 (dated at 1656 ± 9 Ma; Kouyaté et al., 2013) of the Zenaga Inlier (close to Tazenakht City), and doleritic dyke (AMHA069 dated at 1654 ± 16 Ma; Kouyaté et al., 2013) from the Agadir Melloul Inlier. This subalkaline affinity is also evident from Nb/Y ratios that are below 0.67 (cf. Pearce and Cann, 1973; Winchester and Floyd, 1977). Additionally, their non-orogenic tholeiitic character is shown in the Ti vs. V diagram of Shervais (1982) (Fig. 10B) with the data plotting exclusively within the Continental Flood Basalt (CFB) and Mid-Ocean Ridge Basalt (MORB) fields ($20 < \text{V}/\text{Ti} < 50$). When plotted on a multi-element

variation diagram (spidergram; Fig. 11), the dated dykes display patterns similar to some continental tholeiites such as those of the Central Atlantic Magmatic Province (CAMP) with a moderate enrichment in large-ion lithophile elements (LILE) and distinct positive Pb and negative Nb anomalies (Bertrand et al., 1982; Youbi et al., 2003; Marzoli et al., 2004, 2019; Tegner et al., 2019). Nb anomalies have been related to crustal contamination or the influence of a subduction zone, which was either active during magma generation or earlier metasomatized lithospheric mantle with which asthenospheric magma interacted during ascent (e.g. Thompson et al., 1982; Dupuy and Dostal, 1984; Coish and Sinton, 1992; Dorais et al., 2005). Furthermore, the negative Nb anomaly is accompanied by negative P anomaly; this pattern is similar to some low-Ti CFB tholeiitic magmas contaminated by continental crust. This and the relatively high Ti/V ratios (see Fig. 10B) point to a non-orogenic origin of the mafic magmas, which were contaminated with continental crust during magma ascent/emplacement (Ikenne, 1997; Hafid, 1999; Hafid et al., 2001; El Aouli et al., 2004; 2010; El Bahat et al., 2013; Youbi et al., 2013; El Bahat et al., 2017).

4.2. Geochronology: Detrital Zircon Study of the lower Anti-Atlas Supergroup

4.2.1 Metasiltstone of the Tasserda Formation: sample TG-15-1

The sample TG-15-1 (Table 1) is purple, coarse-grained quartz-rich metasiltstone collected from the Tasserda Formation near the village of Taghadout (Figs. 3 and 4). The Tasserda Formation consists of thick- to medium-bedded quartz arenites with few channels filled with conglomerates. The main mineral components of siltstone are quartz and feldspar embedded in a sericite-rich matrix (Fig. 12A).

The zircon grains of this sample exhibit mean lengths and widths of 120 and 60 μm , respectively. In most cases, they are sub-euhedral to euhedral in shape. The internal texture of zircon grains shows oscillatory concentric, sector, and banded zoning (Figure S2 of supplementary data). The Th/U ratio ranges from 0.03 to 1.15 (Fig. 8). About 83 % of the analyzed zircons have Th/U ratios larger than 0.1, potentially indicating that they were derived from intermediate to felsic igneous rocks (cf. Möller et al., 2003; Hoskin and Schaltegger 2003; Linnemann et al., 2007). The other 17 % have a

Th/U ratio below 0.1 and could be of metamorphic origin. From the total of 99 measured spots, 74 yielded less than 10 % discordant ages (Table S2 of supplementary data). About 99 % of the zircon population is of Paleoproterozoic age (26 % are Rhyacian and 73 % are Orosirian), ranging from 2294 $\pm$ 32 Ma to 1831 $\pm$ 35 Ma, with a prominent peak at around 2022 Ma and three minor peaks at about 2294, 2142, and 1870 Ma. One remaining analysis, representing 1% of the concordant ages of this sample, yielded an age of 2653 $\pm$ 32 Ma (Fig. 13A, B). The dominance of the Paleoproterozoic ages indicates that almost all of the detrital zircon grains were derived from the Paleoproterozoic (Eburnean) crust. The seven youngest zircon grains yielded a concordia age of 2049.3 $\pm$ 4.6 Ma, which constrains the maximum depositional age of the Tasserda Formation (Fig. 13A, B).

4.2.2. Quartzites of the Agoumy Formation: sample BA21

The sample BA21 (Table 1) is a medium-grained quartzite collected from the Agoumy Formation at the Tachdamt section (Figs. 3 and 5). The Agoumy Formation consists of thick- to medium-bedded quartzites with few conglomerates nested in channels.

The quartzites are dominated by quartz that is mainly subrounded. Some quartz grains show overgrowth rims. The rock is moderately sorted with a matrix that consists of silt-sized quartz and mica minerals such as sericite. Intergranular voids are filled with secondary Fe-Ti oxides (Fig. 12B).

Zircons from this sample have a mean length and width of 110 and 70 μ m, respectively. Zircon grains are mainly sub-rounded to rounded in shape with some euhedral crystals (Figure S3 of supplementary data). Most of them show oscillatory zoning, suggesting that they are of igneous origin. This is corroborated by their Th/U ratios ranging from 0.19 to 1.50 (Fig. 8). The internal textures include oscillatory zoning, planar zoning, banded zoning, and a minor portion with sector zoning. A total of 55 spots have been analysed (Table S3 of supplementary data), of which 50 yielded 90 to 110% concordant ages. About 84% are Paleoproterozoic (Statherian - 4%, Orosirian - 24%, Rhyacian - 54%, and Siderian - 2%), 12% of all ages are Neoproterozoic and 4% are Mesoproterozoic (2869 $\pm$ 8 Ma and 2857 $\pm$ 32 Ma). The Neoproterozoic ages range from 2735 $\pm$ 11 to

2531 $\pm$ 18 Ma, while the Paleoproterozoic ages range from 2498 $\pm$ 14 to 1786 $\pm$ 20 Ma. The probability-density plot is dominated by peaks at ca. 2205, 2080, 2035, and 1812 Ma. The six youngest grains, ranging from 1834 $\pm$ 24 Ma to 1786 $\pm$ 20 Ma, combined yield a concordia age of 1818 $\pm$ 6.4 Ma (MSWD = 1.5; probability = 0.22), which likely represents the maximum depositional age for the Agoummy Formation (Fig. 13C, D).

4.2.3. Quartzites of the Tachdamt Formation: sample ALBL29

The sample ALBL29 (Table 1) is a medium- to coarse-grained, poorly sorted quartz arenite from the lowermost part of the Tachdamt Formation at the Bleida section (Figs. 3 and 5). This formation consists of basaltic lava flows and interbedded volcanic breccias, lapilli tuffs, and volcanoclastic sediments. The mineral assemblage mainly consists of quartz with a small amount of feldspar and a matrix of silt-sized quartz and mica minerals (Fig. 12C).

Zircon grains from this sample exhibit a mean length and width of 160 and 90 μ m, respectively. The grains are sub-rounded and euhedral (Figure S4 of supplementary data). The internal textures predominantly show concentric zoning, with few grains lacking any zoning. The Th/U ratio ranges from 0.01 to 1.26, with a dominant population of likely igneous origin (98%; 0.1 to 1.26) and a minor population with a potentially metamorphic origin (2%; 0.01 to 0.1; Fig. 8). Out of 96 measured spots (Table S4 of supplementary data), 92 yielded less than 10% discordant ages (Fig. 14A, B). About 54% of the ages are Mesoproterozoic (Calymmian - 22%, Ectasian - 17%, and Stenian - 17%), 41.3% of the ages are Paleoproterozoic (Siderian - 4%, Rhyacian - 8%, Orosirian - 18%, and Statherian - 11%), 2% of the ages are Neoproterozoic (Tonian), 1% of the ages are Mesoarchean, and 1% of the ages are Neoarchean. Neoarchean and Mesoarchean ages are each represented by one analysis at 2577 $\pm$ 53 and 2969 $\pm$ 48 Ma, respectively. Paleoproterozoic ages range from 2482 $\pm$ 66 to 1603 $\pm$ 175 Ma with two prominent peaks at ca. 1978 and 1752 Ma. The Mesoproterozoic ages range from 1572 $\pm$ 55 to 1043 $\pm$ 12 Ma with four prominent peaks at ca. 1525, 1280, 1208, and 1060 Ma. The Neoproterozoic age at 822 $\pm$ 6 Ma is only 92% concordant and has a very low Th/U ratio of 0.01 indicating some overprint of the zircon grain. This grain is

therefore not used to calculate a maximum depositional age. The inferred maximum age of deposition is set by a concordia age derived from the youngest population of concordant grains at 1060 ± 5.5 Ma (four grains), and 930 ± 12 Ma, a 100% concordant single grain analysis with a Th/U ratio of 1.10 (Fig.14A, B).

4.2.4. Metasandstone of the Bleida Formation: sample BA24

The sample BA24 (Table 1) is a fine-grained, moderately sorted metasandstone from the middle part of the Bleida Formation (Figs. 3 and 5) at the Bleida section. The Bleida Formation consists of up to 600 m thick siliciclastic deposits interbedded with volcanoclastics. The sample contains quartz, plagioclase, and lithic fragments with matrix bearing clays, sericite, muscovite, and chlorite.

The grains are subangular to subrounded indicating that the sediment is immature (Fig. 12D).

The zircon grains of this sample exhibit a mean length and width of 125 and 85 μm , respectively, and most of them are sub-rounded to sub-angular in shape. The internal texture of the zircons revealed by CL images comprise concentric and sector zoning, and some unzoned grains (Figure S5 of supplementary data). The Th/U ratio ranges from 0.18 to 1.31, indicating a magmatic origin of the zircon grains (Fig. 8). Of 44 measured spots (Table S5 of supplementary data), 40 yielded less than 10% discordant ages. A total of 93% exhibits Paleoproterozoic ages (Siderian - 7%, Rhyacian - 49%, and Orosirian - 37%), Mesoarchean, Mesoproterozoic, and Neoproterozoic populations are each 2%. The Mesoarchean is represented by one age of 2852 ± 7 Ma, while the Paleoproterozoic ages range from 2487 ± 6 to 1841 ± 12 Ma with one prominent peak at ca. 2075 Ma and three minor peaks at ca. 2158, 1960, and 1852 Ma. The only Mesoproterozoic age is 1276 ± 14 Ma and the Neoproterozoic age is 977 ± 10 Ma. Two youngest grains, overlapping within errors, give a concordia age of 1855 ± 6.5 Ma (Fig. 14C, D).

In summary, the inferred maximum depositional ages based on concordia diagrams for the sedimentary succession of the lower Anti-Atlas Supergroup are 2049.3 ± 4.6 Ma for the Tasserda Formation, 1818 ± 6.4 Ma for the Agoumy Formation, 1060 ± 5.5 Ma for the Tachdamt

Formation, and 1855 ± 6.5 Ma for the Bleida Formation (Table 1). However, the latter formations could be younger (see Section 5.3)

5. Discussion

5.1 The 1.68-1.64 Ga Zenaga Event

The SHRIMP U-Pb geochronologic data for the Great Taghdout sill presented here shows a weighted mean average $^{207}\text{Pb}/^{206}\text{Pb}$ age of 1676 ± 37 Ma interpreted to date the magmatic emplacement of the sill (Fig. 9B). This age is similar within error to those obtained for other dolerite dykes and sills in the studied area (Zenaga Inlier), as well as in other inliers of the Anti-Atlas Belt (e.g. Agadir Melloul Inlier). Kouyaté et al. (2013) obtained ages of 1640 ± 2 Ma and 1641 ± 3 Ma (U-Pb baddeleyite) on the sill (DZ 36), which intrudes the Eburnean basement in the Zenaga Inlier (Fig. 4). They also obtained an age of 1656 ± 9 Ma (U-Pb baddeleyite) on another sill (DZ40), which is connected to a sill intruding the lowermost part of the lower Anti-Atlas Supergroup of the Zenaga Inlier (the Tasserda-Taghatine Group) and an age of 1654 ± 16 Ma (U-Pb baddeleyite) on a large NE-trending dyke, which crosscuts the Eburnean basement of the Agadir Melloul Inlier. More recently, Youbi et al. (2019) dated two sills intruding quartzites of the Lkest Group (equivalent of the Oumoula Formation) in the Kerdous Inlier. One of the sills yielded an upper intercept date of ca. 1640 Ma, constrained with ID-TIMS data for three zircon grains. LA-ICP-MS data for the same sample yielded a weighted mean average $^{207}\text{Pb}/^{206}\text{Pb}$ age of ca. 1650 Ma, identical within analytical uncertainties to the ID-TIMS date. Only a single baddeleyite grain has been analyzed from the second sill, producing a similar crystallization age of ca. 1650 Ma. All of these ages overlap within error with the newly obtained age from this study.

These ca. 1.68-1.64 Ga ages match the Rb-Sr dates, which were obtained for granitic rocks from the Anti-Atlas Belt; the Timrhahbine granite (Zenaga Inlier), Tasserhert granite (Kerdous Inlier), and Oued Chaiba granite (Bas Drâa Inlier) dated at 1650 ± 50 , 1680 ± 35 , and 1640 ± 32 Ma, respectively (Charlot, 1976; 1982). This suggests an event of bimodal magmatism during the late

602 Paleoproterozoic, post-dating the ca. 2.04 Ga Tagragera of Tata LIP extensional event (Kouyaté et
 603 al., 2013; Youbi et al., 2013), and confirms the polycyclic character of magmatism in the Anti-Atlas
 604 Belt, previously inferred in Kerdous and Tagragera of Akka inliers (Gasquet et al., 2004; Barbey et
 605 al., 2004). These ca. 1650 Ma ages are interpreted to belong to a single LIP event, which is termed
 606 the Zenaga Event (Kouyaté et al., 2013; Youbi et al., 2013; Ernst, 2014).

607 Globally, magmatism at ca. 1640-1656 Ma is also observed on other cratons (Fig. 15). In eastern
 608 and southwestern Greenland, the Melville Bugt dyke swarm extends for about 1000 km and is
 609 precisely dated at 1622-1635 Ma by the U-Pb method (1622 ± 3 , 1629 ± 1 , 1632 ± 1 , and 1635 ± 3
 610 Ma; Halls et al., 2011). Following upon previous work on the ca. 1630 Ma Melville Bugt dyke
 611 swarm along the NW coast of Greenland, Klausen and Nilsson (2019) confirmed its proposed
 612 continuation to the SE Greenland, indicating an extraordinary >2000 km long Trans-Greenlandic
 613 dyke swarm of Laurentia. This correlation was not only based on the swarm's continuous pattern,
 614 but also on a matching 1630 ± 4 Ma U-Pb baddeleyite age determination (Klausen and Nilsson,
 615 2019) and similar whole-rock compositions as well as distinctive outcrop characteristics. In
 616 Laurentia, the bimodal Narakay Volcanic Complex in the Dease Arm of the Great Bear Lake,
 617 associated with deposition in the intracontinental Hornby Bay Basin, NWT, Canada, yielded TIMS
 618 U-Pb zircon age of 1663 ± 8 Ma (Bowring and Ross, 1985).

619 Corresponding magmatism on Baltica includes the ca. 1.64 Ga NW-trending dyke swarms of Häme,
 620 Sipoo, and Suomenniemi and the related rapakivi granite complexes exposed in the Svecofennian
 621 Domain of the Fennoscandian Shield in southeast Finland (Vaasjoki et al., 1991; Salminen et al.,
 622 2017). The most extensive of these dyke swarms is the 1642 ± 2 Ma Häme diabase dyke swarm (U-
 623 Pb ID-TIMS baddeleyite age; Salminen et al., 2017).

624 In Southern Siberia, the Nersa Sill Complex from the Biryusa Block of the Irkutsk Promontary has
 625 a 1641 ± 8 Ma U-Pb baddeleyite age (Ernst et al., 2016). Other potential matches are rejected. The
 626 Khibelen dykes of the Baikal Uplift, Siberia have a 1674 ± 29 Ma Sm-Nd pyroxene, plagioclase,
 627 and whole-rock isochron age (Gladkochub et al., 2010). However, the Khibelen dykes are on a

trend from the SW-trending Chaya dyke subswarm, which has a 1752 ± 3 Ma U-Pb baddeleyite age (Gladkochub et al. 2010; Ernst et al. 2016), and so the Chaya and Khibelen dykes likely represent a single ca. 1750 Ma swarm. The type section for the Early Riphean, a major Russian stratigraphical division, in the South Urals Mountains, southeast Baltica, contains poorly-dated trachybasalts of the Navysh Subformation of the Ai Formation, which were originally thought to be of similar age. However, based on more recent high-precision SHRIMP U-Pb zircon dating these volcanics are 1752 ± 11 Ma in age (Krasnobaev et al., 2013a, b; Puchkov et al. 2014).

5.2. Stratigraphic Implications Arising From the U-Pb Dating of the Great Taghdout Sill

Based on the previously published data, the age of the lower Anti-Atlas Supergroup is bracketed between ca. 2032 and 647 Ma (see section 2.3.). The new U-Pb SHRIMP zircon age of 1676 ± 37 Ma obtained for the herein named Taghdout sill provides a new constraint on the minimum age for the lowermost part of the lower Anti-Atlas Supergroup in the Zenaga Inlier. The Taghdout sill intrudes into the lower Anti-Atlas Supergroup and is consequently younger than the intruded sedimentary rocks (that is the Tasserda and Ifrane-n-Taghatine formations) and most likely older than the unconformably overlying Oumoula Formation. The new age indicates that the lowermost part of the lower Anti-Atlas Supergroup was likely deposited during the late Paleoproterozoic (2.03-1.64 Ga). It could potentially be related to extension during the post-Eburnean orogenic collapse and if so should be close in age to ca. 2.0 Ga. In contrast, the much younger ca. 1.67 Ga Taghdout sill, likely related to the ca. 1.65 Ga Zenaga LIP, is a part of the mantle plume event/s that occurred at the late stage of the Nuna / Columbia supercontinent assembly (Fig. 15).

The new 1676 ± 37 Ma age for the sill in the Zenga Inlier is equivalent within uncertainty to the ca. 1706 ± 7 Ma age obtained for mafic intrusions crosscutting basal quartzites in the Iggherm Inlier (Ikenne et al., 2017) and the ca. 1650 Ma age of the thick sills intruding the basal part of the lower Anti-Atlas Supergroup in the Kerdous Inlier (Youbi et al., 2019). Together, these ages indicate associated extension. Therefore, the post-Eburnean hiatus of one billion years on the WAC is no longer valid and the pre-Pan-African sediments and magmatic events are preserved along the

northern margin of the craton. The 1640-1680 Ma (this study; Youbi et al., 2019) and the ca. 1706 Ma (Ikenne et al., 2017) magmatic events indicate that extension predated deposition of the thick quartzites of the Oumoula Formation, but postdated accumulation of the Tasserda and Ifrane-n-Taghatine formations. Further geochronological work is required to determine whether the ca. 1650 Ma, ca. 1676 Ma, and ca. 1706 Ma intrusions represent separate intraplate events on the WAC or belong to a single long-lived 1.70-1.65 Ga LIP event across the WAC and potentially representing an event extending into Baltica and Laurentia (Fig. 15). Regardless, all these ages for mafic sills are additional evidence that the lower part of the lower Anti-Atlas Supergroup has an upper Paleoproterozoic age and, therefore, nearly one billion years older than previously thought.

5.3. Constraints From Detrital Zircons on the Depositional Age of the Lower Anti-Atlas Supergroup

The youngest detrital zircons from the sample TG-15-1 (the Tasserda Formation) are ca. 2049 Ma old (Fig. 13A, B), providing a maximum depositional age for the formation. This is in agreement with the geological context of the lower Anti-Atlas Supergroup: (i) the age of the succession is bracketed between 2032 ± 5 Ma and 647 ± 2 Ma; (ii) the 1676 ± 37 Ma Taghdout sill indicates that the intruded sedimentary rocks (that is the Tasserda and Ifrane-n-Taghatine formations) are older than 1676 ± 37 Ma, consistent with the minimum age constraints of ca. 1650 Ma in the Kerdous Inlier (Youbi et al., 2019) and ca. 1710 Ma in the Igherm Inlier (Ikenne et al., 2017).

The six youngest zircon grains in the sample BA21 (the Agoummy Formation) yielded a late Paleoproterozoic concordia age of 1818 ± 6.4 Ma (Fig. 13C, D), providing the maximum age for deposition of this formation. The U-Pb detrital zircon data obtained from this sample are consistent with the U-Pb zircon data for quartzites of the Mimount Formation in the Siroua Inlier (*sensu* Thomas et al., 2002), which is the stratigraphic equivalent of the Oumoula Formation (*sensu* Bouougri and Saquaque, 2004) of the lower Anti-Atlas Supergroup. For the Mimount Formation, Abati et al. (2010) obtained LA-ICP-MS $^{207}\text{Pb}/^{206}\text{Pb}$ zircon ages from ca. 2936 to ca. 1809 Ma, with

major peaks at ca. 2200 and 2086 Ma and a number of minor peaks (ca. 2005, 1950, and 1818 Ma). The youngest zircon in the Oumoula Formation gave a concordant age of 1809 ± 15 Ma (Abati et al., 2010), while the youngest population age is ca. 1818 Ma, providing a maximum age for deposition of the Oumoula Formation. For the Mimount Formation, Walsh et al. (2012) obtained a range of SHRIMP $^{207}\text{Pb}/^{206}\text{Pb}$ zircon ages from about 2182 to 1987 Ma, with a major peak at ca. 2050 Ma. The similarity between detrital zircon age distribution in our sample BA21 from the Agoumy Formation and that of the Mimount Formation in Abati et al. (2010) and Walsh et al. (2012) indicates a similar provenance for the Mimount (= Oumoula) and Agoumy formations. The lack of ca. 1.67 Ga zircon grains in samples from the Mimount and Agoumy formations is worth a note given the prominence of the 1650 Ma Zenaga event (see Section 5.1). However, mafic melts are usually poor in zircons; this may explain their relative absence in these formations. The youngest detrital zircon grains from the Tachdamt Formation which overlap within errors are ca. 1060 Ma old (Fig. 14A, B). However, there are two younger single zircon grains with ages at 930 ± 12 and 822 ± 6 Ma. The latter age has a concordance of only 92% and a very low Th/U ratio, therefore the maximum age of sedimentation is constrained by ca. 1060 Ma and ca. 930 dates, consistent with ca. 883 Ma U-Pb SHRIMP zircon ages for two tuff horizons interbedded in the basal part of the Tachdamt Formation in the Siroua Inlier (Youbi et al., 2018, Bouougri et al., 2020). The youngest concordant zircon dates that overlap within errors in the sample BA24 from the Bleida Formation yield a concordia age of 1855 ± 6.5 Ma. The two youngest zircon grains provide dates at 1276 ± 14 Ma and 977 ± 10 Ma (Fig. 14C, D). However, Letsch (2018) reported a pronounced bimodal age distribution for a greywacke from the Bleida Formation in the Bou Azzer Inlier with a major Paleoproterozoic population (peaked at ca. 2060 Ma) and a minor Neoproterozoic one (with a prominent mode at ca. 690 Ma); the latter defines, along with the 697.3 ± 2.2 Ma maximum depositional age of the metasiltstone from the most basal part of the Bleida Formation (Bouougri et al., 2020), a Cryogenian maximum depositional age for this formation

(although the minimum depositional age is younger than 883 Ma). These data imply a significant hiatus (697 to <883 Ma) in time of deposition between the Tachdamt and Bleida formations.

Both the Bleida and Tachdamt formations contain Mesoproterozoic detrital zircon ages (Fig. 14), however these ages are less abundant in the Bleida Formation, suggesting a change in the provenance consistent with a major hiatus in the time of deposition for these units. Importantly, detrital zircon ages indicate that the provenance for the lower Anti-Atlas Supergroup was overwhelmingly Paleoproterozoic and Mesoproterozoic in age with a much smaller contribution from Archean and Neoproterozoic sources.

5.4. Interpretation of Detrital Zircon Age Clusters

Four major detrital zircon age clusters have been obtained in this study: 3.0-2.4, 2.3-1.74, 1.7-1.4, and 1.25-0.95 Ga (Fig. 16). The few Archean zircon grains, forming a poorly defined cluster, were likely derived from the Reguibat and Leo-Man shields (Rocci et al., 1991; Thiéblemont et al., 2004; Key et al., 2008; Koffi et al., 2020). The second cluster (2.30-1.74 Ga) reflects contribution from mafic and felsic magmatic intrusions emplaced during the Eburnean Orogeny and its aftermath along the northern margin of the WAC (Abouchami et al., 1990; Boher et al., 1992; Egal et al., 2002; Schofield et al., 2006; Kouyaté et al., 2013; Youbi et al., 2013 Baratoux et al., 2011; 2019; McFarlane et al., 2019). The third cluster (1.7-1.4 Ga) is likely linked to the emplacement of dykes and sills accompanying the protracted rifting and the breakup of the Nuna / Columbia supercontinent (Youbi et al., 2013; Evans et al., 2013). Zircons of the fourth cluster (1.25-0.95 Ga) are common in the early Neoproterozoic sedimentary rocks. It is thought that these zircons were derived from the 1.25-0.95 Ga Grenvillian arcs (Rino et al., 2008) that underwent uplift and erosion during and in the aftermath of the amalgamation of the Rodinia supercontinent, and they may have been transported for thousands of kilometres inside the supercontinent by rivers (Rainbird et al., 1997). Interestingly, metamorphic and intrusive units with these ages are common in the Sunsás Orogen (e.g. Sadowski and Bettencourt, 1996; Santos et al., 2008), which borders the Andean side of the Amazonian Craton (Fig. 17), and the Sveconorwegian Orogen of the northwestern part of

Baltica (e.g. Andersson et al., 2007; Christoffel et al., 1999; Cosca et al., 1998) that were adjacent to the WAC in the Rodinia supercontinent configuration.

5.5. Speculations on the Provenance and Paleogeographic Reconstruction

The main feature of our detrital zircon populations is the general absence or scarcity of Mesoproterozoic ages (Figs. 13 and 14), which is consistent with the earlier studies that argued for a tectonomagmatically quiescent period between ca. 1.7 and 1.0 Ga (e.g. Ennih and Liégeois, 2008).

The absence or scarcity of Mesoproterozoic ages has been considered diagnostic in distinguishing the WAC provenance as a possible source (e.g. Nance and Murphy, 1994; Friedl et al., 2000; Linnemann et al., 2004; Abati et al., 2010; Bahlburg et al., 2010; Gärtner et al., 2013, 2016, 2017). How could this be reconciled with the U-Pb geochronological evidence for major mafic magmatic events at ca. 1.75, 1.65, and 1.41-1.38 Ga (e.g. Youbi et al., 2013) ? The simplest explanation is that the mafic magmatism does not produce a significant detrital zircon population, and hence these major LIP events are not reflected in the detrital zircon record.

Samples from the upper part of the lower Anti-Atlas Supergroup (the Tachdamt and Bleida formations) contain Mesoproterozoic detrital zircons. Considering that in most reconstructions of the Nuna-Columbia and Rodinia supercontinents, the Amazonian Craton and Baltica were adjacent to the WAC (e.g. Hoffman, 1991; Dalziel, 1991, 1997; Weil et al., 1998; Evans, 2013; Li et al., 2013; Johansson, 2009; 2014; Merdith et al., 2017) and had their sources of Mesoproterozoic zircons far from their borders with the WAC (Fig. 15), both of these cratons could have been potential sources of detrital zircons as long as a long-distance fluvial transport was involved. However, the Amazonian Craton (Tassinari and Macambira, 1999) is characterized by a continuous range of Mesoproterozoic ages since it undergone several late Paleoproterozoic to Mesoproterozoic orogenic events, and igneous and metamorphic units with the ages ranging from 2000 to 1000 Ma are common there (e.g. Santos et al., 2000; Tassinari et al., 2000). While the sample ALB29 is indeed characterized by a continuous range of Mesoproterozoic ages, the sample BA24 has only few Mesoproterozoic zircons. Tantalizingly, similar to the sample ALB29 range of

Mesoproterozoic detrital zircon ages was observed for the ca. 1.1 Ga Polissya Series in the western part of the Ukrainian Shield (SW part of Baltica (Sarmatia); Shumlyanskyy et al., 2015) that was positioned adjacent to the WAC in the configuration of Fig. 15, but equally lacks sources for the Mesoproterozoic detrital zircons. A long-distance transport from NW Baltica has thus been inferred for this mode in detrital zircon ages based on the combined ages and Hf isotope systematics. Mesoproterozoic detrital zircon ages were also found in the Neoproterozoic sedimentary units in the Mauritanides and in the Taoudeni Basin (Straathof, 2011; Nicoll et al., 2010; Bradley et al., 2015; see for compilation Gärtner et al., 2017), indicating long-distance fluvial transport of Mesoproterozoic zircons in the aftermath of the Grenville Orogeny (cf. Rainbird et al., 1997).

5.6. Refining Proterozoic Stratigraphy of the Anti-Atlas Belt

Figure 18 shows a tentative correlation of the lower Anti-Atlas Supergroup (*sensu* Thomas et al., 2004) among the Anti-Atlas inliers. Based on our own geochronologic data and on published U-Pb geochronology, as well as on lithostratigraphy, we propose to abandon the Lkest-Taghdout Group name since it combines genetically unrelated successions deposited in different, superimposed basins recording different tectonic settings and bracketed by long-lasting hiatuses. Instead, we propose to use the lower Anti-Atlas Supergroup as the highest stratigraphic hierarchy unit for the Proterozoic, pre-Pan-African successions of the Anti-Atlas Belt that can be further subdivided in five unconformity-bounded lithostratigraphic units.

(1) The lower lithostratigraphic unit is Paleoproterozoic in age (ca. 2030-1706 Ma) and comprises the Tasserda and Ifrane-n-Taghatine formations, named here the Tasserda-Taghatine Group. Our maximum depositional age for the Tasserda Formation (<2049 Ma) agrees well with the only other detrital zircon study for the Tasserda and Ifrane-n-Taghatine formations by Letsch (2018), who inferred <2038-1975 Ma maximum depositional age for this succession. The lower group is herein related to the Eburnean Orogen collapse at ca. 2.0 Ga and deposition in extensional intracratonic basins.

(2) The second lithostratigraphic unit is also Paleoproterozoic in age (ca. 1745-1650 Ma) and comprises the Oumoula (= Mimount) Formation. In the Ifni Inlier, the stratigraphic equivalent of the lower Anti-Atlas Supergroup, the “Groupe des Quartzites” (Jeannette et al., 1981), consists of quartzites and metasiltstones that have been subdivided into three units that unconformably overlie the Eburnean basement. U-Pb dating of the Eburnean basement in this inlier yielded 2042 ± 5 Ma and 2026 ± 7 Ma ages for the Alouzad and Sahel granites, respectively (Schulte et al., 2016). The detrital zircons from the Ifni Inlier quartzites yielded ages ranging between ca. 2020 and 1745 Ma with no evidence of the Mesoproterozoic ages (Benziane et al., 2016). We tentatively correlate the Ifni Inlier succession with the Oumoula Formation based on a similar detrital zircon distribution patterns.

In the Agadir Melloul and south Siroua inliers, the stratigraphic equivalents of the lower Anti-Atlas Supergroup are the Iguiguil-Tizi n'Taghatine and Ahmamad formations, that both unconformably overlie the Eburnean granites and diorites with ca. 2.04-2.03 Ga U-Pb zircon ages (Choubert et al., 1992; Blein et al., 2014b). The Iguiguil-Tizi n'Taghatine Formation (Fig. 18) contains the Jbel Iguiguil quartzites that are sedimentologically similar to the quartzites of the Oumoula Formation (*sensu* Bouougri and Saquaque, 2004) and Mimount Formation (*sensu* Thomas et al., 2002; 2004). The Ahmamad Formation (Fig. 18) consists of quartzites and locally developed carbonate rocks exposed along the strike-slip faults in the southeastern part of the Azaghar-Izazen Inlier (Blein et al., 2014b and references therein). Both the Iguiguil-Tizi n'Taghatine and Ahmamad formations appear to have a similar age, although spatial (and structural) relationships are not clear (Choubert et al., 1992; Blein et al., 2014b). The Ahmamad Formation is similar to the Oumoula Formation in Jbel Iguiguil. The Iguiguil-Tizi n'Taghatine Formation is intruded by 570 ± 7 Ma dyke swarms and sill complexes of dolerites and gabbros (U-Pb zircon age; Blein et al., 2014b).

The Agadir Melloul Inlier has several ridges with nearly vertical conglomeratic quartzite beds sandwiched in Paleoproterozoic granites. Detrital zircon dates from the quartzites yielded the major

807 peak at ca. 2.0 Ga and the youngest ages at ca. 1.8-1.7 Ga (Soulaïmani et al., 2019), similar to the
808 distribution trend shown by detrital zircons from the the Oumoula Formation.

809 In the Kerdous Inlier, the stratigraphic equivalent of the lower Anti-Atlas Supergroup is the Jbel
810 Lkest Group and includes the whole section of the “PII” quartzites in the Anzi District
811 (Hassenforder, 1987; O’Connor et al., 2010). The thick, hematite-rich quartzite sequence is not
812 subdivided because of its monotonous nature and the lack of preservation of original textures due to
813 extensive recrystallization, however it seems to correspond to the same depositional facies as the
814 Oumoula Formation (*sensu* Bouougri and Saquaque, 2004) and Mimount Formation (*sensu* Thomas
815 et al., 2002; 2004) quartzites. The great thickness of this formation and its homogeneity imply a
816 constant supply of sediment sustained for a long period of time balanced by a continuous
817 subsidence. Letsch (2018) constrained the maximum depositional age for the quartzite to be 1780
818 Ma based on U-Pb LA-ICP-MS detrital zircon data. As noted above (Section 5.2.), Youbi et al.,
819 (2019) recently dated two mafic sills emplaced into the quartzite. One of the sills yielded an upper
820 intercept date of ca. 1640 Ma, constrained by ID-TIMS data on three zircon fractions. LA-ICP-MS
821 data for the same sample yielded a weighted mean average $^{207}\text{Pb}/^{206}\text{Pb}$ date of ca. 1650 Ma,
822 identical within analytical uncertainties to the ID-TIMS date. Only a single fraction of baddeleyite
823 has been analyzed from the second sill, indicating a similar crystallization age of ca. 1650 Ma.

824 The unconformably bounded Oumoula Formation is thus bracketed in age between ca. 1745 and
825 1650 Ma (Fig. 18). The data suggest that there was a pronounced change in the provenance across
826 the unconformity between the Oumoula and the underlying Tasserda-Taghatine Group.
827 Furthermore, the underlying units show a different structural pattern from the Oumoula Formation
828 in the Zenaga Inlier (cf. Choubert et al., 1951; Brabers, 1988; Letsch, 2018), suggesting that they
829 were deposited in different basins. Discontinuous distribution of the lower group formations and the
830 Oumoula Formation, combined with geochemical composition of the 1.75-1.64 Ga mafic dykes and
831 sills akin to continental flood basalts (Section 4.1.4.), indicate deposition of the Oumoula Formation
832 at ca. 1.75-1.64 Ga in the compartmentalized extensional basins (rifts).

833 (3) The third lithostratigraphic unit is Paleoproterozoic to Neoproterozoic in age ($< \text{ca. } 1706\text{-}1640$
834 Ma to $> \text{ca. } 883 \text{ Ma}$). It incorporates the Ifarkhs-n-Tirsal, Wanimzi, Tamgarda, Agoumy, and Imi
835 n-Tizi formations. This lithostratigraphic unit is named here the Tizi n'Taghatine Group. Our
836 detrital zircon ages for the Agoumy Formation provide the first maximum depositional age for
837 this succession ($< 1818 \text{ Ma}$), but unfortunately it does not improve independent age constraints for
838 this group. $\text{Ca. } 1650 \text{ Ma}$ sills and dykes that intrude the underlying Oumoula Formation, but not the
839 Tizi n'Taghatine Group provide the maximum depositional age and $\text{ca. } 883$ volcanics in the
840 unconformably overlying Tachdamt Formation set the minimum depositional age for this
841 succession. The succession was likely deposited in the intracratonic basin related to the rifting
842 within the assembling Rodinia supercontinent.

843 (4) The fourth lithostratigraphic unit is the $\text{ca. } 883 \text{ Ma}$ Tachdamt Formation. The Tachdamt
844 Formation consists of a succession of basaltic lava flows interbedded with subordinate pyroclastic
845 rocks. Youbi et al. (2018) and Bouougri et al., (2018; 2020) dated two tuff horizons, interbedded in
846 the basal part of the Tachdamt Formation in the Siroua Inlier, at $885 \pm 5.9 \text{ Ma}$ and $883 \pm 2.3 \text{ Ma}$ (U-
847 Pb SHRIMP on zircon). The NE30°-trending Iguerda-Taïfast mafic dyke swarm in the Zenaga and
848 Iguerda-Taïfast inliers and the coeval volcanics of the Tachdamt Formation are part of the $\text{ca. } 885\text{-}$
849 883 Ma Iguerda-Taïfast Large Igneous Province (LIP) in the Anti-Atlas Belt of Morocco (Kouyaté
850 et al., 2013; Söderlund et al., 2013; Youbi et al., 2013, 2018; Bouougri et al., 2020), related to the
851 rifting within the Rodinia supercontinent. The formation was likely deposited in the rift basin at the
852 late stage in the assembly of the supercontinent Rodinia, but it remains uncertain whether this basin
853 evolved into a passive continental margin setting along the northern margin of the WAC. Similar-
854 age LIP events have been recognized in the southern part of the WAC (Baratoux et al., 2019), and
855 on the Tarim Terrane of North China (Wang et al., 2015; Zhang et al., 2019) and the Congo Craton
856 (Armstrong et al., 2005; Johnson et al., 2007).

857 (5) The uppermost unit, the $\text{ca. } 700 \text{ Ma}$ Bleida Formation is characterized by fine-grained and thin-
858 bedded siliciclastic turbidites, interbedded with minor basaltic lava flows, exhalative facies (jasper

and ironstone), and black shales. Depositional setting of the Bleida Formation and the minimum age of its detrital zircons point to deposition in a foreland basin developed in the lead of the ca. 660-680 Ma Pan-African Orogeny (Thomas et al., 2004; Bouougri et al., 2020).

5.7. Potential correlation with the Taoudeni Basin succession in Mauritania

Our study indicates that the lower Anti-Atlas Supergroup records Mid-Proterozoic protracted and episodic deposition in intracontinental basins developed in association with extension and LIPs events on the WAC. Considering that these conditions, likely linked to far-distant tectonic stresses and emplacement of mantle plumes, could also have affected other parts of the WAC, we explore here correlation with another Mid-Proterozoic basin on the WAC, the Taoudeni Basin, located approximately 500 km to the south of the Anti-Atlas Belt (Fig. 1). It contains three unconformity-bounded successions in the ascending order: the Char, Atar, and Assabet el Hassiane groups.

The lower part of the sedimentary fill of the Taoudeni Basin, the Hodh Supergroup (Super-groupe du Hodh), is subdivided into the three unconformity-bound groups: the Char, Atar, and de l'Assabet el Hassiane (Moussine-Pouchkine and Bertrand-Sarfati, 1997; Benan and Deynoux, 1998). The mixed carbonate-siliclastic Char Group was deposited in a fault-bounded, compartmentalized basin developed on the deeply weathered Eburnean basement with aeolean and fluvial redbeds grading upsection to tidally influenced, marine siliciclastic and carbonate deposits with some evidence (halite pseudomorphs, mudcracks and mudchips, dessication cracks) for warm and arid climate. The depositional and tectonic setting of the Char Group is similar to that of the Tasserda-Taghatine Group (lithostratigraphic unit 1 in the Anti-Atlas Belt). Similarly, both successions carry detrital zircons with youngest ages corresponding to the Eburnean basement with an additional contribution in case of the Zenaga Inlier of the Anti-Atlas Belt of slightly younger, ca. 2042-1975 Ma provenance (Bradley et al., 2015; Straathof, 2011; Letsch, 2018; this study). We provisionally correlate these successions and link the development of their corresponding basins to the post-Eburnean orogen collapse.

884 We recognize no correlative stratigraphic interval in the Taoudeni Basin to the ca. 1745-1650 Ma
 885 Oumoula Formation (lithostratigraphic unit 2 of the Anti-Atlas Belt), which we link to extension
 886 and LIP emplacement on the WAC at the final stage in the assembly of the Nuna supercontinent.
 887 Importantly, ca. 1750 and 1650 Ma mafic magmatism, linked with deposition of the Oumoula
 888 Formation, was not so far recognized in the Taoudeni Basin and the surrounding Reguibat Craton.
 889 The next potentially correlative stratigraphic intervals in the Anti-Atlas Belt and the Taoudeni Basin
 890 are the Tizi n'Tghatine (lithostratigraphic unit 3) and Atar groups, respectively. Both are mixed
 891 carbonate-siliciclastic successions deposited in intracratonic basins in warm and arid climate
 892 (Leblanc and Moussine-Pouchkine, 1994; Moussine-Pouchkine and Bertrand-Sarfati, 1997). Our
 893 youngest detrital zircon ages for the Agoumy Formation (ca. 1818 Ma) correspond well with the
 894 youngest detrital zircon ages reported for the Atar Group (1804 ± 33 Ma, Straathof, 2011; Nicoll et
 895 al., 2010; Bradley et al., 2015). Re-Os isochron ages for black shales from the Atar Group yielded
 896 the late Mesoproterozoic depositional age (1106 ± 12 Ma, 1109 ± 22 Ma, and 1105 ± 37 Ma;
 897 Rooney et al., 2010). These Mesoproterozoic ages are confirmed by the study of microfossils
 898 (Beghin et al., 2017) and chemostratigraphy (Kah et al., 2012). Carbonates of the the Tizi
 899 n'Tghatine Group have highly negative to near zero carbon isotope values (-5.0 to $+1.5$ ‰; Alvaro
 900 et al., 2014). In comparison, carbonates of the Atar Group show large carbon isotope oscillations
 901 from -6.0 to $+4.0$ ‰ (Fairchild et al., 1990; Shields et al., 2007; Kah et al., 2012) that are
 902 characteristic of the latest Mesoproterozoic to early Neoproterozoic seawater.
 903 It is uncertain whether there is a stratigraphic equivalent of either the ca. 883 Ma Tachdamt
 904 Formation / Bleida formation (lithostratigraphic units 4 and 5) in the Taoudeni Basin, but the
 905 siliciclastic Assabel el Hassiane Group yielded youngest ages for detrital zircons of 952 ± 38 Ma
 906 and 965 ± 29 Ma (Straathof, 2011; Bradley et al., 2015), which are consistent with the youngest
 907 ages for detrital zircons from the Tachdamt Formation (see Fig. 18). More importantly, both the
 908 Tachdamt Formation and the Assabel el Hassiane Group show influx of Mesoproterozoic detrital
 909 zircons in the aftermath of the Grenville Orogeny. Similar pattern for the Laurentia has been related

to giant fluvial systems that delivered material from the Grenville Orogen inside the Rodinia supercontinent (Rainbird et al., 1997, 2012; Rainbird and Young, 2009). Kuznetsov et al. (2017) suggested the Rodinia supercontinent configuration ringed by continental arcs and with predominantly internal runoff that seems to fit well to the detrital zircon patterns of the WAC.

6. Concluding remarks

The main conclusions of our study are summarized below.

(1) The lower Anti-Atlas Supergroup was considered as a late Neoproterozoic passive continental margin sequence deposited during rifting and breakup on the northern margin of the WAC, culminating with opening of an oceanic basin between the northern edge of the WAC and an unknown terrane at ca. 760 Ma. Based on U-Pb geochronology data (this work and literature data) and lithostratigraphic constraints, we propose a new lithostratigraphic framework for the Proterozoic lower Anti-Atlas Supergroup of the Anti-Atlas Belt. U-Pb geochronologic data indicate that deposition of the lower Anti-Atlas Supergroup lasted from the middle Paleoproterozoic to late Neoproterozoic with significant hiatuses along unconformities. Five major, unconformity-bounded lithostratigraphic units are distinguished in the lower Anti-Atlas Supergroup. The lower unit is the ca. 2030-1706 Ma Tasserda-Taghatine Group and comprises the Tasserda and Ifrane-n-Taghatine formations. The second unit is the ca. 1745-1650 Ma Oumoula (Mimount) Formation. The third unit is Paleoproterozoic to Neoproterozoic in age (the ca. 1706-1640 Ma to > ca. 883 Ma Tizi 'n'Taghatine Group that incorporates the Ifarkhs-n-Tirsal, Wanimzi, Tamgarda, Agoumy, and Imi n-Tizi formations. The fourth unit is the ca. 883 Ma Tachdamt Formation and the uppermost unit is the ca. 700 Ma Bleida Formation.

Using this lithostratigraphic framework in combination with ages of LIPs on the WAC and herein developed correlation with the Taoudeni Basin succession highlights protracted and yet episodic mid-Proterozoic sedimentation in intracratonic basins on the WAC in association with extensional events and far-distance plate-boundary stresses that lasted from the early stage in the Nuna / Columbia assembly to the early stage in the breakup of the Rodinia supercontinent. Deposition of

the Tasserda-Taghatine Group is potentially linked to the post-Eburnean orogenic collapse. The Oumoula (Mimount) Formation might reflect the intracontinental extension within the Nuna /Columbia supercontinent in response to either ca. 1750 Ma or 1650 Ma LIP event. The Tizi ‘n’Taghatine Group might be correlative to the ca. 1.1 Ga Atar Group (Taoudeni Basin). The Tachdamt Formation records rifting within the Rodinia supercontinent, but it is uncertain whether this event led to the development of the passive continental margin along the northern boundary of the WAC. Finally, the Bleida Formation deposited in the foreland basin records the early stage in the assembly of the Gondwana supercontinent. The WAC formed the core of the Nuna / Columbia supercontinent and transited to the Rodinia supercontinent directly unaffected by the supercontinent breakup and subsequent assembly.

(2) The maximum depositional age and provenance of the lower Anti-Atlas Supergroup are constrained by new detrital zircon ages from the lower, middle, and upper parts of the sedimentary successions at the Zenaga and Bou Azzer El Graara inliers. The inferred maximum depositional ages are: ca. 2049 Ma for the Tasserda Formation, ca. 1818 Ma for the Agoumy Formation, ca. 1060 Ma for the Tachdamt Formation, and ca. 700 Ma for the Bleida Formation. Four detrital zircon age clusters have been defined in this study: ca. 3.0-2.4, 2.3-1.74, 1.7-1.4, and 1.25-0.95 Ga. The provenance of detrital zircon was mainly on the WAC (that is the Reguibat and Leo-Man shields), but other cratons such as Amazonia, Baltica, and the Sahara Metacraton (Hoggar, Mali) cannot be excluded as possible subsidiary source areas. Importantly, we find that subsequent to the Grenville Orogeny, long-distance transport by fluvial systems delivered onto the WAC Mesoproterozoic zircons either from Fennoscandia or Baltica or Sunsas Belt of Amazonia (see Fig. 15). Transport from either of these two areas requires giant fluvial systems directed inward of the Rodinia supercontinent.

(3) The Tachdamt Formation (lithostratigraphic unit 4) consists mainly of a succession of basaltic lava flows. A ca. 885 Ma mafic dyke swarm in the Zenaga and Iguerda-Taïfast inliers and the coeval volcanics of the Tachdamt Formation constitute a LIP in the Anti-Atlas Belt of Morocco,

named the Iguerda-Taïfast LIP. Similarly aged LIPs also occur on other cratons such as the Tarim Terrane (North China) and Congo Craton (Central Africa) and might reflect rifting at the late stage in the assembly of the Rodinia supercontinent.

Acknowledgements

Most of this work was carried out at the Department of Geology of the Faculty of Sciences-Semlalia, Cadi Ayyad University of Marrakech and the “Centro de Pesquisas Geocronológicas (CPGeo), Instituto de Geociências (IG), Universidade de São Paulo-USP, São Paulo (SP), Brazil” within the framework of the scientific and technical agreement between the São Paulo University and the Cadi Ayyad University (responsibles Colombo Celso Gaeta Tassinari and Nasrddine Youbi). We acknowledge Academy Hassan II for Science and Technology for funding studentship for Abdelhak Ait Lahna in 2016 to 2018. Financial support for this work was also provided by several research projects: (i) Academy Hassan II for Science and Technology (HIIAST/SDU/2016.02) to Nasrddine Youbi and co-workers; (ii) Medyna; and (iii) FCT (Portugal)-CNRST (Morocco) to Nasrddine Youbi and João Mata. Richard E. Ernst was partially supported from Mega-Grant 14.Y26.31.0012 of the Russian Federation. João Mata acknowledges the financial FCT support through project UIDB/50019/2020 – IDL. The Editor Prof. Damien Delvaux is gratefully acknowledged for his comments, patience, and support. We greatly appreciate thoughtful comments of reviewers Prof. David Evans and Dr. Shuan-Hong Zhang that helped to improve the manuscript.

Figure captions

Fig. 1. (A) Location of Northwestern Africa; (B) Geological map of Northwestern Africa (after Thiéblemont, 2016 and Brahimi et al., 2018); (C) Distribution of main tectonic elements (after Brahimi et al., 2018); and (D) Geological map of the Anti-Atlas Belt in southern Morocco showing

the studied area in the Zenaga and Bou Azzer El Graara inliers (after Gasquet et al., 2008).

Abbreviations: G.- Groups; J.- Jbel; Paleoprot.- Paleoproterozoic; T.-Tizi

Fig. 2. Generalized lithostratigraphic column for the Anti-Atlas Pan-African orogen. After Thomas et al., (2004), modified by Gasquet et al., (2005; 2008) and Youbi et al., (2013), and taken from Youbi et al., (2013). The conventional viewpoint has been that there is a gap in magmatic activity between 1.7 and 1.0 Ga in the Anti-Atlas Belt (Mesoproterozoic gap). However, this interpretation was challenged with the results of Youbi et al. (2013), El Bahat et al. (2013), Kouyaté et al. (2013), Söderlund et al. (2013), which all indicated important intraplate magmatic events at ca. 1.75, 1.65, and 1.38-1.41 Ga. See the text for more explanation. Note that this figure is the outdated correlation scheme for the Anti-Atlas Belt since it uses outdated chronology of the Anti-Atlas cover successions. Abbreviations: PI (XI), PII (XII), PII-III (XII-III) and PIII (XIII) are the classical stratigraphic symbols used on Anti-Atlas geological maps (Choubert, 1963). Adoud.- Adoudounien (or Adoudounian); Calc. Sup.- “Calcaires Supérieures” (Upper Limestones) ; Cambr.- Cambrian; HKCA- High-K calc-alkaline (granitoids); L. d. Vin - Lie de Vin (wine-coloured) ; Paleoproteroz.- Paleoproterozoic.

Fig. 3. Lithostratigraphic subdivision, sequence stratigraphy, and paleogeographic data for the lower Anti-Atlas Supergroup (modified after Bouougri and Saquaque, 2004). We use the name of the Ifrane-n-Taghatine Formation instead of the Taghdout Formation to avoid confusion with the widely used name of the group - “the Taghdout Group” in the Zenaga Inlier. Abbreviations: systems tracts consist of (LST) lowstand system tracts, (TST) transgressive system tracts, and (HST) highstand system tracts. U1 to U5 are unconformities. DS1 to DS5 are depositional systems.

Fig. 4. Geology of the Zenaga Inlier (modified after Choubert, 1963; Ennih and Liégeois, 2001; Thomas et al., 2002; Kouyaté et al., 2013; Youbi et al., 2013). The Taghdout Group corresponds to

the lower Anti-Atlas Supergroup (Thomas et al., 2004). Inset shows a rose diagram that summarizes mafic dyke trends measured from the map and Google Earth; n is the number of measurements. The U-Pb ages obtained in this study are indicated with asterisks; other age constraints are from literature (Kouyaté et al., 2013; Youbi et al., 2013).

Fig. 5. Simplified geological map of the Bou Azer-El Graara Inlier showing the main lithotectonic units. The localities (Tachdamt and Bleida) mentioned in the text are indicated, as well as the location of samples used for U-Pb detrital geochronology (after Blein et al., 2014a). The Bleida Group corresponds to the lower Anti-Atlas Supergroup (Blein et al., 2014a)

Fig. 6. (A) General view of the unconformity between the Eburnean basement and the overlying quartzites of Oumoula Formation (O Fm.) of the lower Anti-Atlas Supergroup at Tizi n'Taghatine locality. The Eburnean basement (EB: gneiss and schists) is crosscut by pegmatite dyke (PE). The arrow indicates the road from Taliwine to Tazenakht. Geologists are Fida Medina, Maria Giuditta Fellin, Claudio Faccenna, and Riccardo Lanari; (B) View of the unconformity between the schists of the Bleida Formation (B Fm.) and the overlying volcanic rocks of the Ouarzazate Supergroup (OS); (C) View of the Taghdout sill (TS) crosscutting the mixed siliciclastic-carbonate succession of the Ifrane-n-Taghatine Formation (IT Fm.) in the Zenaga Inlier near the Taghdout Town. Note hornfels (h) at the baked margin of the sill. The Ifrane-n-Taghatine Formation is overlain by thick quartzites of Oumoula Formation (O Fm.); and (D) Detailed view of the upper contact between the Taghdout dolerite sill (TS) and limestones of the Ifrane-n-Taghatine Formation (IT Fm.) transformed into hornfels.

Fig. 7. Thin-section photomicrographs of the Great Taghdout sill. (A) coarse-grained dolerite in the centre of the sill with plagioclase (Pl) and clinopyroxene (Cpx) minerals and oxides (Ox), crossed nicols; (B) fine-grained, microlitic porphyric texture at the border of the sill with microlites (tiny

crystals) of plagioclase (Pl) and clinopyroxene (Cpx) minerals and oxides (Ox). Epidote (Ep) is filling vesicles; crossed nicols.

Fig. 8. Th/U ratios versus concordant U-Pb ages of igneous and detrital zircon grains of the lower Anti-Atlas Supergroup.

Fig. 9. Zircon U-Pb SHRIMP geochronology: (A) Wetherill concordia plots of U-Pb zircon data for the Taghdout sill (sample TA) of the Zenaga Inlier. (B) Weighted mean average $^{207}\text{Pb}/^{206}\text{Pb}$ age of analysis with low Th and U concentrations (spots 2.1, 7.1, and 13.1).

Fig. 10. (A) Total alkali vs. silica (TAS) diagram for the chemical classification and nomenclature (after Le Bas et al., 1986). Subalkaline vs. alkaline series boundary (dotted line) is from Irvine and Baragar (1971). (B) Ti vs. V variations in samples from the Zenaga, Agadir Melloul, Iguerda-Taïfast, and Tagragra of Akka inliers including four datasets reported by Mortaji (1989, unpublished thesis), Hafid (1999, unpublished thesis), Hafid et al., (2001) and El Aouli et al., (2010); diagram is after Shervais (1982). The studied samples TA1 (centre of the sill dated at 1676 ± 37 Ma; this work) and TA2 (border of the sill) and the 1.6 Ga dolerite dykes and sills from the Zenaga and Agadir Melloul inliers collected by Kouyaté et al., (2013) are also shown on this diagram.

Fig. 11. Primitive mantle-normalized trace element 'spidergram' for the dated mafic sill and dykes of the Zenaga and Agadir Melloul inliers. Primitive mantle values used for normalization are from Sun and McDonough (1989).

Fig; 12. Thin section photomicrographs of the sedimentary samples. (A) Metasiltstones of the Tasserda Formation: sample TG-15-1; purple, coarse-grained and quartz-rich metasiltstone with

quartz (Qz) and feldspar (Pl) embedded in a sericite-rich matrix (Mx); crossed nicols. (B) Quartzites of the Agoumy Formation: sample BA21, with quartz grains that are mostly subrounded. (C) Quartzites of the Tachdamt Formation: sample ALBL29, with a mineral assemblage comprised mainly of quartz (Qz) associated to a small amount of feldspars (Pl) with a matrix (Mx) of silt-sized quartz and mica minerals. (D) Metasandstone of the Bleida Formation: sample BA24, fine-grained, moderately sorted metasandstone with quartz (Qz), plagioclase (Pl), and lithic fragments in mica matrix (Mx).

Fig. 13. U-Pb ages for detrital zircon grains from samples TG-15-1 (metasiltstones of the Tasserda Formation) and BA21 (quartzites of the Agoumy Formation) of the lower Anti-Atlas Supergroup. (A) and (C) are concordia diagrams; (B) and (D) are combined binned frequency and probability density plots of detrital zircon grains. Abbreviations: Conc.-Concordance; Decay-const. errs.- Decay-constant errors

Fig. 14. U-Pb ages for detrital zircon grains from samples ALBL29 (quartzites of the Tachdamt Formation) and BA24 (metasandstone of the Bleida Formation) of the lower Anti-Atlas Supergroup. (A) and (C) are concordia diagrams; (B) and (D) are combined binned frequency and probability density plots of detrital zircon grains. Abbreviations: Conc.-Concordance; Decay-const. errs.- Decay-constant errors

Fig. 15. Paleomagnetically and geologically informed reconstruction of WAC, Amazonia, and Baltica cratons (SAMBA connection) along with Laurentia and Siberia in the core of the Nuna supercontinent (modified after Evans and Mitchell, 2011); additional ca. 1640-1680 Ma dyke swarms include the 1641 ± 8 Ma Nersa sill complex of Southern Siberia (U-Pb baddeleyite TIMS age; Metelkin et al, 2011; Ernst et al., 2016), 1635.0 ± 2.7 to 1622.1 ± 3.2 Ma Melville Bugt dykes of western Greenland (U-Pb baddeleyite TIMS age; Halls et al., 2011; Klausen and Nilsson, 2019),

1663 ± 8 Ma bimodal Narakay Volcanic Complex of Canada (U-Pb zircon TIMS age; Bowring and Ross, 1985), and 1642 ± 2 Ma dyke swarms of Häme, Sipoo, and Suomenniemi of the Fennoscandian Shield in southeast Finland (U-Pb ID-TIMS baddeleyite age; Vaasjoki et al., 1991; Salminen et al., 2017). Star locates mantle plume centre, proposed for the Melville Bugt swarm by Klausen and Nilsson (2019).

Fig. 16. Age compilation for detrital zircon grains from four samples collected from the lower Anti-Atlas Supergroup in the Zenaga and Bou Azzer El Graara inliers. Ages older than 1.7 Ga indicate the West African provenance (modified from Linnemann et al., 2010 and Abubaker et al., 2017). See the text for more explanation. Abbreviations: Conc.-Concordance

Fig. 17. Distribution of U-Pb ages for detrital zircon grains analyzed in this study from the lower Anti-Atlas Supergroup in the Zenaga and Bou Azzer El Graara inliers (all grouped together) and potential sources compiled from the literature. Gray stripes: detrital zircons; black stripes: zircons from magmatic rocks. Detrital zircon data sources: Pyrenees (Margalef et al., 2016); Lybia (Altumi et al., 2013); Algeria, Ahaggar (Linnemann et al., 2011); Meguma, Ganderia, and Avalonia detrital zircons (Fyffe et al., 2009; Gärtner et al., 2016; Henderson et al., 2016; Murphy et al., 2004a, 2004b; Waldron et al., 2009 and references therein); Anti-Atlas and Sardinia Cambrian (Avigad et al., 2012); Anti-Atlas Neoproterozoic (Abati et al., 2010; Karaoui et al., 2015). Anti-Atlas mafic rocks (Youbi et al., 2013 and references therein). Modified after Linnemann et al., (2011), Pratt et al., (2015), and Marzoli et al., (2017), and references therein. Abbreviations: ANS = Arabian-Nubian Shield; CAMP-Central Atlantic Magmatic Province; Camb.-Cambrian; Dev.- Devonian; Jur., Jurassic; Neoprot.- Neoproterozoic; Ord.- Ordovician; Sil.-Silurian

Fig. 18. A tentative correlation scheme of the lower Anti-Atlas Supergroup and its stratigraphic equivalents in the Anti-Atlas inliers. U1 to U8 are unconformities. U-Pb geochronology data are

from (1) this study, (2) Letsch (2018), (3) Youbi et al., (2018) and Bouougri et al., (2018; 2020), (4) Abati et al., (2010), (5) Walsh et al., (2012), (6) Thomas et al., (2002), (7) Soulaïmani et al., (2013a), Hafid et al., (2013), Blein et al., (2014b), (8) Soulaïmani et al., (2019), (9) Walsh et al., (2002), (10) Ikenne et al., (2017), (11) Aït Malek et al., (1998), (12) Youbi et al., (2019), (13) Benziane et al., (2016), and (14) Schulte et al., (2016).

Table captions

Table 1. Site location and summary of the data for the studied samples. The names of units and formations are sensu Bouougri and Saquaque (2004).

Table 2. Representative whole-rock major and trace element geochemical data for dykes and sills from the Zenaga, Agadir Melloul, Iguerda-Taïfast, and Tagragra of Akka inliers including the dated dykes, samples TA1 from the Zenaga Inlier (1676 ± 37 Ma; this work), DZ 36 (1640 ± 2 Ma and 1641 ± 3 Ma; Kouyaté et al., 2013) and DZ 40 (1656 ± 9 Ma; Kouyaté et al., 2013) from the Zenaga Inlier (Kouyaté et al, 2013), and AMHA069 (1654 ± 16 Ma; Kouyaté et al, 2013) from the Agadir Melloul Inlier. The geochemical data are from unpublished theses (Hafid, 1999; Mortaji, 1989) and published papers (Hafid et al., 2001; El Aouli et al., 2010; Youbi et al., 2013). Fe_2O_{3t} = total Fe expressed as Fe_2O_3 . Loss-on-ignition (L.O.I.) determined by sample weight loss at 1000°C .

Appendix: Supplementary data

Figure S1 of supplementary data. Cathodoluminescence (CL) images of zircon grains obtained from the sample TA 1 showing their internal texture, spots (yellow circles), and apparent ages in Ma (1σ error).

Figure S2 of supplementary data. Cathodoluminescence (CL) images of zircon grains obtained from the sample TG-15-1 (metasiltstone of the Tasserda Formation) of the lower Anti-Atlas Supergroup showing their internal texture, laser spots (yellow circles=concordant ages; red circles=discarded ages due to high discordance and/or U/Th/Pb contents), and apparent ages in Ma (1σ error).

Figure S3 of supplementary data. Cathodoluminescence (CL) images of zircon grains obtained from the sample BA21 (quartzites of the Agoumy Formation) of the lower Anti-Atlas Supergroup showing their internal texture, laser spots (yellow circles=concordant ages; red circles= discarded ages due to high discordance and/or U/Th/Pb contents), and apparent ages in Ma (1σ error).

Figure S4 of supplementary data. Cathodoluminescence (CL) images of zircon grains obtained from the sample ALBL29 (quartzites of the Tachdamt Formation) of the lower Anti-Atlas Supergroup showing their internal texture, laser spots (yellow circles=concordant ages; red circles= discarded ages due to high discordance and/or U/Th/Pb contents), and apparent ages in Ma (1σ error).

Figure S5 of supplementary data. Cathodoluminescence (CL) images of zircon grains obtained from the sample BA24 (metasandstone of the Bleida Formation) of the lower Anti-Atlas Supergroup showing their internal texture, laser spots (yellow circles=concordant ages; red circles= discarded ages due to high discordance and/or U/Th/Pb contents), and apparent ages in Ma (1σ error).

Table S1 of supplementary data. SHRIMP zircon U-Th-Pb data for the Taghdout Sill (Sample TA1).

Table S2 of supplementary data. LA-ICP-MS zircon U-Th-Pb data for the sample TG-15-1 (metasiltstone of the Tasserda Formation) of the lower Anti-Atlas Supergroup.

Table S3 of supplementary data. SHRIMP zircon U-Th-Pb data for the sample BA21 (quartzites of the Agoumy Formation) of the lower Anti-Atlas Supergroup.

Table S4 of supplementary data. LA-ICP-MS zircon U-Th-Pb data for the sample ALBL29 (quartzites of the Tachdamt Formation) of the lower Anti-Atlas Supergroup.

Table S5 of Supplementary data. SHRIMP zircon U-Th-Pb data for the sample BA24 (metasandstone of the Bleida Formation) of the lower Anti-Atlas Supergroup.

References

- Abati, J., Aghzer A.M., Gerdes A., Ennih N., 2010. Detrital zircon ages of Neoproterozoic sequences of the Moroccan Anti-Atlas belt. *Precambrian Research*, v. 181(1), p.115-128. Doi:10.1016/j.precamres.2010.05.018.
- Abati, J., Aghzer, A. M., Gerdes, A., Ennih, N., 2012. Insights on the crustal evolution of the West African craton from Hf isotopes in detrital zircons from the Anti-Atlas belt. *Precambrian Research*, 212-213, 263–274. doi:10.1016/j.precamres.2012.06.005.
- Abouchami, W., Boher, M., Michard, A. and Albarede, F., 1990. A Major 2.1 Ga event of mafic magmatism in West Africa: An early stage of crustal accretion. *Journal of Geophysical Research*, v. 95, p. 17605-17629. Doi:10.1029/jb095ib11p17605
- Abubaker, A., Hofmann, M., Gärtner, A., Linnemann, U., Elicki, O., 2017. First U–Pb geochronology on detrital zircons from Early-Middle Cambrian strata of the Torgau-Doberlug

- 1194 Syncline (eastern Germany) and palaeogeographic implications. *International Journal of Earth*
 1195 *Sciences (Geologische Rundschau)*, v. 106, p. 2445–2459. Doi:10.1007/s00531-016-1440-y
- 1196 Admou, H., and Juteau, T., 1998. Découverte d'un système hydrothermal océanique fossile dans
 1197 l'ophiolite antécambrienne de Khzama (massif du Siroua, Anti-Atlas marocain). *Comptes*
 1198 *Rendus de l'Académie Des Sciences - Series IIA - Earth and Planetary Science*, 327(5), 335–
 1199 340. doi:10.1016/s1251-8050(98)80052-9.
- 1200 Ahmed, A., Arai, S., Abdelaziz, Y., Rahimi, A., 2005. Spinel composition as a petrogenetic
 1201 indicator of the mantle section in the Neoproterozoic Bou Azzer ophiolite, Anti Atlas, Morocco.
 1202 *Precambrian Research*, 138(3-4), 225–234. Doi:10.1016/j.precamres.2005.05.004
- 1203 Aït Malek, H.A., Gasquet, D., Bertrand, J.M., Leterrier, J., 1998. Geochronologie U–Pb sur zircon
 1204 de granitoïdes éburnéens et panafricains dans les boutonnières protérozoïques d'Igherm, du
 1205 Kerdous et du Bas Draâ (Anti-Atlas occidental, Maroc). *Comptes Rendus de l'Académie des*
 1206 *Sciences, Serie II. Sciences de la Terre et des Planetes*: v. 327 (12), p. 819–826.
 1207 Doi:10.1016/s1251-8050(99)80056-1
- 1208 Altumi, M.M., Elicki, O., Linnemann, U., Hofmann, M., Sagawe, A., Gärtner, A., 2013. U–Pb LA-
 1209 ICP-MS detrital zircon ages from the Cambrian of Al Qarqaf Arch, central-western Libya:
 1210 provenance of the West Gondwanan sand sea at the dawn of the early Palaeozoic. *Journal of*
 1211 *African Earth Sciences*, v. 79, p. 74–97. Doi:10.1016/j.jafrearsci.2012.11.007
- 1212 Alvaro, J. J., Pouclet, A., Ezzouhairi, H., Soulaïmani, A., Bouougri, E. H., Imaz, A. G., Fekkak, A.,
 1213 2014. Early Neoproterozoic rift-related magmatism in the Anti-Atlas margin of the West African
 1214 craton, Morocco. *Precambrian Research*, v 255, 433–442. Doi:10.1016/j.precamres.2014.10.008
- 1215 Andersson, U. B., Rutanen, H., Johansson, Å., Mansfeld, J., Rimša, A., 2007. Characterization of
 1216 the Paleoproterozoic Mantle beneath the Fennoscandian Shield: Geochemistry and Isotope
 1217 Geology (Nd, Sr) of ~ 1.8 Ga Mafic Plutonic Rocks from the Transscandinavian Igneous Belt in
 1218 Southeast Sweden. *International Geology Review*, 49(7), 587–625. Doi:10.2747/0020-
 1219 6814.49.7.587

- Armstrong, R. A., Master, S., Robb, L. J., 2005. Geochronology of the Nchanga Granite, and constraints on the maximum age of the Katanga Supergroup, Zambian Copperbelt. *Journal of African Earth Sciences*, 42(1-5), 32–40. doi:10.1016/j.jafrearsci.2005.08.012
- Avigad, D., Gerdes A., Morag, N., Bechstädt, T., 2012. Coupled U–Pb–Hf of detrital zircons of Cambrian sandstones from Morocco and Sardinia: Implications for provenance and Precambrian crustal evolution of North Africa. *Gondwana Research*, v. 21, p.690–703. Doi:10.1016/j.gr.2011.06.005
- Bahlburg, H., Vervoort, J.D., DuFrane, S.A., 2010. Plate tectonic significance of Middle Cambrian and Ordovician siliciclastic rocks of the Bavarian Facies, Armorican Terrane Assemblage, Germany: U–Pb and Hf isotope evidence from detrital zircons. *Gondwana Research*, v. 17, p.223–235. Doi:10.1016/j.gr.2009.11.007
- Baratoux, L., Metelka, V., Naba, S., Jessell, M. W., Grégoire, M., Ganne, J., 2011. Juvenile Paleoproterozoic crust evolution during the Eburnean orogeny (~2.2–2.0Ga), western Burkina Faso. *Precambrian Research*, 191(1-2), 18–45. Doi:10.1016/j.precamres.2011.08.010
- Baratoux, L., Söderlund, U., Ernst, R. E., De roever, E., Jessell, M. W., Kamo, S., Naba, S., Perrouy, S., Metelka, V., Yatte, D., Grenholm, M., Diallo, D. P., Ndiaye, P. M., Dioh, E., Cournède, C., Benoit, M., Baratoux, D., Youbi, N., Rousse S., Bendaoud, A., 2019. New U–Pb Baddeleyite Ages of Mafic Dyke Swarms of the West African and Amazonian Cratons: Implication for Their Configuration in Supercontinents Through Time. Springer Nature Singapore Pte Ltd. 2019 R. K. Srivastava et al. (eds.), *Dyke Swarms of the World: A Modern Perspective*, Springer Geology, p. 263-314, DOI https://doi.org/10.1007/978-981-13-1666-1_7.
- Barbey, P., Oberli, F., Burg, J-P., Nachit, H, Pons, J., Meier, M., 2004. The Palaeoproterozoic in western Anti-Atlas (Morocco): a clarification. *Journal of African Earth Sciences*, v. 39, p. 39–245. Doi:10.1016/j.jafrearsci.2004.07.044

- 1245 Beghin, J., Storme, J.-Y., Blanpied, C., Gueneli, N., Brocks, J. J., Poulton, S. W., Javaux, E. J.,
 1246 2017. Microfossils from the late Mesoproterozoic-early Neoproterozoic Atar/El Mreïti Group,
 1247 Taoudeni Basin, Mauritania, northwestern Africa. *Precambrian Research*, 291, 63–82.
 1248 doi:10.1016/j.precamres.2017.01.009.
- 1249 Benan, C. A. A., Deynoux, M., 1998. Facies analysis and sequence stratigraphy of neoproterozoic
 1250 Platform deposits in Adrar of Mauritania, Taoudeni basin, West Africa. *Geologische Rundschau*,
 1251 87(3), 283–302. Doi:10.1007/s005310050210
- 1252 Benziane, F., Yazidi, A., Schulte, B., Boger, S., Stockhammer, S., Lehmann, A., Saadane, A.,
 1253 Yazidi, M., 2016. Carte géologique du Maroc, feuille au 1/50 000 Tlata Al Akhçaç. Notes et
 1254 Mém. Serv. Géol. Maroc- Rabat n° 543bis.
- 1255 Benziane, F., 2007. Lithostratigraphie et évolution géodynamique de l’Anti-Atlas (Maroc) du
 1256 Paléoprotérozoïque au Néoprotérozoïque: exemples de la boutonnière de Tagragra de Tata et
 1257 du Jbel Saghro. Ph.D. Univ. Savoie, France.
- 1258 Benziane, F., Yazidi, A., Saadane, A., Yazidi, M., El Fahssi, A., Stone, B.D., Walsh, G.J., Burton,
 1259 W.C., Aleinikoff, J.N., Ejjaouani, H., Kalai, M., 2008. Carte géologique au 1/50 000, Feuille
 1260 Qalâat Mgouna. Notes et Mémoires du Service Géologique du Maroc n° 468, 1–139.
- 1261 Bertrand, H., Dostal, J., Dupuy, C., 1982. Geochemistry of early Mesozoic tholeiites from
 1262 Morocco. *Earth and Planetary Science Letters*, 58(2), 225–239. Doi:10.1016/0012-
 1263 821x(82)90196-0
- 1264 Black, L.P., Kamo, S.L., Allen, C.M., Davis, D.W., Aleinikoff, J.N., Valley J.W., Mundil R.,
 1265 Campbell I.H., Korsch R.J., Williams I.S., Foudoulis C., 2004. Improved $^{206}\text{Pb}/^{238}\text{U}$
 1266 microprobe geochronology by the monitoring of a trace-element related matrix effect:
 1267 SHRIMP, ID-TIMS, ELA-ICP-MS and oxygen isotope documentation for a series of zircon
 1268 standards. *Chemical Geology* 205, p. 115–140. Doi:10.1016/j.chemgeo.2004.01.003
- 1269 Blein, O., Baudin, T., Chèvremont, P., Soulaïmani, A., Admou, H., Gasquet, P., Cocherie, A., Egal,
 1270 E., Youbi, N., Razin, P., Bouabdelli, M., Gombert, P., 2014a. Geochronological constraints on

- 1271 the polycyclic magmatism in the Bou Azzer-El Graara inlier (Central Anti-Atlas Morocco).
 1272 Journal of African Earth Sciences, v. 99, p.287–306. Doi:10.1016/j.jafrearsci.2014.04.021
- 1273 Blein, O., Baudin, T., Soulaïmani, A., Cocherie, A., Chèvremont, P., Admou, H., Ouanaimi, H.,
 1274 Hafi, A., Razin, P., Bouabdelli, M., and Roger, J., 2014b. New geochemical, geochronological
 1275 and structural constraints on the Ediacaran evolution of the south Sirwa, Agadir-Melloul and
 1276 Iguerda inliers, Anti-Atlas, Morocco. Journal of African Earth Sciences, v. 98, p. 47–71,
 1277 <https://doi.org/10.1016/j.jafrearsci.2014.06.019>
- 1278 Block, K. A., Steiner, J. C., Puffer, J. H., Jones, K. M., Goldstein, S. L., 2015. Evolution of late
 1279 stage differentiates in the Palisades Sill, New York and New Jersey. Lithos, 230, 121–132.
 1280 doi:10.1016/j.lithos.2015.05.018
- 1281 Bodinier, J. L., Dupuy, C., Dostal, J., 1984. Geochemistry of Precambrian ophiolites from Bou
 1282 Azzer, Morocco. Contributions to Mineralogy and Petrology, 87(1), 43–50. 87(1), 43–50.
 1283 Doi:10.1007/bf00371401
- 1284 Bogdanova, S. V., Bingen, B., Gorbatshev, R., Kheraskova, T. N., Kozlov, V. I., Puchkov, V. N.,
 1285 Volozh, Y. A., 2008. The East European Craton (Baltica) before and during the assembly of
 1286 Rodinia. Precambrian Research, 160 (1-2), 23–45. Doi:10.1016/j.precamres.2007.04.024
- 1287 Bogdanova, S.V., Gorbatshev, R., Garetsky, R.G., 2016. Europe/East European Craton. Reference
 1288 Module in Earth Systems and Environmental Sciences, Elsevier, pp. 1-18. Doi:10.1016/b978-0-
 1289 12-409548-9.10020-x
- 1290 Boher, M., Abouchami, W., Michard, A., Albarede, F. and Arndt, N. T., 1992. Crustal growth in
 1291 West Africa at 2.1 Ga. Journal of Geophysical Research, v. 97, p. 345-369.
 1292 Doi:10.1029/91jb01640
- 1293 Bouougri, E.H., Saquaque, A., 2004. Lithostratigraphic framework and correlation of the
 1294 Neoproterozoic northern Western African Craton passive margin sequence (Siroua-Zenaga Bou
 1295 Azzer El Graarara Inliers, Central Anti-Atlas, Morocco): an intergrated approach. Journal of
 1296 African Earth Sciences, v. 39, p. 227–238. Doi:10.1016/j.jafrearsci.2004.07.045

- 1297 Bouougri, E. H., Ait Lahna, A., Tassinari, C. C. G., Basei, M. A. S., Youbi, N., Admou, H.,
 1298 Maacha, L., 2020. Time constraints on Early Tonian Rifting and Cryogenian Arc terrane-
 1299 continent convergence along the northern margin of the West African craton: Insights from
 1300 SHRIMP and LA-ICP-MS zircon geochronology in the Pan-African Anti-Atlas belt (Morocco).
 1301 Gondwana Research. Volume 85, p. 169-188. doi.org/10.1016/j.gr.2020.03.011
- 1302 Bowring, S. A., Ross, G. M., 1985. Geochronology of the Narakay Volcanic Complex: implications
 1303 for the age of the Coppermine Homocline and Mackenzie igneous events. Canadian Journal of
 1304 Earth Sciences, 22(5), 774–781. Doi:10.1139/e85-082
- 1305 Brabers, P. M., 1988. A plate tectonic model for the Panafrican Orogeny in the Anti-Atlas,
 1306 Morocco. Lecture Notes in Earth Sciences, 61–80. Doi:10.1007/bfb0011586
- 1307 Bradley, D.C., O’Sullivan, P., Cosca, MA., Motts, HA., Horton, JD., Taylor, CD., Beaudoin, G.,
 1308 Lee G.K., Ramezani, J., Bradley, D.B., Jones, J.V., Bowring, S., 2015. Synthesis of Geological,
 1309 Structural, and Geochronologic Data (Phase V, Deliverable 53): Chapter A of Taylor CD (ed.),
 1310 Second Projet de Renforcement Institutionnel du Secteur Minier de la République Islamique de
 1311 Mauritanie (PRISM-II). U.S. Geological Survey Open-File Report 2013-12080-A, p. 328.
 1312 <https://doi.org/10.3133/ofr20131280A>
- 1313 Brahimi, S., Liégeois, J.-P., Ghienne, J.-F., Munsch, M., Bourmatte, A., 2018. The Tuareg shield
 1314 terranes revisited and extended towards the northern Gondwana margin: Magnetic and
 1315 gravimetric constraints. Earth-Science Reviews, 185, 572–599.
 1316 Doi:10.1016/j.earscirev.2018.07.002
- 1317 Cahen, L., Snelling, N.J., Delhal, J., Vail, J.R., 1984. The Geochronology and Evolution of Africa:
 1318 Clarendon Press, Oxford, United Kingdom, p. 512.
- 1319 Chacko, T., De, S. K., Creaser, R. A., Muehlenbachs, K., 2000. Tectonic setting of the Taltson
 1320 magmatic zone at 1.9–2.0 Ga: a granitoid-based perspective. Canadian Journal of Earth Sciences,
 1321 37 (11), 1597–1609. Doi:10.1139/e00-029

- Charlot, R., 1976. The precambrian of the Anti-Atlas (Morocco): a geochronologic synthesis: Precambrian Research, v. 3, p. 273–299. Doi:10.1016/0301-9268(76)90013-9
- Charlot, R., 1982. Caracterisation des evenements eburneens et panafricains dans l’Anti-Atlas marocain; apport de la methode geochronologique Rb–Sr : Notes et Mémoires du Service Géologique du Maroc. v. 313, 106p.
- Choubert, G., 1962. Réflexions sur les parallélismes probables des formations quaternaires atlantiques du Maroc avec celles de la Méditerranée: Quaternaria, v. 5, p.137-175.
- Choubert, G., 1963. Histoire géologique de l’Anti-Atlas de l’archéen à l’aurore des temps primaires. Notes et Mémoires du Service Géologique du Maroc, v. 62, p. 350–352.
- Choubert, G., Faure-Muret, A., Dahmani, M., Mouatani, E.A., Horrenberger, J.C., Salem, M., 1992. Carte géologique du Maroc à 1/100 000 feuille Agadir Melloul. Notes et Mémoires n° 359. Ministère de l’Energie et des Mines, Rabat.
- Choubert, G., Termier, H., Termier, G., 1951. Les calcaires précambriens de Taghdout et leurs organismes problématiques. Notes et Mémoires du Service géologique du Maroc 85 (5), 9-34.
- Christoffel, C. A., Connelly, J. N., Åhäll, K.-I., 1999. Timing and characterization of recurrent pre-Sveconorwegian metamorphism and deformation in the Varberg–Halmstad region of SW Sweden. Precambrian Research, 98(3-4), 173–195. Doi:10.1016/s0301-9268(99)00046-7
- Clauer, N., 1976. Géochimie isotopique du strontium des milieux sédimentaires. Application à la géochronologie du craton ouest-africain. Sciences géologiques, Strasbourg, v. 45, 256 p.
- Coish, R. A., and Sinton, C. W., 1992. Geochemistry of mafic dikes in the Adirondack Mountains: Implications for late Proterozoic continental rifting. Contributions to Mineralogy and Petrology, v. 110, p. 500–514. Doi:10.1007/bf00344084
- Corrigan, D., Hajnal, Z., Németh, B., Lucas, S. B., 2005. Tectonic framework of a Paleoproterozoic arc-continent to continent-continent collisional zone, Trans-Hudson Orogen, from geological and seismic reflection studies. Canadian Journal of Earth Sciences, 42 (4), 421–434. Doi:10.1139/e05-025

- Corrigan, D., Pehrsson, S., Wodicka, N., de Kemp, E., 2009. The Palaeoproterozoic Trans-Hudson Orogen: a prototype of modern accretionary processes. Geological Society, London, Special Publications, 327(1), 457–479. Doi:10.1144/sp327.19
- Cosca, M. A., Mezger, K., Essene, E. J., 1998. The Baltica-Laurentia Connection: Sveconorwegian (Grenvillian) Metamorphism, Cooling, and Unroofing in the Bamble Sector, Norway. The Journal of Geology, 106(5), 539–552. Doi:10.1086/516040
- Dalziel, I.W.D., 1991. Pacific margins of Laurentia and East Antarctica–Australia as a conjugate rift pair: evidence and implications for an Eocambrian supercontinent. Geology 19, p. 598–601. Doi:10.1130/0091-7613(1991)019<0598:pmolae>2.3.co;2
- Dalziel, I.W.D., 1997. Neoproterozoic–Paleozoic geography and tectonics: review, hypothesis and environmental speculation. Bulletin of the Geological Society of America, v. 109, p. 16–42. Doi:10.1130/0016-7606(1997)109<0016:onpgat>2.3.co;2
- Deynoux, M., Affaton, P., Trompette, R., Villeneuve, M., 2006. Pan-African tectonic evolution and glacial events registered in Neoproterozoic to Cambrian cratonic and foreland basins of West Africa. J. Afr. Earth Sci. 46, 397–426. Doi:10.1016/j.jafrearsci.2006.08.005
- Doblas, M., Lopez-Ruiz, J., Cebria, J.M., Youbi, N., Degroote, E., 2002. Mantle insulation beneath the West African Craton during the Precambrian-Cambrian transition. Geology 30, p. 839–842. Doi:10.1130/0091-7613(2002)030<0839:mibtwa>2.0.co;2
- Dorais, M.J., Harper, M., Larson, S., Nugroho, H., Richardson, P., Roosmawati, N., 2005. A comparison of Eastern North America and Coastal New England magma suites: implications for subcontinental mantle evolution and the broad terrane hypothesis. Canadian Journal of Earth Sciences, v. 42, p. 1571–1587. Doi:10.1139/e05-056
- Dupuy, C., and Dostal, J., 1984. Trace element geochemistry of some continental tholeiites. Earth and Planetary Science Letters, 67(1), 61–69. doi:10.1016/0012-821x(84)90038-4
- Egal, E., Thiéblemont, D., Lahondère, D., Guerrot, C., Adi Coseta, C., Iliescu, D., Delor, C., Goujou, J-C., Lafon, JM., Tegye, M., Diaby, S., Kolié, P., 2002. Late Eburnean granitization

and tectonics along the western and northwestern margin of the Archean Kénéma-Man domain (Guinea, West African Craton). *Precambrian Research*, v. 117, p.57–84. Doi:10.1016/s0301-9268(02)00060-8

El Aouli, E.H., Ikenne, M., Amaouain, H., 2010. Petrographic and geochemical characterization of Cryogenian mafic dykes of the Iguerda-Taïfast inlier (Central Anti-Atlas, Morocco). *International Journal of Geomatics and Geosciences*, v. 1, p. 355–371.

El Aouli, E.H., Ikenne, M., Beraaouz, E.H., Mortaji, A., 2004. Les roches basiques des boutonnières d'Agadir Melloul et d'Iguerda-Taïfast: témoins de l'histoire préorogénique de la chaîne panafricaine de l'Anti-Atlas (Maroc). *Estudios Geologicos* 60, p. 11–20. <https://doi.org/10.3989/egeol.04601-269>

El Bahat, A., Ikenne, M., Cousens, B., Söderlund, U., Ernst, R., Klausen, M., B., Youbi, N., 2017. New constraints on the geochronology and Sm-Nd isotopic characteristics of Bas-Drâa mafic dykes, Anti-Atlas of Morocco. *Journal of African Earth Sciences*, 127, 77-87. Doi:10.1016/j.jafrearsci.2016.09.003

El Bahat, A., Ikenne, M., Söderlund, U., Cousens, B., Youbi N., Ernst R., Soulaïmani A., Janati M., Ahmid Hafid A., 2013. U–Pb baddeleyite ages and geochemistry of dolerite dykes in the Bas Drâa Inlier of the Anti-Atlas of Morocco: Newly identified 1380 Ma event in the West African Craton. *Lithos*, V. 174, p. 85–98. Doi:10.1016/j.lithos.2012.07.022

El Hadi, H., Simancas, J.F., Martínez-Poyatos, D., Azor, A., Tahiri, A., Montero, P., Fanning, C.M., Bea, F., González-Lodeiro, F., 2010. Structural and geochronological constraints on the evolution of the Bou Azzer Neoproterozoic ophiolite (Anti-Atlas, Morocco). *Precambrian Research*, v. 182, p. 1–14. Doi:10.1016/j.precamres.2010.06.011

Ennih, N., and Liégeois, J. P., 2008. The boundaries of the West African craton, with special reference to the basement of the Moroccan metacratonic Anti-Atlas belt. Geological Society, London, Special Publications, v. 297(1), p. 1-17.

- 1399 Ennih, N., Liégeois, J. P., 2001. The Moroccan Anti-Atlas: the West African craton passive margin
1400 with limited pan-african activity. Implications for the northern limit of the craton. *Precambrian*
1401 *Research*, v. 112, p. 289-302. [https://doi.org/10.1016/S0301-9268\(01\)00195-4](https://doi.org/10.1016/S0301-9268(01)00195-4)
- 1402 Ernst, R.E., Hamilton, M.A., Söderlund, U., Hanes, J.A., Gladkochub, D.P., Okrugin, A.V.,
1403 Kolotilina, T., Mekhonoshin, A.S., Bleeker, W., LeCheminant, A.N., Buchan, K.L.,
1404 Chamberlain, K.R., Didenko, A.N., 2016. Long-lived connection between southern Siberia and
1405 northern Laurentia in the Proterozoic. *Nature Geoscience*, v. 9, p. 464-469.
1406 [Doi:10.1038/ngeo2700](https://doi.org/10.1038/ngeo2700)
- 1407 Ernst, R.E. 2014. *Large Igneous Provinces*. Cambridge University Press, 653 p.
1408 <https://doi.org/10.1017/CBO9781139025300>
- 1409 Ernst, R.E., and Buchan, K.L., 2001. Large mafic magmatic events through time and links to mantle
1410 plume heads: In: Ernst, R.E., Buchan, K.L. (Eds.), *Mantle Plumes: Their Identification Through*
1411 *Time*, Geological Society of America, Special Paper, v. 352, p. 483–575.
1412 <https://doi.org/10.1130/0-8137-2352-3.483>
- 1413 Ernst, R.E., and Youbi, N., 2017. How large igneous provinces affect global climate, sometimes
1414 cause mass extinctions, and represent natural markers in the geological record.:
1415 *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 478, p. 30–52.
1416 [Doi:10.1016/j.palaeo.2017.03.014](https://doi.org/10.1016/j.palaeo.2017.03.014)
- 1417 Evans, D.A.D. 2013. Reconstructing pre-Pangean supercontinents. *Geological Society of America*
1418 *Bulletin*, 125, 1735–1751. [Doi:10.1130/b30950.1](https://doi.org/10.1130/b30950.1)
- 1419 Evans, D.A.D., Li, Z.X., Murphy, J.B., 2016. Four-dimensional context of Earth's supercontinents.
1420 In: Li, Z.X., Evans, D.A.D., Murphy, J.B. (Eds.), *Supercontinent Cycles Through Earth History*.
1421 Geological Society, London, Special Publications, pp. 1–14. [Doi:10.1144/sp424.12](https://doi.org/10.1144/sp424.12)
- 1422 Evans, D.A.D., Mitchell, R.N., 2011. Assembly and breakup of the core of Paleoproterozoic–
1423 Mesoproterozoic supercontinent Nuna. *Geology* 39, p. 443–446. [doi:10.1130/g31654.1](https://doi.org/10.1130/g31654.1)

- 1424 Fairchild, I.J., Marshall, J.D., Bertrand-Sarfati, J., 1990. Stratigraphic shifts in carbon isotopes from
 1425 Proterozoic stromatolitic carbonates (Mauritania): influence of primary mineralogy and
 1426 diagenesis. *Am. J. Sci.* 290A:46–79
- 1427 Fekkak, A., Pouclet, A., Badra, L., 2002. The Pre-Pan-African rifting of Saghro (Anti-Atlas,
 1428 Morocco): example of the middle Neoproterozoic Basin of Boumalne. *Bulletin de la Société*
 1429 *géologique de France*, v. 173(1), p. 25–35. Doi:10.2113/173.1.25
- 1430 Fekkak, A., Pouclet, A., Ouguir, H., Ouazzani, H., Badra, L., Gasquet, D., 2001. Géochimie et
 1431 signification géotectonique des volcanites du Cryogénien inférieur du Saghro (Anti-Atlas
 1432 oriental, Maroc) / Geochemistry and geotectonic significance of Early Cryogenian volcanics of
 1433 Saghro (Eastern Anti-Atlas, Morocco). *Geodinamica Acta*, 14(6), 373–385.
 1434 doi:10.1080/09853111.2001.10510730.
- 1435 Fodor, R. V., 2009. Diorite Segregations in Gabbro: Geochemical Characteristics and Conditions
 1436 for Origin Assessed at Diorite-Gabbro Contacts. *The Journal of Geology*, 117(2), 109–125.
 1437 doi:10.1086/596506
- 1438 Friedl, G., Finger, F., McNaughton, N.J., Fletcher, I.R., 2000. Deducing the ancestry of terranes:
 1439 SHRIMP evidence for South America-derived Gondwana fragments in central Europe. *Geology*
 1440 28, p.1035-1038. Doi:10.1130/0091-7613(2000)28<1035:dtat>2.0.co;2
- 1441 Fyffe, L.L., Barr, S.M., Johnson, S.C., McLeod, M.J., McNicoll, V.J., Valverde-Vaquero, P., van
 1442 Staal, C.R., White, C.E., 2009. Detrital zircon ages from Neoproterozoic and Early Paleozoic
 1443 conglomerate and sandstone units of New Brunswick and coastal Maine: implications for the
 1444 tectonic evolution of Ganderia. *Atlantic Geology*, v. 45, p. 110–144.
 1445 Doi:10.4138/atlgel.2009.006
- 1446 Gärtner, A., Villeneuve, M., Linnemann, U., El Archi, A., Bellon, H., 2013. An exotic terrane of
 1447 Laurussian affinity in the Mauritanides and Souttoudides (Moroccan Sahara). *Gondwana*
 1448 *Research*, v. 24, p. 687–699. Doi:10.1016/j.gr.2012.12.019

- 1449 Gärtner, A., Villeneuve, M., Linnemann, U., Gerdes, A., Youbi, N., Hofmann, M., 2016. Similar
 1450 crustal evolution in the western units of the Adrar Souttoug Massif (Moroccan Sahara) and the
 1451 Avalonian terranes: insights from Hf isotope data. *Tectonophysics*, v. 681, p. 305–317.
 1452 Doi:10.1016/j.tecto.2015.11.030
- 1453 Gärtner, A., Youbi, N., Villeneuve, M., Linnemann, U., Sagawe, A., Hofmann, M., Zieger, J.,
 1454 Mahmoudi, A., Boumehdi, M. A., 2018. Provenance of detrital zircon from siliciclastic rocks of
 1455 the Sebkha Gezmayet unit of the Adrar Souttoug Massif (Moroccan Sahara) – Palaeogeographic
 1456 implications. *Comptes Rendus Geoscience*. Doi:10.1016/j.crte.2018.06.004
- 1457 Gärtner, A., Youbi, N., Villeneuve, M., Sagawe, A., Hofmann, M., Mahmoudi, A., Boumehdi
 1458 M.A., Linnemann U. 2017. The zircon evidence of temporally changing sediment transport—the
 1459 NW Gondwana margin during Cambrian to Devonian time (Aoucert and Smara areas, Moroccan
 1460 Sahara). *International Journal of Earth Sciences* 106 (8): 2747–2769. Doi:10.1007/s00531-017-
 1461 1457-x
- 1462 Gasquet, D., Chevremont, P., Baudin, T., Chalot-Prat, F., Guerrot, C., Cocherie, A., Roger, J.,
 1463 Hassenforder, B., Cheilletz, A., 2004. Polycyclic magmatism in the Tagragra and Kerdous-
 1464 Tafeltast inliers (Western Anti-Atlas, Morocco). *Journal of African Earth Sciences*, v. 39, p.
 1465 267–275. Doi:10.1016/j.jafrearsci.2004.07.062
- 1466 Gasquet, D., Ennih, N., Liégeois, J.-P., Soulaïmani, A., Michard, A., 2008. The Pan-African belt,
 1467 In: Michard, A., Saddiqi, O., Chalouan, A., Frizon de Lamotte, D. (Eds.), *Continental Evolution:
 1468 The Geology of Morocco*. Lecture Notes in Earth Sciences 116, Springer-Verlag, Berlin
 1469 Heidelberg, p. 33–64, Chapter 2. Doi:10.1007/978-3-540-77076-3_2
- 1470 Gasquet, D., Levresse, G., Cheilletz, A., Azizi-Samir, M.R., Mouttaqi A., 2005. Contribution to a
 1471 geodynamic reconstruction of the Anti-Atlas (Morocco) during Pan-African times with the
 1472 emphasis on inversion tectonics and metallogenic activity at the Precambrian-Cambrian
 1473 transition. *Precambrian Research*, v. 140, p. 157–182. Doi:10.1016/j.precamres.2005.06.009

- 1474 Geyer, G., Landing, E., 2006. Ediacaran–Cambrian depositional environments and stratigraphy of
 1475 the western Atlas regions. In: Geyer, G., Landing, E. (Eds.), Morocco 2006 – Ediacaran–
 1476 Cambrian depositional environments and stratigraphy of the western Atlas regions. UCL
 1477 Maghreb Petroleum Research Group, Infracambrian/Early Palaeozoic Field Guide Series 1 und
 1478 Beringeria Special Issue 6, pp. 47–120.
- 1479 Geyer, G., Landing, E., 2020. Cambrian deposition in northwestern Africa: Relationship of Tamlelt
 1480 massif (Moroccan–Algerian border region) succession to the Moroccan Meseta. *Journal of*
 1481 *African Earth Sciences*, 103772. Doi:10.1016/j.jafrearsci.2020.103772
- 1482 Gladkochub, D.P., Pisarevsky, S.A., Donskaya, T.V., Ernst, R.E., Wingate, M.T.D., Söderlund, U.,
 1483 Mazukabzov, A.M., Sklyarov, E.V., Hamilton, M.A., Hanesg, J.A., 2010. Proterozoic mafic
 1484 magmatism in Siberian craton: an overview and implications for paleocontinental reconstruction.
 1485 *Precambrian Research* 183, p. 660–668. Doi:10.1016/j.precamres.2010.02.023
- 1486 Grenholm, M., Jessell, M., Thébaud, N., 2019. A geodynamic model for the Paleoproterozoic (ca.
 1487 2.27–1.96 Ga) Birimian Orogen of the southern West African Craton – Insights into an evolving
 1488 accretionary-collisional orogenic system. *Earth-Science Reviews*.
 1489 Doi:10.1016/j.earscirev.2019.02.006
- 1490 Hafid, A., 1999. Magmatisme basique filonien Néoproterozoïque (Précambrien II inferieur) dans
 1491 l’Anti-Atlas central et occidental. Minéralogie, géochimie, pétrogenèse et implication
 1492 géodynamique. Unpublished Thèse de Doctorat d’Etat, Faculté des Sciences-Semlalia.
 1493 Université Cadi Ayyad.
- 1494 Hafid, A., Sagon, J.P., Julivert, M., Arboleya, M.L., Saquaque, A., El Boukhari, A., Saidi, A., Soler,
 1495 J.M.F., 2001. Neoproterozoic basic dykes of the Zenaga Inlier, central Anti-Atlas, Morocco:
 1496 petrology, geochemistry and geodynamic significance. *Journal of African Earth Sciences* 32,
 1497 707–721. Doi.org/10.1016/S0899-5362(02)00050-7
- 1498 Hafid, A., Blein, O., Admou, H., Soulaïmani, A., Razin, P.H., Simon, B., Ouanaïmi, H., El janati,
 1499 M., Chevemont, P.H., Baudin, T., Bouabdelli, M., Abia, E.H., Beni Akhy, R., 2013. Notice

- 1500 explicative carte géol. Maroc (1/50 000), feuille Assaragh, Notes et Mémoires Serv. Géol.
 1501 Maroc, n°550, MEM/BRGM.
- 1502 Halls, H.C., Hamilton, M., Denyszyn, S.W., 2011. The Melville Bugt dyke swarm of Greenland: A
 1503 connection to the 1.5 - 1.6 Ga Fennoscandian rapakivi granite province?. In: Srivastava (ed.)
 1504 Dyke Swarms: keys for Geodynamic Interpretation, p. 509-535. Doi:10.1007/978-3-642-12496-
 1505 9_27
- 1506 Hassenforder, B., 1987. La tectonique panafricaine et varisque de l'Anti-Atlas dans le massif de
 1507 Kerdous (Maroc). Ph.D. thesis. Université Strasbourg, Strasbourg, France, 249p.
- 1508 Hefferan, K., et al. 2014. A reconsideration of Pan African orogenic cycle in the Anti-Atlas
 1509 Mountains, Morocco. Journal of African Earth Sciences, v. 98, p. 34-46.
 1510 Doi:10.1016/j.jafrearsci.2014.03.007
- 1511 Henderson, B.J., Collins, W.J., Murphy, J.B., Gutierrez-Alonso, G., Hand, M., 2016. Gondwanan
 1512 basement terranes of the Variscan–Appalachian orogen: Baltican, Saharan and West African
 1513 hafnium isotopic fingerprints in Avalonia, Iberia and the Armorican Terranes. Tectonophysics,
 1514 v. 681, 278–304. Doi:10.1016/j.tecto.2015.11.020
- 1515 Hindermeyer, J., 1953. Le Précambrien III du Sarhro. Comptes Rendus Hebdomadaires des Séances
 1516 de l'Académie des Sciences, 237 (17), 1024–1026.
- 1517 Hodel, F., Macouin, M., Triantafyllou, A., Carlut, J., Berger, J., Rousse, S., Ennih, N., Trindade, R.
 1518 I. F., 2017. Unusual massive magnetite veins and highly altered Cr-spinels as relics of a Cl-rich
 1519 acidic hydrothermal event in Neoproterozoic serpentinites (Bou Azzer ophiolite, Anti-Atlas,
 1520 Morocco). Precambrian Research, 300, 151–167. Doi:10.1016/j.precamres.2017.08.005
- 1521 Hoffman, P. F., 1988. United Plates of America, the Birth of a Craton: Early Proterozoic Assembly
 1522 and Growth of Laurentia. Annual Review of Earth and Planetary Sciences, 16 (1), 543–603.
 1523 Doi:10.1146/annurev.ea.16.050188.002551
- 1524 Hoffman, P.F., 1991. Did the breakout of Laurentia turn Gondwana inside out?. Science, v. 252, p.
 1525 1409–1412. Doi:10.1126/science.252.5011.1409

- 1526 Hoskin, P.W.O., Schaltegger, U., 2003. The composition of zircon and igneous and metamorphic
 1527 petrogenesis. In: Hanchar JM, Hoskin PWO (eds) Zircon; Reviews in Mineralogy and
 1528 Geochemistry, v. 53, p. 27–62. Doi:10.2113/0530027
- 1529 Ikenne, M., 1997. La boutonnière précambrienne du bas Draa (Anti-Atlas occidental, Maroc):
 1530 caractérisation pétrologique et géochimique des roches magmatiques et métamorphiques et leurs
 1531 relations avec la déformation. Ph.D. thesis. University of Agadir, 259p.
- 1532 Ikenne, M., Söderlund, U., Ernst, R.E., Pin, C., Youbi, N., El Aouli, E., Hafid, A., 2017. A c. 1710
 1533 Ma mafic sill emplaced into a quartzite and calcareous series from Ighrem, Anti-Atlas –
 1534 Morocco: Evidence that the Taghdout passive margin sedimentary group is nearly 1 Ga older
 1535 than previously thought. Journal of African Earth Sciences, v. 127, p. 62-76.
 1536 Doi:10.1016/j.jafrearsci.2016.08.020
- 1537 Inglis, J.D., Hefferan, K., Samson, S.D., Admou, H., Saquaque, A., 2016. Determining age of Pan-
 1538 African metamorphism using Sm-Nd garnet-whole rock geochronology and phase equilibria
 1539 modeling in the Tasriwine ophiolite, Sirwa, Anti-Atlas Morocco. Journal of African Earth
 1540 Sciences. v. 127, p. 88–98. Doi:10.1016/j.jafrearsci.2016.06.021
- 1541 Inglis, J.D., MacLean, J.S., Samson, S.D., D’Lemos R.S., Admou, H., Hefferan, K., 2004. A precise
 1542 U–Pb zircon age for the Bleïda granodiorite, Anti-Atlas, Morocco: implications for the timing
 1543 of deformation and terrane assembly in the eastern Anti-Atlas. Journal of African Earth
 1544 Sciences, v. 39, p.277–283. doi:10.1016/j.jafrearsci.2004.07.041
- 1545 Irvine, T.N. and Baragar, W.R.A., 1971. A guide to the chemical classification of the common
 1546 volcanic rocks. Canadian Journal of Earth Science, 8, 523-548. [https://doi.org/10.1139/e71-](https://doi.org/10.1139/e71-055)
 1547 055
- 1548 Jackson, S. E., Pearson, N. J., Griffin, W. L., Belousova, E. A., 2004. The application of laser
 1549 ablation-inductively coupled plasma-mass spectrometry to in situ U–Pb zircon geochronology.
 1550 Chemical Geology, 211(1-2), 47–69. doi:10.1016/j.chemgeo.2004.06.017

- 1551 Jeannette, D., Benziane F., Yazidi A., 1981. Lithostratigraphie et datation du Proterozoïque de la
 1552 boutonnière d'Ifni (Anti-Atlas, Maroc). *Precambrian Research*; 14(3):363-378.
 1553 Doi:10.1016/0301-9268(81)90045-0
- 1554 Jessell, M.W., Begg, G.C., Miller, M.S., 2016. The geophysical signatures of the west African
 1555 craton. *Precambrian Res.* 274, 3–24. Doi:10.1016/j.precamres.2015.08.010
- 1556 Johansson, Å., 2009. Baltica, Amazonia and the SAMBA connection - 1000 million years of
 1557 neighbourhood during the Proterozoic?. *Precambrian Research*, v. 175, p. 221–234.
 1558 Doi:10.1016/j.precamres.2009.09.011
- 1559 Johansson, Å., 2014. From Rodinia to Gondwana with the 'SAMBA' model – A distant view from
 1560 Baltica towards Amazonia and beyond. *Precambrian Research* 244, 226-235.
 1561 Doi:10.1016/j.precamres.2013.10.012
- 1562 Kah, L. C., Bartley, J. K., Teal, D. A., 2012. Chemostratigraphy of the Late Mesoproterozoic Atar
 1563 Group, Taoudeni Basin, Mauritania: Muted isotopic variability, facies correlation, and global
 1564 isotopic trends. *Precambrian Research*, 200-203, 82–103.
 1565 Doi:10.1016/j.precamres.2012.01.011
- 1566 Karaoui, B., Breitzkreuz, C., Mahmoudi, A., Youbi, N., 2014. Physical volcanology, geochemistry
 1567 and basin evolution of the Ediacaran volcano-sedimentary succession in the Bas Draâ Inlier
 1568 (Ouarzazate Supergroup, Western Anti-Atlas, Morocco). *Journal of African Earth Sciences*. v.
 1569 99(2), p. 307-331. Doi:10.1016/j.jafrearsci.2014.06.022
- 1570 Karaoui, B., Breitzkreuz, C., Mahmoudi, A., Youbi, N., Hofmann, M., Gärtner, A., Linnemann, U.,
 1571 2015. U-Pb zircon ages from volcanic and sedimentary rocks of the Ediacaran Bas Draa inlier
 1572 (Anti-Atlas Morocco): chronostratigraphic and provenance implications. *Precambrian Research*.
 1573 v. 263, p. 43-58. Doi:10.1016/j.precamres.2015.03.003
- 1574 Key, R.M., Loughlin, S.C., Horstwood, M.S.A., Gillespie, M., Pitfield, P.E.J., Henney, P.J.,
 1575 Crowley, Q.G., Del Rio, M., 2008. Two Mesoarchaean terranes in the Reguibat shield of NW

- 1576 Mauritania. In: Ennih, N., Liégeois, J.-P. (Eds.), *Boundaries of the West African Craton*, v. 297.
 1577 Special Publication of the Geological Society, London, p. 33–52. Doi:10.1144/sp297.3
- 1578 Klausen, M.B. Nilsson, M.K.M. 2019. The Melville Bugt Dyke Swarm across SE Greenland: A
 1579 closer link to Mesoproterozoic AMCG-complexes. Volume 329, pages 88-107.
 1580 Doi:10.1016/j.precamres.2018.06.001
- 1581 Koffi, G., R-S., Kouamelan, A. N., Allialy, M. E., Coulibaly, Y., Peucat, J.-J., 2020. Re-evaluation
 1582 of Leonian and Liberian events in the geodynamical evolution of the Man-Leo Shield (West
 1583 African Craton). *Precambrian Research*, 105582. Doi:10.1016/j.precamres.2019.105582
- 1584 Kouyaté, D., Söderlund, U., Youbi, N., Ernst, R., Hafid, A., Ikenne, M., Soulaïmani, A., Bertrand,
 1585 H., El Janati, M., Chaham, K.R., 2013. U–Pb baddeleyite and zircon ages of 2040 Ma, 1650 Ma
 1586 and 885 Ma on dolerites in the West African Craton (Anti-Atlas Inlier): possible links to break-
 1587 up of Precambrian supercontinents. *Lithos*, v. 174, p. 71–84. doi:10.1016/j.lithos.2012.04.028
- 1588 Krasnobaev, A., Kozlov, V.I., Puchkov, V.N., Busharina, S.V., Sergeeva, N.D., Paderin I.P., 2013a.
 1589 Zircon geochronology of the Mashak volcanic rocks and the problem of the age of the lower
 1590 middle Riphean boundary (Southern Urals). *Stratigraphy and Geological Correlation* 21, 465–
 1591 481. Doi:10.1134/s0869593813050055
- 1592 Krasnobaev, A., Puchkov, V.N., Kozlov, V.I., Sergeeva, N.D., Busharina, S.V., Lepekhina E.N.,
 1593 2013b. Zirconology of Navysh Volcanic Rocks of the Ai Suite and the Problem of the Age of the
 1594 Lower Riphean Boundary in the Southern Urals. *Doklady Earth Sciences* 448, 185–190.
 1595 Doi:10.1134/s1028334x13020050
- 1596 Kuznetsov, A. B., Bekker, A., Ovchinnikova, G. V., Gorokhov, I. M., Vasilyeva, I. M., 2017.
 1597 Unradiogenic strontium and moderate-amplitude carbon isotope variations in early Tonian
 1598 seawater after the assembly of Rodinia and before the Bitter Springs Excursion. *Precambrian*
 1599 *Research*, 298, 157–173. Doi:10.1016/j.precamres.2017.06.011

- 1600 LE Bas, M. J., Maitre, R. W. L., Streckeisen, A., Zanettin, B., 1986. A Chemical Classification of
 1601 Volcanic Rocks Based on the Total Alkali-Silica Diagram. *Journal of Petrology*, 27(3), 745–750.
 1602 Doi:10.1093/petrology/27.3.745
- 1603 Leblanc, M., 1975. Ophiolites précambriennes et gîtes arsénites de cobalt (Bou Azzer, Maroc).
 1604 Ph.D. thesis, université. Paris VI, p. 329.
- 1605 Leblanc, M., 1981. The late Proterozoic ophiolites of bou Azzer (Morocco): evidence for Pan-
 1606 African plate tectonics. In: Kroner, A. (Ed.), *Precambrian Plate Tectonics*. Elsevier, Amsterdam,
 1607 p. 435–451. Doi:10.1016/s0166-2635(08)70022-7
- 1608 Leblanc, M., and Moussine-Pouchkine, A., 1994. Sedimentary and volcanic evolution of a
 1609 Neoproterozoic continental margin (Bleida, Anti-Atlas, Morocco). *Precambrian Research*, 70(1-
 1610 2), 25–44. doi:10.1016/0301-9268(94)90019-1.
- 1611 Letsch, D., 2018. The Anti-Atlas belt (Morocco) during the Proterozoic – a sedimentary
 1612 perspective. Ph.D. thesis, ETH Zurich, 201p. <https://doi.org/10.3929/ethz-b-000296186>
- 1613 Li, Z. X., D. A. D. Evans, and G. P. Halverson. 2013. Neoproterozoic glaciations in a revised global
 1614 palaeogeography from the breakup of Rodinia to the assembly of Gondwanaland. *Sedimentary*
 1615 *Geology*, 294, 219–232. Doi:10.1016/j.sedgeo.2013.05.016
- 1616 Liégeois, J.-P., Abdelsalam, M.G., Ennih, N., Ouabadi, A., 2013. Metacraton: nature, genesis and
 1617 behavior. *Gondwana Res.* 23, 220–237. Doi:10.1016/j.gr.2012.02.016
- 1618 Liégeois, J.-P.; Fekkak, A.; Bruguier, O.; Errami, E.; Ennih, N, 2006. The Lower Ediacaran (630–
 1619 610 Ma) Saghro group: An orogenic transpressive basin development during the early
 1620 metacratonic evolution of the Anti-Atlas (Morocco). In *Proceedings of the IGCP485 4th*
 1621 *Meeting*, Algiers, Algeria, 2 September 2006; p. 57.
- 1622 Linnemann, U., Gerdes, A., Drost, K., Buschmann, B., 2007. The continuum between Cadomian
 1623 Orogenesis and opening of the Rheic Ocean: constraints from LA-ICP-MS U–Pb zircon dating
 1624 and analysis of plate-tectonic setting (Saxo-Thuringian Zone, NE Bohemian massif, Germany).
 1625 In: Linnemann, U., Nance, D., Kraft, P., Zulauf, G. (Eds.), *The Evolution of the Rheic Ocean:*

- 1626 From Avalonian–Cadomian Active Margin to Alleghenian–Variscan Collision: Memoir of
 1627 Geological Society of America, v. 423, p. 61–96. Doi:10.1130/2007.2423(03)
- 1628 Linnemann, U., Hofmann, M., Romer, R.L., Gerdes, A., 2010. Transitional stages between the
 1629 Cadomian and Variscan Orogenies: Basin development and tectonomagmatic evolution of the
 1630 southern margin of the Rheic Ocean in the Saxo-Thuringian Zone (North Gondwana shelf). In:
 1631 Linnemann U, Romer RL (eds) Pre-Mesozoic Geology of Saxo-Thuringia-From the Cadomian
 1632 Active Margin to the Variscan Orogen. Schweizerbart Science Publishers, Stuttgart, pp 59–98.
- 1633 Linnemann, U., McNaughton, N.J., Romer, R.L., Gehmlich, M., Drost, K., and Tonk, C., 2004.
 1634 West African provenance for Saxo-Thuringia (Bohemian Massif): Did Armorica ever leave pre-
 1635 Pangean Gondwana? U/Pb-SHRIMP zircon evidence and the Nd-isotopic record. International
 1636 Journal of Earth Sciences, v. 93, p. 683–705. doi: 10.1007/s00531-004-0413-8.
- 1637 Linnemann, U., Ouzegane, K., Drareni, A., Hofmann, M., Becker, S., Gärtner, A., Sagawe, A.,
 1638 2011. Sands of West Gondwana: an archive of secular magmatism and plate interactions—a case
 1639 study from the Cambro-Ordovician section of the Tassili Ouan Ahaggar (Algerian Sahara) using
 1640 U–Pb LA-ICP-MS detrital zircon ages. Lithos, v. 123, p. 188–203.
 1641 Doi:10.1016/j.lithos.2011.01.010
- 1642 Lottaroli, F., Craig, J., Thusu, B., 2009. Neoproterozoic–Early Cambrian (Infracambrian)
 1643 hydrocarbon prospectivity of North Africa: a synthesis. In: Craig, J., Thurow, J., Thusu, B.,
 1644 Whitham, A., Abutarruma, Y. (Eds.), Global Neoproterozoic Petroleum Systems: The Emerging
 1645 Potential in North Africa. Vol. 326. Geological Society of London, Special Publication, pp. 137–
 1646 156. Doi:10.1144/sp326.7
- 1647 Ludwig, K.R., 2003. Isoplot 3. In: A Geochronological toolkit for Microsoft Excel: Berkeley
 1648 Geochronology Centre Special Publication 1a.
- 1649 Maloof, A., Schrag, D., Crowley, J., and Bowring, S., 2005. An expanded record of early Cambrian
 1650 carbon cycling from the Anti-Atlas margin, Morocco. Canadian Journal of Earth Sciences, v.
 1651 42(12), p. 2195–2216. Doi:10.1139/e05-062

- 1652 Margalef, A., Castiñeiras, P., Casas, J.M., Navidad, M., Liesa, M., Linnemann, U., Hofmann, M.,
 1653 Gärtner, A., 2016. Detrital zircon from the Ordovician rocks of the Pyrenees: geochronological
 1654 constraints and provenance. *Tectonophysics*, v. 681, p. 121–134.
 1655 Doi:10.1016/j.tecto.2016.03.015
- 1656 Marzoli, A., Bertrand, H., Knight, K., Cirilli, S., Buratti, N., Verati, C., Nomade, S., Renne, P. R.,
 1657 Youbi, N., Martini, R., Allenbach, K., Neuwerth, R., Rapaille, C., Zaninetti, L., Bellieni, G.,
 1658 2004. Synchrony of the Central Atlantic Magmatic province and the Triassic–Jurassic boundary
 1659 climatic and biotic crisis. *Geology*, 32, 973–976. Doi:10.1130/g20652.1
- 1660 Marzoli, A., Bertrand, H., Youbi, N., Callegaro, S., Merle, R., Reisberg, L., Chiaradia, M., Brownlee,
 1661 S., Jourdan, F., Zanetti, A., Davies, J. H F L, Cuppone, T., Mahmoudi, A., Medina, F. Renne,
 1662 P.R., Bellieni, G., Crivellari, S. EL Hachimi, H., Bensalah, M.K., Meyzen, C.M., Tegner, C.,
 1663 2019. The Central Atlantic Magmatic Province (CAMP) in Morocco. *Journal of Petrology*,
 1664 Volume 60, Issue 5, May 2019, Pages 945–996. Doi:10.1093/petrology/egz021
- 1665 Marzoli, A., Davies, J.H.F.L., Youbi, N., Merle, R., Dal Corso, J., Dunkley, D.J., Fioretti, A.M.,
 1666 Bellieni, G., Medina, F., Wotzlaw, J.-F., McHone, G., Font, E., Bensalah, M.K., 2017.
 1667 Proterozoic to Mesozoic evolution of North-West Africa and Peri-Gondwana microplates:
 1668 Detrital zircon ages from Morocco and Canada. *Lithos*, 278–281, 229–239. DOI:
 1669 10.1016/j.lithos.2017.01.016. Doi:10.1016/j.lithos.2017.01.016
- 1670 Massironi, M., Moratti, G., Algouti, A., Benvenuti, M., Dal Piaz, G.V., Eddebbi, A., El Boukhari,
 1671 A., Laftouhi, N., Ouanaimi, H., Schiavo, A., Taj-Eddine, K., Visonà, D., 2007. Carte géologique
 1672 au 1/50 000, Feuille Boumalne. *Not. Mém. Serv. Géol. Maroc* 521, 80.
- 1673 McFarlane, H.B., Ailleres, L., Betts, P., Ganne, J., Baratoux, L., Jessell, M.W., Block S., 2019.
 1674 Episodic collisional orogenesis and lower crust exhumation during the Palaeoproterozoic
 1675 Eburnean Orogeny: Evidence from the Sefwi Greenstone Belt, West African Craton.
 1676 *Precambrian Research*, 325, 88–110. DOI: 10.1016/j.precamres.2019.02.012.

- 1677 Merdith, A. S., Williams, S. E., Müller, R. D., Collins, A. S., 2017. Kinematic constraints on the
 1678 Rodinia to Gondwana transition. *Precambrian Research*, 299, 132–150.
 1679 Doi:10.1016/j.precamres.2017.07.013
- 1680 Metelkin, D.V., Ernst, R.E., Hamilton, M.A., 2011. A ca. 1640 Ma mafic magmatic event in
 1681 southern Siberia, and links with northern Laurentia. *Geological Society of America Annual*
 1682 *Meeting*, abstract, Minneapolis, USA.
- 1683 Michard, A., Saddiqi, O., Chalouan, A. Frizon de La Motte, D. (Eds.) 2008. *Continental Evolution:*
 1684 *the Geology of Morocco*. Springer-Verlag Berlin Heidelberg, p. 404 p. Doi:10.1007/978-3-540-
 1685 77076-3
- 1686 Michard, A., Soulaïmani, A., Ouanaïmi, H., Raddid, Y., Aït Brahim, L., Rjimati, E. ., Baidder, L.,
 1687 Saddiqi, O., 2017. Saghro Group in the Ougnat Massif (Morocco), an evidence for a continuous
 1688 Cadomian basin along the northern West African Craton. *Comptes Rendus Geoscience*, v.
 1689 349(2), p. 81-90. Doi:10.1016/j.crte.2017.01.001
- 1690 Möller, A., O'Brien, P.J., Kennedy, A., Kröner, A., 2003. Linking growth episodes of zircon and
 1691 metamorphic textures to zircon chemistry: an example from the ultrahigh-temperature granulites
 1692 of Rogaland (SW Norway). *Geological Society, London, Special Publications*, v. 220, p. 65-81.
 1693 Doi:10.1144/gsl.sp.2003.220.01.04
- 1694 Mortaji, A. 1989. *La boutonniere précambrienne de Tagragra d'Akka (Anti-Atlas Occidental,*
 1695 *Maroc), Pérologie et géochimie des granitoides. filons basiques et métamorphites associées.*
 1696 *Thèse 3ème Cycle, Université Nancy I*, 211 p.
- 1697 Moussine-Pouchkine, A., and Bertrand-Sarfati, J., 1978. Le Gourma; un aulacogene du Precambrien
 1698 supérieur?. *Bulletin de la Société Géologique de France* ; S7-XX (6): 851–855.
 1699 Doi:10.2113/gssgfbull.s7-xx.6.851
- 1700 Murphy, J.B., Fernández-Suárez, J., Keppie, J.D., Jeffries, T.E., 2004a. Contiguous rather than
 1701 discrete Paleozoic histories for the Avalon and Meguma terranes based on detrital zircon data.
 1702 *Geology*, v. 32, 585–588. Doi:10.1130/g20351.1

- 1703 Murphy, J.B., Fernández-Suárez, J., Keppie, J.D., Jeffries, T.E., 2004b. Lithogeochemical and Sm–
 1704 Nd and U–Pb isotope data from the Silurian–Lower Devonian Arisaig Group clastic rocks,
 1705 Avalon terrane, Nova Scotia: a record of terrane accretion during the Appalachian–Caledonide
 1706 orogen. *Geological Society of America Bulletin*, v. 116, p. 1183–1201. Doi:10.1130/b25423.1
- 1707 Naidoo, D. D., Bloomer, S. H., Saquaque, A., Hefferan, K., 1991. Geochemistry and significance of
 1708 metavolcanic rocks from the Bou Azzer-El Graara ophiolite (Morocco). *Precambrian Research*,
 1709 53(1-2), 79–97. Doi:10.1016/0301-9268(91)90006-v
- 1710 Nance, RD., Murphy, JB., 1994. Contrasting basement isotopic signatures and the palinspastic
 1711 restoration of peripheral orogens: example from the Neoproterozoic Avalonian-Cadomian belt.
 1712 *Geology*, v. 22(7), p.617–620. Doi:10.1130/0091-7613(1994)022<0617:cbisat>2.3.co;2
- 1713 Nicoll, G., Straathof, G., Tait, J., Lo K., Ousmane, N., El Moctar Dahmada, M., Berndt, J., Key, R.,
 1714 2010. Provenance analysis and tectonic setting of the Neoproterozoic sediments within the
 1715 Taoudeni Basin, Northern Mauritania. *Geophys Res Abstr* 12:EGU2010-7094-2.
- 1716 O'Connor, E.A., Barnes, R.P., Beddoe-Stephens, B., Fletcher, T., Gillespie, M.R., Hawkins, M.P.,
 1717 Loughlin, S.C., Smith, M., Smith, R.A., Waters, C.N., Williams, M., 2010. Geology of the Drâa,
 1718 Kerdous, and Boumalne Districts, Anti Atlas, Morocco. Keyworth, Nottingham British
 1719 Geological Survey, p. 324.
- 1720 Ouguir, H., Macaudière, J., Dagallier, G., 1996. Le Protérozoïque supérieur d'Imiter, Saghro
 1721 oriental, Maroc : un contexte géodynamique d'arrière-arc. *Journal of African Earth Sciences*, v.
 1722 22, p.173–189. Doi:10.1016/0899-5362(96)00002-4
- 1723 Pearce, J. A., and Cann, J. R., 1973. Tectonic setting of basic volcanic rocks determined using trace
 1724 element analyses. *Earth and Planetary Science Letters*, 19(2), 290–300. Doi:10.1016/0012-
 1725 821x(73)90129-5
- 1726 Pehrsson, S. J., Berman, R. G., Eglington, B., Rainbird, R., 2013. Two Neoarchean supercontinents
 1727 revisited: The case for a Rae family of cratons. *Precambrian Research*, 232, 27–43.
 1728 Doi:10.1016/j.precamres.2013.02.005

- 1729 Pratt, J.R., Barbeau Jr., D.L., Garver, J.I., Emran, A., Izykowski, T.M., 2015. Detrital zircon
1730 geochronology of Mesozoic sediments in the Rif and Middle Atlas Belts of Morocco:
1731 provenance constraints and refinement of the West African signature. *Journal of Geology*, v.
1732 123, p. 177–200. Doi:10.1086/681218
- 1733 Puchkov, V.N., Krasnobaev, A.A., Sergeeva, N.D., 2014. The New Data on Stratigraphy of the
1734 Riphean Stratotype in the Southern Urals, Russia. *Journal of Geoscience and Environment*
1735 *Protection*, v. 2, p. 108-116. DOI: 10.4236/gep.2014.23015
- 1736 Rainbird, R., Cawood, P.A., and Gehrels, G., 2012. The Great Grenvillian Sedimentation Episode:
1737 Record of Supercontinent Rodinia's Assembly. In Busby, C., and Azor, A., eds., *Recent*
1738 *Advances in the Tectonics of Sedimentary Basins*: Blackwell Publishing Ltd., p. 583 601.
- 1739 Rainbird, R.H., McNicoll, V.J., Theraiault, R.J., Heaman, L.M., Abbott, J.G., Long, D.G.F.,
1740 Thorkelson, D.J., 1997. Pan-continental river systems draining Grenville orogen recorded by U-
1741 Pb and Sm-Nd geochronology of Neoproterozoic quartz arenites and mudrock, northwestern
1742 Canada. *Journal of Geology*, v. 105, p. 1e10. Doi:10.1086/606144
- 1743 Rainbird, R.H., Young, G.M., 2009. Colossal rivers, massive mountains and supercontinents. *Earth*
1744 54, 52-61.
- 1745 Rino, S., Kon, Y., Sato, W., Maruyama, S., Santosh, M., Zhao, D., 2008. The Grenvillian and Pan-
1746 African Orogenies: World's largest orogenies through geologic time, and their implications on
1747 the origin of the superplume. *Gondwana Research*, v. 14, p. 51–72.
1748 Doi:10.1016/j.gr.2008.01.001
- 1749 Rocci, G., Bronner, G., Deschamps, M., 1991. Crystalline Basement of the West African Craton. In:
1750 Dallmeyer, R.D., Lécorché, J.P. (Eds.), *the West African Orogens and Circum-Atlantic*
1751 *Correlatives*: Springer Verlag, Berlin, p. 31–61. Doi:10.1007/978-3-642-84153-8_3
- 1752 Rooney, A.D., Selby, D., Houzay, J.-P., Renne, P.R., 2010. Re–Os geochronology of a
1753 Mesoproterozoic sedimentary succession, Taoudeni basin, Mauritania: implications for basin

- 1754 wide correlations and re-Os organic-rich sediments systematics. *Earth Planet. Sci. Lett.* 289,
 1755 486–496. Doi:10.1016/j.epsl.2009.11.039
- 1756 Sadowski, G.R., Bettencourt, J.S., 1996. Mesoproterozoic tectonic correlations between eastern
 1757 Laurentia and the western border of the Amazon craton. *Precambrian Research*, v. 76, 213-227.
 1758 Doi:10.1016/0301-9268(95)00026-7
- 1759 Salminen, J., Klein, R., Veikkolainen, T., Mertanen, S., Mänttari, I., 2017. Mesoproterozoic
 1760 geomagnetic reversal asymmetry in light of new paleomagnetic and geochronological data for
 1761 the Häme dyke swarm, Finland: Implications for the Nuna supercontinent. *Precambrian*
 1762 *Research*, 288, 1–22. Doi:10.1016/j.precamres.2016.11.003
- 1763 Samson, S.D., Inglis, J.D., D'Lemos, R.S., Admou, H., Blichert-Toft, J., Hefferan, K., 2004.,
 1764 Geochronological, geochemical, and Nd–Hf isotopic constraints on the origin of Neoproterozoic
 1765 plagiogranites in the Tasriwine ophiolite, Anti-Atlas orogen, Morocco: . *Precambrian Research*,
 1766 v. 135, p. 133–147. Doi:10.1016/j.precamres.2004.08.003
- 1767 Santos, J.O.S., Hartmann, L.A., Gaudette, H.E., Groves, D.I., Mcnaughton, N.J., and Fletcher, I.R.,
 1768 2000. A new understanding of the provinces of the Amazon craton based on integration of field
 1769 mapping and U–Pb and Sm–Nd geochronology. *Gondwana Research*, v. 3, no. 4, p.453–488.
 1770 Doi:10.1016/s1342-937x(05)70755-3
- 1771 Santos, J.O.S., Rizzotto, G.J., Potter, P.E., McNaughton, N.J., Matos, R.S., Hartmann, L.A.,
 1772 Chemale Jr., F., Quadros, M.E.S., 2008. Age and autochthonous evolution of the Sunsás Orogen
 1773 in West Amazon Craton based on mapping and U–Pb geochronology. *Precambrian Research*
 1774 165, 120–152. Doi:10.1016/j.precamres.2008.06.009
- 1775 Saquaque, A., Admou, H., Karson, J., Hefferan, K., Reuber, I., 1989. Precambrian accretionary
 1776 tectonics in the Bou Azzer-El Graara region, Anti-Atlas, Morocco. *Geology*, 17(12), 1107.
 1777 Doi:10.1130/0091-7613(1989)017<1107:patitb>2.3.co;2

- 1778 Schiavo, A., Taj-Eddine, K., Algouti, A., Benvenuti, M., Dal Piaz, G.V., Eddebbi, A., El Boukhari,
 1779 A., Laftouhi, N., Massironi, M., Moratti, G., Ouanaïmi, H., Pasquarè, G., Visonà, D., 2007.
 1780 Carte géologique au 1/50 000, Feuille Imtir. Not. Mém. Serv. Géol. Maroc 518, 96.
- 1781 Schofield, D.I., Horstwood, M.S.A., Pitfield, P.E.J., Crowley, Q.G., Wilkinson, A.F., Sidaty,
 1782 H.C.O, 2006. Timing and kinematics of Eburnean tectonics in the central Reguibat Shield.
 1783 Mauritania: Journal of the Geological Society, v. 163, p.549–560. Doi:10.1144/0016-764905-
 1784 097
- 1785 Schulte B., Benziane F., Yazidi A., Boger S., Stockhammer S., Lehmann A., Saadane A., Yazidi
 1786 M., 2016. Carte géologique du Maroc, feuille au 1/50 000 Arbaa Sahel. Notes et Mém. Serv.
 1787 Géol. Maroc- Rabat n° 543bis.
- 1788 Shervais, J.W., 1982. Ti–V plots and the petrogenesis of modern and ophiolitic lavas. Earth and
 1789 Planetary Science Letters, v. 59, p. 101–118. Doi:10.1016/0012-821x(82)90120-0
- 1790 Shields, G. A., Deynoux, M., Strauss, H., Paquet, H., Nahon, D., 2007. Barite-bearing cap
 1791 dolostones of the Taoudéni Basin, northwest Africa: Sedimentary and isotopic evidence for
 1792 methane seepage after a Neoproterozoic glaciation. Precambrian Research, 153(3-4), 209–235.
 1793 Doi:10.1016/j.precamres.2006.11.011
- 1794 Shumlyanskyy, L., Hawkesworth, C., Dhuime, B., Billström, K., Claesson, S., Storey, C., 2015.
 1795 ²⁰⁷Pb/²⁰⁶Pb ages and Hf isotope composition of zircons from sedimentary rocks of the
 1796 Ukrainian shield: Crustal growth of the south-western part of East European craton from
 1797 Archaean to Neoproterozoic. Precambrian Research, 260, 39–54.
 1798 Doi:10.1016/j.precamres.2015.01.007
- 1799 Sircombe, K.N., 2004. AgeDisplay: an EXCEL workbook to evaluate and display univariate
 1800 geochronological data using binned frequency histograms and probability density distributions.
 1801 Computational Geosciences, v. 30, p. 21–31. Doi:10.1016/j.cageo.2003.09.006
- 1802 Söderlund, U., Ibanez-Mejia, M., El Bahat, A., Ernst, R.E., Ikenne, M., Soulaïmani, A., Youbi, N.,
 1803 Cousens, B., El Janati, M., Hafid, A., 2013. Reply to Comment on “U–Pb baddeleyite ages and

- 1804 geochemistry of dolerite dykes in the Bas Drâa inlier of the Anti-Atlas of Morocco: Newly
 1805 identified 1380 Ma event in the West African Craton” by André Michard and Dominique
 1806 Gasquet: *Lithos*, v. 174, p.101–108. Doi:10.1016/j.lithos.2013.04.003
- 1807 Soulaïmani, A., Ouanaimi, H., Michard, A., Montero, P., Bea, F., Corsini, M., Molina, J. F.,
 1808 Rjimatif, E. C., Saddiqi, O., Hefferan, K., 2019. Quartzite crests in Paleoproterozoic granites
 1809 (Anti-Atlas, Morocco); a hint to Pan-African deformation of the West African Craton margin.
 1810 *Journal of African Earth Sciences*, 157, 103501. Doi:10.1016/j.jafrearsci.2019.05.009
- 1811 Soulaïmani, A., Ouanaimi, H., Saddiqi, O., Baidder, L., Michard, A., 2018. The Anti-Atlas Pan-
 1812 African Belt (Morocco): Overview and pending questions. *Comptes Rendus Geoscience*. 350,
 1813 279-288. Doi:10.1016/j.crte.2018.07.002
- 1814 Soulaïmani, A., Blein, O., Chèvremont, P., Ouanaimi, H., Hafid, A., Admou, H., Baudin, T.,
 1815 Bouabdelli, M., Razin, P.H., Abia, E.H., Beni Akhy, R., 2013a. Notice explicative carte géol.
 1816 Maroc (1/50 000), feuille Agadir Melloul, Notes et Mémoires Serv. Géol. Maroc, n°549,
 1817 MEM/BRGM.
- 1818 Spencer, C.J., Kirkland, C.L., Taylor, R.J.M., 2016. Strategies towards statistically robust
 1819 interpretation of in situ U-Pb zircon geochronology. *Geoscience Frontiers* 7, 581–589.
 1820 doi.org/10.1016/j.gsf.2015.11.006
- 1821 Stacey, J. S., and Kramers, J. D., 1975. Approximation of terrestrial lead isotope evolution by a two
 1822 stage model. *Earth and Planetary Science Letters*, 26(2), 207–221. doi:10.1016/0012-
 1823 821x(75)90088-6.
- 1824 Straathof, G.B., 2011. Neoproterozoic Low Latitude Glaciations: An African Perspective. Ph.D.
 1825 thesis, University of Edinburgh, 285p.
- 1826 Sun, S.S., and McDonough, W.F., 1989. Chemical and isotopic systematics of oceanic basalts;
 1827 implications for mantle composition and processes. In: *Magmatism in the ocean basins*.
 1828 Saunders, A.D. and Norry, M.J. (Editors), Geological Society of London, London, v. 42, p. 313-
 1829 345. Doi:10.1144/gsl.sp.1989.042.01.19

- 1830 Tapsoba, B., Lo, C.-H., Wenmenga, U., Iizuka, Y., Chung, S.-L., Shellnutt, G., 2018. Chemical and
 1831 Sr-Nd compositions and $^{40}\text{Ar}/^{39}\text{Ar}$ ages of NW-trending dolerite dikes of Burkina Faso:
 1832 Evidence for a Mesoproterozoic magmatism in the West African Craton. *Geoscience Frontiers*.
 1833 doi:10.1016/j.gsf.2017.12.015
- 1834 Tassinari, C.C.G, Macambira, M.J., 1999. Geochronological provinces of the Amazonian Craton.
 1835 International Union of Geological Sciences, v. 22, p. 174–182.
 1836 <https://doi.org/10.18814/epiiugs/1999/v22i3/004>
- 1837 Tassinari, C.C.G., Bettencourt, J.S., Geraldès, M.C., Macambira, M.J.B., Lafon, J.M., Cordani, U.,
 1838 Milani, E.J., Thomaz Filho, A., and Campos, D.A., 2000. The Amazonian Craton, in Cordani,
 1839 U.G., Milani, E.J., Thomaz Filho, A., and Campos, D.A., eds., *Tectonic evolution of South*
 1840 *America: Rio de Janeiro, 31st International Geological Congress*, p.41–95.
- 1841 Tegner, C., Michelis, S.A.T, McDonald, I., Brown, E.L., Youbi, N., Callegaro, S., Lindström, S.,
 1842 Marzoli, A., 2019. Mantle Dynamics of the Central Atlantic Magmatic Province (CAMP):
 1843 Constraints from Platinum Group, Gold and Lithophile Elements in Flood Basalts of Morocco.
 1844 *Journal of Petrology*. Doi:10.1093/petrology/egz041
- 1845 Thiéblemont, D., 2016. An updated geological map of Africa at 1/10 000 000 scale. In: Presented at
 1846 the 35th International Geological Congress: IGC 2016. [https://hal-brgm.archives-](https://hal-brgm.archives-ouvertes.fr/hal-01326139)
 1847 [ouvertes.fr/hal-01326139](https://hal-brgm.archives-ouvertes.fr/hal-01326139)
- 1848 Thiéblemont, D., Goujou, J.C., Egal, E., Cocherie, A., Delor, C., Lafon, J.M., Fanning, C.M., 2004.
 1849 Archean evolution of the Leo Rise and its Eburnean reworking. *Journal of African Earth*
 1850 *Sciences*, v. 39, p. 97–104. Doi:10.1016/j.jafrearsci.2004.07.059
- 1851 Thomas, R. J., Chevakier, L. P., Gresse, P. G., Harmer, R. E., Eglington, B. M., Armstrong, R. A.,
 1852 De Beer, C. H., Martini, J. E. J., De Kock, G. S., Macey, P. H., and Ingram, B. A., 2002.
 1853 Precambrian evolution of the Sirwa Window, Anti-Atlas Orogen, Morocco: Precambrian
 1854 Research, v. 118, p. 1-57. Doi:10.1016/s0301-9268(02)00075-x

- 1855 Thomas, R.J., Fekkak, A., Ennih, N., Errami, E., Loughlin, S.C., Gresse, P.G., Chevakier, L.P. and
 1856 Liegeois J.P., 2004. A new lithostratigraphic framework for the Anti-Atlas Orogen, Morocco.
 1857 Journal of African Earth Sciences, v. 39, p. 217-226. Doi:10.1016/j.jafrearsci.2004.07.046
- 1858 Thompson, R.N., Dickin, A.P., Gibson, I.L., Morrison, M.A., 1982. Elemental fingerprints of
 1859 isotopic contamination of Hebridean Palaeocene mantle-derived magmas by Archaean sial.
 1860 Contributions to Mineralogy and Petrology, v. 79, p. 159–168. Doi:10.1007/bf01132885
- 1861 Toummite, A., Liégeois, J.P., Gasquet, D., Bruguier, O., Beraaouz, E.H., Ikenne, M., 2013. Field,
 1862 geochemistry and Sr-Nd isotopes of the pan-African granitoids from the Tifnoute Valley (Sirwa,
 1863 anti-atlas, Morocco): a post-collisional event in a metacratonic setting. Mineral. Petrol. 107,
 1864 739–763. Doi:10.1007/s00710-012-0245-3
- 1865 Triantafyllou, A., 2015. Evolution géodynamique d'un arc insulaire néoproterozoïque de l'Anti -
 1866 Atlas marocain : caractérisation des processus de croissance intra-océanique et d'accrétion. PhD
 1867 thesis. Université de Nantes, Nantes, France, 390 p.
- 1868 Triantafyllou, A., Berger, J., Baele, J.M., Bruguier, O., Diot, H., Ennih, N., Monnier, C., Plissart,
 1869 G., Vandycke, S., Watlet, A., 2018. Intra-oceanic arc growth driven by magmatic and tectonic
 1870 processes recorded in the Neoproterozoic Bougmane Arc complex (Anti-Atlas, Morocco).
 1871 Precambrian Res. 304, 39–63. Doi:10.1016/j.precamres.2017.10.022
- 1872 Triantafyllou, A., Berger, J., Baele, J.-M., Diot H., Ennih, N., Plissart, G., Monnier, C., Watlet, A.,
 1873 Bruguier, O., Spagna, P., Vandycke S., 2016. The Tachakoucht-Iriri-Tourtit arc complex
 1874 (Moroccan Anti-Atlas): Neoproterozoic records of polyphased subduction-accretion dynamics
 1875 during the Pan-African orogeny. Journal of Geodynamics, v. 98, p.81–103.
 1876 Doi:10.1016/j.jog.2015.07.004
- 1877 Triantafyllou, A., Berger, J., Baele, J.-M., Mattielli, N., Ducea, M. N., Sterckx, S., Samson, S.,
 1878 Florent Hodel, F., Ennih, N., 2020. Episodic magmatism during the growth of a
 1879 Neoproterozoic oceanic arc (Anti-Atlas, Morocco). Precambrian Research, 105610.
 1880 Doi:10.1016/j.precamres.2020.105610

- 1881 Triantafyllou, A., Hodel, F., Berger, J., Macouin, M., Baele, J.M., Mattielli, N., Monnier, C., Ducea
 1882 Mihai, N., Poujol, M., Langlade, J., Trindade, R. I. F., 2019. The Bou azzer and Sirwa ophiolites
 1883 (Anti-Atlas, Morocco): insight into polyphased subduction-accretion dynamics during
 1884 Neoproterozoic times. GSA Annual Meeting, Phoenix, Arizona, USA. Doi:
 1885 10.1130/abs/2019AM-336212
- 1886 Tuduri, J., Chauvet, A., Barbanson, L., Labriki, M., Dubois, M., Trapy, P.-H., Lahfid, A., Poujol,
 1887 M., Melleton, J., Badra, L., Ennaciri, A., Maacha, L., 2018. Structural control, magmatic-
 1888 hydrothermal evolution and formation of hornfels-hosted, intrusion-related gold deposits: Insight
 1889 from the Thaghassa deposit in Eastern Anti-Atlas, Morocco. Ore Geology Reviews, 97, 171–
 1890 198. doi:10.1016/j.oregeorev.2018.04.023
- 1891 Vaasjoki, M., Tapani Rämö, O., Sakko, M., 1991. New U-Pb ages from the Wiborg rapakivi area:
 1892 constraints on the temporal evolution of the rapakivi granite-anorthosite-diabase dyke
 1893 association of southeastern Finland. Precambrian Research, 51(1-4), 227–243.
 1894 Doi:10.1016/0301-9268(91)90102-g
- 1895 Villeneuve, M., 2005. Paleozoic basins in West Africa and the Mauritanide thrust belt. Journal of
 1896 African Earth Sciences, 43(1-3), 166–195. Doi:10.1016/j.jafrearsci.2005.07.012
- 1897 Waldron, J.W.F., White, C.E., Barr, S.M., Simonetti, A., Heaman, L.M., 2009. Provenance of the
 1898 Meguma terrane, Nova Scotia: rifted margin of early Paleozoic Gondwana. Canadian Journal
 1899 of Earth Sciences, v. 46, p. 1–8. Doi:10.1139/e09-004
- 1900 Walsh, G.J., Aleinikoff, J.N., Benziane, F., Yazidi, A., Armstrong, T.R., 2002. U-Pb zircon
 1901 geochronology of the Paleoproterozoic Tagragra de Tata inlier and its Neoproterozoic cover,
 1902 western Anti-Atlas, Morocco. Precambrian Research, v. 117, p. 1–20. Doi:10.1016/s0301-
 1903 9268(02)00044-x
- 1904 Walsh, G.J., Benziane, F., Aleinikoff, J.N., Harisson, R.W., Yazidi, A., Burton, W.C., Quick, J.E.,
 1905 Saadane, A., 2012. Neoproterozoic tectonic evolution of the Jebel Saghro and Bou Azer-El

- 1906 Graara inliers, eastern and central Anti-Atlas, Morocco. *Precambrian Research*, v. 216–219, p.
1907 23–62. Doi:10.1016/j.precamres.2012.06.010
- 1908 Wang, C., Zhang, J.-H., Li, M., Li, R.-S., Peng, Y., 2015. Generation of ca. 900–870Ma bimodal
1909 rifting volcanism along the southwestern margin of the Tarim Craton and its implications for the
1910 Tarim–North China connection in the early Neoproterozoic. *Journal of Asian Earth Sciences*,
1911 113, 610–625. doi:10.1016/j.jseaes.2015.08.002
- 1912 Weil, A.B., Van der Voo, R., MacNiocall, C., Meert, J.G., 1998. The Proterozoic supercontinent
1913 Rodinia: paleomagnetically derived reconstructions for 1100–800 Ma. *Earth Planet: Science*
1914 *Letters*, v. 154, p. 13–24. Doi:10.1016/s0012-821x(97)00127-1
- 1915 Williams, I.S., 1998. U-Th-Pb geochronology by ion microprobe. In: McKibben, M.A., Shanks,
1916 W.C., Riley, W.I. (Eds.), *Applications of Microanalytical Techniques to Understanding*
1917 *Mineralising Processes. Reviews in Economic Geology*, v.7, p. 1–35.
1918 <https://doi.org/10.5382/Rev.07.01>
- 1919 Willner, A.P., Barr, S.M., Gerdes, A., Massonne, H.-J., White, C.E., 2013. Origin and evolution of
1920 Avalonia: evidence from U–Pb and Lu–Hf isotopes in zircon from the Mira terrane, Canada, and
1921 the Stavelot–Venn Massif, Belgium. *Journal of the Geological Society of London*: v. 170, p.
1922 769–784. Doi:10.1144/jgs2012-152
- 1923 Winchester, J. A., and Floyd, P. A., 1977. Geochemical discrimination of different magma series
1924 and their differentiation products using immobile elements. *Chemical Geology*, 20, 325–343.
1925 doi:10.1016/0009-2541(77)90057-2
- 1926 Youbi N., Söderlund U., Ernst R. E., Ait Malek M., Barzouk A., Ait Lahna A., Palassia M., Gong
1927 Z., Evans D., Wen B., Jing X-Q., Boumehdi M. A., Ikenne M., Bensalah M. K., Admou H.,
1928 Bajddi A., Maacha L, 2019. Ca. 1.65 Ga mafic sills emplaced into the quartzite of jbel Lkest
1929 from Kerdous Inlier, Anti-Atlas, West African craton, Morocco: additional evidence that the
1930 basal part of Taghdout group is nearly 1 Ga older than previously thought. *The 7 International*
1931 *Science Conference Abstracts*. Tomsk: p.165.

- 1932 Youbi, N., Bouougri, El H., Ait Lahna, A., Kouyaté, D., Tassinari, C.C.G., Admou, H., Mata, J.,
 1933 Boumehdi, M.A., Bensalah, M.K., Basei, M., A., S., Sato., K., Ernst, R.E., Söderlund, U., Chaib,
 1934 L., Bairouk L., Gärtner, A., Linnemann, U., Ikenne M., Bodinier, J-L., 2018. A grande província
 1935 ígnea de Iguerda-Taifast (ca. 885 Ma) no Anti-Atlas (Marrocos-Cratão Oeste africano): os
 1936 enxames de diques máficos de Ifzwane e os vulcanitos máficos do grupo Tizi n'Taghatine. X
 1937 Congresso National de geologia, Açores, Portugal, Vulcânica, v.2, p.21.
- 1938 Youbi, N., Ernst, R.E., Söderlund, U., Boumehdi, M.A., Ait Lahna, A., Tassinari, C.C.G., El
 1939 Moume, W., Bensalah, M.K., 2020. The Central Iapetus magmatic province: An updated review
 1940 and link with the ca. 580 Ma Gaskiers glaciation. In: Thierry Adatte; David P.G. Bond; Gerta
 1941 Keller (Editors). Mass Extinctions, Volcanism, and Impacts: New Developments. Geological
 1942 Society of America Special Papers, Volume 544. Published: December 31, 2019 DOI:
 1943 [https://doi.org/10.1130/2020.2544\(02\)](https://doi.org/10.1130/2020.2544(02))
- 1944 Youbi, N., Kouyaté, D., Söderlund, U., Ernst, R.E., Soulaïmani, A., Hafid, A., Ikenne, M., El Bahat,
 1945 A., Bertrand, H., Chaham, K.R., Ben Abbou, M., Mortaji, A., El Ghorfi, M., Zouhair, M., El
 1946 Janati, M., 2013. The 1750 Ma Magmatic Event of the West African Craton (Anti-Atlas,
 1947 Morocco). *Precambrian Research*, v. 236, p.106–123. Doi:10.1016/j.precamres.2013.07.003
- 1948 Youbi, N., Martins, L. T., Munhá, J. M., Ibouh, H., Madeira, J., Ait Chayeb, H., El Boukhari, A.,
 1949 2003. The Late Triassic-Early Jurassic Volcanism of Morocco and Portugal in the Framework of
 1950 the Central Atlantic Magmatic province: An Overview. In: Hames, W. E., Machone, J. G.,
 1951 Renne, P. R., Ruppel, C. (Eds.) *The Central Atlantic Magmatic Province: Insights from*
 1952 *Fragments of Pangea*. American Geophysical Union Geophysical Monograph Series, 136, 179-
 1953 207. Doi:10.1029/136gm010
- 1954 Zhang, C.-L., Ye, X.-T., Ernst, R. E., Zhong, Y., Zhang, J., Li, H.-K., Long, X.-P., 2019. Revisiting
 1955 the Precambrian evolution of the Southwestern Tarim terrane: implications for its role in
 1956 Precambrian supercontinents. *Precambrian Research*. Doi:10.1016/j.precamres.2019.01.018
- 1957

Table 1. Site location and summary of the data for the studied samples. The names of units and formations are sensu Bouougri and Saquaque (2004).

Sample ID	Coordinates (GPS WGS84)	Inlier/ Subinlier	Lithology	Units/Formations/Suites	Methodes/ Mineral	Number of spots	U-Pb zircon Age (Ma)
BA24	N 30°23'7.90", W 6°30'33.40"	Bou Azzer El Graara/Bleida	Fine-grained sandstone	Upper Unit/Bleida Formation	SHRIMP II/Zircon	44 spots	1855 ± 6.5 - 977 ± 10*
ALBL29	N 30°20'19.28", W 6°24'48.70"	Bou Azzer El Graara/Bleida	Medium to coarse-grained quartzite	Middle Unit/Tachdamt Formation	LA-ICP-MS/Zircon	96 spots	1060 ± 5.5 - 930 ± 12*
BA21	N 30°29'11.40", W 6°52'44.00"	Bou Azzer El Graara/Tachdamt	Medium-grained quartzite	Lower Unit/Agoumy Formation	SHRIMP II/Zircon	55 spots	1818 ± 6.4*
TA1	N 30°36'43.6", W 7°17'13.4"	Zenaga	Coarse-grained to a fine-grained microlitic porphyritic dolerite	Ifzwane Suite	SHRIMP II/Zircon	18 spots	1676 ± 37**
TG-15-1	N 30°35'11.76", W 7°20'06.92"	Zenaga	Purple coarse-grained and quartz rich metasiltstone	Lower Unit/Tasserda Formation	LA-ICP-MS/Zircon	99 spots	2049.3 ± 4.6*

* U-Pb detrital zircon, maximum depositional age

** U-Pb age emplacement

SHRIMP II: Sensitive High Resolution Ion Microprobe

LA-ICP-MS: Laser Ablation Inductively Coupled Plasma Mass Spectrometry

ID: Identification

GPS WGS84: The Global Positioning System (GPS) uses the World Geodetic System (WGS84) as its reference coordinate system.

Table 1. Site location and data summary of the investigated samples. The names of Units and Formations are sensu Bouougri and Saquaue (2004).

Sample ID	Coordinates (GPS WGS84)	Inlier/ Subinlier	Lithology	Units/Formations/Suites	Methodes/ Mineral	Number of spots	U-Pb zircon Age (Ma)
BA 24	N 30°23'7.90", W 6°30'33.40"	Bou Azzer El Graara/Bleida	Fine-grained sandstone	Upper Unit/Bleida Formation	SHRIMP/II/ Zircon	44 spots	1957 ± 63*
ALBL29	N 30°20'19.28", W 6°24'48.70"	Bou Azzer El Graara/Bleida	Medium to coarse-grained quartzarenite	Middle Unit/Tachdamt Formation	LA-ICP-MS/Zircon	96 spots	1060 ± 5.5*
BA 21	N 30°29'11.40", W 6°52'44.00"	Bou Azzer El Graara/Tachdamt	Medium-grained quartzarenite	Lower Unit/Agoummy Formation	SHRIMP/II/ Zircon	54 spots	1818 ± 6.4*
TA 1	N 30°36'43.6", W 7°17'13.4"	Zenaga	Coarse-grained to a fine-grained microlitic porphyric dolerite	Ifzwane Suite	SHRIMP/II/ Zircon	18 spots	1639 ± 34 **
TG-15-1	N 30°35'11.76", W 7°20'06.92"	Zenaga	Purple coarse-grained and quartz rich siltstone	Lower Unit/Tassarda Formation	LA-ICP-MS/Zircon	99 spots	2047 ± 9.7*

* U-Pb detrital zircon, maximum depositional age

** U-Pb age emplacement

SHRIMP/II: Sensitive High Resolution Ion Microprobe

LA-ICP-MS: Laser Ablation Inductively Coupled Plasma Mass Spectrometry

ID: Identification


GPS WGS84: The Global Positioning System (GPS) uses the World Geodetic System (WGS84) as its reference coordinate system.


Table 2. Representative whole-rock major and trace element geochemical data for dykes and sills from the Zenaga, Agadir Melloul, Iguerda-Taïfast, and Tagragra of Akka inliers including the dated dykes, samples TA1 from the Zenaga Inlier (1676 ± 37 Ma; this work), DZ 36 (1640 ± 2 Ma and 1641 ± 3 Ma; Kouyaté et al., 2013) and DZ 40 (1656 ± 9 Ma; Kouyaté et al., 2013) from the Zenaga Inlier (Kouyaté et al, 2013), and AMHA069 (1654 ± 16 Ma; Kouyaté et al, 2013) from the Agadir Melloul Inlier. The geochemical data are from unpublished theses (Hafid, 1999; Mortaji, 1989) and published papers (Hafid et al., 2001; El Aouli et al., 2010; Youbi et al., 2013). Fe₂O₃t= total Fe expressed as Fe₂O₃. Loss-on-ignition (L.O.I.) determined by sample weight loss at 1000°C.


Sample	TA1	TA2	DZ36	DZ40	AMHA69	ZEN-4	ZEN-13	ZEN-1	ZEN-11	IT-5	IT-7	IT-12	IT-13	NC40	NC379	NC314aMor	NC 361
Inlier	Zenaga	Zenaga	Zenaga	Zenaga	Agadir Melloul	Zenaga	Zenaga	Zenaga	Zenaga	Iguerda-Taïfast	Iguerda-Taïfast	Iguerda-Taïfast	Iguerda-Taïfast	Tagragra of Akka	Tagragra of Akka	Tagragra of Akka	Tagragra of Akka
Latitude N	30°36'43.6"	30°36'43.7"															
Longitude W	7°17'13.4"	7°17'13.3"															
Orientation	N45°	N45°	N125°	N140°		N170°	N125°	N85°	N170°		N00°	N30°	N45°	N50°-N110°	N50°-N110°	N30°-N160°	N30°-N160°
Ages (Ma)	1676 ± 37		1640 ± 2/ 1641 ± 3	1656 ± 9	1654 ± 16												
Reference	(this work)	(this work)	Kouyaté (2013)	Kouyaté (2013)	Kouyaté (2013)	Hafid (1999)	Hafid (1999)	Hafid (1999)	Hafid (1999)	Hafid (1999)	Hafid (1999)	Hafid (1999)	Hafid (1999)	Mortaji (1989)	Mortaji (1989)	Mortaji (1989)	Mortaji (1989)
SiO ₂	49.31	49.73	50.66	50.84	45.78	48.56	49.71	48.38	48.30	49.12	49.98	49.00	47.98	50.85	47.58	44.35	49.18
TiO ₂	1.64	1.58	0.85	0.73	0.79	2.66	2.09	1.98	1.64	2.05	2.52	2.17	2.06	0.91	1.93	2.86	2.99
Al ₂ O ₃	13.79	13.48	14.55	14.61	13.73	13.58	14.08	13.97	13.97	13.50	13.67	13.06	13.20	14.72	13.62	15.25	12.75
Fe ₂ O ₃ t	14.51	14.35	10.27	9.78	9.50	14.43	14.83	13.71	13.16	15.17	15.02	14.87	15.54	12.60	15.63	16.53	16.06
MnO	0.23	0.23	0.16	0.18	0.25	0.17	0.19	0.16	0.20	0.26	0.17	0.23	0.26	0.19	0.20	0.18	0.12
MgO	6.34	6.68	7.31	7.55	9.87	6.61	5.76	6.55	7.73	5.28	5.24	5.91	6.24	6.69	6.29	5.95	5.21
CaO	6.70	6.20	10.01	11.27	12.18	8.71	8.51	10.18	10.34	8.52	6.98	10.04	8.40	7.22	8.82	7.16	4.36
Na ₂ O	3.26	4.06	2.17	1.94	2.49	2.40	2.59	1.86	2.10	2.46	2.45	1.91	2.73	3.36	2.04	2.58	2.63
K ₂ O	0.96	0.32	0.74	0.62	0.29	0.54	0.78	0.32	0.64	1.15	1.79	0.36	0.95	0.14	0.77	1.04	1.31
P ₂ O ₅	0.14	0.19	0.07	0.08	0.06	0.24	0.33	0.18	0.15	0.43	0.36	0.19	0.17	0.14	0.24	0.69	0.50
L.O.I.	2.47	3.04	2.48	2.38	4.37	2.35	2.18	2.67	2.02	1.46	1.73	2.07	1.99	3.67	2.69	3.80	5.00
Total	99.34	99.84	99.49	100.25	99.68	100.25	101.05	99.96	100.25	99.40	99.91	99.81	99.52	100.49	99.81	100.39	100.11
V	343.24	317.91	240.00	233.70	223.00	326.90	317.14	306.30	332.92	290.97	339.52	400.00	413.57	278.00	367.00	189.00	367.00
Cr	18.95	18.83	307.70	410.10	802.70	189.80	96.76	197.10	181.59	114.14	126.86	113.12	180.31	139.00	114.00	180.00	118.00
Co	51.53	54.75	46.00	44.67	57.36	43.13	50.47	40.43	55.85	44.43	46.07	44.10	43.62	57.00	54.00	46.00	46.00
Ni	39.42	37.22	143.00	185.20	178.30	117.00	69.71	94.16	121.74	44.20	64.28	79.64	92.72	55.00	95.00	109.00	65.00
Zn	126.00	158.84	90.36	92.06	228.20	150.20	152.60	119.80	161.27	131.25	129.48	134.44	155.27		116.00	153.00	62.00
Rb	35.95	8.84	26.89	20.34	10.88	16.85	20.90	8.44	19.70	63.40	68.05	24.02	45.85	10.00	43.00	29.00	81.00
Sr	192.75	220.51	215.20	196.60	191.10	391.10	410.50	269.90	256.15	372.90	370.30	253.56	200.55	593.00	185.00	247.00	120.00
Y	23.11	23.50	16.27	14.16	12.11	28.80	30.91	27.12	22.87	37.00	28.97	29.81	28.79	19.46	30.85	41.74	26.22
Zr	88.44	93.50	69.43	56.11	40.35	151.90	123.40	87.81	75.23	142.31	178.72	111.29	108.45		122.00	293.00	167.00
Nb	6.80	7.02	4.56	3.64	2.99	13.10	7.60	4.62	3.77	8.57	15.77	7.74	8.29		10.00	36.00	20.00
Ba	238.53	148.42	146.60	342.30	43.29	169.70	482.90	70.51	113.54	762.92	700.14	224.48	425.88	103.00	274.00	528.00	418.00


La	8.50	9.31	9.45	7.77	3.82	18.75	25.71	8.90	8.06	26.09	30.90	7.84	8.38	10.47	9.79	31.79	18.87
Ce	20.35	22.54	20.54	16.79	8.92	50.90	63.10	23.08	19.89	59.41	70.09	21.36	21.78	27.83	28.15	70.15	45.07
Pr	2.87	3.24	2.51	2.06	1.24	5.75	6.45	3.18	2.75	7.76	8.92	3.23	3.28				
Nd	13.34	15.24	11.26	9.54	6.45	25.95	26.89	14.22	12.43	33.49	37.03	16.27	15.26	13.43	17.19	40.17	27.38
Sm	3.65	4.05	2.74	2.29	1.84	6.44	6.11	4.55	3.81	7.32	7.46	4.56	4.44	3.60	5.26	8.75	7.06
Eu	1.36	1.31	0.87	0.78	0.68	2.21	2.07	1.56	1.31	1.96	2.42	1.68	1.53	1.01	1.52	1.87	2.32
Gd	3.98	4.24	2.81	2.42	2.11	6.61	6.10	4.47	3.55	6.89	7.19	5.96	5.53	3.40	5.49	7.93	6.94
Tb	0.66	0.69	0.47	0.39	0.35	1.07	0.99	0.84	0.67	1.06	1.05	0.96	0.93				
Dy	4.27	4.38	2.87	2.44	2.14	5.68	5.75	5.02	4.05	6.69	5.91	5.58	5.70	3.35	5.65	7.94	5.55
Ho	0.89	0.93	0.57	0.49	0.43	1.11	1.16	1.03	0.83	1.41	1.18	1.13	1.19				
Er	2.38	2.41	1.61	1.39	1.19	2.68	3.00	2.66	2.24	3.66	2.99	3.04	3.08	1.99	2.97	4.35	2.55
Tm	0.34	0.34	0.24	0.21	0.18	0.37	0.44	0.39	0.32	0.56	0.42	0.45	0.46				
Yb	2.21	2.26	1.62	1.38	1.14	2.39	2.70	2.45	1.88	3.34	2.53	2.68	2.77	2.01	2.82	4.12	2.04
Lu	0.34	0.36	0.25	0.22	0.18	0.35	0.39	0.36	0.29	0.51	0.37	0.40	0.42	0.37	0.46	0.67	0.36
Cu	279.90	20.75	67.27	60.43	193.30	22.64	61.31	68.13	10.68	73.30	20.24	200.74	18.26	15.00	148.00	33.00	18.00
U	0.28	0.32	0.53	0.41	0.14	0.54	0.47	0.24	0.25	0.40	1.65	0.18	0.21				
Th	0.81	0.92	1.69	1.33	0.37	1.77	2.42	0.89	0.79	1.70	3.65	0.64	0.80				
Ta	0.51	0.54	0.36	0.28	0.23	0.82	0.24	0.19	0.20	0.48	1.00	0.59	0.59				
Hf	2.45	2.65	2.00	1.61	1.16	3.86	3.03	2.29	1.95	3.99	4.92	3.36	3.29				
Sc	48.93	48.88				32.09	41.73	39.63	48.49	38.97	28.38	38.58	41.10		45.00	35.00	30.00
Ga	20.82	19.25	16.88	15.52	13.62	21.02	21.94	18.46	19.54	21.78	22.31	19.06	18.76				
Pb	4.00	4.47	8.16	16.85	8.78	7.07	9.65	5.42	5.55	8.50	6.98	5.52	5.90				
Cs	0.87	0.45	1.17	0.65	0.30	1.05	1.15	0.73	1.21	7.09	1.39	1.41	2.00				


Sample	Inlier	Latitude N	Longitude W	Orientation	Ages (Ma)	Reference	SiO2	TiO2	Al2O3	Fe2O3t	MnO	MgO	CaO	Na2O	K2O	P2O5	L.O.I.	Total	V	Cr	Co	Ni	Zn	Rb	Sr	Y	Zr	Nb	Ba	La	Ce	Pr	Nd	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu	Cu	U	Th	Ta	Hf	Sc	Ga	Pb	Cs	
TA1	Zenaga	30°36'43.6"	7°17'13.4"	N45°	1639 ± 34 Ma	(this work)	49.31	1.64	13.79	14.51	0.23	6.34	6.70	3.26	0.96	0.14	2.47	99.34	343.24	18.95	51.53	39.42	126.00	35.95	192.75	23.11	88.44	6.80	238.53	8.50	20.35	2.87	13.34	3.65	1.36	3.98	0.66	4.27	0.89	2.38	0.34	2.21	0.34	279.90	0.28	0.81	0.51	2.45	48.93	20.82	4.00	0.87	
TA2	Zenaga	30°36'43.7"	7°17'13.3"	N30°		(this work)	49.73	1.58	13.48	14.35	0.23	6.68	6.20	4.06	0.32	0.19	3.04	99.84	317.91	18.83	54.75	37.22	158.84	8.84	220.51	23.50	93.50	7.02	148.42	9.31	22.54	3.24	15.24	4.05	1.31	4.24	0.69	4.38	0.93	2.41	0.34	2.26	0.36	20.75	0.32	0.92	0.54	2.65	48.88	19.25	4.47	0.45	
DZ36	Zenaga			N125°	1640 ± 2 Ma/ 1641 ± 3 Ma	Kouyate (2013)	50.7	0.85	14.55	10.27	0.16	7.31	10	2.17	0.74	0.07	2.48	99.49	240	307.7	46	143	90.36	26.9	215.2	16.3	69.43	4.56	146.6	9.45	20.5	2.51	11.3	2.74	0.87	2.81	0.47	2.87	0.57	1.61	0.24	1.62	0.25	67.27	0.53	1.69	0.36	2	16.9	8.1606	1.17		
DZ40	Zenaga			N140°	1656 ± 9 Ma	Kouyate (2013)	50.8	0.73	14.61	9.78	0.18	7.55	11.3	1.94	0.62	0.08	2.38	100.3	233.7	410.1	44.7	185.2	92.06	20.3	196.6	14.2	56.11	3.64	342.3	7.77	16.8	2.06	9.54	2.29	0.78	2.42	0.39	2.44	0.49	1.39	0.21	1.38	0.22	60.43	0.41	1.33	0.28	1.61	15.5	16.847	0.65		
AMHA69	Agadir Melloul				1654 ± 16 Ma	Kouyate (2013)	45.8	0.79	13.73	9.5	0.25	9.87	12.2	2.49	0.29	0.06	4.37	99.68	223	802.7	57.4	178.3	228.2	10.9	191.1	12.1	40.35	2.99	43.29	3.82	8.92	1.24	6.45	1.84	0.68	2.11	0.35	2.14	0.43	1.19	0.18	1.14	0.18	193.3	0.14	0.37	0.23	1.16	13.6	8.7784	0.3		
ZEN-4	Zenaga			N170°		Hafid (1999)	48.6	2.66	13.58	14.43	0.17	6.61	8.71	2.4	0.54	0.24	2.35	100.3	326.9	189.8	43.1	117	150.2	16.9	391.1	28.8	151.9	13.1	169.7	18.8	50.9	5.75	26	6.44	2.21	6.61	1.07	5.68	1.11	2.68	0.37	2.39	0.35	22.64	0.54	1.77	0.82	3.86	32.1	21	7.07	1.05	
ZEN-13	Zenaga			N125°		Hafid (1999)	49.7	2.09	14.08	14.83	0.19	5.76	8.51	2.59	0.78	0.33	2.18	101.1	317.1	96.76	50.5	69.71	152.6	20.9	410.5	30.9	123.4	7.6	482.9	25.7	63.1	6.45	26.9	6.11	2.07	6.1	0.99	5.75	1.16	3	0.44	2.7	0.39	61.31	0.47	2.42	0.24	3.03	41.7	21.9	9.65	1.15	
ZEN-1	Zenaga			N85°		Hafid (1999)	48.4	1.98	13.97	13.71	0.16	6.55	10.2	1.86	0.32	0.18	2.67	99.96	306.3	197.1	40.4	94.16	119.8	8.44	269.9	27.1	87.81	4.62	70.51	8.9	23.1	3.18	14.2	4.55	1.56	4.47	0.84	5.02	1.03	2.66	0.39	2.45	0.36	68.13	0.24	0.89	0.19	2.29	39.6	18.5	5.42	0.73	
ZEN-11	Zenaga			N170°		Hafid (1999)	48.3	1.64	13.97	13.16	0.2	7.73	10.3	2.1	0.64	0.15	2.02	100.3	332.9	181.6	55.9	121.7	161.3	19.7	256.2	22.9	75.23	3.77	113.5	8.06	19.9	2.75	12.4	3.81	1.31	3.55	0.67	4.05	0.83	2.24	0.32	1.88	0.29	10.68	0.25	0.79	0.2	1.95	48.5	19.5	5.55	1.21	
IT-5	Iguerd-Taifast					Hafid (1999)	49.1	2.05	13.5	15.17	0.26	5.28	8.52	2.46	1.15	0.428	1.46	99.4	291	114.1	44.4	44.2	131.3	63.4	372.9	37	142.3	8.57	762.9	26.1	59.4	7.76	33.5	7.32	1.96	6.89	1.06	6.69	1.41	3.66	0.56	3.34	0.51	73.3	0.4	1.7	0.48	3.99	39	21.8	8.5	7.09	
IT-7	Iguerd-Taifast			N00°		Hafid (1999)	50	2.52	13.67	15.02	0.17	5.24	6.98	2.45	1.79	0.362	1.73	99.91	339.5	126.9	46.1	64.28	129.5	68.1	370.3	29	178.7	15.8	700.1	30.9	70.1	8.92	37	7.46	2.42	7.19	1.05	5.91	1.18	2.99	0.42	2.53	0.37	20.24	1.65	3.65	1	4.92	28.4	22.3	6.98	1.39	
IT-12	Iguerd-Taifast			N30°		Hafid (1999)	49	2.17	13.06	14.87	0.23	5.91	10	1.91	0.36	0.189	2.07	99.81	400	113.1	44.1	79.64	134.4	24	253.6	29.8	111.3	7.74	224.5	7.84	21.4	3.23	16.3	4.56	1.68	5.96	0.96	5.58	1.13	3.04	0.45	2.68	0.4	200.7	0.18	0.64	0.59	3.36	38.6	19.1	5.52	1.41	
IT-13	Iguerd-Taifast			N45°		Hafid (1999)	48	2.06	13.2	15.54	0.26	6.24	8.4	2.73	0.95	0.174	1.99	99.52	413.6	180.3	43.6	92.72	155.3	45.9	200.6	28.8	108.5	8.29	425.9	8.38	21.8	3.28	15.3	4.44	1.53	5.53	0.93	5.7	1.19	3.08	0.46	2.77	0.42	18.26	0.21	0.8	0.59	3.29	41.1	18.8	5.9	2	
NC40	Tagragra of Akka			N50°-N110°		Mortaji (1989)	50.9	0.91	14.72	12.6	0.19	6.69	7.22	3.36	0.14	0.14	3.67	100.5	278	139	57	55		10	593	19.5		103	10.5	27.8					13.4	3.6	1.01	3.4		3.35	1.99		2.01	0.37	15								
NC379	Tagragra of Akka			N50°-N110°		Mortaji (1989)	47.6	1.93	13.62	15.63	0.2	6.29	8.82	2.04	0.77	0.24	2.69	99.81	367	114	54	95	116	43	185	30.9	122	10	274	9.79	28.2				17.2	5.26	1.52	5.49		5.65	2.97		2.82	0.46	148				45				
NC314aMor	Tagragra of Akka			N30°-N160°		Mortaji (1989)	44.4	2.86	15.25	16.53	0.18	5.95	7.16	2.58	1.04	0.69	3.8	100.4	189	180	46	109	153	29	247	41.7	293	36	528	31.8	70.2				40.2	8.75	1.87	7.93		7.94	4.35		4.12	0.67	33				35				
NC 361	Tagragra of Akka			N30°-N160°		Mortaji (1989)	49.2	2.99	12.75	16.06	0.12	5.21	4.36	2.63	1.31	0.5	5	100.1	367	118	46	65	62	81	120	26.2	167	20	418	18.9	45.1				27.4	7.06	2.32	6.94		5.55	2.55		2.04	0.36	18				30				

























































Highlights

- ▶ The Taghdout mafic sill of the Ifzwane Suite yielded an age of 1676 ± 37 Ma
- ▶ The maximum depositional age of the lower Anti-Atlas Supergroup ranges from 2049 Ma (bottom) to 700 Ma (top) and the provenance is mainly from the WAC
- ▶ A new 5-part lithostratigraphic framework for the Proterozoic lower Anti-Atlas Supergroup of the Anti-Atlas Belt is proposed
- ▶ The 1650-1640 Ma Zenaga and 885-883 Ma Iguerda-Taïfast LIPs provide important stratigraphic constraints
- ▶ Correlation between the lower Anti-Atlas Supergroup and the Taoudeni Basin succession in Mauritania

Declaration of interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: