[Click here to view linked References](#)

**Killing two birds with one stone:
how budding yeast Mps1 controls chromosome segregation and spindle
assembly checkpoint through phosphorylation of a single kinetochore protein**

Giorgia Benzi and Simonetta Piatti

CRBM, University of Montpellier, CNRS

1919 Route de Mende

34293 Montpellier (France)

Contact: simonetta.piatti@crbm.cnrs.fr

Keywords: mitosis; chromosome biorientation; spindle assembly checkpoint;

kinetochore; Mps1; Aurora B

Abstract

During mitosis, the identical sister chromatids of each chromosome must attach through their kinetochores to microtubules emanating from opposite spindle poles. This process, referred to as chromosome biorientation, is essential for equal partitioning of the genetic information to the two daughter cells. Defects in chromosome biorientation can give rise to aneuploidy, a hallmark of cancer and genetic diseases. A conserved surveillance mechanism called spindle assembly checkpoint (SAC) prevents the onset of anaphase until biorientation is attained. Key to chromosome biorientation is an error correction mechanism that, by disengaging faulty kinetochore-microtubule connections, allows kinetochores to establish proper bipolar attachments. Error correction relies on the Aurora B and Mps1 kinases that also promote SAC signalling, raising the possibility that they are part of a single sensory device responding to improper attachments and concomitantly controlling both their disengagement and a temporary mitotic arrest.

In budding yeast Aurora B and Mps1 promote error correction independently from one another, but while the substrates of Aurora B in this process are at least partially known, the mechanism underlying the involvement of Mps1 in the error correction pathway is unknown. Through the characterization of a novel *mps1* mutant and an unbiased genetic screen for extragenic suppressors, we recently gained evidence that a common mechanism based on Mps1-dependent phosphorylation of the Knl1/Spc105 kinetochore scaffold and subsequent recruitment of the Bub1 kinase is critical for the function of Mps1 in chromosome biorientation as well as for SAC activation (Benzi et al., EMBO Rep. 2020 Apr 19:e50257. doi: 10.15252/embr.202050257).

Introduction

Chromosome segregation is a vulnerable, error-prone process that must be tightly regulated in time and space. Several requirements must be fulfilled in order for daughter cells to get an equal chromosome complement. First, chromosomes must be faithfully duplicated into identical sister chromatids through DNA replication. Second, sister chromatids must be glued together by sister chromatid cohesion, which allows cells to distinguish genetically identical from distinct chromosomes. Third, sister chromatids need to attach via their kinetochores to microtubules emanating from opposite spindle poles, a process referred to as chromosome biorientation (reviewed in Bloom and Yeh, 2010). If this fails, a surveillance mechanism called Spindle Assembly Checkpoint (SAC) temporarily halts cell cycle progression in metaphase to allow error correction. Conversely, when all chromosomes are bipolarly attached, SAC is satisfied and anaphase can shortly ensue.

SAC detects the lack of attachment between spindle microtubules and kinetochores (reviewed in Maresca and Salmon, 2010), which are large protein assemblies residing at the centromere of each chromosome (reviewed in Musacchio and Desai 2017). Unattached kinetochores generate an alert signal that ultimately leads to the formation of a Mitotic Checkpoint Complex (MCC), made by Bub3, Mad2, BubR1 (Mad3 in yeast) and Cdc20, that binds to and inhibits the E3 ubiquitin ligase Anaphase Promoting Complex bound to its activator Cdc20 (APC^{Cdc20}). In turn, stabilisation of APC^{Cdc20} targets, such as securin and cyclin B, prevents sister chromatid separation and mitotic exit, thereby imposing a reversible metaphase arrest that provides the time for correction of improper kinetochore-microtubule connections (reviewed in Musacchio, 2015 ; Sacristan and Kops, 2015).

1 The dual-specificity mitotic kinase Mps1 (Monopolar spindle 1) is a key player in SAC
2 signaling. Mps1 was first identified in budding yeast (Winey et al. 1991) due to its
3 involvement in the duplication of spindle pole bodies (SPBs), the functional
4 equivalents of centrosomes in animal cells.
5

6 While the involvement of Mps1 in centrosome duplication in eukaryotic cells other
7 than yeast is controversial (Stucke et al. 2002; Fisk et al. 2003; Kwiatkowski et al.
8 2010), Mps1 is almost universally implicated in SAC activation (reviewed in Pachis
9 and Kops, 2018). Mps1 works at the apex of SAC signaling through phosphorylation
10 of the kinetochore protein Knl1/Spc105 on its N terminal MELT repeats, which in turn
11 is essential for the kinetochore recruitment of the Bub1-Bub3 complex (London et al.
12 2012; Shepperd et al. 2012; Yamagishi et al. 2012; Primorac et al. 2013). Mps1 also
13 phosphorylates Bub1 and Mad1, allowing their interaction and kinetochore
14 recruitment of Mad2 that acts as a catalyzer for MCC assembly (Faesen et al., 2017;
15 Ji et al., 2018; London and Biggins, 2014; Mora-Santos et al., 2016; Moyle et al.,
16 2014).
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

39 **Coupling chromosome biorientation with the Spindle Assembly Checkpoint**

40 During the SAC induced-metaphase arrest, cells rectify the tensionless kinetochore-
41 microtubule attachments with the aim of establishing chromosome biorientation. An
42 error correction mechanism involves the continuous detachment of improper
43 attachments to provide kinetochores with further opportunities to capture
44 microtubules (reviewed in Lampson and Grishchuk, 2017). When tension across
45 kinetochores is finally established, bipolar attachments are stabilized and
46 chromosomes congress to the cell's equator, due to the equilibrium between pulling
47 and cohesive forces (by microtubules and sister chromatid cohesion, respectively).
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 A key player in the error correction pathway is the Chromosomal Passenger Complex
2 (CPC). The CPC is composed of the Aurora B kinase, the inner centromere protein
3 INCENP, Survivin and Borealin (Ipl1, Slp1, Bir1 and Nbl1, respectively, in budding
4 yeast) (reviewed in Carmena et al., 2012). Aurora B detaches improper kinetochore-
5 microtubule connections by progressively phosphorylating Ndc80/Hec1, the subunit
6 of the NDC80 complex that directly interacts with microtubules (Cheeseman et al.
7 2006; DeLuca et al. 2006; Ciferri et al. 2008; Alushin et al. 2010; Tooley et al. 2011;
8 Zaytsev et al. 2014, 2015). Moreover, Aurora B/Ipl1 phosphorylates the DAM1
9 complex in budding yeast and its functional homolog SKA complex in mammalian
10 cells (Chan et al., 2012; Cheeseman et al., 2002; Kalantzaki et al., 2015; Lampert et
11 al., 2010; Tien et al., 2010), which couples chromosome movement to microtubule
12 depolymerization (Abad et al., 2014; Asbury et al., 2006; Grishchuk et al., 2008;
13 Lampert et al., 2010; Schmidt et al., 2012; Tien et al., 2010; Welburn et al., 2009;
14 Westermann et al., 2006).

15 Besides correcting improper kinetochore-microtubule connections, Aurora B is also
16 part of SAC signaling (reviewed in Krenn and Musacchio, 2015). On one side, it
17 generates unattached kinetochores that are sensed by SAC (Pinsky et al. 2006). On
18 the other, it facilitates the rapid recruitment of Mps1 to kinetochores (Santaguida et
19 al., 2011; Saurin et al., 2011) and phosphorylates the RVSF motif in KNL1 (Liu et al.
20 2010). The latter hampers the binding of the PP1 phosphatase, which is required for
21 SAC silencing (Liu et al., 2010; Pinsky et al., 2006; Rosenberg et al., 2011;
22 Vanoosthuyse and Hardwick, 2009). Thus, Aurora B contributes to chromosome
23 biorientation by both promoting error correction and sustaining SAC signaling.
24 Similarly, The Mps1 kinase has been involved in the correction of faulty kinetochore-
25 microtubule attachments, in addition to triggering SAC activity (Jones et al., 2005;

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
Maure et al., 2007; Santaguida et al., 2010; Hewitt et al., 2010; Maciejowski et al.,
2010). However, its role in the error correction pathway is far from being understood.
While in mammalian cells Mps1 seems to work in concert with Aurora B (Vigneron et
al. 2004; Jelluma et al. 2008; Saurin et al. 2011; Santaguida et al. 2011), in yeast the
two kinases appear to play independent roles (Maure et al. 2007; Storchová et al.
2011). In addition, while in mammalian cells the SKA complex has been proposed to
be a relevant Mps1 substrate in the error correction pathway (Maciejowski et al.,
2017), in the budding yeast *S. cerevisiae* the critical targets of Mps1 in this process
remain to be identified. Indeed, although *S.c.*Mps1 phosphorylates Dam1 and Ndc80
on several residues, none of these phosphorylations is required for chromosome
biorientation (Shimogawa et al. 2006; Kemmler et al. 2009; Kalantzaki et al. 2015).
The identification of the critical targets of *S.c.*Mps1 in chromosome biorientation is
not an easy task, and is complicated by the essential function of *S.c.*Mps1 in SPB
duplication and spindle assembly (Weiss and Winey 1996). Indeed, the initial
discovery that Mps1 is involved in the error correction pathway was based on a
complex experimental set-up (Maure et al. 2007).

41 **A novel genetic tool to dissect the function of Mps1 in chromosome**

42 **biorientation**

43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
We have recently characterized a novel temperature-sensitive mutant, named *mps1-3*, that bears a single substitution of serine 635 to phenylalanine. The *mps1-3* mutant is lethal at temperatures above 32°C, and at temperatures ranging from 32°C to 34°C is severely defective in chromosome biorientation but proficient in SPB duplication and spindle elongation (Benzi et al. 2020), unlike the majority of *mps1* mutants (Weiss and Winey 1996; Schutz and Winey 1998; Castillo et al. 2002; Araki et al.

2010). Contextually, *mps1-3* cells are also SAC-deficient, which allows them to progress through the cell cycle in the face of chromosome misalignment, thus accumulating massive aneuploidy. We could attribute lack of chromosome biorientation in *mps1-3* cells to faulty correction of improper attachments, rather than to a failure to establish kinetochore-microtubule connections, in agreement with earlier conclusions on Mps1 inhibition through an analogue-sensitive mutant (Jones et al. 2005; Maure et al. 2007). Consistent with defects in the error correction pathway, in *mps1-3* cells sister chromatids co-segregate preferentially toward the bud (Benzi et al. 2020), where the old SPB usually migrates during anaphase (Pereira et al. 2001). This phenotype had been previously described for *ip11* (Aurora B) mutants and is likely due to a delay in the gain of microtubule-nucleating activity of the new versus the old SPB, thereby biasing kinetochore attachment to the old, mature SPB (Tanaka et al. 2002).

To our surprise, the *mps1-3* mutation, which hits the kinase domain of the protein (aa 440-720), increases the *in vitro* kinase activity of Mps1 relative to the wild type protein. However, the Mps1-3 does not localize at kinetochores at restrictive temperatures. Although Mps1 can sustain SAC signaling away from kinetochores under non-physiological conditions (Maciejowski et al., 2017; Fraschini et al., 2001; Yuan et al., 2017), its kinetochore recruitment clearly contributes to proficient chromosome biorientation and SAC signaling in normal cells (Jelluma et al. 2010; Saurin et al. 2011; Heinrich et al. 2012; Nijenhuis et al. 2013; Zhu et al. 2013; Aravamudhan et al. 2016). Thus, the SAC and chromosome segregation defects of *mps1-3* mutant cells likely stem from the lack of phosphorylation of critical Mps1 substrates at kinetochores. Consistently, phosphorylation of known Mps1 substrates at kinetochores, such as Spc105/Knl1, Bub1, and Mad1 (Shepperd et al. 2012;

1 Yamagishi et al. 2012; Primorac et al. 2013; London and Biggins 2014; Moyle et al.
2 2014; Mora-Santos et al. 2016; Faesen et al. 2017; Qian et al. 2017; Ji et al. 2018) is
3
4 markedly compromised in *mps1-3* mutant cells. Additionally, artificial anchoring of
5
6 Mps1 to the kinetochore protein Mtw1 partially restores proper chromosome
7
8 segregation in *mps1-3* cells (Benzi et al. 2020).
9
10

11 The domain responsible for Mps1 kinetochore localization lies at the N-terminus of
12
13 the protein, far from the catalytic domain (Ji et al.; Liu et al. 2003; Stucke et al. 2004;
14
15 Kemmler et al. 2009; Araki et al. 2010; Hached et al. 2011; Nijenhuis et al. 2013;
16
17 Hiruma et al. 2015; Maciejowski et al. 2017). Thus, it is unlikely that the *mps1-3*
18
19 mutation interferes directly with Mps1 binding to its kinetochore receptor(s).
20
21

22
23 Nonetheless, Mps1 has been proposed to accelerate its own kinetochore turnover
24
25 through autophosphorylation (Sliedrecht et al. 2010; Kwiatkowski et al. 2010; Hewitt
26
27 et al. 2010; Santaguida et al. 2010; Jelluma et al. 2010; Wang et al. 2014; Dou et al.
28
29 2015; Koch et al. 2019; Hayward et al. 2019). Furthermore, activation of the Mps1
30
31 kinase leads to extensive conformational changes in the protein (Combes et al.
32
33 2018), which in turn could mask its kinetochore-binding regions. Thus, the increased
34
35 kinase activity of Mps1-3 could reduce its retention time at kinetochores, thereby
36
37 impairing SAC signaling and chromosome biorientation. We attempted to test directly
38
39 this hypothesis by introducing into the *mps1-3* allele the analog-sensitive *mps1-as1*
40
41 mutation, which inhibits the kinase in the presence of the ATP analog inhibitor 1-NM-
42
43 PP1 (Jones et al. 2005). However, the double mutant turned out to be lethal even in
44
45 the absence of the analog (our unpublished data), making this approach
46
47 inconclusive. The characterization of the internal suppressors that we identified (see
48
49 below) might provide us with alternative tools to test the contribution of the
50
51 hyperactivity of the Mps1-3 kinase in its own kinetochore turnover.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

The Spc105/Knl1 kinetochore scaffold as a critical target of Mps1 in chromosome biorientation beyond SAC signaling

Out of the two processes that are affected by the *mps1-3* mutation, i.e. SAC signaling and chromosome biorientation, only the latter is essential for cell viability and accounts for the lethality of *mps1-3* cells at high temperatures. This makes the *mps1-3* mutant a valuable genetic tool to underpin the molecular bases underlying the involvement of Mps1 in the error correction pathway. We therefore exploited the potential of budding yeast as a powerful genetic tool to identify spontaneous suppressors that enable *mps1-3* cells to divide at 34°C. This unbiased genetic screen was instrumental to our current model envisioning a common Mps1 substrate (Spc105/Knl1, i.e. the stone) in the control of both chromosome biorientation and SAC (i.e. the two birds).

Through our genetic screen we isolated several intragenic suppressors that carried second site mutations within the *mps1-3* allele. These were all missense mutations in the catalytic kinase domain. Their future characterization might reveal important insights into the control of Mps1 activity and kinetochore recruitment. We also isolated extragenic suppressors that bore, instead, a suppressing mutation in genes other than *MPS1*. Strikingly, all of them rescued both chromosome segregation and SAC defects of *mps1-3* cells, suggesting that a common mechanism underlies the role of Mps1 in these processes. Among these suppressors, we found mutations in *SPC105*, which codes for the kinetochore protein Spc105/Knl1, and *GLC7*, which codes for the catalytic subunit of the protein phosphatase PP1. Importantly, the *spc105* suppressing mutations hit the motif (RVSF) for Spc105 binding to PP1, while the *GLC7* suppressors carried a missense mutation of Phe256, which resides in the

hydrophobic groove involved in PP1 interaction with partners carrying the RV/IXF consensus motif (Wu and Tatchell 2001). As mentioned above, Mps1-dependent phosphorylation of Spc105/Knl1 at its MELT repeats recruits the Bub3-Bub1 complex and is critical for SAC signaling (London et al. 2012; Shepperd et al. 2012; Yamagishi et al. 2012; Primorac et al. 2013). Upon chromosome biorientation, Spc105/Knl1 phosphorylation is reversed by the phosphatase PP1, which binds the RVSF motif and silences the SAC (Pinsky et al. 2009; Vanoosthuyse and Hardwick 2009; Liu et al. 2010; Rosenberg et al. 2011; Meadows et al. 2011; London et al. 2012; Moura et al. 2017). Importantly, the *spc105* and *GLC7* suppressors that we isolated did not restore Mps1 kinetochore localization, strongly suggesting that PP1 opposes Mps1 activity at kinetochores for both proper chromosome biorientation and SAC signaling. The antagonism between Mps1 and PP1 has been well characterized for SAC, while its involvement in the error correction pathway had not been established. Altogether, our data have two important implications: first, the same Mps1-dependent sensory device may be used for both correction of improper kinetochore-microtubule attachments and SAC signaling, in line with previous proposals (Musacchio 2011; Caldas et al. 2013); second, PP1 is expected to stabilize kinetochore-microtubule attachments in yeast, as it does in human cells (Liu et al. 2010).

On the basis of our results, we wondered if Spc105 phosphorylation and subsequent Bub1 recruitment could be the main function of Mps1 in the error correction pathway, as it is in SAC activation. Consistent with this idea, Bub1 is no longer recruited to kinetochores in *mps1-3* mutant cells, while it regains kinetochore binding in the suppressors. Furthermore, artificial tethering of Bub1 to Spc105 is sufficient to restore balanced chromosome segregation and partial SAC signaling in *mps1-3* cells (Benzi et al. 2020). Thus, our data strongly imply that Mps1 promotes the error

1 correction pathway and SAC activity via a common mechanism that relies on
2 Spc105/Knl1 phosphorylation (Fig. 1). Along this line, phosphorylation of other Mps1
3
4 kinetochore targets, such as Dam1 and Ndc80, turned out to be dispensable for
5
6 equal chromosome segregation (Shimogawa et al. 2006; Kemmler et al. 2009;
7
8 Kalantzaki et al. 2015).
9

10
11 The main conclusion of our paper is consistent with several published observations: a
12
13 *spc105-6A* yeast mutant with unphosphorylatable MELT repeats is defective in Bub1
14
15 kinetochore recruitment and displays chromosome missegregation (London et al.
16
17 2012); phospho-mimicking mutations in the MELT repeats of Spc7 (the fission yeast
18
19 homologue of Spc105/Knl1) partially suppress the chromosome biorientation defects
20
21 of fission yeast *mps1Δ* cells, though in fission yeast Mps1 only modestly contributes
22
23 to chromosome biorientation (Yamagishi et al. 2012). Finally, Knl1 depletion in
24
25 human cells impairs Aurora B activity and chromosome alignment on the metaphase
26
27 plate (Caldas et al. 2013).
28
29
30
31
32
33
34
35

36 **Is Bub1 kinetochore recruitment the only function of Mps1 in chromosome** 37 38 **biorientation?** 39

40
41 Bub1 has been extensively involved in chromosome biorientation through
42
43 phosphorylation of T120 of histone H2A (S121 in yeast), which in turn recruits
44
45 shugoshin proteins (Sgo1 in budding yeast) to pericentromeres (Fernius and
46
47 Hardwick 2005; Kawashima et al. 2010; Yamagishi et al. 2010; Liu et al. 2015).
48
49 Shugoshins, in turn, promote recruitment of the CPC and biorientation of sister
50
51 chromatids (Indjeian et al. 2005; Kawashima et al. 2007; Huang et al. 2007; Kiburz et
52
53 al. 2008; Tsukahara et al. 2010; Yamagishi et al. 2010; Verzijlbergen et al. 2014; Liu
54
55 et al. 2015). Consistent with Bub1 being required for Sgo1 accumulation at
56
57
58
59
60
61
62
63
64
65

pericentromeres, localization of Sgo1 at kinetochores is compromised in *mps1-3* mutant cells. However, the synthetic lethality between the *mps1-3* mutation and *SGO1* deletion suggests that Mps1 could have at least one additional function in chromosome biorientation, besides recruiting Bub1 to kinetochores. Indeed, Bub1 and Sgo1 are not essential proteins, in contrast to Mps1. Furthermore, artificial tethering of Bub1 to Spc105 cannot fully rescue the chromosome biorientation defects and temperature-sensitivity of *mps1-3* cells (Benzi et al. 2020), suggesting that other Mps1 substrates and/or other factors binding to Spc105 might intervene. One intriguing hypothesis, which would be supported by our genetic data, is that Mps1 could keep PP1 inhibited at kinetochores. PP1, in turn, stabilizes bipolar attachments (Liu et al. 2010).

Another interesting candidate that might play a function complementary to Bub1 in chromosome biorientation, downstream of Mps1, is the microtubule rescue factor Stu1, which is a member of the CLASP family (Al-Bassam and Chang 2011). At kinetochores, Stu1 stabilizes microtubules, and, consistently, it gets sequestered at unattached kinetochores to facilitate their own capturing (Funk et al. 2014). Stu1 sequestration at unattached kinetochores requires Mps1 and Spc105 phosphorylation of its MELT repeats, but not Bub1 (Kolenda et al. 2018), suggesting that Stu1 and Bub1 might cooperate at Spc105 to achieve successful bipolar attachment. Stu1 recruits Stu2, a microtubule-associated protein of the XMAP215 family, to polymerize kinetochore-microtubules (Vasileva et al. 2017). Remarkably, Stu2 has been recently proposed to mediate an error correction pathway independent of Aurora B/Ipl1 in budding yeast (Miller et al. 2016, 2019). Whether and how Mps1 modulates Stu1 and/or Stu2 function at kinetochores to achieve

1 chromosome biorientation is an important question that deserves future
2 investigations.

3
4 Another important question that remains open is whether an interplay between
5 S.c.Mps1 and the CPC in the error correction pathway can be completely ruled out.
6

7 As already mentioned, in budding yeast Mps1 and Ipl1 were proposed to operate
8 independently from one another (Maure et al. 2007; Storchová et al. 2011). Our data
9 are consistent with this view. First, phosphorylation of the known kinetochore
10 substrate of Ipl1/Aurora B Dam1 is not affected in *mps1-3* cells (Benzi et al. 2020).
11

12 Similarly, Mps1 inhibition does not perturb phosphorylation of Aurora B substrates in
13 human cells (Hewitt et al. 2010; Maciejowski et al. 2010; Santaguida et al. 2010).
14

15 Second, the temperature-sensitivity of *mps1-3* cells cannot be suppressed by either
16 the *NDC80-13D* phospho-mimicking mutant allele where all Mps1-dependent
17 phosphorylation sites of Ndc80 have been replaced by aspartate (Kemmler et al.
18 2009), or the *DAM1-S221F* allele (also known as *DAM1-765*, (Shimogawa et al.
19 2006), which was previously found to suppress the temperature-sensitivity of *ipl1-321*
20 mutant cells (Shimogawa et al. 2010; our unpublished data). Third, mutations in the
21 CPC subunit Sli15 that activate Ipl1/Aurora B and rescue the sickness of *bir1Δ*,
22 *bub1Δ* and *sgo1Δ* cells (Campbell and Desai 2013) do not suppress the lethality and
23 chromosome segregation defects of *mps1-3* cells at high temperature (Benzi et al.
24 2020). Thus, we favour the hypothesis that Mps1 sets in motion an error correction
25 mechanism that does not involve Aurora B/Ipl1. However, it remains possible that,
26 while differing in their respective upstream targets, the Mps1- and Ipl1-dependent
27 error correction pathways ultimately converge on shared downstream effectors to
28 destabilise flawed attachments. Future work will certainly shed light on this exciting
29 possibility.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

ACKNOWLEDGMENTS

We thank all members of Piatti's lab for useful discussions. Work in Piatti's lab is supported by Fondation pour la Recherche Médicale (DEQ20150331740 to S.P.), Agence Nationale de la Recherche (ANR-18-CE13-0015-01 to S.P.) and by Labex EpiGenMed (PhD fellowship to G.B.).

REFERENCES

- Abad MA, Medina B, Santamaria A, et al (2014) Structural basis for microtubule recognition by the human kinetochore Ska complex. *Nat Commun* 5:2964. <https://doi.org/10.1038/ncomms3964>
- Al-Bassam J, Chang F (2011) Regulation of microtubule dynamics by TOG-domain proteins XMAP215/Dis1 and CLASP. *Trends Cell Biol* 21:604–614. <https://doi.org/10.1016/j.tcb.2011.06.007>
- Alushin GM, Ramey VH, Pasqualato S, et al (2010) The Ndc80 kinetochore complex forms oligomeric arrays along microtubules. *Nature* 467:805–810. <https://doi.org/10.1038/nature09423>
- Araki Y, Gombos L, Migueleti SPS, et al (2010) N-terminal regions of Mps1 kinase determine functional bifurcation. *J Cell Biol* 189:41–56. <https://doi.org/10.1083/jcb.200910027>
- Aravamudhan P, Chen R, Roy B, et al (2016) Dual mechanisms regulate the recruitment of spindle assembly checkpoint proteins to the budding yeast kinetochore. *Molecular Biology of the Cell* 27:3405–3417. <https://doi.org/10.1091/mbc.e16-01-0007>
- Asbury CL, Gestaut DR, Powers AF, et al (2006) The Dam1 kinetochore complex harnesses microtubule dynamics to produce force and movement. *Proceedings of the National Academy of Sciences* 103:9873–9878. <https://doi.org/10.1073/pnas.0602249103>
- Benzi G, Camasses A, Atsunori Y, et al (2020) A common molecular mechanism underlies the role of Mps1 in chromosome biorientation and the spindle assembly checkpoint. *EMBO Rep*. <https://doi.org/10.15252/embr.202050257>
- Bloom K, Yeh E (2010) Tension Management in the Kinetochore. *Curr Biol* 20:R1040–R1048. <https://doi.org/10.1016/j.cub.2010.10.055>
- Caldas GV, DeLuca KF, DeLuca JG (2013) KNL1 facilitates phosphorylation of outer kinetochore proteins by promoting Aurora B kinase activity. *The Journal of Cell Biology* 203:957–969. <https://doi.org/10.1083/jcb.201306054>
- Campbell CS, Desai A (2013) Tension sensing by Aurora B kinase is independent of survivin-based centromere localization. *Nature* 497:118–121. <https://doi.org/10.1038/nature12057>
- Carmena M, Wheelock M, Funabiki H, Earnshaw WC (2012) The chromosomal passenger complex (CPC): from easy rider to the godfather of mitosis. *Nature Reviews Molecular Cell Biology* 13:789–803. <https://doi.org/10.1038/nrm3474>
- Castillo AR, Meehl JB, Morgan G, et al (2002) The yeast protein kinase Mps1p is required for assembly of the integral spindle pole body component Spc42p. *J Cell Biol* 156:453–465. <https://doi.org/10.1083/jcb.200111025>
- Chan YW, Jeyaprakash AA, Nigg EA, Santamaria A (2012) Aurora B controls kinetochore-microtubule attachments by inhibiting Ska complex-KMN network interaction. *J Cell Biol* 196:563–571. <https://doi.org/10.1083/jcb.201109001>
- Cheeseman IM, Anderson S, Jwa M, et al (2002) Phospho-regulation of kinetochore-microtubule attachments by the Aurora kinase Ipl1p. *Cell* 111:163–172. [https://doi.org/10.1016/s0092-8674\(02\)00973-x](https://doi.org/10.1016/s0092-8674(02)00973-x)

- 1 Cheeseman IM, Chappie JS, Wilson-Kubalek EM, Desai A (2006) The Conserved KMN
2 Network Constitutes the Core Microtubule-Binding Site of the Kinetochore. *Cell*
3 127:983–997. <https://doi.org/10.1016/j.cell.2006.09.039>
- 4 Ciferri C, Pasqualato S, Screpanti E, et al (2008) Implications for kinetochore-microtubule
5 attachment from the structure of an engineered Ndc80 complex. *Cell* 133:427–439.
6 <https://doi.org/10.1016/j.cell.2008.03.020>
- 7
8 Combes G, Barysz H, Garand C, et al (2018) Mps1 Phosphorylates Its N-Terminal Extension
9 to Relieve Autoinhibition and Activate the Spindle Assembly Checkpoint. *Current*
10 *Biology* 28:872-883.e5. <https://doi.org/10.1016/j.cub.2018.02.002>
- 11 DeLuca JG, Gall WE, Ciferri C, et al (2006) Kinetochore microtubule dynamics and
12 attachment stability are regulated by Hec1. *Cell* 127:969–982.
13 <https://doi.org/10.1016/j.cell.2006.09.047>
- 14
15 Ditchfield C, Johnson VL, Tighe A, et al (2003) Aurora B couples chromosome alignment
16 with anaphase by targeting BubR1, Mad2, and Cenp-E to kinetochores. *J Cell Biol*
17 161:267–280. <https://doi.org/10.1083/jcb.200208091>
- 18
19 Dou Z, Liu X, Wang W, et al (2015) Dynamic localization of Mps1 kinase to kinetochores is
20 essential for accurate spindle microtubule attachment. *Proc Natl Acad Sci USA*
21 112:E4546-4555. <https://doi.org/10.1073/pnas.1508791112>
- 22
23 Egloff MP, Johnson DF, Moorhead G, et al (1997) Structural basis for the recognition of
24 regulatory subunits by the catalytic subunit of protein phosphatase 1. *EMBO J*
25 16:1876–1887. <https://doi.org/10.1093/emboj/16.8.1876>
- 26
27 Faesen AC, Thanasoula M, Maffini S, et al (2017) Basis of catalytic assembly of the mitotic
28 checkpoint complex. *Nature* 542:498–502. <https://doi.org/10.1038/nature21384>
- 29
30 Famulski JK, Chan GK (2007) Aurora B kinase-dependent recruitment of hZW10 and hROD
31 to tensionless kinetochores. *Curr Biol* 17:2143–2149.
32 <https://doi.org/10.1016/j.cub.2007.11.037>
- 33
34 Fernius J, Hardwick KG (2005) Bub1 Kinase Targets Sgo1 to Ensure Efficient Chromosome
35 Biorientation in Budding Yeast Mitosis. *PLoS Genet* preprint:e213.
36 <https://doi.org/10.1371/journal.pgen.0030213.eor>
- 37
38 Fisk HA, Mattison CP, Winey M (2003) Human Mps1 protein kinase is required for
39 centrosome duplication and normal mitotic progression. *PNAS* 100:14875–14880.
40 <https://doi.org/10.1073/pnas.2434156100>
- 41
42 Funk C, Schmeiser V, Ortiz J, Lechner J (2014) A TOGL domain specifically targets yeast
43 CLASP to kinetochores to stabilize kinetochore microtubules. *J Cell Biol* 205:555–
44 571. <https://doi.org/10.1083/jcb.201310018>
- 45
46 Grishchuk EL, Spiridonov IS, Volkov VA, et al (2008) Different assemblies of the DAM1
47 complex follow shortening microtubules by distinct mechanisms. *Proc Natl Acad Sci*
48 *USA* 105:6918–6923. <https://doi.org/10.1073/pnas.0801811105>
- 49
50 Hached K, Xie SZ, Buffin E, et al (2011) Mps1 at kinetochores is essential for female mouse
51 meiosis I. *Development* 138:2261–2271. <https://doi.org/10.1242/dev.061317>
- 52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 Hayward D, Bancroft J, Mangat D, et al (2019) Checkpoint signaling and error correction
2 require regulation of the MPS1 T-loop by PP2A-B56. *The Journal of Cell Biology*
3 218:3188–3199. <https://doi.org/10.1083/jcb.201905026>
- 4 Heinrich S, Windecker H, Hustedt N, Hauf S (2012) Mph1 kinetochore localization is crucial
5 and upstream in the hierarchy of spindle assembly checkpoint protein recruitment to
6 kinetochores. *J Cell Sci* 125:4720–4727. <https://doi.org/10.1242/jcs.110387>
- 8 Hendrickx A, Beullens M, Ceulemans H, et al (2009) Docking motif-guided mapping of the
9 interactome of protein phosphatase-1. *Chem Biol* 16:365–371.
10 <https://doi.org/10.1016/j.chembiol.2009.02.012>
- 12 Hewitt L, Tighe A, Santaguida S, et al (2010) Sustained Mps1 activity is required in mitosis to
13 recruit O-Mad2 to the Mad1–C-Mad2 core complex. *J Cell Biol* 190:25–34.
14 <https://doi.org/10.1083/jcb.201002133>
- 16 Hiruma Y, Sacristan C, Pachis ST, et al (2015) Competition between MPS1 and microtubules
17 at kinetochores regulates spindle checkpoint signaling. *Science* 348:1264–1267.
18 <https://doi.org/10.1126/science.aaa4055>
- 20 Huang H, Feng J, Famulski J, et al (2007) Tripin/hSgo2 recruits MCAK to the inner
21 centromere to correct defective kinetochore attachments. *Journal of Cell Biology*
22 177:413–424. <https://doi.org/10.1083/jcb.200701122>
- 24 Indjeian VB, Stern BM, Murray AW (2005) The Centromeric Protein Sgo1 Is Required to
25 Sense Lack of Tension on Mitotic Chromosomes. *Science* 307:130–133.
26 <https://doi.org/10.1126/science.1101366>
- 28 Jelluma N, Brenkman AB, van den Broek NJF, et al (2008) Mps1 phosphorylates Borealin to
29 control Aurora B activity and chromosome alignment. *Cell* 132:233–246.
30 <https://doi.org/10.1016/j.cell.2007.11.046>
- 32 Jelluma N, Dansen TB, Slidrecht T, et al (2010) Release of Mps1 from kinetochores is
33 crucial for timely anaphase onset. *J Cell Biol* 191:281–290.
34 <https://doi.org/10.1083/jcb.201003038>
- 36 Ji Z, Gao H, Jia L, et al (2018) A sequential multi-target Mps1 phosphorylation cascade
37 promotes spindle checkpoint signaling. *eLife* 6:. <https://doi.org/10.7554/eLife.22513>
- 39 Ji Z, Gao H, Yu H Kinetochore attachment sensed by competitive Mps1 and microtubule
40 binding to Ndc80C. 6
- 42 Jones MH, Huneycutt BJ, Pearson CG, et al (2005) Chemical Genetics Reveals a Role for
43 Mps1 Kinase in Kinetochore Attachment during Mitosis. *Current Biology* 15:160–165.
44 <https://doi.org/10.1016/j.cub.2005.01.010>
- 46 Kalantzaki M, Kitamura E, Zhang T, et al (2015) Kinetochore-microtubule error correction is
47 driven by differentially regulated interaction modes. *Nat Cell Biol* 17:421–433.
48 <https://doi.org/10.1038/ncb3128>
- 50 Kawashima SA, Tsukahara T, Langeegger M, et al (2007) Shugoshin enables tension-
51 generating attachment of kinetochores by loading Aurora to centromeres. *Genes Dev*
52 21:420–435. <https://doi.org/10.1101/gad.1497307>

- 1 Kawashima SA, Yamagishi Y, Honda T, et al (2010) Phosphorylation of H2A by Bub1
2 Prevents Chromosomal Instability Through Localizing Shugoshin. *Science* 327:172–
3 177. <https://doi.org/10.1126/science.1180189>
- 4 Kemmler S, Stach M, Knapp M, et al (2009) Mimicking Ndc80 phosphorylation triggers
5 spindle assembly checkpoint signalling. *The EMBO Journal* 28:1099–1110.
6 <https://doi.org/10.1038/emboj.2009.62>
- 7
8 Kiburz BM, Amon A, Marston AL (2008) Shugoshin promotes sister kinetochore biorientation
9 in *Saccharomyces cerevisiae*. *Molecular biology of the cell* 19:1199–1209
- 10
11 Koch LB, Opoku KN, Deng Y, et al (2019) Autophosphorylation is sufficient to release Mps1
12 kinase from native kinetochores. *Proc Natl Acad Sci USA* 116:17355–17360.
13 <https://doi.org/10.1073/pnas.1901653116>
- 14
15
16 Kolenda C, Ortiz J, Pelzl M, et al (2018) Unattached kinetochores drive their own capturing
17 by sequestering a CLASP. *Nature Communications* 9:.
18 <https://doi.org/10.1038/s41467-018-03108-z>
- 19
20
21 Krenn V, Musacchio A (2015) The Aurora B Kinase in Chromosome Bi-Orientation and
22 Spindle Checkpoint Signaling. *Frontiers in Oncology* 5:.
23 <https://doi.org/10.3389/fonc.2015.00225>
- 24
25 Kwiatkowski N, Jelluma N, Filippakopoulos P, et al (2010) Small Molecule Kinase Inhibitors
26 Provide Insight into Mps1 Cell Cycle Function. *Nat Chem Biol* 6:359–368.
27 <https://doi.org/10.1038/nchembio.345>
- 28
29
30 Lampert F, Hornung P, Westermann S (2010) The Dam1 complex confers microtubule plus
31 end-tracking activity to the Ndc80 kinetochore complex. *The Journal of Cell Biology*
32 189:641–649. <https://doi.org/10.1083/jcb.200912021>
- 33
34 Lampson MA, Grishchuk EL (2017) Mechanisms to Avoid and Correct Erroneous
35 Kinetochore-Microtubule Attachments. *Biology* 6:1.
36 <https://doi.org/10.3390/biology6010001>
- 37
38 Liu D, Vleugel M, Backer CB, et al (2010) Regulated targeting of protein phosphatase 1 to
39 the outer kinetochore by KNL1 opposes Aurora B kinase. *The Journal of Cell Biology*
40 188:809–820. <https://doi.org/10.1083/jcb.201001006>
- 41
42 Liu H, Qu Q, Warrington R, et al (2015) Mitotic Transcription Installs Sgo1 at Centromeres to
43 Coordinate Chromosome Segregation. *Mol Cell* 59:426–436.
44 <https://doi.org/10.1016/j.molcel.2015.06.018>
- 45
46
47 Liu S-T, Chan GKT, Hittle JC, et al (2003) Human MPS1 kinase is required for mitotic arrest
48 induced by the loss of CENP-E from kinetochores. *Mol Biol Cell* 14:1638–1651.
49 <https://doi.org/10.1091/mbc.02-05-0074>
- 50
51
52 London N, Biggins S (2014) Mad1 kinetochore recruitment by Mps1-mediated
53 phosphorylation of Bub1 signals the spindle checkpoint. *Genes & Development*
54 28:140–152. <https://doi.org/10.1101/gad.233700.113>
- 55
56 London N, Ceto S, Ranish JA, Biggins S (2012) Phosphoregulation of Spc105 by Mps1 and
57 PP1 Regulates Bub1 Localization to Kinetochores. *Current Biology* 22:900–906.
58 <https://doi.org/10.1016/j.cub.2012.03.052>
- 59
60
61
62
63
64
65

- 1 Maciejowski J, Drechsler H, Grundner-Culemann K, et al (2017) Mps1 Regulates
2 Kinetochore-Microtubule Attachment Stability via the Ska Complex to Ensure Error-
3 Free Chromosome Segregation. *Developmental Cell* 41:143-156.e6.
4 <https://doi.org/10.1016/j.devcel.2017.03.025>
- 5 Maciejowski J, George KA, Terret M-E, et al (2010) Mps1 directs the assembly of Cdc20
6 inhibitory complexes during interphase and mitosis to control M phase timing and
7 spindle checkpoint signaling. *J Cell Biol* 190:89-100.
8 <https://doi.org/10.1083/jcb.201001050>
- 9
10 Maresca TJ, Salmon ED (2010) Welcome to a new kind of tension: translating kinetochore
11 mechanics into a wait-anaphase signal. *J Cell Sci* 123:825-835.
12 <https://doi.org/10.1242/jcs.064790>
- 13
14 Maure J-F, Kitamura E, Tanaka TU (2007) Mps1 Kinase Promotes Sister-Kinetochore Bi-
15 orientation by a Tension-Dependent Mechanism. *Current Biology* 17:2175-2182.
16 <https://doi.org/10.1016/j.cub.2007.11.032>
- 17
18 Maure J-F, Komoto S, Oku Y, et al (2011) The Ndc80 loop region facilitates formation of
19 kinetochore attachment to the dynamic microtubule plus end. *Curr Biol* 21:207-213.
20 <https://doi.org/10.1016/j.cub.2010.12.050>
- 21
22 Meadows JC, Shepperd LA, Vanoosthuyse V, et al (2011) Spindle checkpoint silencing
23 requires association of PP1 to both Spc7 and kinesin-8 motors. *Dev Cell* 20:739-750.
24 <https://doi.org/10.1016/j.devcel.2011.05.008>
- 25
26 Miller MP, Asbury CL, Biggins S (2016) A TOG Protein Confers Tension Sensitivity to
27 Kinetochore-Microtubule Attachments. *Cell* 165:1428-1439.
28 <https://doi.org/10.1016/j.cell.2016.04.030>
- 29
30 Miller MP, Evans RK, Zelter A, et al (2019) Kinetochore-associated Stu2 promotes
31 chromosome biorientation in vivo. *PLoS Genet* 15:.
32 <https://doi.org/10.1371/journal.pgen.1008423>
- 33
34 Mora-Santos MDM, Hervas-Aguilar A, Sewart K, et al (2016) Bub3-Bub1 Binding to
35 Spc7/KNL1 Toggles the Spindle Checkpoint Switch by Licensing the Interaction of
36 Bub1 with Mad1-Mad2. *Curr Biol* 26:2642-2650.
37 <https://doi.org/10.1016/j.cub.2016.07.040>
- 38
39 Moura M, Osswald M, Leça N, et al (2017) Protein Phosphatase 1 inactivates Mps1 to
40 ensure efficient Spindle Assembly Checkpoint silencing. *eLife* 6:.
41 <https://doi.org/10.7554/eLife.25366>
- 42
43 Moyle MW, Kim T, Hattersley N, et al (2014) A Bub1-Mad1 interaction targets the Mad1-
44 Mad2 complex to unattached kinetochores to initiate the spindle checkpoint. *J Cell*
45 *Biol* 204:647-657. <https://doi.org/10.1083/jcb.201311015>
- 46
47 Musacchio A (2015) The Molecular Biology of Spindle Assembly Checkpoint Signaling
48 Dynamics. *Current Biology* 25:R1002-R1018.
49 <https://doi.org/10.1016/j.cub.2015.08.051>
- 50
51 Musacchio A (2011) Spindle assembly checkpoint: the third decade. *Philos Trans R Soc*
52 *Lond B Biol Sci* 366:3595-3604. <https://doi.org/10.1098/rstb.2011.0072>
- 53
54 Musacchio A, Desai A (2017) A Molecular View of Kinetochore Assembly and Function.
55 *Biology (Basel)* 6:.
56 <https://doi.org/10.3390/biology6010005>

- 1 Nijenhuis W, von Castelmur E, Littler D, et al (2013) A TPR domain–containing N-terminal
2 module of MPS1 is required for its kinetochore localization by Aurora B. *J Cell Biol*
3 201:217–231. <https://doi.org/10.1083/jcb.201210033>
- 4 Pachis ST, Kops GJPL (2018) Leader of the SAC: molecular mechanisms of Mps1/TTK
5 regulation in mitosis. *Open Biol* 8:180109. <https://doi.org/10.1098/rsob.180109>
- 6
7 Pereira G, Tanaka TU, Nasmyth K, Schiebel E (2001) Modes of spindle pole body
8 inheritance and segregation of the Bfa1p–Bub2p checkpoint protein complex. *EMBO*
9 *J* 20:6359–6370. <https://doi.org/10.1093/emboj/20.22.6359>
- 10
11 Pinsky BA, Kung C, Shokat KM, Biggins S (2006) The Ipl1-Aurora protein kinase activates
12 the spindle checkpoint by creating unattached kinetochores. *Nat Cell Biol* 8:78–83.
13 <https://doi.org/10.1038/ncb1341>
- 14
15
16 Pinsky BA, Nelson CR, Biggins S (2009) Protein Phosphatase 1 Regulates Exit from the
17 Spindle Checkpoint in Budding Yeast. *Current Biology* 19:1182–1187.
18 <https://doi.org/10.1016/j.cub.2009.06.043>
- 19
20
21 Primorac I, Weir JR, Chirolì E, et al (2013) Bub3 reads phosphorylated MELT repeats to
22 promote spindle assembly checkpoint signaling. *eLife* 2:e01030.
23 <https://doi.org/10.7554/eLife.01030>
- 24
25 Qian J, García-Gimeno MA, Beullens M, et al (2017) An Attachment-Independent
26 Biochemical Timer of the Spindle Assembly Checkpoint. *Molecular Cell* 68:715–
27 730.e5. <https://doi.org/10.1016/j.molcel.2017.10.011>
- 28
29 R. F, A. B, G. L, S. P (2001) Role of the kinetochore protein Ndc10 in mitotic checkpoint
30 activation in *Saccharomyces cerevisiae*. *Molecular Genetics and Genomics* 266:115–
31 125. <https://doi.org/10.1007/s004380100533>
- 32
33
34 Rosenberg JS, Cross FR, Funabiki H (2011) KNL1/Spc105 Recruits PP1 to Silence the
35 Spindle Assembly Checkpoint. *Current Biology* 21:942–947.
36 <https://doi.org/10.1016/j.cub.2011.04.011>
- 37
38 Sacristan C, Kops GJPL (2015) Joined at the hip: kinetochores, microtubules, and spindle
39 assembly checkpoint signaling. *Trends Cell Biol* 25:21–28.
40 <https://doi.org/10.1016/j.tcb.2014.08.006>
- 41
42 Santaguida S, Tighe A, D'Alise AM, et al (2010) Dissecting the role of MPS1 in chromosome
43 biorientation and the spindle checkpoint through the small molecule inhibitor
44 reversine. *J Cell Biol* 190:73–87. <https://doi.org/10.1083/jcb.201001036>
- 45
46
47 Santaguida S, Vernieri C, Villa F, et al (2011) Evidence that Aurora B is implicated in spindle
48 checkpoint signalling independently of error correction: Aurora B is directly implicated
49 in spindle checkpoint. *The EMBO Journal* 30:1508–1519.
50 <https://doi.org/10.1038/emboj.2011.70>
- 51
52
53 Saurin AT, van der Waal MS, Medema RH, et al (2011) Aurora B potentiates Mps1 activation
54 to ensure rapid checkpoint establishment at the onset of mitosis. *Nat Commun* 2:316.
55 <https://doi.org/10.1038/ncomms1319>
- 56
57 Schmidt JC, Arthanari H, Boeszoermenyi A, et al (2012) The kinetochore-bound Ska1
58 complex tracks depolymerizing microtubules and binds to curved protofilaments. *Dev*
59 *Cell* 23:968–980. <https://doi.org/10.1016/j.devcel.2012.09.012>
- 60
61
62
63
64
65

- Schutz AR, Winey M (1998) New Alleles of the Yeast MPS1 Gene Reveal Multiple Requirements in Spindle Pole Body Duplication. *Mol Biol Cell* 9:759–774
- Shepherd LA, Meadows JC, Sochaj AM, et al (2012) Phosphodependent recruitment of Bub1 and Bub3 to Spc7/KNL1 by Mph1 kinase maintains the spindle checkpoint. *Curr Biol* 22:891–899. <https://doi.org/10.1016/j.cub.2012.03.051>
- Shimogawa MM, Graczyk B, Gardner MK, et al (2006) Mps1 phosphorylation of Dam1 couples kinetochores to microtubule plus ends at metaphase. *Curr Biol* 16:1489–1501. <https://doi.org/10.1016/j.cub.2006.06.063>
- Shimogawa MM, Wargacki MM, Muller EG, Davis TN (2010) Laterally attached kinetochores recruit the checkpoint protein Bub1, but satisfy the spindle checkpoint. *Cell Cycle* 9:3619–3628. <https://doi.org/10.4161/cc.9.17.12907>
- Slidrecht T, Zhang C, Shokat KM, Kops GJPL (2010) Chemical Genetic Inhibition of Mps1 in Stable Human Cell Lines Reveals Novel Aspects of Mps1 Function in Mitosis. *PLoS One* 5:. <https://doi.org/10.1371/journal.pone.0010251>
- Storchová Z, Becker JS, Talarek N, et al (2011) Bub1, Sgo1, and Mps1 mediate a distinct pathway for chromosome biorientation in budding yeast. *Molecular Biology of the Cell* 22:1473–1485. <https://doi.org/10.1091/mbc.e10-08-0673>
- Stucke VM, Baumann C, Nigg EA (2004) Kinetochores localization and microtubule interaction of the human spindle checkpoint kinase Mps1. *Chromosoma* 113:1–15. <https://doi.org/10.1007/s00412-004-0288-2>
- Stucke VM, Silljé HHW, Arnaud L, Nigg EA (2002) Human Mps1 kinase is required for the spindle assembly checkpoint but not for centrosome duplication. *EMBO J* 21:1723–1732. <https://doi.org/10.1093/emboj/21.7.1723>
- Tanaka TU, Rachidi N, Janke C, et al (2002) Evidence that the Ipl1-Sli15 (Aurora Kinase-INCENP) Complex Promotes Chromosome Bi-orientation by Altering Kinetochores-Spindle Pole Connections. *Cell* 108:317–329. [https://doi.org/10.1016/S0092-8674\(02\)00633-5](https://doi.org/10.1016/S0092-8674(02)00633-5)
- Tien JF, Umbreit NT, Gestaut DR, et al (2010) Cooperation of the Dam1 and Ndc80 kinetochores complexes enhances microtubule coupling and is regulated by aurora B. *The Journal of Cell Biology* 189:713–723. <https://doi.org/10.1083/jcb.200910142>
- Tooley JG, Miller SA, Stukenberg PT (2011) The Ndc80 complex uses a tripartite attachment point to couple microtubule depolymerization to chromosome movement. *Mol Biol Cell* 22:1217–1226. <https://doi.org/10.1091/mbc.E10-07-0626>
- Tsukahara T, Tanno Y, Watanabe Y (2010) Phosphorylation of the CPC by Cdk1 promotes chromosome bi-orientation. *Nature* 467:719–723. <https://doi.org/10.1038/nature09390>
- Vanoosthuyse V, Hardwick KG (2009) A novel protein phosphatase 1-dependent spindle checkpoint silencing mechanism. *Curr Biol* 19:1176–1181. <https://doi.org/10.1016/j.cub.2009.05.060>
- Vasileva V, Gierlinski M, Yue Z, et al (2017) Molecular mechanisms facilitating the initial kinetochores encounter with spindle microtubules. *J Cell Biol* 216:1609–1622. <https://doi.org/10.1083/jcb.201608122>

- 1 Verzijlbergen KF, Nerusheva OO, Kelly D, et al (2014) Shugoshin biases chromosomes for
2 biorientation through condensin recruitment to the pericentromere. *Elife* 3:e01374
- 3 Vigneron S, Prieto S, Bernis C, et al (2004) Kinetochore Localization of Spindle Checkpoint
4 Proteins: Who Controls Whom? *Mol. Biol. Cell* 15:13
- 5
6 Wang X, Yu H, Xu L, et al (2014) Dynamic Autophosphorylation of Mps1 Kinase Is Required
7 for Faithful Mitotic Progression. *PLOS ONE* 9:e104723.
8 <https://doi.org/10.1371/journal.pone.0104723>
- 9
10 Weiss E, Winey M (1996) The *Saccharomyces cerevisiae* spindle pole body duplication gene
11 MPS1 is part of a mitotic checkpoint. *The Journal of cell biology* 132:111–123
- 12
13 Welburn JPI, Grishchuk EL, Backer CB, et al (2009) The human kinetochore Ska1 complex
14 facilitates microtubule depolymerization-coupled motility. *Dev Cell* 16:374–385.
15 <https://doi.org/10.1016/j.devcel.2009.01.011>
- 16
17 Westermann S, Wang H-W, Avila-Sakar A, et al (2006) The Dam1 kinetochore ring complex
18 moves processively on depolymerizing microtubule ends. *Nature* 440:565–569.
19 <https://doi.org/10.1038/nature04409>
- 20
21 Winey M, Goetsch L, Baum P, Byers B (1991) MPS1 and MPS2: novel yeast genes defining
22 distinct steps of spindle pole body duplication. *The Journal of Cell Biology* 114:745–
23 754. <https://doi.org/10.1083/jcb.114.4.745>
- 24
25 Wu X, Tatchell K (2001) Mutations in Yeast Protein Phosphatase Type 1 that Affect
26 Targeting Subunit Binding [†]. *Biochemistry* 40:7410–7420.
27 <https://doi.org/10.1021/bi002796k>
- 28
29 Yamagishi Y, Honda T, Tanno Y, Watanabe Y (2010) Two histone marks establish the inner
30 centromere and chromosome bi-orientation. *Science* 330:239–243.
31 <https://doi.org/10.1126/science.1194498>
- 32
33 Yamagishi Y, Yang C-H, Tanno Y, Watanabe Y (2012) MPS1/Mph1 phosphorylates the
34 kinetochore protein KNL1/Spc7 to recruit SAC components. *Nature Cell Biology*
35 14:746–752. <https://doi.org/10.1038/ncb2515>
- 36
37 Yuan I, Leontiou I, Amin P, et al (2017) Generation of a Spindle Checkpoint Arrest from
38 Synthetic Signaling Assemblies. *Current Biology* 27:137–143.
39 <https://doi.org/10.1016/j.cub.2016.11.014>
- 40
41 Zaytsev AV, Mick JE, Maslennikov E, et al (2015) Multisite phosphorylation of the NDC80
42 complex gradually tunes its microtubule-binding affinity. *Mol Biol Cell* 26:1829–1844.
43 <https://doi.org/10.1091/mbc.E14-11-1539>
- 44
45 Zaytsev AV, Sundin LJR, DeLuca KF, et al (2014) Accurate phosphoregulation of
46 kinetochore-microtubule affinity requires unconstrained molecular interactions. *J Cell*
47 *Biol* 206:45–59. <https://doi.org/10.1083/jcb.201312107>
- 48
49 Zhu T, Dou Z, Qin B, et al (2013) Phosphorylation of Microtubule-binding Protein Hec1 by
50 Mitotic Kinase Aurora B Specifies Spindle Checkpoint Kinase Mps1 Signaling at the
51 Kinetochore. *J Biol Chem* 288:36149–36159.
52 <https://doi.org/10.1074/jbc.M113.507970>
- 53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 1

Benzi and Piatti, Fig. 1