Discussion on ‘Palaeoseismic structures in Quaternary sediments, related to an assumed fault zone north of the Permian Peissen-Gnutz salt structure (NW Germany) – Neotectonic activity and earthquakes from the Saalian to the Holocene’ (Grube, 2019)

Pascal Bertran, Kevin Manchuel, Deborah Sicilia

► **To cite this version:**

Pascal Bertran, Kevin Manchuel, Deborah Sicilia. Discussion on ‘Palaeoseismic structures in Quaternary sediments, related to an assumed fault zone north of the Permian Peissen-Gnutz salt structure (NW Germany) – Neotectonic activity and earthquakes from the Saalian to the Holocene’ (Grube, 2019). *Geomorphology*, 2020, 365, pp.106704. 10.1016/j.geomorph.2019.03.010 . hal-03136518

HAL Id: hal-03136518

<https://hal.science/hal-03136518>

Submitted on 9 Feb 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L’archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d’enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Manuscript Number: GEOMOR-8414R1

Title: Discussion on 'Palaeoseismic structures in Quaternary sediments, related to an assumed fault zone north of the Permian Peissen-Gnutz salt structure (NW Germany) - Neotectonic activity and earthquakes from the Saalian to the Holocene' (Grube, 2019)

Article Type: Reply

Keywords: Palaeoseismology; Permafrost; Brittle deformation

Corresponding Author: Dr. Pascal Bertran, PhD

Corresponding Author's Institution: INRAP / PACEA

First Author: Pascal Bertran, PhD

Order of Authors: Pascal Bertran, PhD; Kevin Manchuel, Ph.D.; Deborah Sicilia, Ph.D.

Abstract: We discuss the significance of deformation structures in Quaternary sediments observed by Grube (2019) in the Peissen quarries (NW Germany) in light of the geological context. Evidence for polygonal patterns visible in aerial images in the study area shows that the wedge structures interpreted by Grube (2019) as earthquake-induced sand blows may rather correspond to thermal contraction cracks filled with aeolian sand in a permafrost environment. In the study sites, brittle deformations caused by (i) the rise of a salt diapir, (ii) salt dissolution, (iii) the development of Pleistocene permafrost and (iv) possibly, water circulation under pressure in the Scandinavian ice sheet margin may have coexisted. We support the idea that, while the morphology of deformation generally makes it possible to determine the stress state to which the sediments have been subjected and the quantity of water available in the system at the time of deformation, the nature of the factors causing the stresses remains difficult to identify. In the end, we highlight other useful criteria that should be privileged for palaeoseismic research in such complex geological settings.

Discussion on 'Palaeoseismic structures in Quaternary sediments, related to an assumed fault zone north of the Permian Peissen-Gnutz salt structure (NW Germany) – Neotectonic activity and earthquakes from the Saalian to the Holocene' (Grube, 2019)

Pascal Bertran^{1*}, Kevin Manchuel², Deborah Sicilia²

¹ Inrap / PACEA, bâtiment B2, allée Geoffroy-Saint-Hilaire, 33615 Pessac, France

² EDF, TEGG/Service Géologique Géotechnique, 905 Avenue du Camp de Menthe, 13097 Aix-en-Provence, France

* Corresponding author. Email address: pascal.bertran@inrap.fr

Abstract

We discuss the significance of deformation structures in Quaternary sediments observed by Grube (2019) in the Peissen quarries (NW Germany) in light of the geological context. Evidence for polygonal patterns visible in aerial images in the study area shows that the wedge structures interpreted by Grube (2019) as earthquake-induced sand blows may rather correspond to thermal contraction cracks filled with aeolian sand in a permafrost environment. In the study sites, brittle deformations caused by (i) the rise of a salt diapir, (ii) salt dissolution, (iii) the development of Pleistocene permafrost and (iv) possibly, water circulation under pressure in the Scandinavian ice sheet margin may have coexisted. We support the idea that, while the morphology of deformation generally makes it possible to determine the stress state to which the sediments have been subjected and the quantity of water available in the system at the time of deformation, the nature of the factors causing the stresses remains difficult to identify. In the end, we highlight other useful criteria that should be privileged for palaeoseismic research in such complex geological settings.

Keywords: Palaeoseismology; Permafrost; Brittle deformation

1. Introduction

Grube's (2019) article on brittle and ductile deformations affecting Quaternary sediments in the Peissen region (northwest Germany) provides interesting data on potential traces of palaeoearthquakes and is part of a recent effort to extend seismicity catalogues in Europe beyond the period for which instrumental data and historical texts are available. These deformation

structures complement the many other structures already described in northern Europe, particularly by [Hoffmann and Reicherter \(2012\)](#), [Brandes and Winsemann \(2013\)](#), [van Loon and Pisarska-Jamroz \(2014\)](#) and [van Loon et al. \(2016\)](#), which were also interpreted as earthquake-induced. However, the geographical proximity of all these structures does not constitute proof of a common origin. The main reasons for this relate to the complex local geomorphological context, marked by (1) the development of permafrost during the last glaciation, (2) the proximity of the Scandinavian ice sheet during the last Glacial Maximum, and (3) the presence of a salt diapir in the immediate vicinity of the study sites. These aspects will be developed in detail here. The outcome of this analysis is that, while the morphology of deformation generally makes it possible to determine the stress state to which the sediments have been subjected and the quantity of water available in the system at the time of deformation, the nature of the factors causing the stresses remains difficult to identify. Laboratory experiments and field observations indicate that similar brittle and/or ductile deformations may occur in relation to various geomorphological processes. Therefore, the multiplicity of potential factors that may have been involved in the sector studied by Grube (2019) does not make it possible to determine confidently the origin of the deformation structures.

2. Sand blows and sand wedges in a permafrost context

The wedge-shaped structures filled with sand described by Grube (2019) and interpreted as sand blows due to ground fluidization caused by seismic shaking are very similar to the wedges caused by thermal contraction of the ground in periglacial environments and filled with aeolian sand (sand wedges) or a mixture of sand and ice (composite wedge pseudomorphs) (e.g., [Murton, 2013](#); [Andrieux et al., 2016a, b](#)). These wedges have a V shape, sand veins extending from the base (apophyses), a depth (up to 1.7 m) and an internal organization typical of periglacial structures. Vertical lamination is preserved in one of the wedges illustrated in Fig. 11e of Grube (2019). The lamination, common in thermal contraction wedges, is related to repeated cracking of the ground during winters and the infiltration of aeolian sand into the crack ([Fig. 1](#)). When the filling is composed of sand and ice (composite wedges), melting of ice causes the lamination to vanish and the wedge to deform. Coarser material from the host sediment can fill the depression. [Ghysels and Heyse \(2006\)](#) and [Buylaert et al \(2009\)](#) have described similar examples in Belgium OSL-dated to the Weichselien and Saalien. Creep of fine-grained host material during melting or repeated thaw cycles can also lead to the formation of globular structures comparable to those shown in Grube's (2019) Fig. 11b ([Fig. 1D](#)).

One of the most relevant criteria for discriminating these structures from sand blows is their organization into large polygons, which reach 10 to 30 m in diameter. Unfortunately, Grube (2019)

has not documented this aspect. However, some data indicate that the presence of sand wedges or composite wedge pseudomorphs is highly probable in the study area. These data are (1) polygons visible on satellite photographs accessible in Google Earth; one of the identified sites is located in the immediate vicinity of Pit 3, another of Pit 2 (Fig. 2); (2) the mention by Christensen (1978) of abundant fossil thermal contraction polygons visible in aerial photographs in Denmark and northern Germany; the geo-referencing of Christensen's map shows that the study sites are located in an area where more than 5% of agricultural land is affected by polygons; and (3) the concomitance of continuous permafrost during the last glaciation (Vandenberghe et al., 2014) and coversand deposition (Kasse, 1997; Zeeberg, 1998), which is a pattern highly favourable to the formation of sand wedges over large surfaces. Therefore, the interpretation of these structures as periglacial sand wedges rather than seismogenic sand blows seems most likely in the current state of the analysis and a seismic origin cannot therefore be retained without more supporting evidence.

3. Brittle deformation

Unconsolidated and well-drained sediments under stress deform in a brittle manner. The stress can be caused by wide range of geological processes. Many experiments using analog models under stress have been described in the literature. These models use cohesionless granular materials, most often sand, sometimes interlayered with ductile levels (cohesive wet clays, or purely viscous silicone paste). The results constitute a data set that can be used to understand the brittle deformation of non-lithified Quaternary sediments.

Experiments by Komuro (1987) and Walter and Troll (2001) reproduced the growth of a lenticular body (a putty ball or an inflated chamber) under a granular cover. The increase in volume leads to the formation of a dome and the development of subvertical radial cracks at the top (Fig. 3A,B). As the dome grows, cracks are formed that cut at right angles the radial cracks and are arranged in a more or less concentric pattern. At the top of the dome, the blocks bounded by the cracks collapse, leading to the formation of a polygonal central depression limited by normal faults. Walter and Troll (2001) reproduced the succession of growth phases followed by collapse of the cover. The interaction between the fractures created during swelling and those related to collapse generates blocks of variable size and a high degree of material fragmentation (Fig. 3F).

This type of structure accurately reflects the fracturing of the sediment or peat layer that covers intrusive ice mounds (pingos, frost blisters) or segregated ice mounds (lithalsas) in a periglacial context. Examples have been described in the Canadian Arctic by Mackay (1988, 1998) and Pollard (1991) (Fig. 4A). Davison et al. (2000) and Marco et al. (2002) also observed a system of radial and tangential cracks in the sedimentary cover at the periphery of salt diapirs (Fig. 4B). In the example

detailed by [Marco et al. \(2002\)](#), many structures correspond to clastic dykes and probably result from the filling of cracks opened from the surface, which were formed in connection with the rise of the diapir.

Experiments that reproduce the rise of a block of substratum, and those that simulate the collapse of an unconsolidated cover over a cavity, provide comparable results. In both cases, the faults are concentrated to a limited area at the edge of the raised block or the cavity. [Sanford's \(1959\)](#) experiments consisted in lifting a rigid block limited by vertical edges under a cover of sand or sand and clay. Reverse curved faults propagate towards the surface and the subsided area ([Fig. 3C](#)). As pointed out by [Sanford \(1959\)](#), the reverse faults, which are subvertical at the base and evolve into thrusts towards the surface, are not related to horizontal compression in the model, but to vertical movements at depth. Similar results were also obtained in a series of experiments designed to reproduce the roof collapse over a cavity ([Roche et al., 2001; Walter and Troll, 2001; Geyer et al., 2006; Coumans and Stix, 2016](#)). In these experiments, bell-shaped reverse fractures form above the cavity together with annular extension fractures starting from the surface at the periphery. Progressive roof collapse occurs in connection with the propagation of the bell-shaped fractures up to the surface ([Fig. 3D,E](#)). [Coumans and Stix \(2016\)](#) reproduced the situation in which the thickness of the cover is not homogeneous. In this case, subsidence associated with reverse faulting occurs where the sediment thickness above the cavity is lowest, whereas normal faults develop at the opposite side ([Fig. 3G](#)). The final depression has an asymmetrical shape.

Many examples of curved reverse faults and associated normal faults have been described in natural environments. They comprise the cover of salt diapirs undergoing dissolution ([Simon and Soriano, 1986; Davison et al., 1996](#)) ([Fig. 4B](#)), sediments affected by the collapse of a karstic cavity ([Soriano et al., 2012; Simon et al., 2014; Luzón et al., 2012](#)) ([Fig. 4E](#)), subglacial deposits (eskers) deformed by glacier melt-out ([McDonald and Shilts, 1975](#)), and lahars or fluvio-glacial deposits (jökulhlaup) deformed by melting of ice-blocks ([Branney and Gilbert, 1995; Fay, 2000](#)). [Calmels et al. \(2008\)](#) identified reverse ice-filled faults dipping 50 to 90° in a segregated ice mound (lithalsa) in northern Quebec. According to [Calmels et al. \(2008\)](#), these faults formed during the growth of ice lenses during permafrost build-up. Reverse faults have also been identified in ramparts created by the melting of Pleistocene ice mounds ([Payette and Séguin, 1979; Kasse and Bohncke, 1992; Pissart, 2000; Bertran et al., 2018](#)) and in thermokarst lake deposits ([Murton, 1996; Bertran et al., 2018](#)) ([Fig. 4C](#)).

Other experiments have reproduced deformations created by shortening ([Bonini et al., 2000; Bonini, 2007](#)). In these models, detachment occurs at the substratum - cover interface and reverse faults

with low dip (thrusts) form in the cover to accommodate the shortening (Fig. 3E). Thrusting leads to the formation of an anticline at the top of the ramp and conjugate reverse faults delimiting pop-up structures develop. The frontal bulge can collapse along normal faults that form as shortening and bulging progress.

This type of deformation is typically observed in the frontal bulge of landslides (Coombs and Norris, 1981; McCalpin and Thakkar, 2003) (Fig. 4D) and at the front of emerging deep-seated reverse faults (Philip et al., 1992; McCalpin and Thakkar, 2003).

In summary, the available models show that it is possible to identify the stress state that caused brittle deformation from the fracture geometry. However, the factors underlying stress remain more difficult to determine and observations in natural environments indicate that many geological processes are able to generate similar deformations. Comparable conclusions were also drawn for soft-sediment deformations (e.g., Moretti et al., 2016). Analysis of the fracture pattern can provide insight into the factors that may be involved. However, the usually limited extent (i.e., a few tens of metres) of the outcrops is one of the main limitations for documenting accurately the general fracture pattern. An exception is fracturing due to thermal contraction of the ground in a periglacial context, where the growth of ice or sand wedges generates easily identifiable polygonal patterns.

The site studied by Grube (2019) is located in a complex geological environment, where deformations created by the rise of a salt diapir, salt dissolution, the development of Pleistocene permafrost and possibly, by water circulation under pressure in the Scandinavian ice sheet margin (see Boulton et al., 1993; Murton, 2005; Ravier et al., 2015) have overlapped. Consequently, this site seems unfavourable to the detection of palaeoearthquakes, insofar as the structures observed at the scale of the outcrops do not allow the factors potentially involved to be discriminated against. The association of normal and reverse faults (Fig. 12 of Grube (2019)) may well reflect the deformation associated with the rise and dissolution of the underlying salt diapir, or that created by the growth and melting of ice bodies during the Weichselian, rather than earthquake-induced processes. Diapir uplift caused by loading of the surrounding land by the Weichselian ice sheet likely occurred as demonstrated by Lang et al. (2014).

4. Conclusion and prospects

The arguments proposed by Grube (2019) for a seismic origin of the structures observed in the Peissen quarries are not convincing when considering the context. In such geological settings, the types of criteria that should be favoured in palaeoseismic research are (1) surface lineation identifiable by the relief or an offset in geological structures, which may reflect the emergence of deep-seated faults (although possibly non-seismogenic), and (2) fractured pebbles associated with a

164 fault or in the associated damage zone, which provide evidence for the seismogenic nature of the
165 fault.

166 The first criterion is traditionally used in palaeoseismology (e.g., [Chardon et al., 2005](#); [Camelbeeck et al., 2007](#); [Baize et al., 2019](#)) and serves as a guide for trenching to study the fault's history. This
167 approach ensures that the structures analysed in cross-section are effectively related to a deep-
168 seated fault. The persistence of relief, however, implies that the fault has been active recently.

170 Pebble fracturing along seismogenic faults is attested by some authors, particularly [Kübler et al \(2018\)](#), in settings where the lithostatic stress is null (subsurface). Fragmentation results from the
171 development of stress higher than pebble strength (especially in the case of poorly resistant
172 lithologies such as argillites or sandstones) caused by shearing along the fault. This stress remains
173 significantly lower than that required for quartz grain cataclasis (typically in deep fault gouges, e.g.,
174 [Cashman and Cashman, 2000](#); [Torabi et al., 2007](#); [Mair and Abe, 2008](#); [Kristensen et al., 2013](#)) and
175 requires the pebbles to be in contact with each other (clast-supported material). According to [Radjai et al. \(1998\)](#), the stress transmitted along the load-bearing network is much higher than the average
176 vertical stress, allowing pressures higher than pebble strength to develop during shear. Seismic
177 compression waves and waves released by the bursting of neighbouring pebbles are likely to be
178 involved in fracturing ([Davies et al., 2012](#)), as long as fracturing is not observed along non-
179 seismogenic faults.
180
181

182 Cataclasis could thus provide a reliable indicator that can be used in palaeoseismological analysis,
183 particularly when it affects many pebbles in the damage zone around a fault ([Fig. 5A, B](#)). Other
184 factors that can cause fracturing of subsurface pebbles include gelifraction ([Matsuoka, 2001, 2008](#);
185 [Jia et al., 2017](#)) and mass flow of debris. Gelifraction is caused by crack expansion resulting from ice
186 growth in fissured rocks and segregation ice growth due to water migration in weak and highly
187 porous rocks such as chalk. Gelifraction is a common feature in cold environments and mainly affects
188 limestones, shales and all kinds of fractured rocks ([Fig. 5C, D](#)). It remains ineffective on compact rocks
189 such as most alluvial pebbles, for which the impacts caused by the fluvial transport have eliminated
190 the least resistant and most fissured parts. Fractured pebbles are actually rare (but not totally
191 absent) in Quaternary alluvial deposits and are scattered throughout the deposits. Cataclasis typifies
192 the sedimentary flows involving a large volume of debris, i.e., rock avalanches ([Siebert, 1984](#);
193 [Yarnold, 1993](#); [Bertran, 2003](#)). It develops within the whole flow and does not concentrate along
194 identifiable faults. It gives rise to so-called "jigsaw" structures ([Fig. 5E, F](#)). In smaller flows (debris
195 flows, snow avalanches), flaking of the transported blocks dominates. In most cases,

sedimentological criteria and the non-localized nature of cataclasis make it possible to identify the factor involved.

Fractured pebbles associated with faults and embedded in unconsolidated or weakly cemented fluvial deposits were mentioned by several authors ([Jorda, 1982](#); [Carbon et al., 1993](#); [Baize et al., 2002](#); [Guignard et al., 2005](#)) in southeastern France, the most seismic region of the country during historical times, and provide reliable indices for palaeo-earthquakes. To our knowledge, similar such structures have not been reported in areas further north and should focus research.

Acknowledgements

We acknowledge O. Bellier for showing us sites with fractured pebbles in SE France. The funds have been provided by Electricité de France (EDF) and Institut de Recherches en Archéologie Préventive (Inrap) through the SISMOGEL project. M. Moretti and an anonymous reviewer are also thanked for their constructive remarks on the manuscript.

References

- [Andrieux, E., Bertran, P., Saito, K., 2016a. Spatial analysis of the French Pleistocene permafrost by a GIS database. *Permafrost and Periglacial Processes* 27, 17-30.](#)
- [Andrieux, E., Bertran, P., Antoine, P., Deschodt, L., Lenoble, A., Coutard, S., 2016b. Database of pleistocene periglacial features in France: description of the online version. *Quaternaire* 27, 329-339.](#)
- [Baize, S., Cushing, M., Lemeille, F., Granier, T., Grellet, B., Carbon, D., Combes, P., Hibschi, C., 2002. Inventaire des indices de rupture affectant le Quaternaire, en relation avec les grandes structures connues, en France métropolitaine et dans les régions limitrophes. *Mémoires de la Société Géologique de France, Nouvelle Série*, 175, 141 p.](#)
- [Baize, S., Reicherter, K., Avagyan, A., Belyashov, A., Pestov, E., Vittori, E., Arakelyan, A., Decker, K., 2019. First assessment of recent tectonics and paleoearthquakes along the Irtysh fault \(eastern Kazakhstan\). *Geomorphology* 326, 90-106.](#)
- [Ballard, J.F., Brun, J.P., Van den Driessche, J., Allemand, P., 1987. Propagation des chevauchements au-dessus des zones de décollement : modèles expérimentaux. *Courrier de l'Académie des Sciences de Paris* 305, série II, 1249-1253.](#)

225 Bertran, P., 2003. The rock-avalanche of February 1995 at Claix (French Alps). *Geomorphology* 54,
226 339-346.

227 Bertran, P., Andrieux, E., Bateman, M., Font, M., Manchuel, K., Sicilia, D., 2018. Features caused by
228 ground ice growth and decay in Late Pleistocene fluvial deposits, Paris Basin, France. *Geomorphology*
229 310, 84-101.

230 Bonini, M., 2007. Deformation patterns and structural vergence in brittle-ductile thrust wedges: An
231 additional analogue modelling perspective. *Journal of Structural Geology* 29, 141-158.

232 Bonini, M., Sokoutis, D., Mulutega, G., Katrivano,s E., 2000. Modelling hanging wall accomodation
233 above rigid thrust ramps. *Journal of Structural Geology* 22, 1165-1179.

234 Boulton, G.S., Slot, T., Blessing, K., Glasbergen, P., Leijnse, T., van Gijssel, K., 1993. Deep circulation of
235 groundwater in overpressured subglacial aquifers and its geological consequences. *Quaternary*
236 *Science Reviews* 12, 739-745.

237 Brandes, C., Winsemann, J., 2013. Soft-sediment deformation structures in NW Germany caused by
238 Late Pleistocene seismicity. *International Journal of Earth Sciences* 102, 2255-2274.

239 Branney, M.J., Gilbert, J.S., 1995. Ice-melt collapse pits and associated features in the 1991 lahar
240 deposits of Volcan Hudson, Chile: criteria to distinguish eruption-induced glacier melt. *Bulletin of*
241 *Volcanology* 57, 293-302.

242 Buylaert, J.P., Ghysels, G., Murray, A.S., Thomsen, K.J., Vandenberghe, D., De Corte, F., Heyse, I., Van
243 den Haute, P., 2009. Optical dating of relict sand wedges and composite-wedge pseudomorphs in
244 Flanders, Belgium. *Boreas* 38, 160–175.

245 Calmels, F., Allard, M., Delisle, G., 2008. Development and decay of a lithalsa in Northern Quebec: a
246 geomorphological history. *Geomorphology* 97, 287-299.

247 Camelbeeck, T., Vanneste, K., Alexandre, P., Verbeeck, K., Petermans, T., Rosset, P., Everaerts, M.,
248 Warnant, R., Van Camp, M., 2007. Relevance of active faulting and seismicity studies to assessments
249 of long-term earthquake activity and maximum magnitude in intraplate northwest Europe, between
250 the Lower Rhine Embayment and the North Sea. In: Stein, S., Mazzotti, S. (ed.), *Continental Intraplate*
251 *Earthquakes: Science, Hazard, and Policy Issues*. The Geological Society of America Special Paper 425,
252 pp. 193-223.

253 Carbon, D., Combes, P., Cushing, M., Granier, T., 1993. Enregistrement d'un paléoséisme dans des
254 sédiments du pléistocène supérieur dans la vallée du Rhône : quantification de la
255 déformation. *Géologie Alpine* 69, 33-48.

256 Cashman, S., Cashman, K., 2000. Cataclasis and deformation-band formation in unconsolidated
 257 marine terrace sand, Humboldt County, California. *Geology* 28, 111-114.

258 Chardon, D., Hermitte, D., Nguyen, F., Bellier, O., 2005. First paleoseismological constraints on the
 259 strongest earthquake in France (Provence) in the twentieth century. *Geology* 33, 901- 904.

260 Coombs, D.S., Norris, R.J., 1981. The east Abbotsford, Dunedin, New Zealand, landslide of august 8,
 261 1979, an interim report. *Bulletin de Liaison des Ponts-et-Chaussées*, n° spécial X, 27-34.

262 Coumans, J.P., Stix, J., 2016. Caldera collapse at near-ridge seamounts: an experimental investigation.
 263 *Bulletin of Volcanology* 78, 70, doi 10.1007/s00445-016-1065-9.

264 Christensen, L., 1978. Waterstress conditions in cereals used in recognizing fossil ice-wedge
 265 polygonal patterns in Denmark and northern Germany. *Proceedings of the Third International*
 266 *Conference on Permafrost*, National research Council of Canada, Edmonton, Canada, vol. 1, pp. 254-
 267 261.

268 Davison, I., Bosence, D., Alsop, G.I., Al-Aawah, M.H., 1996. Deformation and sedimentation around
 269 active Miocene salt diapirs on the Tihama Plain, northwest Yemen. In: Alsop, G.I., Blundell, D.J.,
 270 Davison, I. (eds.), *Salt tectonics*. Geological Society, London, Special Publication 100, pp. 23-39.

271 Davison, I., Alsop, G.I., Evans, N.G., Safaricz, M., 2000. Overburden deformation patterns and
 272 mechanisms of salt diapir penetration in the Central Graben, North Sea. *Marine and Petroleum*
 273 *Geology* 17, 601-618.

274 Davies, T.R.H., McSaveney, M.J., Boulton, C.J., 2012. Elastic strain energy release from fragmenting
 275 grains: effects on fault rupture. *Journal of Structural Geology* 38, 265-277.

276 Fay, H., 2000. Formation of kettle holes following a glacial outburst flood (jökulhlaup),
 277 Skeidararsandur, southern Iceland. *IAHS Publ.* 271, 205-210.

278 Geyer, A., Folch, A., Marti, J., 2006. Relationship between caldera collapse and magma chamber
 279 withdrawal: an experimental approach. *Journal of Volcanology and Geothermal Research* 157, 375-
 280 386.

281 Guignard, P., Bellier, O., Chardon, D., 2005. Géométrie et cinématique post-oligocène des failles d'Aix
 282 et de la moyenne Durance (Provence, France). *Comptes-Rendus Géoscience* 337, 375-384.

283 Guysels, G., Heyse, I., 2006. Composite-wedge pseudomorphs in Flanders, Belgium. *Permafrost and*
 284 *Periglacial Processes* 17, 145-161.

285 Hoffmann, G., Reicherter, K., 2012. Soft-sediment deformation of Late Pleistocene sediments along
 286 the southwestern coast of the Baltic Sea (NE Germany). *International Journal of Earth Sciences* 101,
 287 351-363.

288 Jia, H., Leith, K., Krautblatter, M., 2017. Path-dependent frost-wedging experiments in fractured, low-
 289 permeability granite. *Permafrost and Periglacial Processes* 28, 698-709.

290 Jorda, M., 1992. La tectonique plio-quaternaire des Préalpes de Digne et ses prolongements récents.
 291 L'enseignement des hautes nappes alluviales. In: *Le Villafranchien méditerranéen*, Université des
 292 Sciences et Techniques de Lille, 1982, 425-440.

293 Kasse, C., 1997. Cold-climate Aeolian Sand-sheet formation in North-Western Europe (c. 14-12.4 ka);
 294 a response to permafrost degradation and increased Aridity. *Permafrost and Periglacial Processes* 8,
 295 295-311.

296 Kasse, K., Bohncke, S., 1992. Weichselian Upper Pleniglacial aeolian and ice-cored morphology in the
 297 southern Netherlands (Noort-Brabant, Groote Peel). *Permafrost and Periglacial Processes* 3, 327–
 298 342.

299 Komuro, H., 1987. Experiments on cauldron formation: a polygonal cauldron and ring fractures.
 300 *Journal of Volcanology and Geothermal Research* 31, 139-149.

301 Kübler, S., Friedrich, A.M., Gold, R.D., Strecker, M.R., 2018. Historical coseismic surface deformation
 302 of fluvial gravel deposits, Schafberg fault, Lower Rhine Graben, Germany. *International Journal of*
 303 *Earth Science* 107, 571-585.

304 Kristensen, M.B., Child, C., Olesen, N.Ø., Korstgård, J.A., 2013. The microstructure and internal
 305 architecture of shear bands in sand-clay sequences. *Journal of Structural Geology* 46, 129-141.

306 Lang, J., Hampel, A., Brandes, C., Winsemann, J., 2014. Response of salt structures to ice loading:
 307 implications for ice-marginal and subglacial processes. *Quaternary Science Reviews* 101, 217-233.

308 Luzón, A., Rodríguez-López, J.P., Pérez, A., Soriano, M.A., Gil, H., Pocovi, A., 2012. Karst subsidence as
 309 a control on the accumulation and preservation of aeolian deposits: A Pleistocene example from a
 310 proglacial outwash setting, Ebro Basin, Spain. *Sedimentology* 59, 2199-2225.

311 Mackay, J.R., 1988. Pingo collapse and paleoclimatic reconstruction. *Canadian Journal of Earth*
 312 *Sciences* 25, 495-511.

313 Mackay, J.R., 1998. Pingo growth and collapse, Tuktoyaktuk Peninsula area, western Arctic coast,
 314 Canada: A long-term field study. *Géographie Physique et Quaternaire* 52, 271-323.

315 Mair, K., Abe, S., 2008. 3D numerical simulations of fault gouge evolution during shear: Grain size
316 reduction and strain localization. *Earth and Planetary Science Letters* 274, 72-81.

317 Marco, S., Weinberger, R., Agnon, A., 2002. Radial clastic dykes formed by a salt diapir in the Dead
318 Sea Rift, Israel. *Terra Nova* 14, 288-294.

319 Matsuoka, N., 2001. Microgelivation versus macrogelivation: towards bridging the gap between
320 laboratory and field frost weathering. *Permafrost and Periglacial Processes* 12, 299-313.

321 Matsuoka, N., 2008. Frost weathering and rockwall erosion in the southeastern Swiss Alps: Long-term
322 (1994–2006) observations. *Geomorphology* 99, 353-368.

323 McCalpin, J.P., Thakkar, M.G., 2003. 2001 Bhuj-Kachchh earthquake: surface faulting and its relation
324 with neotectonics and regional structures, Gujarat, Western India. *Annals of Geophysics* 46, 937-956.

325 McDonald, B.C., Shilts, W.W., 1975. Interpretation of faults in glaciofluvial sediments. In: Jopling,
326 A.V., McDonald, B.C. (eds.), *Glaciofluvial and glaciolacustrine sedimentation*. Society of Economic
327 Paleontologists and Mineralogists, Tulsa, USA, Special Publication 23, pp. 123-131.

328 Moretti, M., Alfaro, P., Owen, G., 2016. The environmental significance of soft-sediment deformation
329 structures: key signatures for sedimentary and tectonic processes. *Sedimentary Geology* 344, 1-4.

330 Murton, J.B., 1996. Thermokarst-lake-basin sediments, Tuktoyaktuk Coastlands, western arctic
331 Canada. *Sedimentology* 43, 737-760.

332 Murton, J.B., 2005. Ground-ice stratigraphy and formation at North Head, Tuktoyaktuk Coastlands,
333 Western Arctic Canada: a product of glacier–permafrost interactions. *Permafrost and Periglacial*
334 *Processes* 16, 31-50.

335 Murton, J.B., 2013. Ice wedges and ice wedge casts. In: Elias, S.A., Mock, C.J. (Eds.), *Encyclopedia of*
336 *Quaternary Science*. Elsevier, Amsterdam, pp. 436-451.

337 Payette, S., Séguin, M.K., 1979. Les buttes minérales cryogènes dans les basses terres de la Rivière-
338 aux-Feuilles, Nouveau Québec. *Géographie Physique et Quaternaire* XXIII, 339-358.

339 Philip, H., Rogozhin, E., Cisternas, A., Bousquet, J.C., Borisov, B., Karakhanian, A., 1992. The Armenian
340 earthquake of 1988 December 7: faulting and folding, neotectonics and palaeoseismicity.
341 *Geophysical Journal International* 110, 141-158.

342 Pissart, A., 2000. Remnants of lithalsas of the Hautes Fagnes, Belgium: a summary of present-day
343 knowledge. *Permafrost and Periglacial Processes* 11, 327–355.

344 Pollard, W.H., 1991. Seasonal frost-mounds. *The Canadian Geographer* 35, 2014-2018.

345 Radjai, F., Wolf, D.E., Jean, M., Moreau, J.J., 1998. Bimodal Character of Stress Transmission in
346 Granular Packings. *Physical Review Letters* 80, 61-64.

347 Ravier, E., Buoncristiani, J.F., Menzies, J., Guiraud, M., Portier, E., 2015. Clastic injection dynamics
348 during ice front oscillations: a case example from Sólheimajökull (Iceland). *Sedimentary Geology* 323,
349 92–109.

350 Roche, O., van Wyk de Vries, B., Druitt, T.H., 2001. Sub-surface structures and collapse mechanisms
351 of summit pit. *Journal of Volcanology and Geothermal Research* 105, 1-18.

352 Sanford, A.R., 1959. Analytical and experimental study of simple geologic structures. *Bulletin of the*
353 *Geological Society of America* 70, 19-52.

354 Siebert, L., 1984. Large volcanic debris avalanches: characteristics of source areas, deposits, and
355 associated eruptions. *Journal of Volcanology and Geothermal Research* 22, 163–197.

356 Simón, J.L., Soriano, A., 1986. Diapiric deformations in the Quaternary deposits of the central Ebro
357 Basin, Spain. *Geological Magazine* 123, 45-57.

358 Simón, J.L., Soriano, M.A., Pérez, A., Luzón, A., Pocovi, A., Gil H., 2014. Interacting tectonic faulting,
359 karst subsidence, diapirism and continental sedimentation in Pleistocene deposits of the central Ebro
360 Basin (Spain). *Geological Magazine* 151, 1115-1134.

361 Soriano, M.A., Luzón, A., Yuste, A., Pocovi, A., Pérez, A., Simón, J.L., Gil H., 2012. Quaternary alluvial
362 sinkholes: record of environmental conditions of karst development, examples from the Ebro Basin,
363 Spain. *Journal of Cave and Karst Studies* 74, 173-185.

364 Torabi, A., Braathen, A., Cuisiat, F., Fossen, H., 2007. Shear zones in porous sand: insights from ring-
365 shear experiments and naturally deformed sandstones. *Tectonophysics* 437, 37-50.

366 Vandenberghe, J., French, H., Gorbunov, A., Marchenko, S., Velichko, A.A., Jin, H., Cui, Z., Zhang, T.,
367 Wan, X., 2014. The Last permafrost Maximum (LPM) map of the Northern Hemisphere: permafrost
368 extent and mean annual air temperatures, 25-17 ka. *Boreas* 43, 652-666.

369 Van Loon, A.J., Pisarska-Jamrozy, M., 2014. Sedimentological evidence of Pleistocene earthquakes in
370 NW Poland induced by glacio-isostatic rebound. *Sedimentary Geology* 300, 1-10.

371 Van Loon, A.J., Pisarska-Jamrozy, M., Nartiss, M., Krievans, M., Soms, J., 2016. Seismites resulting
372 from high-frequency, high-magnitude earthquakes in Latvia caused by Late Glacial glacio-isostatic
373 uplift. *Journal of Palaeogeography* 5, 360-380.

Walter, T.R., Troll, V.R., 2001. Formation of caldera periphery faults: an experimental study. *Bulletin of Volcanology* 63, 191-203.

Yarnold, J.C., 1993. Rock-avalanche characteristics in dry climates and the effect of flow into lakes: insights from mid-Tertiary sedimentary breccias near Artillery Peak, Arizona. *Geological Society of America Bulletin* 105, 345– 360.

Zeeberg, J., 1998. The European sand belt in Eastern Europe and comparison of Late Glacial dune orientation with GCM simulation results. *Boreas* 27, 127–139.

Figure captions

Figure 1. (A) Pleistocene sand wedge, Salaunes (SW France), (B) Close up view of the vertical lamination in a sand wedge from La Louverie (Loire valley, France), (C) Sand wedge with apophyses, Saint-Amand-les-Eaux (N France), (D) Globular structure (DSW) caused by the deformation of a sand wedge, Saint-André-de-Cubzac (SW France); the sand wedge (SW) is visible on the bottom of the trench below the dotted line.

Figure 2. (A) Ground thermal contraction polygons, Peissen (X = 9.5878°E, Y = 54.0414°N); (B) Polygons, Beldorf (X = 9.3540°E, Y = 54.1213°N) (Google Earth, photos 2009).

Figure 3. Brittle deformation of sandy soil models. (A) Radial fractures and central polygonal depression formed by the rise of a ball under a granular cover, redrawn from Komuro (1987); (B) Fracturing of a sand cover following chamber inflation, from Walter and Troll (2001); (C) Formation of curved reverse faults in sand due to the lifting of a substratum block, from Sanford (1959); (D) Bell-shaped reverse faults above a cavity, from Geyer et al. (2006); (E) Annular fracture above a cavity, from Walter and Troll (2001) (the chamber is indicated by the grey dashed line); (F) Deformation after swelling followed by emptying of a chamber, from Walter and Troll (2001); (G) Asymmetric collapse above a cavity in the presence of a relief, from Coumans and Stix (2016); (H) Deformation of a cover as a result of horizontal shortening, from Ballard et al. (1987).

Figure 4. (A) Radial cracks on a pingo (ice-cored mound), Tuktoyaktuk, Canada (Google Earth); (B) Radial and tangential faults around a salt diapir, redrawn from Davison et al. (2000); (C) Reverse faults in Pleistocene thermokarst lake deposits, Gourgauçon (Paris Basin, France); (D) Thrust planes at the toe of a landslide, Les Leches (SW France); (E) Bell-shaped reverse faults above a karstic cavity, Mérignac (SW France).

404 Figure 5. (A, B) Fractured pebbles in the Lower Pleistocene Valensole II Formation, near Sisteron (SE
405 France); the finer-grained material to the left of photo B is a fault gouge; the largest pebble is 10 cm
406 long; (C) GelifRACTED pebble in an active layer above permafrost, Tuktoyaktuk (Canada); (D)
407 GelifRACTED sandy limestones, French Pyrenees (France); the largest pebbles are 20 cm in diameter;
408 (E) Jigsaw structure in the Mont Granier rock avalanche (French Alps); knife for scale; (F) Jigsaw
409 structure in a volcanic rock avalanche, Guadeloupe Island (French West Indies); the section is 1.5 m
410 high.

Figure (Color)
[Click here to download high resolution image](#)

Figure (Color)
[Click here to download high resolution image](#)

Figure (Color)
[Click here to download high resolution image](#)

Figure (Color)
[Click here to download high resolution image](#)

Figure (Color)
[Click here to download high resolution image](#)