Non-invasive estimation of myocardial work in aortic stenosis from modelling approach

K. R. Owashi, A. Hubert, Elena Galli, Erwan Donal, A. Hernandez, V Le Rolle

► To cite this version:

K. R. Owashi, A. Hubert, Elena Galli, Erwan Donal, A. Hernandez, et al.. Non-invasive estimation of myocardial work in aortic stenosis from modelling approach. *European Heart Journal*, 2020, 41 (Supplement_2), pp.51-51. 10.1093/ehjci/ehaa946.0051 . hal-03155845

HAL Id: hal-03155845

<https://hal.science/hal-03155845>

Submitted on 11 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Non-invasive estimation of Myocardial Work in Aortic Stenosis from Modelling Approach

Kimi P. Owashi¹, Arnaud Hubert¹, Elena Galli¹, Erwan Donal¹, Alfredo I. Hernandez¹ and Virginie Le Rolle¹

¹ Univ Rennes, Inserm, LTSI - UMR 1099, F-35000 Rennes, France

Background: The noninvasive assessment of myocardial work by pressure-strain loops (PSL) analysis is a recently introduced tool to estimate myocardial performance. PSL analysis cannot be applied to patients having an obstacle to LV ejection, as in the case of aortic stenosis (AS), because of the difficulty to estimate left ventricular (LV) pressure. The objective of this work is to propose a non-invasive model-based estimation of the left ventricular pressure curve in AS patients in order to evaluate myocardial work indices.

Methods: Twelve patients with moderate-to-severe AS underwent cardiac catheterization to acquire the LV pressure (P_{lv}^{exp}). All patients underwent a standard trans-thoracic echocardiography to extract regional myocardial strain curves and estimate the aortic valve area (AVA). Systolic and diastolic arterial pressures were also measured. The proposed cardiovascular system (CVS) model is composed of four main coupled sub-models simulating: *i*) cardiac electrical activities, *ii*) mechanical cardiac cavity, *iii*) the systemic and pulmonary circulation, and *iv*) cardiac valves (Figure 1). A 2-step parameter identification strategy, based evolutionary algorithms, was implemented to learn LV parameters from P_{lv}^{exp} and to estimate patient-specific model-based LV pressure curves (P_{lv}^{model}) starting from parameters obtained from non-invasive hemodynamics. Global myocardial constructive work (GCW) and myocardial wasted work (GWW) were calculated using P_{lv}^{exp} and P_{lv}^{model} .

Figure 1: Clinical Dataset (left). Parameter identification strategy (middle). CVS model (right).

Results: A close match was observed between P_{lv}^{exp} and P_{lv}^{model} , with a mean total relative error of 12.27% (range 5.9% to 17.40%). When considering GCW and GWW, global correlation was equal to 0.92 ($p < 0.0001$). In BA analysis, mean bias was -2.9 mmHg.%, which corresponds to relative bias equal to 0.11% with respect to mean value of work indices (Figure 2).

Figure 2: Work indices comparison for the 12 patients. Scatter and BA plots for GWW (blue) and GCW (red).

Conclusions: A model-based approach can be used to estimate LV pressure and myocardial work indices in patients with AS, and be provide a promising tool for the assessment of myocardial performance in patients with AS.