Earth and Space Science

COMMENT

10.1029/2020EA001298

This article is a comment on Le Mouél et al. (2020), <https://doi.org/10.1029/2019EA000671>.

Key Points:

- No compelling results showing a significant covariability between sunspot amounts and temperature
- The radiative forcing associated with slow solar components (60 years, trend) is very weak over the past 150 years
- The global temperature and slow solar sunspots number components have opposite trend in the last 30 years

Correspondence to:

Y. Cuypers,
Yannis.Cuypers@ocean-ipsl.upmc.fr

Citation:

Cuypers, Y., Codron, F., & Crepon, M. (2021). Comment on “Characteristic time scales of decadal to centennial changes in global surface temperatures over the past 150 years” by J. L. Le Mouél, F. Lopes, and V. Courtillot. *Earth and Space Science*, 8, e2020EA001298. <https://doi.org/10.1029/2020EA001298>

Received 17 JUN 2020

Accepted 19 FEB 2021

© 2021. The Authors. Earth and Space Science published by Wiley Periodicals LLC on behalf of American Geophysical Union.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs License](#), which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Comment on “Characteristic Time Scales of Decadal to Centennial Changes in Global Surface Temperatures Over the Past 150 Years” by J. L. Le Mouél, F. Lopes, and V. Courtillot

Y. Cuypers¹ , F. Codron¹ , and M. Crepon¹

¹Laboratoire d’Océanographie et du Climat: Expérimentations et approches numériques, Unité Mixte de Recherche 7159 CNRS/IRD/Sorbonne Université/MNH, Institut Pierre Simon Laplace, Paris, France

Abstract Based on Singular Spectral Analysis (SSA) analysis of global earth surface temperature and solar activity (sunspots), Le Mouél et al. (2020, <https://doi.org/10.1029/2019EA000671>) suggest that the variability in Earth surface temperature observed since 1850 is natural and controlled by the Sun. We cannot agree with their conclusions for several reasons: the lack of compelling results from the Fourier spectra and SSA estimates which are provided without confidence intervals, the small radiative forcing associated with the sunspot variability, and finally the simple evidence that the slowly varying components of the temperature and sunspots time show opposite trends in the last 30 years.

First of all, the standard Fourier spectra of International Sunspots Numbers (ISSN; WDC-SILSO, Royal Observatory of Belgium, Brussels; SILSO World Data Center, 2021), ISSN and HadCRUT4 temperature (simply temperature hereafter) time series (Morice et al., 2012) provided in Figures 1 and 11 by Le Mouél et al. (2020) look very different and do not seem at first sight to share any common significant peak. Moreover, the authors did not provide any confidence interval to allow meaningful conclusions. For this reason, we have recomputed the frequency spectra of temperature and ISSN sunspots to compare their relative amplitudes at different periods and provide a confidence interval (Figures 1a and 1b). We used the Thomson multitapered method (Ghil et al., 2002) with a time–bandwidth product of 3, a method which is well adapted to a spectrum which exhibits both continuous and singular components. Each spectrum was normalized by its signal variance to ease the comparison. The only significant peaks that exceed the 95% confidence interval level are the well-known 11-year peak and its harmonic at 5.5 years in the ISSN time series; but there is no corresponding significant peak at the 11- and 5.5-year periods or any other one in the temperature time series. The 60-year period shows unsurprisingly more energy in the temperature signal (the temperature, as many geophysical time series, is a “red signal” with more energy at low frequencies). Still, we could not isolate a clear distinctive peak at this period on the temperature spectrum; nor on the ISSN one. This may be a limit of the spectral analysis, as the length of the time series is too short to single out a component at a period so close to its total length, but so far the frequency analysis did not suggest any significant common periods between the two time series.

Can we learn more from the Singular Spectral Analysis (SSA) performed by Le Mouél et al. (2020)? Unfortunately, the SSA eigenvalues of the temperature and ISSN time series computed by Le Mouél et al. (2020) suggest the same conclusion (Figures 1c and 1d): the dominant frequencies are completely different between the two time series; the temperature SSA is dominated by a 60-year eigenvalue which is simply absent in the ISSN SSA. The ISSN SSA spectrum is dominated by the 11-year component which emerges only at rank 15 in the temperature SSA spectrum (such a high rank is usually associated with noise). In addition, we do not have any indication whether these weak 22-, 11- and 5.5-year components are even significant in the temperature SSA (by construction, the SSA method will produce peaks at single frequencies even in a purely random noise time series). Although the SSA significance is not straightforward to assess, it could be have been tested using a Monte Carlo SSA test (Ghil et al., 2002). Since this crucial information is lacking, we can attempt to select the significant components by locating the slope break in the temperature SSA (Ghil et al., 2002): the idea being that there is a sharp break between the highest (significant) eigenvalues and the

Figure 1. (a) Power spectrum of International Sunspots Number (ISSN, WDC-SILSO, Royal Observatory of Belgium, Brussels) using the multitaper method with a time–bandwidth product of 3; (b) same for the HadCRUT4 temperature time series. The red bars in each case represent the 95% confidence. Vertical lines indicate the main periods (60, 22, 11, and 5.5 years) studied in Le Mouél et al. (2020). (c) ISSN and (d) temperature SSA components from Le Mouél et al. (2020) (Figures 3 and 13) reproduced here for the sake of convenience. SSA, Singular Spectral Analysis.

lowest insignificant ones. This suggests that all components shorter than 60-year period are not significant for the temperature, a conclusion that is consistent with the frequency spectral analysis.

So, if these short period components associated with the solar activity are likely not significant in the temperature series, is there a more robust signal for the longer periods? The main problem here is that the 60-year period that dominates the temperature SSA is absent in the ISSN SSA, as acknowledged in the Le Mouél et al. (2020) paper. Le Mouél et al. (2020) managed to extract a 60-year component in the ISSN sunspot series by processing the SSA, as described in the beginning of Section 4 (“The 60 year component does not show as such explicitly in the eigenvalue spectrum of ISSN... It is actually extracted from a re-analysis of the first component, i.e. the trend, again by changing the L value – L is the length of the time series – to one that is better adapted to long periods but not to shorter ones”). This methodology seems extremely obscure: the adapted L value is not given, making the result impossible to interpret or reproduce, and once again a confidence estimate is missing. In addition, the fact that the solar and temperature 60-year components are in phase before 1950, but out of phase in the 2000s (as noticed by Le Mouél et al. [2020]), makes it unlikely that one could cause the other. Finally, and not surprisingly since we are dealing with a “further singular spectrum analysis of the trend,” the extracted 60-year component in the ISSN is extremely small (0.3 solar sunspot peak to peak) which literally corresponds to the thickness of the line if compared to the original time series (Figure 11a). It is interesting in this respect to compare Figures 16 and 14 upper panels showing the reconstructed time series at 11- and 60-year periods for ISSN and temperature. Le Mouél et al. (2020) suggest that sunspot oscillations of amplitude of ± 80 sunspots at the 11-year period induce temperature fluctuations of $\pm 0.02^\circ\text{C}$, while much smaller oscillations of amplitude of ± 0.15 sunspot at 60 years would induce much larger temperature fluctuations of $\pm 0.15^\circ\text{C}$. This would represent a huge gain increase of the

global temperature response of a factor of $\sim 5,000$ for a solar forcing at a 60-year periods compared to the forcing at 11-year period, which seems physically unrealistic. This discrepancy was not discussed by Le Mouél et al. (2020).

Let us now consider the second point. The energy fluxes associated with the solar periodicities are unable to drive Earth's surface temperature variations due to their low amplitude. The strongest sunspot variability (0–250 sunspots) occurs for a 11-year period and corresponds to a variation in the solar radiation of the order of 1 W m^{-2} (Lean, 2018) which implies a variability of the solar flux absorbed by the planet of 0.2 W m^{-2} after taking the sphericity of the planet and its albedo into account. This is small compared with the anomalous energy fluxes due to anthropogenic greenhouse gases increase, which have been of the order of 2 W m^{-2} since the preindustrial era (IPCC, 2014). This discrepancy only worsens when considering longer-term changes: the ISSN 60-year component has a very small peak to peak amplitude of at most 0.3 sunspots (Figure 15); a one sunspot difference corresponds to $0.2/250 = 0.0008 \text{ W m}^{-2}$, therefore the amplitude of this component is at most $\sim 0.3 \times 0.0008 = 0.00024 \text{ W m}^{-2}$, that is about 4 orders of magnitude lower than the anthropogenic greenhouse effect. The sunspot trend is a bit larger: it reaches about a 16 sunspots difference over the 1850–2017 period, which corresponds to a small radiative addition of 0.013 W m^{-2} , but this is still 2 orders of magnitude lower than the anthropogenic greenhouse gas effect over the same period. Moreover, the long-term trend of the ISSN sunspots was computed over 1700–2017 while the temperature trend over 1850–2017 (see Figure 20 caption) making any conclusions incorrect.

No physical mechanism is provided by Le Mouél et al. (2020) to explain how radiative forcing of such vastly different amplitudes whether comparing long-term solar to greenhouse effect, or to the 11-year solar cycle—could have comparable impacts on the global temperature. The Occam razor's principle—invoked by Le Mouél et al. (2020) would naturally lead to think that the radiative forcing from anthropogenic greenhouse gas increase, being orders of magnitude larger than the one from solar variability, would dominate the long-term temperature trends. Indeed, general climate models that include detailed radiative transfer calculations, and are forced by both observed changes in greenhouse gas concentrations and in solar forcing, show a very good agreement with observed temperature changes. As shown in the recent study by Hausfather et al. (2020), since the earliest ones developed around the 1970s, climate models have predicted the recent temperature trends with a correct amplitude and consistently attributed it predominantly to greenhouse gases.

The main point of Le Mouél et al. (2020) was that the solar activity and global temperature time series shared low-frequency components, and therefore—in a big leap of faith—the former should cause the latter: to quote them “When components are shared by two different time series, it is reasonable to assume that there is a common source to these variations, without having to know the full theory of that source” and in the conclusions “It could therefore be argued, as an application of Occam's razor, that most of the variability of surface temperatures could be natural and primarily controlled by the Sun.” Obviously, correlation does not imply causality (or even common cause), and Le Mouél et al. (2020) provide no possible mechanism. But more than that, we showed the followings:

- From a statistical viewpoint, no common significant frequency peak can be found in the temperature and sunspot time series, whether analyzed with the SSA technique or with a standard spectral transform.
- From a quantitative, physical viewpoint, the sunspot fluctuations over longer periods (60 years up to long-term trend) are small and lead to radiative forcing 2–4 orders of magnitude smaller than the anthropogenic greenhouse effect.

As a concluding sanity check, we go back and compare in Figure 2 the raw time series of global-mean temperature and ISSN sunspot numbers. A quick glance immediately reveals that the two raw series behave completely differently: the temperature is slowly increasing (faster near the end of the record), while the solar activity is dominated by the 11-year cycle. An additional 11-year smoothing is then applied to the sunspot series to isolate its slowly varying component. It is clear that while the filtered sunspot series tracks the temperature for ~ 70 years between 1900 and 1970, a period when both are increasing, this is not true before 1900 (decreasing sunspots) and especially since 1970 when both are moving steadily in opposite directions. This opposite trend was noted by Lockwood and Fröhlich (2007) and has further increased in the last two decades. From this figure, the fit in the first half of the twentieth century clearly looks like a coincidence,

Figure 2. ISSN (International Solar Sunspots) time series (black dashed line) with 11-year sliding average (blue line); HadCRUT4 observed temperature time series (orange).

that does not even solve the energy flux amplitude discrepancy, or the lack of 11-year signal in the temperature time series.

Data Availability Statement

The ISSN time series can be downloaded from <http://www.sidc.be/silso/datafiles>. The HadCRUT4 temperature time series can be downloaded from <https://crudata.uea.ac.uk/cru/data/temperature/>.

References

- Ghil, M., Allen, M. R., Dettinger, M. D., Ide, K., Kondrashov, D., Mann, M. E., et al. (2002). Advanced spectral methods for climatic time series. *Reviews of Geophysics*, 40(1), 1003. <https://doi.org/10.1029/2000RG000092>
- Hausfather, Z., Drake, H. F., Abbott, T., & Schmidt, G. A. (2020). Evaluating the performance of past climate model projections. *Geophysical Research Letters*, 47, e2019GL085378. <https://doi.org/10.1029/2019GL085378>
- IPCC. (2014). Climate change 2014: Synthesis report. In Core Writing Team, R. K. Pachauri, & L. A. Meyer (Eds.), *Contribution of Working Groups I, II and III to the fifth assessment report of the Intergovernmental Panel on Climate Change* (p. 151). Geneva, Switzerland: IPCC.
- Lean, J. L. (2018). Estimating solar irradiance since 850. *Earth and Space Science*, 5, 133–149. <https://doi.org/10.1002/2017EA000357>
- Le Mouél, J. L., Lopes, F., & Courtillot, V. (2020). Characteristic time scales of decadal to centennial changes in global surface temperatures over the past 150 years. *Earth and Space Science*, 7(4), e2019EA000671. <https://doi.org/10.1029/2019EA000671>
- Lockwood, M., & Fröhlich, C. (2007). Recent oppositely directed trends in solar climate forcings and the global mean surface air temperature. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 463(2086), 2447–2460.
- Morice, C. P., Kennedy, J. J., Rayner, N. A., & Jones, P. D. (2012). Quantifying uncertainties in global and regional temperature change using an ensemble of observational estimates: The HadCRUT4 dataset. *Journal of Geophysical Research*, 117, D08101. <https://doi.org/10.1029/2011JD017187>
- SILSO World Data Center. (2021). Sunspot number and long-term solar observations. On-line Sunspot Number catalogue, 1700–2018. Brussel, Belgium: Royal Observatory of Belgium. <https://www.sidc.be/SILSO/>