Journal Pre-proof

Sustained productivity and respiration of degrading kelp detritus in the shallow benthos: detached or broken, but not dead.

Nadia Frontier, Florian de Bettignies, Andy Foggo, Dominique Davoult

PII: S0141-1136(21)00026-X

DOI: <https://doi.org/10.1016/j.marenvres.2021.105277>

Reference: MERE 105277

To appear in: *Marine Environmental Research*

Received Date: 18 October 2020

Revised Date: 28 January 2021

Accepted Date: 4 February 2021

Please cite this article as: Frontier, N., de Bettignies, F., Foggo, A., Davoult, D., Sustained productivity and respiration of degrading kelp detritus in the shallow benthos: detached or broken, but not dead., *Marine Environmental Research*, <https://doi.org/10.1016/j.marenvres.2021.105277>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier Ltd.

CRedit AUTHOR STATEMENT

Nadia Frontier: Conceptualisation, Methodology, Data curation, Formal analysis, Writing – Original Draft. **Florian de Bettignies:** Conceptualisation, Methodology, Writing- Review and Editing. **Andrew Foggo:** Data curation, Formal analysis, Writing- Review and Editing. **Dominique Davault:** Funding acquisition, Resources, Writing- Review and Editing.

Research paper

**Sustained productivity and respiration of degrading kelp detritus in the
shallow benthos: detached or broken, but not dead.**

Nadia Frontier^{a, b†*}, Florian de Bettignies^{a†}, Andy Foggo^b, Dominique Davoult^a

^a Sorbonne Université, CNRS, UMR 7144 AD2M, Station Biologique de Roscoff, Place
Georges Teissier, F-29680 Roscoff, France

^b Marine Biology and Ecology Research Centre, School of Biological and Marine Sciences,
University of Plymouth, Drake Circus, Plymouth PL4 8AA, UK

Running title: Kelp detritus can sustain primary producer functions

*Corresponding author: frontier.nadia@gmail.com

Postal address: 29A Eythrope Road, Stone, Aylesbury, Buckinghamshire, HP17 8PH

† The authors contributed equally to this work

Running header: Kelp detritus fulfilling primary producer functions

All revisions have been highlighted in yellow

22

23 **ABSTRACT**

24 Temperate kelp forests contribute significantly to marine primary productivity and
25 fuel many benthic and pelagic food chains. A large proportion of biomass is exported
26 from kelp forests as detritus into recipient marine ecosystems, potentially contributing
27 to Blue Carbon sequestration. The degradation of this organic material is slow and
28 recent research has revealed the preservation of photosynthetic functions over time.
29 However, the physiological correlates of detrital breakdown in *Laminaria* spp. have
30 not yet been studied. The warming climate threatens to reshuffle the species
31 composition of kelp forests and perturb the dynamics of these highly productive
32 ecosystems. The present study compares the physiological response of degrading
33 detritus from two competing North East Atlantic species; the native Boreal *Laminaria*
34 *hyperborea* and the thermally tolerant Boreal-Lusitanian *L. ochroleuca*. Detrital
35 fragment degradation was measured by a mesocosm experiment across a gradient of
36 spectral attenuation (a proxy for depth) to investigate the changes in physiological
37 performance under different environmental conditions. Degradation of fragments was
38 quantified over 108 days by measuring the biomass, production and respiration (by
39 respirometry) and efficiency of Photosystem II (by PAM fluorometry). Data indicated
40 that whilst degrading, the photosynthetic performance of the species responded
41 differently to simulated depths, but fragments of both species continued to produce
42 oxygen for up to 56 days and sustained positive net primary production. This study
43 reveals the potential for ostensibly detrital kelp to contribute to Blue Carbon fixation
44 through sustained primary production which should be factored into Blue Carbon
45 management. Furthermore, the physiological response of kelp detritus is likely

dependent upon the range of habitats to which it is exported. In the context of climate change, shifts in species composition of kelp forests and their detritus are likely to have wide-reaching effects upon the cycling of organic matter in benthic ecosystems.

Keywords: Climate change; Detritus; North East Atlantic; Oxygen production; PAM fluorescence; Respiration; Trophic transfer

1. INTRODUCTION

Within the temperate to sub-polar regions of the globe, kelps are of paramount importance to marine ecosystem functioning. They contribute to the functional integrity of coastal ecosystems as habitat forming species (HFS) via their biogenic structure, through wave energy dissipation and via cycling of carbon (Christie *et al.* 2003; Laffoley & Grimsditch, 2009; Teagle *et al.* 2017). As highly productive primary producers, kelps fix carbon dioxide to produce organic matter via photosynthesis (Bartsch *et al.* 2008). The accumulated biomass is eventually released or eroded and exported as detritus in a variety of forms into recipient benthic ecosystems (Duggins *et al.* 1989; Krumhansl and Scheibling 2012; Pessarrodona *et al.* 2018a). This transfer of carbon helps to sustain adjacent food webs. The breakdown of detrital material by erosion and grazing activity provides a source of particulate organic matter (POM) available to filter-feeding invertebrates (Renaud *et al.* 2015; de Bettignies *et al.* 2020a; Vilas *et al.* 2020) which in turn can play a major role in mediating carbon fluxes along the sediment-water interface (Queirós *et al.* 2019; Pedersen *et al.* 2020).

A significant proportion of primary production from marine primary autotrophs is exported as detritus rather than entering the consumer food web (Moore *et al.* 2004). In a global context,

as much as 82% of annual kelp biomass may be transferred as detrital subsidies (Krumhansl and Scheibling 2012; de Bettignies *et al.* 2013). Knowing the fate of such detritus is fundamental in understanding the dynamics, connectivity and functioning of coastal ecosystems (Polis *et al.* 1997). Indeed, detritus enhances both primary and secondary production (Polis *et al.* 1997; Marczak *et al.* 2007) whilst contributing to the structure and stability of food webs in recipient ecosystems (Huxel *et al.* 2002). Furthermore, accumulation of detritus can physically modify the habitat structure (Arroyo and Bonsdorff 2016). However, the influences upon recipient ecosystems are dependent upon the timing, frequency and residence time of the detrital deposition (Yang *et al.* 2008).

Macroalgae represent an as yet under-estimated contributor to the oceanic carbon cycle and in particular to carbon sequestration and long-term storage. Their contribution to global carbon assimilation and sequestration is currently not considered by the IPCC (Ciais *et al.* 2013), and whilst the Global Carbon Project (GCP) identifies coastal habitats as carbon stocks (Friedlingstein *et al.* 2019), their full contribution to the global marine carbon sink remains neglected. Unlike seagrasses or phytoplankton, macroalgae are not routinely considered as contributors to the ‘Blue Carbon’ budget (McLeod *et al.* 2011; Howard *et al.* 2017). Marine macrophytes collectively fix around 1.8 Gt C yr^{-1} (of which macroalgae contribute $\sim 1.5 \text{ Gt C yr}^{-1}$) (Duarte *et al.* 2013; Duarte and Krause-Jensen 2017; Krause-Jensen and Duarte 2016) in contrast to the $\sim 58 \text{ Gt C yr}^{-1}$ of phytoplankton production (Buitenhuis *et al.* 2013; Middelburg 2019). However, the lability of phytoplankton carbon and the high rate of remineralisation through consumption and degradation mean that only in the region of $0.23 \text{ Gt C yr}^{-1}$ (0.4%) of phytoplankton production becomes sequestered into carbon stores or sinks, whilst the lower rate of consumption and refractory chemical nature of macroalgal detritus mean that $\sim 0.17 \text{ Gt C yr}^{-1}$ (11.4%) is sequestered. Thus, including

seagrasses, marine macrophytes potentially contribute a comparable mass to direct global carbon sequestration to that of phytoplankton.

Species of the order Laminariales are particularly important contributors to coastal primary production (Kirk 1994) and form key constituents of shallow subtidal kelp forests across four continents (Steneck *et al.* 2002; Teagle *et al.* 2017). Along North East Atlantic coastlines, the stipitate canopy-forming *Laminaria hyperborea* dominates shallow subtidal rocky ecosystems (Smale and Moore 2017). Detrital production by *L. hyperborea* is unique because the majority of the previous-season's post-meristematic growth (the 'growth collar') is shed, often intact, between March and May (Bartsch *et al.* 2008); the release of this pulse of biomass contributes significantly to the total detritus production (Pessarrodona *et al.* 2018a). *L. hyperborea*'s detritus is a trophic resource connecting habitats which becomes increasingly accessible to consumers as it degrades (Norderhaug *et al.* 2003) and is an especially important dietary subsidy during periods of low plankton productivity (Leclerc *et al.* 2013a). As well as accumulating in coastal embayments and on beaches, a significant fraction of the carbon fixed by *L. hyperborea* flows into seafloor depressions, low-energy habitats and deep subtidal regions (Filbee-Dexter and Scheibling 2016; Filbee-Dexter *et al.* 2018). Research on *L. hyperborea* has already revealed that its degradation is slow relative to other species and accumulations in shallow subtidal ecosystems are able to maintain photosynthetic activity across several months (de Bettignies *et al.* 2020b), acting as net producers for a sustained period of time dependent upon the illumination regime.

The conspicuous thermally tolerant congeneric kelp (Franco *et al.* 2017; Hargrave *et al.* 2017) of Lusitanian origin, *L. ochroleuca* (Bachelot de la Pylaie), has expanded its range northwards from Morocco and across the English Channel in response to the changing

climate over the last century. Today, *L. ochroleuca* is expanding its leading edge eastwards and northwards at around 5.4 and 2.5 km per year, respectively (Straub *et al.* 2016). Its range now overlaps substantially with the native assemblage's dominant kelp species, *L. hyperborea* (Smale *et al.* 2015; Hargrave *et al.* 2017) which has undergone a ~250 km range contraction at its warm leading-edge since 1970 (Assis *et al.* 2016). In comparison to the pulsed detrital production from *L. hyperborea*, detrital production by *L. ochroleuca* is greater in volume, more frequent and degrades faster between May and October (Pessarrodona *et al.* 2018b). The expanding thermal envelope and subsequent spread of *L. ochroleuca* is modifying both the taxonomic and functional composition of HFS, with resulting effects on detrital composition. The re-arrangement of HFS has been shown to have marked ecological implications; extending to adjacent communities which rely on these subsidies of allochthonous material (Bishop *et al.* 2010; Straub *et al.* 2016).

The ecological function of detritus varies between species. Recent studies (Pessarrodona *et al.* 2018b) comparing *L. ochroleuca* and *L. hyperborea* have demonstrated interspecific differences in seasonal detrital decomposition rates, with *L. ochroleuca* exhibiting significantly faster rates (6.5 x) compared to *L. hyperborea* (Pessarrodona *et al.* 2018b) in spring. In addition, the species exhibit differences in timing of detrital production: the May cast from *L. hyperborea* accounting for ~40-60% of annual lamina erosion (Pessarrodona *et al.* 2018a) compared with the continual release of detritus from the fronds of *L. ochroleuca* (Pessarrodona *et al.* 2018b). Such alterations to the supply and cycling of organic matter will likely be amplified as sea surface temperatures increase. In the North East Atlantic, a rise of > 2°C is forecast in the next 100 years (Philippart *et al.* 2011; Masson-Delmotte *et al.* 2018), which will likely result in the gradual replacement of the native cool-water *L. hyperborea* by the warm-water tolerant *L. ochroleuca*.

145

146 Degradation is dynamic and testing the response of detritus across different environmental
147 conditions is important to contextualise this process. The depth distribution of photosynthetic
148 kelp-forest communities is strongly influenced by light availability (Kirk 1994; Gorman *et al.*
149 2013; Bajjouk *et al.* 2015) and the photosynthetic mechanisms of some kelp species have
150 been demonstrated to respond to changes in underwater light (e.g. in *Saccharina latissima*;
151 and *L. digitata*) (Gevaert *et al.* 2002; Delebecq *et al.* 2011). *L. hyperborea* and *L. ochroleuca*
152 forests have similar natural depth range but little is known about the photosynthetic
153 efficiency of these two species within their depth thresholds. Investigating the degradation of
154 kelp detritus across a gradient of depth will further elucidate its role in the marine
155 environment.

156

157 The aim of this study therefore was to monitor the break-down dynamics and physiological
158 performance of material from *L. hyperborea* and *L. ochroleuca* across a gradient of light
159 attenuation to investigate the influence of depth across time. We examined biomass
160 dynamics, oxygen production, respiration and photosynthetic performance and efficiency of
161 photosystem II of detritus from the two species. We hypothesized that *Laminaria* species
162 have the capacity to resist degradation, maintaining primary production function depending
163 upon the illumination regime. Following the findings of Pessarrodona *et al.* (2018a) we
164 hypothesised that the two species should exhibit differential responses with respect to
165 degradation, with a quicker decomposition and breakdown of physiological capacity for *L.*
166 *ochroleuca* compared to *L. hyperborea*. We also speculated that there would be an effect of
167 the amount of light, and for the first time, empirically tested the influence of depth-related
168 light attenuation as a potential driver of variability in the degradation of kelp material.

169

170 **2. MATERIAL AND METHODS**

171 The study was conducted in the Roscoff Biological Station close to the Bay of Morlaix
172 (France). In May 2018, *L. hyperborea* was harvested at La Veille (48°42'36.22"N,
173 3°54'04.81"W) and *L. ochroleuca* at An Nehou (48°41'34.27"N, 3°56'25.52"W) from ~ 8-10
174 m below chart datum. Five typical mature canopy-forming individuals of each kelp species
175 were harvested by severing the stipe just above the holdfast. The mean total length of *L.*
176 *hyperborea* individuals was 348 cm compared to *L. ochroleuca* individuals with a mean
177 length of 169 cm. Within two hours of collection, the sporophytes were submerged in the
178 dark in a 500 L flow-through tank of aerated ambient sea water pumped directly from the Bay
179 and left to acclimate for 48 hours.

180

181 *2.1 Experimental design*

182

183 For each adult individual, three blade fragments of equal length (20 cm) were cut at a
184 standardised distance (20 cm) above the meristem, away from the distal section of the kelp
185 blade which is subject to greater epiphytic colonisation. Fragments were tagged with a unique
186 ID number, their mass recorded and were randomly allocated to a depth (0, 15 and 30 m)
187 treatment. Thus there were five replicates per species within each depth treatment.

188

189 To simulate different light intensities across a depth gradient, light attenuation optical filters
190 (LEE Filters, Andover, UK) were constructed to completely cover the aquaria (Fig. 1). A
191 photon flux simulating light availability at 15 and 30 m was recreated using an average light
192 extinction coefficient (*k*) of 0.14 (from Boutler *et al.* 1974; spring and summer surveys

around Roscoff) incorporated into equation 1, where light availability at a specified depth equates to:

$$I(t) = I(z_0) \exp(-kz) \quad \text{equation 1.}$$

The irradiance at depth ($I(t)$) depends on the light at the surface $I(z_0)$, the depth (z) and the extinction coefficient of the water in the column (k). The 0 m treatment, representing strandline and floating detritus, were submerged in 8 cm of water hence $z_0 = 0.08$ m. The percentage of light available at 0, 15 and 30 m was 98.9%, 12.3% and 1.5% respectively; these values were utilised when assembling the relevant filters. We did not, however, account for the change in spectral character as a function of depth (Saulquin *et al.* 2013).

Fig. 1. Experimental set-up composed of filters for 15 and 30 m aquaria, constant air bubbling and HOBO loggers. Two unlabelled aquaria were used for experiments that are outside the scope of this paper. Each aquarium contained five replicate fragments. The brown squares represent aquaria with *L. hyperborea* fragments and yellow squares contain *L. ochroleuca* fragments.

The fragments acclimated for a further 48 hours before being transferred to 8 L aquaria corresponding to their respective species by depth treatment where they remained for the

duration of the experiment. Each aquaria was installed with a wide mesh basket which held the fragments in 8 cm of water and raised them 5 cm from the bottom. The baskets facilitated water circulation whilst constant aeration promoted diffusion across the boundary layer (Noisette and Hurd 2018). The six aquaria were held in a circulating water bath system supplied with a continual flow of unfiltered seawater (renewal rate of 10 times the aquarium volume per hour) located outside (Fig. 1). The fragments were therefore exposed to ambient temperature and irradiance in order to re-create their natural environment. Two aquaria were fitted with temperature/light loggers (HOBO Pendant Temperature/Light Weatherproof Pendant Data Logger 16K) recording at 10-minute intervals. Data was regularly downloaded to ensure conditions did not deviate significantly from outside conditions. Physiological measurements (see below) began 9 days after the fragments were transferred into the experimental system and observations were repeated at 7-day intervals thereafter across five-time periods, (T1-T5) with a 19 day gap between T5 and T6, for a total duration of 56 days.

2.2 Reproductive tissues and biomass change

Photographs of each fragment were taken at each time point to monitor visual changes in tissue surface (Figure S1), for example, the presence of sorus (reproductive) tissue (Figure S2). Repeated measures of wet biomass were taken on the same fragment at each time point. Biomass changes were calculated as a percentage of the starting mass remaining at each time point. Completely degraded fragments were removed and their biomass from the preceding time was recorded as mass lost. This aspect of the trial was extended to 108 days and final biomass measurements were recorded after all the fragments from the 30 m treatment had degraded.

2.3 Production and respiration measurements

Production and respiration rates were measured using a closed bottle respirometry technique (Migné *et al.* 2002; Biscéré *et al.* 2019). Twenty incubation chambers, each comprising of a 1.2 L transparent glass jar closed by a watertight seal, were distributed across four 50 L experimental units. The experimental units were each fitted with optical filters replicating light conditions in the corresponding aquaria and were replenished with the same ambient unfiltered seawater after each set of production and respiration incubations. Fragments were transferred in the dark from their aquaria into the corresponding depth treatment of the experimental unit with each jar containing an individual fragment. Two different incubations were conducted consecutively, measuring both oxygen production and consumption (respiration) respectively. Production was measured during a 60-minute incubation (long enough to allow detection of a change in oxygen concentration whilst avoiding oxygen saturation) under the experimental aquarium conditions. The jars were then opened to allow water exchange for 30 minutes and the containers were covered by dark tarpaulin to halt photosynthetic activity. The jars were subsequently re-sealed and the respiration incubations were conducted on the same fragments for 60 minutes in complete darkness. Dissolved oxygen concentrations were measured before and after both incubations (production and respiration) using a portable multi-meter (HQ40d, Hach®, Loveland, USA) coupled with a luminescent/optical dissolved oxygen probe (Intellical™ LDO101, Hach®, accuracy ± 0.2 mg L⁻¹). At the end of the incubations, fragments were retrieved from the jars, gently blotted dry and weighed (WW in g). Production and respiration rates were estimated by calculating the difference between initial and final oxygen O₂ concentrations after being corrected for temperature change (Aminot and Kérouel, 2004). The rates were expressed in mgO₂ kg WW⁻¹ hr⁻¹. The production rate is a measure of Net Primary Production (NPP), which represents the sum of photosynthesis and respiration in tandem. Meanwhile, Gross Primary Production

(GPP) was calculated by adding the dark respiration rate (R) to the NPP to derive the total oxygen produced via photosynthesis. The seawater temperature and incident photosynthetically active radiation (PAR, 400-700 nm, $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, Li-Cor QuantumSA-190, LI-COR®, Lincoln, USA), were recorded every minute during the trials.

2.4 Photosynthetic efficiency measurement

To reveal to the extent to which fragments' photosynthetic apparatuses were active, the operating efficiency of photosystem II (ΦPSII) and the maximal quantum yield of PSII photochemistry (F_v/F_m) were measured using a portable pulse-amplitude-modulated fluorometer (PAM, Heinz Walz, Effeltrich, Germany). *In vivo* ΦPSII was measured in ambient light whereas F_v/F_m measurements were obtained from fragments that had been dark adapted for 15 minutes (following Hargrave *et al.* 2017), before being flashed with a 0.8 s saturating white light pulse ($2500 \mu\text{mol photons.m}^{-2} \text{s}^{-1}$). F_v/F_m values normally range between 0.7-0.8 for Phaeophyceae and values below are indicative of a stress response (Bischof *et al.* 1999; Hanelt, 2018)

2.5 Statistical analyses

2.5.1 Biomass change

As a consequence of bimodality in the bounded biomass data, a Euclidean distance-based permutational ancova (PERMANCOVA) was used in PRIMER ver 6.1 (Primer-E, Plymouth) to test effects of species and depth (fixed factors) over time (continuous covariate), with fragment identity (the plant from which the fragment was cut) as a random factor. P values

were obtained from type III sums of squares. MDS plots were inspected as a way of identifying potential issues relating to heterogeneities of dispersion, though no formal tests were performed due to the complexity of the model.

2.5.2 Production, respiration and photosynthetic parameters

The effects of species and depths across time on net and gross oxygen production (NPP and GPP), dark respiration (R) and combining NPP with 12 h of R to represent a full day's rate of oxygen flux (NPP-R), were tested with linear mixed effect models using Maximum Likelihood (package *lme4* in R 3.6.1) (Bates *et al.* 2015; R Core Team 2019). To account for variation in abiotic conditions across the experiment, mean PAR was used as a continuous covariate for oxygen production and mean temperature for dark respiration (PAR and temperature were strongly correlated). Fragment identity was included as a random factor and p values were obtained from type III Wald χ^2 tests. Plots of residuals versus fits were used to check assumptions of analyses. Data describing the photosynthetic parameters Fv/Fm and Φ_{PSII} were analysed by similar model structure using generalised least squares (glS) fitting (package *nlme*, Pinheiro *et al.* 2019) due to heterogeneities of variance; PAR was again used as a continuous covariate; no random factor could be included in the glS model, between-plant variability being accounted for in the overall model fit. P values were derived as described above.

2.5.3 Omnibus PERMANCOVA

To examine the *in toto* physiological response of the two species, a multivariate approach was employed. A z-transformation was used to normalise the physiological variables (GPP,

R, Fv/Fm and Φ PSII), and a Euclidean distance – based multivariate similarity matrix was produced in PRIMER ver 6.1. This was then used as the basis for a mixed model PERMANCOVA with 999 permutations of a reduced model employing the same structure as the linear models described above; p values were obtained from type III sums of squares using a pseudo-F statistic.

3. RESULTS

3.1 Degradation dynamics

The analysis of biomass change with depth indicated strong heterogeneity of response to the environmental factor over time in the different species (3-way interaction pseudo- $F_{2,144} = 12.857$, $p < 0.001$; Fig. 2); no evidence of confounding heterogeneity of dispersion was apparent in the MDS plots. The interspecific differences were most apparent in the 15 and 30 m treatments, in both cases some of the *L. hyperborea* fragments degraded more quickly than *L. ochroleuca*. Under surface conditions neither species degraded markedly until after 56 days. No intact fragments of *L. ochroleuca* remained at the end of the 108 days, while *L. hyperborea* fragments were generally still intact at this point. With light attenuated to levels at 15 m, *L. hyperborea* began to degrade after 23 days, whilst *L. ochroleuca* persisted until the end of the study in the majority of cases. At 30 m most *L. hyperborea* fragments had completely broken down after 36 d, in contrast to *L. ochroleuca* where degradation was significantly slower (Fig. 2, Table 1).

3.2 Reproductive tissues and biomass change

Clear evidence of sorus material was first detected whilst conducting the final physiological measurements, 56 days into the experimental period on two *L. ochroleuca* fragments, one in the 0 m treatment and one in the 15 m treatment. When the study was concluded 108 days later, two different *L. ochroleuca* fragments had evidence of sorus material on the blade surface (Figure S2).

Fig. 2. Remaining biomass at three different simulated depths over time in *Laminaria hyperborea* and *Laminaria ochroleuca* (N = 144). Boxes indicate the interquartile range, horizontal line inside the bars indicates the median.

Table 1: Results of permanova analysis (type III SS) of % biomass remaining in the different species and depths (fixed factors) over 108 days (continuous covariate). Significant terms (P<0.05) are indicated in bold.

Source	Df	Pesudo-F	P
Time	1	119.37	0.001

Species	1	2.977	0.104
Depth	2	12.686	0.002
Time:Species	1	18.33	0.001
Species:Depth	2	2.893	0.066
Time: Depth	2	14.448	0.001
Time:Species:Depth	2	12.857	0.001
Fragment ID	24	6.331	0.001

349

350 *3.3 Production and respiration measurements*

351 As expected, the availability of light was the dominant factor affecting GPP; both in terms of
352 PAR at the surface ($\chi^2_{1,161} = 15.735$, $p < 0.001$) and as a main effect of photo-attenuation
353 related to depth ($\chi^2_{2,161} = 92.036$, $p < 0.001$). Contrary to expectation, however, across the
354 degradation period GPP did not decline as expected, and whilst the two species responded
355 differently to the passage of time ($\chi^2_{1,161} = 5.122$, $p = 0.024$), it was apparent that overall
356 oxygen evolution was indicative of the metabolic integrity of surviving tissues remaining
357 remarkably intact often beyond 36 days post-detachment (Fig. 3a, Table 2).

358 Respiration (R) was independent of temperature, but differed strongly between species ($\chi^2_{1,161} = 22.555$,
359 $p < 0.001$), and as a function of time ($\chi^2_{1,161} = 8.462$, $p = 0.004$) and depth ($\chi^2_{2,161} = 10.276$,
360 $p = 0.006$), modifying the species effect. Notwithstanding the declining
361 number of fragments persisting to the end of the trial (Fig. 3), it was clear that R did not
362 display a simple response over time; only the data for *L. hyperborea* under surface light
363 levels are suggestive of a simple linear increase over time (Fig. 3b, Table 2).

364 Most importantly, Net Primary Production (NPP) during lit periods remained at, or recovered
365 to, positive net production levels in the majority of fragments until the end of the 56 day trial
366 (Fig. 3c, Table 2). As with GPP and R, NPP differed strongly at different depths ($\chi^2_{1,161} =$
367 88.31 , $p < 0.001$). NPP did not differ statistically between the two species, but the nature of
368 this experiment restricts these conclusions to brightly lit daylight hours. Assuming

369 equinoctial fully light and dark conditions, with NPP and an additional 12 h of R representing
370 a full day's oxygen flux, suggests a significant effect of species with an additional modifying
371 effect of depth upon this ($\chi^2_{2,161} = 6.244$, $p = 0.044$). Under this '24 h' model, a more realistic
372 interpretation of differential GPP responses between the species can be detected (Fig. 3d,
373 Table 2).
374

Fig. 3. (a) Gross Primary Production (GPP), (b) Respiration (R), and (c) Net Primary Production (NPP) and (d) '24 hour' model of oxygen flux (NPP-R) rates, expressed as oxygen consumption or production, at three different simulated depths over time for *Laminaria hyperborea* and *Laminaria ochroleuca* (N = 161). Boxes indicate the interquartile range, horizontal line inside the bars indicates the median.

Table 2: Results of likelihood ratio tests based upon linear mixed effect models examining responses of (a) Gross Primary Production (GPP), (b) Respiration, (c) Net Primary Production (NPP) and (d) '24 hour' model of oxygen flux (NPP-R) rates across time (56 days), between kelp species and depth (fixed factors) with PAR or Temperature as a covariate. Significant terms ($P < 0.05$) are indicated in bold.

Source	df	Chisq	P
(a) GPP			
PAR	1	15.735	< 0.001
Time (days)	1	0.566	0.452
Species	1	0.000	0.996
Depth	2	92.036	< 0.001
Time: Species	1	5.122	0.024
Species: Depth	2	1.157	0.561
Time: Depth	2	6.316	0.043
Time:Species:Depth	2	4.058	0.131
Fragment ID	1	9.664	0.002
(b) R			
Temp	1	0.098	0.754
Time (days)	1	6.110	0.013
Species	1	22.555	< 0.001
Depth	2	2.234	0.327
Time:Species	1	8.462	0.004
Species: Depth	2	10.276	0.006
Time: Depth	2	1.369	0.504
Time: Species: Depth	2	0.047	0.977
Fragment ID	1	3.727	0.054
(c) NPP			
PAR	1	9.160	0.002

Time (days)	1	3.833	0.050
Species	1	2.032	0.154
Depth	2	88.311	< 0.001
Time:Species	1	1.246	0.262
Time: Depth	2	7.147	0.028
Species: Depth	2	3.589	0.166
Time: Species: Depth	2	3.704	0.157
Fragment ID	1	11.22	0.001
(d) NPP- R			
PAR	1	4.050	0.044
Time (days)	1	7.924	0.005
Species	1	6.215	0.013
Depth	2	74.259	< 0.001
Time:Species	1	0.007	0.933
Time: Depth	2	6.861	0.032
Species: Depth	2	6.244	0.044
Time: Species: Depth	2	2.893	0.235
Fragment ID	1	11.226	< 0.001

3.4 Photosynthetic response

PAR was a significant covariate for the Φ_{PSII} response ($\chi^2_1 = 7.901$, $p < 0.01$). Despite time exerting a non-statistically significant effect on Φ_{PSII} (as it did upon NPP and GPP), the response of detrital fragments clearly declined over the degradation period (Fig. 4a) but became highly variable at increasing depth, swamping any trend. Fig. 4b shows that F_v/F_m displayed a similarly noisy decline over time. There was no significant difference in the photosynthetic response between the species. Both parameters were significantly affected by depth. No interactions were significant and therefore the photosynthetic behaviour of detrital resources from different species was not affected by time or by depth (Figs 4 a-b, Table 3).

397

398 **Fig. 4.** (a) ΦPSII and (b) F_v/F_m measurements at three different simulated depths over time
 399 for *Laminaria hyperborea* and *Laminaria ochroleuca* (N = 161). Boxes indicate the
 400 interquartile range, horizontal line inside the bars indicates the median.

401 **Table 3:** Results of likelihood ratio testing based upon generalized least squares fitting to
 402 examine responses of (a) ΦPSII (b) F_v/F_m across time (56 days), between kelp species and
 403 depths (fixed factors) with PAR as a covariate. Significant terms ($P < 0.05$) are indicated in
 404 bold.

Source	<i>df</i>	Chisq	<i>P</i>
(a) Φ PSII			
PAR	1	7.901	0.005
Time	1	0.221	0.638
Species	1	0.935	0.334
Depth	2	51.403	<0.001
Time:Species	1	0.694	0.405
Species: Depth	2	2.700	0.260
Time:Species:Depth	2	1.331	0.514
(b) F_v/F_m			
PAR	1	13.418	0.001
Time	1	1.620	0.203
Species	1	0.230	0.632
Depth	2	5.981	0.050
Time:Species	1	0.428	0.513
Species: Depth	2	2.654	0.265
Time:Species:Depth	2	1.641	0.440

405

406 *3.5 Omnibus permanova*

407 The combined response of GPP, R, Φ PSII and F_v/F_m variables was strongly affected by the
408 light availability in terms of PAR ($\chi^2_1 = 11.176$, $p < 0.001$) and Depth ($\chi^2_1 = 21.585$, $p <$
409 0.001). A realistic picture of detrital degradation revealed species responded differently to the
410 passage of time ($\chi^2_1 = 9.327$, $p < 0.001$) but not depth (Table 4).

411

412 **Table 4:** Results of omnibus PERMANOVA analysis of standardized data describing GPP,
413 R, Φ PSII and F_v/F_m across time (56 days), between kelp species and depth (fixed factors)
414 with PAR as a covariate. Significant terms ($P < 0.05$) are indicated in bold.

415

Source	<i>df</i>	Pesudo-F	<i>P</i>
--------	-----------	----------	----------

GPP, R, ΦPSII, F_v/F_m			
PAR	1	11.176	0.001
Time	1	4.837	0.012
Species	1	8.729	0.001
Depth	2	21.585	0.001
Time:Species	1	9.327	0.001
Species: Depth	2	2.269	0.081
Time:Species:Depth	2	0.848	0.519

4. DISCUSSION

4.1. Production and respiration measurements

For the first time, the primary producer functions of algal detritus have been quantified across a simulated depth spectrum. Detritus continued to sustain net production of oxygen across a 56-day period, underscoring its capacity to maintain key functions such as carbon fixation. Detrital tissues slightly increased in biomass during the early stages of degradation, demonstrating that measured oxygen production is reflective of fresh material being generated. Over time, the respiration of detrital fragments, as a proxy for stress or associated bacterial activity, increased in some contexts (Sosik and Simenstad, 2013). Evidence of some degradation was further reinforced by a measured decline in photosynthetic performance (Φ PSII and F_v/F_m), in accordance with the initial hypothesis. At the end of the first time period, values between 0.6-0.8 at 15 m and 30 m reflected an optimum performance at depths at which the kelps inhabit (Gorman *et al.*, 2013). Low photosynthetic parameters at 0 m follow reported trends in the literature of decreases in both measurements with increasing irradiances (for *Laminaria digitata*, Delebecq *et al.* 2011).

Respiration is a central process in kelp decomposition (de Bettignies *et al.* 2020b) and measured increases in the rate of oxygen consumption were mirrored by a decline in the

436 detrital biomass. However, it is important to note that the measurements reflect the holobiont
 437 respiration of *Laminaria* tissue together with its ensemble of microorganisms (Baedke *et al.*
 438 2020). The results of this study reveal that the respiration response differed between the two
 439 species across time. Detritus from *L. hyperborea* exhibited higher rates of respiration
 440 compared to *L. ochroleuca* over time in the 30 m treatment and only under surface conditions
 441 was a linear increase in respiration detected for *L. hyperborea*. This differential response
 442 between the species may be indicative of stress or higher biofilm activity associated with *L.*
 443 *hyperborea* fragments. *L. hyperborea* degraded faster than *L. ochroleuca* at 15 m and 30 m,
 444 suggesting that low light conditions may impede the regeneration of damaged tissue for *L.*
 445 *hyperborea* which points to the role of light in determining the detrital fate of some species
 446 (Hader *et al.* 1998; Swanson and Fox 2007). Furthermore, in some instances, during the
 447 initial stages of degradation at 0 and 15 m, *L. ochroleuca* positively increased in biomass
 448 compared to *L. hyperborea* (Fig. 2). However, this trend was reversed after 56 days in the 0
 449 m treatment when all *L. ochroleuca* fragments were completely degraded in sharp contrast to
 450 the relatively intact fragments of *L. hyperborea*. 4.2 Considering the first 40 days of this trial,
 451 this result is comparable, yet contrary to the findings of Pessarrodona *et al.*'s (2018b) *in situ*
 452 study comparing the biomass loss from the same study species. Pessarrodona *et al.* (2018b)
 453 report that *L. ochroleuca* lost biomass at a faster rate than *L. hyperborea* in spring, prior to
 454 the latter shedding of old fronds. The present study separated the change in mass across
 455 regular time intervals and elongated the temporal scale of degradation to 108 days, as well as
 456 eliminating physical erosion as a mechanism of degradation, casting light on the purely
 457 organic dimension of how kelp matter breaks down. Our results suggest that *L. hyperborea*
 458 has the potential to be a more persistent spatial subsidy and thus may reside in the ecosystem
 459 for longer under attenuated illumination. However, hydrodynamics and physical processes

such as abrasion and grazing may further confound elucidation of the decomposition process.
(Nielsen *et al.* 2004; Braeckman *et al.* 2019).

4.2. Contextualizing the findings

These differences in persistence will become increasingly important as *L. ochroleuca* gradually replaces *L. hyperborea*, in habitats where the two species coexist, (Teagle and Smale 2018) and will be further exacerbated by the seasonal modification of detrital production between the species; ultimately affecting the quality and supply of organic kelp derived material entering the detrital pathway (Bishop *et al.* 2010). The extent to which benthic ecosystems in the North East Atlantic are dependent on detritus derived from *L. hyperborea* remains unquantified. However, modifications in the turnover of organic material, transport (Pedersen *et al.* 2005), trophic connectivity (Leclerc *et al.* 2013b), organic matter content (Abdullah *et al.* 2017) and a homogenization of the nutrient supply from kelp detritus derived from different species will likely have ramifying effects across food webs.

Extending the investigation to consider the degradation process across environmental contexts revealed that other measured physiological processes of kelp detritus (NPP, GPP and Φ_{PSII} and F_v/F_m) are not only species specific but also highly dependent on light availability. Depth significantly affected 24-hour oxygen flux and the Species response to the effect of Depth was significantly heterogeneous in the case of both respiration and the 24-hour model of oxygen flux (Fig. 3b and d). Therefore, considering the dynamic nature of detrital transport, the depth to which detritus is exported significantly determines the carbon fixation, respiration and photosynthetic performance of kelp material (Filbee-Dexter and Scheibling, 2016). Furthermore, detritus was clearly less productive as a function of decreasing irradiance

from 0 m > 15 m > 30 m (Fig. 4a). Photosynthetic capacity was critically impaired at 0 m, probably as a result of photoinhibition (Delebecq *et al.* 2011), but remained mostly invariant between 15 and 30 m. Although the activity of the photosynthetic apparatus was maintained in low light conditions, the decline in oxygen production is indicative that detritus is a less effective primary producer at 30 m. The differing responses of respiration in the two species over time and at different depths highlight that the process of degradation needs to be considered across different environmental conditions with a temporal dimension. Species' respiration fluctuated across time and their response was dependent on light attenuation, which indicates that one species may have the physiological capacity to persist for longer than its congener at a specific depth.

4.3. Re-defining kelp detritus

An unexpected result was that *L. ochroleuca* fragments developed visible sorus tissue at the tissue surface indicating that such fragments remain capable of maintaining reproductive functions (Fig S1). Moreover, this was detected first for two fragments during the final time period (T6) in July. Visible sorus remained at the blade surface two independent *L. ochroleuca* fragments when the experiment ended 108 days post fragment preparation in September. This time window is in accordance with the reproductive period for *L. ochroleuca* (Pereira *et al.* 2019), further revealing the metabolic competency of the kelps post-detachment. Further research would be necessary to determine spore viability from such tissues. de Bettignies *et al.* (2020b) reported a similar phenomenon for *L. hyperborea* in October after subjecting detrital fragments to 5 months of degradation *in situ*. In terms of reproductive capacity, some of the energy generated by detrital photosynthesis is invested into the generation of reproductive tissue which potentially has considerable implications for the dispersal of these species. If current-borne kelp detritus can still

reproduce, then kelp populations structure is likely to be far more fluid than would be indicated by typical spore or gamete dispersal, which is low for individual sporophytes within a kelp forest (~ 5-200 m, Fredriksen *et al.* 1995). This phenomenon compliments the functional role of herbivores in increasing fertilisation success and dispersing kelp spores which germinate post-digestion (Ruz *et al.* 2018).

Another important outcome of the present study and other recent work on physiological viability of macroalgal ‘detritus’ (de Bettignies *et al.* 2020b) is a questioning of the rationale behind defining this material as ‘detritus’. Currently, cast or detached macroalgal fragments are considered as ‘non-living’ organic matter generated by the growth and production of living organisms (Hagen *et al.* 2012): a viewpoint that is based upon terrestrial botanical science. Although such tissues may be no longer attached, and as our data suggests, may be in the process of slowly entering the detrital food web, they remain ‘alive’ and physiologically competent for much longer than would be typical for tissues of land plants. This ‘productive necromass’ – physiologically persistent and viable, unattached or fragmented algal biomass – continues to photosynthesize and, in the right conditions, can continue to fix carbon for months after it would otherwise be inaccurately defined as detrital. Our study has demonstrated that kelp detritus can sustain productivity for long periods of time and therefore detrital contribution to Blue Carbon is likely to be much greater than has been previously accounted for. Accordingly, incorporating productivity from macroalgal detritus could account for important increases to the contribution of macroalgae in the Blue Carbon economy.

4.4 The role of kelp detritus in benthic ecosystems

The vast majority of kelp production enters the detrital pathway (about 82% according to Krumhansl and Scheibling 2012), therefore fundamental variability in the respiration and photosynthetic performance of kelp material across time and under different environmental conditions are important for understanding the persistence and productivity of detrital resources. The composition of kelp forests and thus their detritus, will likely be modified as climate warming persists (Smale et al. 2015), with the gradual increase of HFS with high thermal affinities and gradual local extinctions of native HFS with cooler thermal affinities (Wiens 2016). These replacements will likely impact many aspects of the cycling of organic material, from the spatial subsidies exported kelp matter provides, to the fitness of grazers and detritivores feeding upon it, and the rate of carbon sequestration into longer residence forms in the benthos. Pessarrodona *et al.* (2018b) speculated about an alteration in the functional importance of this detrital material due to a variation in the rate of supply from different kelp species. Here we have shown that the continued biological activity of fragmented kelp tissues also differs between species, across time and environments, further altering the contribution of kelp mass to spatial subsidies (Leclerc *et al.* 2013a).

This study provides a snapshot of the degradation dynamics of two kelps under controlled conditions. There are numerous potential sources for error; for example a change in detrital respiration could be attributed an upregulation of metabolic processes in *L. hyperborea* or to an increase in oxygen consuming saprophytic bacteria and fungi (Fenchel and Jørgensen, 1977; Williams *et al.* 2004). Furthermore, variation in species physiology cannot be generalized across the entirety of benthic ecosystems in the North East Atlantic because despite advances in areas of low-moderate wave exposure (Smale *et al.* 2015), *L. ochroleuca* may not be able to fully replace *L. hyperborea* in sites exposed to high wave exposure (Pessarrodona *et al.* 2018b) and therefore the composition of detritus from such assemblages may remain relatively unchanged. Also, the role of macrofaunal colonization and consumption (de Bettignies *et al.* 2020; Ramirez-Llodra *et al.* 2016) of

the kelps in question could change the degradation dynamics completely. Hargrave *et al.* (2016) reported lower concentrations of grazer-deterrent polyphenolics and flavonoids in *L. ochroleuca* than in another common NE Atlantic congener: *L. digitata*, with higher grazing rate by herbivorous gastropods in the former. Meanwhile, Pessarrodona *et al.* (2018b) found a similar herbivore preference for *L. ochroleuca* over *L. hyperborea*, recording a faster degradation of the former. If detritivore preference for the more palatable *L. ochroleuca* mirrors that of grazers, then the reconciliation of the slower degradation of *L. ochroleuca* in the earlier stages of the present study with the faster *in situ* degradation of *L. ochroleuca* observed by Pessarrodona *et al.* (2018b) could be attributed to mechanical effects, especially of saprovores aiding conversion of frond tissue into coarse particulate organic matter and facilitating ingress by saprophytic micro-organisms. Future study should therefore aim to evaluate the importance of the ecological consequences of environmentally realistic deposits of macroalgal detritus with their attendant bacterial and detritivorous assemblages and also continue to compare the detrital compositions from species at risk of local extinction from a variety of range-expanding or invasive algae (Sorte *et al.* 2010), taking into account different environmental contexts in which kelp degradation may occur, testing the degradation response *in situ* across a gradient of turbulence, and explicitly focusing on detritivores as a driver of tissue fragmentation and a facilitator of micro-organismal colonization. These questions have indeed been addressed in other contexts spanning from shallow shorelines to the deep sea whilst the shallow subtidal has often been considered as a mere transitional deposition area. However, we argue that detrital accumulations can exert measurable impacts on ecosystem functioning and deserve enhanced attention.

5. CONCLUSION

Kelp fragments remained physiologically competent for 56 days and possibly reproductively active for up to 108 days after they were cut from the thallus. The fragments' photosynthetic apparatus continued to function adequately to allow short-term (~ 56 days in *L. ochroleuca*) net gains in biomass which was sustained under ambient light levels. Light availability was the greatest determinant of photosynthetic performance, and net and gross primary production, whilst light and interspecific differences dominated the response of respiration. The range-expanding *L. ochroleuca* broke down more slowly under attenuated illumination, challenging previous findings. We believe that a re-evaluation of recently formed macroalgal detritus as 'living material' is due and should certainly be considered when evaluating the functional integrity and spatial subsidies afforded by kelp forests. The predicted expansion of the thermally tolerant species, *L. ochroleuca*, and replacement of the less tolerant *L. hyperborea* across extensive areas is likely to alter the cycling of organic matter. Overall, these findings demonstrate the indispensable need to consider detritus as an important autotrophic resource, even after exportation, depending on the environmental conditions.

DATA AVAILABILITY

Data will be archived in GitHub repository: <https://github.com/nadiafrontier/The-degradation-of-kelp-detritus.git>

FUNDING

This work was supported by the Brittany Regional Council and the French Government through the National Research Agency with regards to the investment expenditure program IDEALG [grant number ANR-10-BTBR-04].

SUPPLEMENTARY INFORMATION

Supplementary tables 1-4 contain abridged statistical tables following permanova, linear mixed effect models and generalised last squares analysis to support the summary statistics quoted in the manuscript.

ACKNOWLEDGEMENTS

We thank the marine operation staff at Roscoff Biological Station (*Service Mer & Observation SBR*) L. Levêque, Y. Fontana, W. Thomas, M. Camusat, N. Guidal and F. Le Ven for their help with scuba diving kelp collection and T. de Bettignies from the Natural History Museum of Paris. We also thank Marine Roscoff Aquarium Services (*FR2424*), especially R. Garnier and G. Schires for their help with the experimental design and set-up. E. Bocher and M. Kervellec for their assistance in the experimental work. NF, FdB and DD conceived the ideas and designed the study. FdB and NF built the experimental system. NF and FdB conducted the experimental manipulation. AF and NF analysed the data. NF and FdB wrote the first draft and AF extensively contributed to subsequent drafts. All authors gave final approval for publication.

REFERENCES

- Abdullah MI, Fredriksen S, Christie H. 2017.** The impact of the kelp (*Laminaria hyperborea*) forest on the organic matter content in sediment of the west coast of Norway. *Marine Biology Research* **13**: 151–160. doi: 10.1080/17451000.2016.1240369
- Aminot A, Kerouel R. 2005.** *Hydrologie des écosystèmes marins: paramètres et analyses. Méthodes d'analyse en milieu marin.* Ifremer, France.
- Arroyo N, Bonsdorff E. 2016.** The Role of Drifting Algae for Marine Biodiversity. In: Olafsson E. *Marine Macrophytes as Foundation Species*. 1st edn. CRC Press: Taylor &

- Francis group, 100-123. doi: 10.4324/9781315370781-6
- Assis J, Lucas AV, Bárbara I, Serrão EÁ. 2016.** Future climate change is predicted to shift long-term persistence zones in the cold-temperate kelp *Laminaria hyperborea*. *Marine Environmental Research* **113**: 174–182. doi: 10.1016/j.marenvres.2015.11.005
- Baedke J, Fábregas-Tejeda A, Nieves Delgado A. 2020.** The holobiont concept before Margulis. *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution*, **334**: 149–155. doi: 10.1002/jez.b.22931
- Bajjouk T, Rochette S, Laurans M, Ehrhold A, Hamdi A, Le Niliot P. 2015.** Multi-approach mapping to help spatial planning and management of the kelp species *L. digitata* and *L. hyperborea*: Case study of the Molène Archipelago, Brittany. *Journal of Sea Research* **100**: 2–21. doi: 10.1016/j.seares.2015.04.004
- Bartsch I, Wiencke C, Bischof K, et al. 2008.** The genus *Laminaria* sensu lato: Recent insights and developments. *European Journal of Phycology* **43**: 1–86. doi: 10.1080/09670260701711376
- Bates D, Maechler, M, Bolker B, Walker, S. 2015.** Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software* **67**: 1-48. doi:10.18637/jss.v067.i01.
- Biscéré T, Zampighi M, Lorrain A, et al. 2019.** High pCO₂ promotes coral primary production. *Biology Letters* **15**. doi: 10.1098/rsbl.2018.0777
- Bischof K, Hanelt D, Wiencke C. 1999.** Acclimation of maximal quantum yield of photosynthesis in the brown alga *Alaria esculenta* under high light and UV radiation. *Plant Biology* **1**: 435–444. doi: 10.1111/j.1438-8677.1999.tb00726.x
- Bishop MJ, Coleman MA, Kelaher BP. 2010.** Cross-habitat impacts of species decline: Response of estuarine sediment communities to changing detrital resources. *Oecologia* **163**: 517–525. doi: 10.1007/s00442-009-1555-y
- Boutler J, Carbioch L, Grall R. 1974.** Quelques observations sur la pénétration de la

- 654 lumière dans les eaux marines au voisinage de Roscoff et ses conséquences écologiques.
 655 *Laboratoire d'Océanographie Physique*. Paris, 129–140.
- 656 **Braeckman U, Pasotti F, Vázquez S, et al.** 2019. Degradation of macroalgal detritus in
 657 shallow coastal Antarctic sediments. *Limnology and Oceanography*, **64**: 1423–1441. doi:
 658 10.1002/lno.11125
- 659 **Buitenhuis ET, Hashioka T, Quéré C Le.** 2013. Combined constraints on global ocean
 660 primary production using observations and models. *Global Biogeochemical Cycles* **27**: 847–
 661 858. doi: 10.1002/gbc.20074
- 662 **Christie H, Jørgensen NM, Norderhaug KM, Waage-Nielsen E.** 2003. Species
 663 distribution and habitat exploitation of fauna associated with kelp (*Laminaria hyperborea*)
 664 along the Norwegian coast. *Journal of the Marine Biological Association of the United*
 665 *Kingdom* **83**: 687–699. doi: 10.1017/S0025315403007653h
- 666 **Ciais P, Sabine C, Bala G, et al.** 2013. Carbon and other Biogeochemical Cycles. In:
 667 *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the*
 668 *Fifth Assessment Report of the Intergovernmental Panel on Climate Change*: 465–570.
- 669 **de Bettignies T, Wernberg T, Lavery PS, Vanderklift MA, Mohring MB.** 2013.
 670 Contrasting mechanisms of dislodgement and erosion contribute to production of kelp
 671 detritus. *Limnology and Oceanography* **58**: 1680–1688. doi: 10.4319/lo.2013.58.5.1680
- 672 **de Bettignies F, Dauby P, Lepoint G. et al.** 2020a. Temporal succession of a macrofaunal
 673 community associated with kelp fragment accumulations in an in situ experiment. *Marine*
 674 *Ecology Progress Series*. in press. doi: 10.3354/meps13391
- 675 **de Bettignies F, Dauby P, Thomas F, et al.** 2020b. Degradation dynamics and processes
 676 associated with the accumulation of *Laminaria hyperborea* (Phaeophyceae) kelp fragments:
 677 an in situ experimental approach. *Journal of Phycology*. in press. doi: 10.1111/jpy.13041
- 678 **Delebecq G, Davoult D, Menu D, et al.** 2011. In situ photosynthetic performance of

- 679 *Laminaria digitata* (phaeophyceae) during spring tides in northern Brittany. *Cahiers de*
680 *Biologie Marine* **52**: 405–414. doi: 10.1080/09670262.2015.1104556
- 681 **Duarte CM, Krause-Jensen D. 2017.** Export from seagrass meadows contributes to marine
682 carbon sequestration. *Frontiers in Marine Science* **4**: 1–7. doi: 10.3389/fmars.2017.00013
- 683 **Duarte CM, Losada IJ, Hendriks IE, Mazarrasa I, Marbà N. 2013.** The role of coastal
684 plant communities for climate change mitigation and adaptation. *Nature Climate Change* **3**:
685 961–968. doi: 10.1038/nclimate1970
- 686 **Duggins DO, Simenstad CA, Estes JA. 1989.** Magnification of secondary production by
687 kelp detritus in coastal marine ecosystems. *Science* **245**: 170–173. doi:
688 10.1126/science.245.4914.170
- 689 **Fenchel TM, Jørgensen BB. 1977.** Detritus Food Chains of Aquatic Ecosystems: The Role
690 of Bacteria. : 1–58. doi: 10.1007/978-1-4615-8219-9_1
- 691 **Filbee-Dexter K, Scheibling RE. 2016.** Spatial Patterns and Predictors of Drift Algal
692 Subsidy in Deep Subtidal Environments. *Estuaries and Coasts* **39**: 1724–1734. doi:
693 10.1007/s12237-016-0101-5
- 694 **Filbee-Dexter K, Wernberg T, Norderhaug KM, Ramirez-Llodra E, Pedersen MF.**
695 **2018.** Movement of pulsed resource subsidies from kelp forests to deep fjords. *Oecologia*
696 **187**: 291–304. doi: 10.1007/s00442-018-4121-7
- 697 **Franco JN, Tuya F, Bertocci I, Rodríguez L, Sousa-pinto BMI, Arenas F. 2017.** The
698 ‘golden kelp ’ *Laminaria ochroleuca* under global change : Integrating multiple eco-
699 physiological responses with species distribution models. *Journal of Ecology* **106**: 47-58. doi:
700 10.1111/1365-2745.12810
- 701 **Fredriksen S, Sjøtun K, Lein TE, Rueness J, 1995.** Spore dispersal in *Laminaria*
702 *hyperborea* (laminariales, phaeophyceae). *Sarsia*, **80**, pp.47-53. doi:
703 10.1080/00364827.1995.10413579

- 704 **Friedlingstein P, Jones MW, O’Sullivan M, et al. 2019.** Global carbon budget 2019. *Earth*
 705 *System Science Data* **11**: 1783–1838. doi: 10.5194/essd-11-1783-201
- 706 **Gevaert F, Creach A, Davoult D, Holl AC, Seuront L, Lemoine Y. 2002.** Photo-inhibition
 707 and seasonal photosynthetic performance of the seaweed *Laminaria saccharina* during a
 708 simulated tidal cycle: Chlorophyll fluorescence measurements and pigment analysis. *Plant,*
 709 *Cell and Environment* **25**: 859–872. doi: 10.1046/j.1365-3040.2002.00869.x
- 710 **Gorman D, Bajjouk T, Populus J, Vasquez M, Ehrhold A. 2013.** Modeling kelp forest
 711 distribution and biomass along temperate rocky coastlines. *Marine Biology* **160**: 309–325.
 712 doi: 10.1007/s00227-012-2089-0
- 713 **Hader DP, Kumar HD, Smith RC and Worrest RC, 1998.** Effects on aquatic ecosystems.
 714 *Journal of photochemistry and photobiology B: Biology.* **46**: 53-68. doi 10.1016/S1011-
 715 **1344(98)00185-7**
- 716 **Hagen EM, McCluney KE, Wyant KA, et al. 2012.** A meta-analysis of the effects of
 717 detritus on primary producers and consumers in marine, freshwater, and terrestrial
 718 ecosystems. *Oikos* **121**: 1507–1515. doi: 10.1111/j.1600-0706.2011.19666.x
- 719 **Hanelt D. 2018.** Photosynthesis assessed by chlorophyll fluorescence. In: Häder D, Erzingen
 720 G, *Bioassays: Advanced Methods and Applications*. Amsterdam: Elsevier Inc, 169-196
- 721 **Hargrave MS, Foggo A, Pessarrodona A, Smale DA. 2017.** The effects of warming on the
 722 ecophysiology of two co-existing kelp species with contrasting distributions. *Oecologia* **183**:
 723 531–543. doi: 10.1007/s00442-016-3776-1
- 724 **Howard J, Sutton-Grier A, Heer D, et al. 2017.** Clarifying the role of coastal and marine
 725 systems in climate mitigation. *Frontiers in Ecology and the Environment* **15**: 42–50. doi:
 726 10.1038/news050808-1
- 727 **Huxel GR, McCann K, Polis GA. 2002.** Effects of partitioning allochthonous and
 728 autochthonous resources on food web stability. *Ecological Research* **17**: 419–432. doi:

- 729 10.1046/j.1440-1703.2002.00501.x
- 730 **Kirk JT, 1994.** *Light and photosynthesis in aquatic ecosystems*, 2nd edn. Cambridge:
- 731 Cambridge university press. doi: 10.1017/CBO9780511623370
- 732 **Krause-Jensen D, Duarte CM. 2016.** Substantial role of macroalgae in marine carbon
- 733 sequestration. *Nature Geoscience* **9**: 737–742. doi: 10.1038/ngeo2790
- 734 **Krumhansl KA, Scheibling RE. 2012a.** Production and fate of kelp detritus. *Marine*
- 735 *Ecology Progress Series* **467**: 281–302. doi: 10.3354/meps09940
- 736 **Krumhansl KA, Scheibling RE. 2012b.** Detrital subsidy from subtidal kelp beds is altered
- 737 by the invasive green alga *Codium fragile* ssp. *fragile*. *Marine Ecology Progress Series* **456**:
- 738 73–85. doi: 10.3354/meps09671
- 739 **Laffoley D, Grimsditch G. 2009.** *The management of natural coastal carbon sinks*. Gland:
- 740 Switzerland: IUCN. doi: 10.1007/s00114-001-0283-x
- 741 **Leclerc J-C, Riera P, Leroux C, Lvque L, Davoult D. 2013a.** Temporal variation in
- 742 organic matter supply in kelp forests: Linking structure to trophic functioning. *Marine*
- 743 *Ecology Progress Series* **494**: 87–105. doi: 10.3354/meps10564
- 744 **Leclerc JC, Riera P, Leroux C, et al. 2013b.** Trophic significance of kelps in kelp
- 745 communities in Brittany (France) inferred from isotopic comparisons. *Marine Biology* **160**:
- 746 3249–3258. doi: 10.1007/s00227-013-2306-5
- 747 **Marczak LB, Thompson RM, Richardson JS. 2007.** Meta-analysis: Trophic level, habitat,
- 748 and productivity shape the food web effects of resource subsidies. *Ecology* **88**: 140–148. doi:
- 749 10.1890/0012-9658(2007)88[140:MTLHAP]2.0.CO;2
- 750 **Masson-Delmotte V, Zhai P, Prtner HO, 2018.** IPCC Summary for Policymakers. Global
- 751 Warming of 1.5C. IPCC Special Report on the impacts of global warming of 1.5C above
- 752 pre-industrial levels and related global greenhouse gas emission pathways, in the context of
- 753 strengthening the global response to the threat of climate change, sustainable development,

and efforts to eradicate poverty.

McLeod E, Chmura GL, Bouillon S, et al. 2011. A blueprint for blue carbon: Toward an improved understanding of the role of vegetated coastal habitats in sequestering CO₂.

Frontiers in Ecology and the Environment **9**: 552–560. doi: 10.1890/110004

Middelburg JJ, 2019. Marine carbon biogeochemistry: a primer for earth system scientists. Springer Briefs in Earth System Sciences. doi: 10.1007/978-3-030-10822-9

Migné A, Davoult D, Spilmont N, et al. 2002. A closed-chamber CO₂-flux method for estimating intertidal primary production and respiration under emerged conditions. *Marine Biology* **140**: 865–869. doi: 10.1007/s00227-001-0741-1

Moore JC, Berlow EL, Coleman DC, et al. 2004. Detritus, trophic dynamics and biodiversity. *Ecology Letters* **7**: 584–600. doi: 10.1111/j.1461-0248.2004.00606.x

Nielsen SL, Banta GT, Pedersen M, 2004. Decomposition Of Marine Primary Producers: Consequences For Nutrient Recycling And Retention In Coastal Ecosystems. In: Nielsen S.L., Banta G.T., Pedersen M.F. (eds) *Estuarine Nutrient Cycling: The Influence of Primary Producers*. Aquatic Ecology Book Series, **vol 2**. Springer, Dordrecht. https://doi.org/10.1007/978-1-4020-3021-5_7

Noisette F, Hurd C. 2018. Abiotic and biotic interactions in the diffusive boundary layer of kelp blades create a potential refuge from ocean acidification. *Functional Ecology* **32**: 1329–1342. doi: 10.1111/1365-2435.13067

Norderhaug KM, Fredriksen S, Nygaard K. 2003. Trophic importance of *Laminaria hyperborea* to kelp forest consumers and the importance of bacterial degradation to food quality. *Marine Ecology Progress Series* **255**: 135–144. doi: 10.3354/meps255135

Pedersen MF, Filbee-Dexter K, Norderhaug KM, et al. 2020. Detrital carbon production and export in high latitude kelp forests. *Oecologia* **192**: 227–239.

Pedersen MF, Stæhr PA, Wernberg T, Thomsen MS. 2005. Biomass dynamics of exotic

- 779 *Sargassum muticum* and native *Halidrys siliquosa* in Limfjorden, Denmark - Implications of
 780 species replacements on turnover rates. *Aquatic Botany* **83**: 31–47. doi: 10.1007/s00442-019-
 781 04573-z
- 782 **Pereira TR, Azevedo IC, Oliveira P, Silva DM, Sousa-Pinto I. 2019.** Life history traits of
 783 *Laminaria ochroleuca* in Portugal: The range-center of its geographical distribution. *Aquatic*
 784 *Botany* **152**: 1–9. doi: 10.1016/j.aquabot.2018.09.002
- 785 **Pessarrodona A, Moore PJ, Sayer MDJ, Smale DA. 2018a.** Carbon assimilation and
 786 transfer through kelp forests in the NE Atlantic is diminished under a warmer ocean climate.
 787 *Global Change Biology* **24**: 4386–4398. doi: 10.1111/gcb.14303 doi: 10.1111/1365-
 788 2745.13053
- 789 **Pessarrodona A, Foggo A, Smale DA. 2018b.** Can ecosystem functioning be maintained
 790 despite climate-driven shifts in species composition? Insights from novel marine forests.
 791 *Journal of Ecology* **107**: 91–104.
- 792 **Philippart CJM, Anadón R, Danovaro R, et al. 2011.** Impacts of climate change on
 793 European marine ecosystems: Observations, expectations and indicators. *Journal of*
 794 *Experimental Marine Biology and Ecology* **400**: 52–69. doi: 10.1016/j.jembe.2011.02.023
- 795 **Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team. 2019.** Package 'nlme': Linear and
 796 Nonlinear Mixed Effects Models_. R package version 3.1-140, URL: [https://CRAN.R-](https://CRAN.R-project.org/package=nlme)
 797 [project.org/package=nlme](https://CRAN.R-project.org/package=nlme).
- 798 **Polis GA, Anderson WB, Holt RD. 1997.** Toward an integration of landscape and food
 799 webs ecology: the dynamics of spatially subsidized food webs. *Annual Review of Ecology*
 800 *and Systematics* **28**: 289–316. doi: 10.1146/annurev.ecolsys.28.1.289
- 801 **Queirós AM, Stephens N, Widdicombe S, et al. 2019.** Connected macroalgal-sediment
 802 systems: blue carbon and food webs in the deep coastal ocean. *Ecological Monographs*. doi:
 803 10.1002/ecm.1366

- 804 **Ramirez-Llodra E, Rinde E, Gundersen H, et al. 2016.** A snap shot of the short-term
 805 response of crustaceans to macrophyte detritus in the deep Oslofjord. *Scientific Reports* **6**: 1–
 806 5. doi: 10.1038/srep23800
- 807 **Renaud PE, Løkken TS, Jørgensen LL, Berge J, Johnson BJ. 2015.** Macroalgal detritus
 808 and food-web subsidies along an Arctic fjord depth-gradient. *Frontiers in Marine Science* **2**:
 809 1–15. doi: 10.3389/fmars.2015.00031
- 810 **Ruz CS, Muth AF, Tala F, Pérez-Matus A. 2018.** The herbivorous fish, *Aplodactylus*
 811 *punctatus*, as a potential facilitator of dispersal of kelp, *Lessonia trabeculata*, in Chile.
 812 *Journal of Experimental Marine Biology and Ecology* **500**: 112–119. doi:
 813 10.1016/j.jembe.2017.12.007
- 814 **Saulquin B, Hamdi A, Gohin F, Populus J, Mangin A. 2013.** Archimer Estimation of the
 815 diffuse attenuation coefficient K dPAR using MERIS and application to seabed habitat
 816 mapping. *Remote Sensing of Environment* **128**: 224–233. doi: 10.1016/j.rse.2012.10.002
- 817 **Smale DA, Moore P. 2017.** Variability in kelp forest structure along a latitudinal gradient in
 818 ocean temperature. *Journal of Experimental Marine Biology and Ecology* **486**: 255–264. doi:
 819 10.1016/j.jembe.2016.10.023
- 820 **Smale DA, Wernberg T, Yunnies A, Vance T. 2015.** The rise of *Laminaria ochroleuca* in
 821 the Western English Channel (UK) and comparisons with its competitor and assemblage
 822 dominant *Laminaria hyperborea*. *Marine Ecology* **36**: 1033–1044. doi: 10.1111/maec.12199
- 823 **Sorte CJB, Williams SL, Carlton JT. 2010.** Marine range shifts and species introductions:
 824 Comparative spread rates and community impacts. *Global Ecology and Biogeography* **19**:
 825 303–316. doi: 10.1111/j.1466-8238.2009.00519.x
- 826 **Sosik EA, Simenstad CA. 2013.** Isotopic evidence and consequences of the role of microbes
 827 in macroalgae detritus-based food webs. *Marine Ecology Progress Series* **494**: 107–119. doi:
 828 10.3354/meps10544

- 829 **Steneck R, Graham B J Bourque D Corbett J M Erlandson MH, Graham MH, et al.**
 830 **2002.** Kelp forest ecosystems: biodiversity, stability, resilience and future. *Environmental*
 831 *Conservation* **29**: 436–459. doi: 10.1017/S0376892902000322
- 832 **Straub S, Thomsen MS, Wernberg T. 2016.** The dynamic biogeography of the
 833 Anthropocene: The speed of recent range shifts in seaweeds. In: Fraser C, Hu Z. *Seaweed*
 834 *Phylogeography: Adaptation and Evolution of Seaweeds under Environmental Change.*
 835 Dordrecht:Springer, 63-69. doi: 10.1007/978-94-017-7534-2
- 836 **Swanson AK, Fox, CH. 2007.** Altered kelp (Laminariales) phlorotannins and growth under
 837 elevated carbon dioxide and ultraviolet-B treatments can influence associated intertidal food
 838 webs. *Global Change Biology*, **13**: 1696–1709. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2486.2007.01384.x)
 839 [2486.2007.01384.x](https://doi.org/10.1111/j.1365-2486.2007.01384.x)
- 840 **Teagle H, Hawkins SJ, Moore PJ, Smale DA. 2017.** The role of kelp species as biogenic
 841 habitat formers in coastal marine ecosystems. *Journal of Experimental Marine Biology and*
 842 *Ecology* **492**: 81–98. doi: 10.1016/j.jembe.2017.01.017
- 843 **Teagle H, Smale DA. 2018.** Climate-driven substitution of habitat-forming species leads to
 844 reduced biodiversity within a temperate marine community. *Diversity and Distributions* **24**:
 845 1367–1380. doi: 10.1111/ddi.12775
- 846 **Vilas D, Coll M, Pedersen T, et al. 2020.** Kelp-carbon uptake by Arctic deep-sea food webs
 847 plays a noticeable role in maintaining ecosystem structural and functional traits. *Journal of*
 848 *Marine Systems* **203**: 103268. doi: 10.1016/j.jmarsys.2019.103268
- 849 **Wiens JJ. 2016.** Climate-related local extinctions are already widespread among plant and
 850 animal species. *PLoS Biology* **14**: 1–18. doi: 10.1371/journal.pbio.2001104
- 851 **Williams PJB, Giorgio PA, King GM, et al. 2004.** *Respiration in aquatic ecosystems.*
 852 Oxford University Press, USA.
- 853 **Yang LH, Bastow JL, Spence KO, Wright AN. 2008.** What can we learn from resource

854 pulses? *Ecology* **89**: 621–634. doi: 10.1890/07-0175.1

855

Journal Pre-proof

Sustained productivity and respiration of degrading kelp detritus in the shallow benthos: detached or broken, but not dead.

Nadia Frontier^{a, b†*}, Florian de Bettignies^{a†}, Andy Foggo^b, Dominique Davoult^a

^a Sorbonne Université, CNRS, UMR 7144 AD2M, Station Biologique de Roscoff, Place

Georges Teissier, F-29680 Roscoff, France

Marine Environmental Research Highlights-

- The term ‘detritus’ should be revised when referring to kelp material
- Detached kelp material continues to photosynthesise and fix carbon for months
- Light availability strongly influences productivity and photosynthetic performance
- Material from warm-water and cold-water kelp species degrade at different rates
- The cycling of organic material will likely be modified in a changing climate

Declaration of interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper. **(TICK)**

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: