Triathlon and Ultra-Endurance Events in Tropical Environments

Clovis Chabert, Eric Hermand, Olivier Hue

► To cite this version:

Clovis Chabert, Eric Hermand, Olivier Hue. Triathlon and Ultra-Endurance Events in Tropical Environments. Heat Stress in Sport and Exercise, Springer International Publishing, pp.283-296, 2019, 10.1007/978-3-319-93515-7_15 . hal-03187448

HAL Id: hal-03187448

<https://hal.science/hal-03187448>

Submitted on 1 Apr 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

15. Triathlon and ultra-endurance events in tropical environments

Chabert C[†], Hermand E[†], Hue O.

Laboratory « Adaptations au Climat Tropical, Exercice et Santé » (ACTES; EA 3596), French West Indies University, Pointe-à-Pitre, Guadeloupe, France.

[†]: Both authors contributed equally to this work.

Abstract

Physical performance in tropical environments, which combine heat and high humidity, is a challenge that requires specific preparation. The high humidity of a tropical climate alters thermoregulatory capacity by limiting the rate of sweat evaporation. Proper management of whole-body temperature is thus essential to complete an endurance event like a long-distance triathlon or an ultramarathon in such an environment. In triathlon and ultra-endurance races, which can last from 8 to 20 h, performance in tropical settings is closely linked to the capacity to maintain hydration status. Indeed, the rate of withdrawal in these longer events has been associated with water intake, with many finishers showing alterations in electrolyte (e.g. sodium) balance. To counterbalance the impact of a tropical climate and maintain performance, several countermeasures can be adopted, such as using hydration and cooling strategies, and heat acclimation.

1 Influence of heat on performance and health

1.1 Specificity of the triathlon

The Ironman and Challenge triathlon series are trademark brands for long-distance triathlons consisting of a 3.8-km swim, 180-km cycling leg, and full marathon (42.195 km). The best

professional triathletes complete the distance in under 7 h and 50 min, whereas age group athletes vary from 9 to 13 h and more, making this one of the longest endurance races. The duration depends on several factors, including terrain (i.e. course route and geography) and environmental conditions. For example, hot and humid climates impact on performance and the rate of withdrawal, as observed in the Kona (Hawaii) Ironman World Championships every October (Giambelluca et al. 2014; Stiefel et al. 2013): top age-groupers were slower at Hawaii than at their qualifier races (Stiefel et al. 2013), and abandon rate can reach 10%, a high proportion considering the fitness level of athletes participating in this particular event, more resistant to heat stress (Cheung and McLellan 1998; Ironman 2018b). Similarly, more than 25% of the athletes did not reach the finish line at the inaugural Ironman Vichy in 2015, which was held in unusually hot conditions (31~35°C), in contrast to North American races with a minimum 95% finish rate (Britt 2011).

The following sections detail how thermal stress can specifically impact each of the triathlon legs: swimming, cycling and running.

1.1.1 Swimming

Most long distance triathlons allow neoprene wetsuits to be worn for the swim leg as long as water temperature does not exceed 24.5°C. Between 24.5°C and 28.8°C, athletes who choose to wear a wetsuit are allowed to participate, but are not eligible for age-group awards or for qualifying slots for the Ironman World Championships, and over 28.8°C wetsuits are prohibited (Ironman 2018). With its enhanced buoyancy, the neoprene wetsuit reduces ventilation and $\dot{V}O_2$ consumption at a given swimming speed, and therefore metabolic heat production (Trappe et al. 1996). However, wetsuits reduce heat dissipation capacity and can lead to high core body temperatures and dehydration (Kerr et al. 1998; Laursen et al. 2006). Hence, although banning wetsuits in warm water might impair swimming performance, it might also decrease the risk of hyperthermia.

Interestingly, Kerr et al. (Kerr et al. 1998) showed in a simulated Olympic triathlon that core temperature did not change by wearing a wetsuit, thought to limit thermoregulatory mechanisms

49 through heat-insulating properties of neoprene, as excess heat was still transferred at the periphery,
50 leading to a higher skin temperature. A subsequent field study confirmed that core temperature
51 remained around 38°C in well-trained athletes wearing a wetsuit in 20.5°C water during a 3.8 km
52 swim (Laursen et al. 2006). This was again confirmed by a case report in which an athlete's core
53 temperature was continuously monitored during an Ironman (Laursen et al. 2009). In 29.5°C water,
54 another study showed that the core temperature of moderately trained athletes increased by an
55 average 0.7°C and remained under 38°C during a non-wetsuit half-Ironman (Figure 1; (Baillot and
56 Hue 2015)). Thus, no study appears to have shown a dangerous elevation in core temperature during
57 the swim leg of long-distance triathlon events. Moreover, while 13 of the 14 deaths in triathlon
58 races from 2006 to 2008 occurred during the swim leg, pre-existing cardiovascular abnormalities
59 seem to have been the major factor, ruling out hyperthermia as a potential cause (Harris et al. 2010).

60 1.1.2 Cycling

61 When cycling, some of the heat produced by the muscles is dissipated via convection. This
62 however, depends on a sufficient temperature gradient between the skin and ambient air.
63 Convection is therefore limited in hot conditions when air temperatures exceeds 35°C. Heat is also
64 dissipated during cycling via the evaporation of sweat, which depends on the water vapor partial
65 pressure gradient between air and skin. Like convection, this mechanism is reduced in humid
66 environments, making performance in a tropical environment challenging as thermoregulatory
67 efficiency is reduced (Nybo 2010). In a tropical climate (31-33°C, 70-75% RH), one-hour of
68 pedalling at a submaximal intensity increases core temperature, heart rate, sweat rate and water loss
69 (Saat et al. 2005a; Saat et al. 2005b; Voltaire et al. 2003). These results can be extrapolated to
70 describe the physiological impact of 4–6-h events, especially if the cycling course includes hills,
71 where the diminished air speed reduces heat dissipation (Nybo 2010). Laboratory-controlled studies
72 showed that increasing ambient temperature impaired cycling capacity when relative humidity was
73 clamped at 70% (Galloway and Maughan 1997; Maughan et al. 2012). During a 9-day (2.5 h/day)

74 cycling race held in Guadeloupe (31.1°C, 75.6% RH), tympanic temperatures measured
75 immediately after the stages never exceeded 2°C of resting pre-race values (Hue et al. 2006). In
76 contrast to the variability associated with stage racing in a peloton, triathletes will adopt a much
77 steadier pace during racing in some cases to “save their legs” for the following marathon (Wu et al.
78 2015). Interestingly, Baillot et al. (Baillot and Hue 2015) showed a negative correlation between
79 body mass and the evolution of core temperature throughout the cycling leg, implying greater heat
80 storage and inertia. A bigger body thus presents a larger surface to preserve heat exchange at high
81 velocity, even in a hot and humid environment (Nielsen 1996).

82 In summary, the high velocities and individual nature of cycling in the triathlon support heat
83 dissipation by convection on the condition that air temperature is lower than skin temperature. In a
84 thermally challenging environment, sweat evaporation is the main avenue for heat dissipation but it
85 is limited by the high humidity of a tropical environment. Nevertheless, triathletes appear to show
86 adequate thermoregulatory capability to prevent excessive hyperthermia, mainly through intensity
87 management and pacing strategies (Baillot and Hue 2015).

88 1.1.3 Running

89 Of the three sports in the triathlon, running may carry the greatest risk of hyperthermia for
90 triathletes competing in a tropical climate. Several factors come into play. Firstly, although running
91 is faster than swimming, air possesses neither the heat capacity nor transfer celerity of water, hence
92 limiting the quantity of heat energy transferred from the body. Secondly, unlike in cycling, the
93 lower speed associated with running does not allow for excess heat dissipation via convection
94 (Nielsen 1996). These factors can have a strong negative impact on running performance (Kenefick
95 et al. 2007), particularly the marathon (Ely et al. 2007). This negative impact is amplified (i) by a
96 heavier body mass, which, at a given pace, requires more energy for running propulsion and
97 therefore produces more heat, and (ii) in heavier athletes running in a hot/humid environment

98 limiting heat exchange by convection and evaporation, who display greater imbalance between heat
99 production and dissipation, even at slower paces (Marino et al. 2000).

100 While the core temperature data of professional triathletes in the Hawaii Ironman or other
101 tropical locations are currently not available, a few studies of moderately to very well trained age
102 group athletes have shown that none of them suffered from heat-related illness and excessive
103 increases in core temperature (Laursen et al. 2006) (Figure 1; (Baillot and Hue 2015; Del Coso et al.
104 2014)). In the report from Laursen et al. (Laursen et al. 2006) 7 out of 10 well trained triathletes
105 performed a high-level sub-10-hour performance at Ironman Western Australia in temperate but
106 humid conditions (23.3°C, 60% RH), and the average core body temperature remained close to
107 38°C. Despite a much hotter and more humid environment in Guadeloupe (27.2°C, 80% RH), the
108 average body temperature of moderately trained athletes during a half-Ironman run was 38.2°C and
109 none of them reached 40°C (Baillot and Hue 2015). These values are not surprising given the
110 relative intensity and absolute workload at which most long-distance triathlon races are performed,
111 which is in contrast to shorter events performed at a higher intensity and thus higher heat production
112 (Siegel et al. 2010; Baillot and Hue 2015). The importance of exercise intensity was confirmed in
113 dryer conditions in a study showing that the fastest athletes exhibited the highest core temperatures
114 (Del Coso et al. 2014).

115

116 1.2 Specificity of ultra-endurance events

117 1.2.1 Ultra-endurance in the heat

118 Ultramarathons last more than 6 h (Zaryski and Smith 2005). In the past decade, trail running
119 races have become a major ultra-endurance sport. Since the first edition of the Ultra Trail du Mont
120 Blanc (160 km and +10,000 m of elevation), the number of finishers has grown from 67 in 2003 to
121 1685 in 2017. This kind of effort involves several physiological and psychological parameters that
122 are modulated by environmental factors. Ultramarathon races are performed with a variety of off-

123 road terrains, elevation profiles (positive: D+ and negative: D- changes), and distances, all of which
124 greatly complicate the attempt to understand how environmental conditions affect performance.
125 Marathon races have more standardized characteristics, facilitating the comparison of performances
126 in order to study the role played by heat (see chapter 12). Based on the analysis of winners' times in
127 several marathons, (Maughan 2010) et al., 2010 determined that the best running temperature is
128 between 10°C to 12°C for elite athletes. Beyond this temperature, the analysis of 136 marathons
129 performed between a wet-bulb globe temperature (WBGT) of 5 and 25°C showed a decrease in
130 performance with an increase in WBGT (Figure 2; (Ely et al. 2007)). Interestingly, these authors
131 also showed that slower runners, who are exposed to heat for a longer time than faster runners, are
132 more affected by temperature. For ultramarathon distances, the same effect of hot environments on
133 performance was observed in a study that examined running times in the Western States Endurance
134 Run (161 km and 6000 m D+) in 2006 and 2007 (Parise and Hoffman 2011). In these two editions
135 (Figure 3), the authors followed 50 runners who had finished the race in both years to investigate
136 the effect of temperature changes from 7–38°C in 2006 to 2–30°C in 2007. Independently of their
137 level, the athletes were 8% slower in the hot condition (2006) than in the cooler environment
138 (2007), and the withdrawal rate was 14% higher in 2006. In this study, slower runners did not
139 appear to be more impacted by the hot environment, but this observation was probably due to the
140 longer time running at a comfortable temperature (i.e. night).

141 1.2.2 Hydration requirements during ultra-endurance exercise in the heat

142 During muscle contraction, only ~25% of the substrate energy is converted to mechanical
143 work, with the other ~75% released as heat. Effective thermoregulation during exercise depends on
144 the balance between the absolute mechanical work generated by the athlete and the heat loss process
145 (Chapter 2). Ultramarathons, characterized by the maintenance of a low relative exercise intensity
146 over very long periods, do not significantly challenge the heat balance in temperate environments.
147 However, maintaining mechanical load over a longer time may challenge heat balance in athletes,

148 especially when competing in hot and/or humid environments. Furthermore, environmental heat
149 stress (i.e. temperature) reduces dry heat loss, due to the small (or even negative) temperature
150 gradient between the skin and air, which progressively places sweat evaporation as the only
151 mechanism available for the organism to dissipate endogenous heat (Brotherhood 2008; Taylor
152 2006). Indeed, at in 30°C conditions a dry heat loss of 75 W is observed whereas at 40°C, it induces
153 a dry heat gain of 75 W, increasing the sweating rate require to maintain body core temperature
154 (Gagnon et al. 2013). Thus, ultramarathons performed in a hot environment require bigger volumes
155 of water intake to compensate for increased water losses. During the Badwater Ultramarathon of
156 2012 (217 km and 4000 m D+), the temperature oscillated from 10.1°C at night to 46.6°C during
157 the day. The follow-up of 4 runners who completed the race in 36 ± 3 h showed a mean water intake
158 of 34 ± 13 L (Brown and Connolly 2015). On average, each athlete drank $0.93 \text{ L}\cdot\text{h}^{-1}$, whereas on
159 completion of the Biel Ultramarathon in Switzerland (100 km and 645 m D+), in a temperate
160 environment varying from 8 to 28°C, the runners consumed $0.65 \text{ L}\cdot\text{h}^{-1}$ (Knechtle et al. 2011). When
161 the distances of these two races were linked to elevation changes (100 m D+ corresponds to 1 km of
162 distance), the corrected distance for the Biel Ultramarathon was 106.45 km and 257 km for
163 Badwater. According to these corrected distances, the Badwater runners drank almost twice as
164 much water as the Biel runners ($0.13 \text{ L}\cdot\text{km}^{-1}$ and $0.069 \text{ L}\cdot\text{km}^{-1}$, respectively), despite the higher
165 mechanical workload developed during the Biel Ultramarathon due to its shorter distance and lower
166 elevation change ($6.6 \text{ min}\cdot\text{km}^{-1}$ vs. $8.5 \text{ min}\cdot\text{km}^{-1}$ for Badwater). The characteristics of the 2007
167 Peninsula Ultra Fun Run (PUFfeR) of South Africa (80 km and 1000 m D+) were close to those of
168 the Biel Ultramarathon, but the participants were subjected to lower environmental temperatures
169 (8–20°C). The runners drank less during this more temperate race than during the Biel
170 Ultramarathon ($0.028 \text{ L}\cdot\text{km}^{-1}$ and $0.069 \text{ L}\cdot\text{km}^{-1}$, respectively) (Tam et al. 2009). Analysis of the
171 hydration status of 16 ultramarathon runners (161 km, 7000 m D+) in a hot environment
172 (4.8–37.8°C) showed that finishers (n=6) had drunk significantly more water at the 48th kilometer
173 than non-finishers (n=10) (Stuempfle et al. 2011). However, the authors were unable to determine

174 whether beverage intake was directly linked to finishing capacity or if it was due to other factors,
175 such as the experience or endurance capacity of the runners. A second study performed during a
176 160-km foot race (positive elevation not known) in temperatures that peaked at 38°C followed only
177 those runners who had completed at least one previous ultramarathon among the top 50% (Glance et
178 al. 2002). The authors showed that, despite the homogenous level and experience of these runners,
179 non-finishers also drank significantly less than finishers (-35%, $p < 0.01$). All these data confirmed
180 the relationship between event temperature and water intake that is crucial to health and
181 performance. A study during the Gwada Run of 2011, a 6-day multi-stage race for a total of 142 km
182 under tropical conditions ($30 \pm 2.4^\circ\text{C}$ and $82 \pm 4\%$ RH) showed a sweat loss of 0.19 L.km^{-1} (Hue et al.
183 2014). Yet water intake ($1.5 \pm 0.3 \text{ L}$ per stage) was probably distorted by the shortness of the stages
184 (from 16 to 21 km), which enabled athletes to tolerate transient dehydration during the races (-
185 $4.2 \pm 0.9 \text{ L}$ per stage).

186

187 2 Countermeasures to optimize performance and health

188 2.1 Hydration

189 Limiting dehydration during triathlons and ultramarathons in hot environments seems to be
190 essential to maintain the exercise workload over many hours. The role of hydration in a tropical
191 climate (see chapter 6) is critical since it has been shown that hyperthermia and dehydration are
192 worse than hyperthermia alone (Sawka et al. 2001; Sawka and Noakes 2007). Despite contradictory
193 observations on influencing thermoregulation and core temperature (Latzka et al. 1997; Ross et al.
194 2012), it seems that hyperhydration can delay the development of dehydration (Latzka and Sawka
195 2000; Latzka et al. 1997; Sawka et al. 1984). However, hyperhydration does not seem to lead to
196 better performances during a laboratory-based 46 km cycling time trial by elite cyclists (Ross et al.
197 2012) or a 60 min run in endurance trained runners (Scheidler 2009). As hyperhydration results in

198 non-negligible added body weight, it may not be an advantage for running performance, which is
199 greatly affected by extra weight (Teunissen et al. 2007). In addition, it does not enhance
200 thermoregulation in a hot and humid climate and augments the risk of hyponatremia (Noakes et al.
201 2005).

202 In the triathlon, although hydration depends very much on the availability of adequate means
203 set up by the race organization, the descriptive papers to date report that no triathlete has suffered
204 from dehydration symptoms in hot and humid events, despite occasional significant water losses
205 (Baillot and Hue 2015; Del Coso et al. 2014; Laursen et al. 2006; Speedy et al. 2001). This suggests
206 that the hydration strategies employed by the athletes meet the body's water requirements (Kenefick
207 2018).

208 Maintaining optimal hydration is more complicated than it would seem in the ultramarathon.
209 Indeed, analysis of the hydration status of ultramarathon runners in 2 different races showed that
210 finishers had drunk significantly more water than non-finishers, even if the runners' experience was
211 taking into account (Glance et al. 2002; Stuenkel et al. 2011). In contrast, slower runners became
212 over-hydrated because they feared dehydration, which may lead to hyponatremia (Hew-Butler et al.
213 2008; Noakes 2007). Hyponatremia however, is not only due to over-hydration, but may be the
214 consequence of insufficient sodium intake during the race (Glance et al. 2002; Stuenkel et al. 2011).
215 In a hot environment, hyponatremia affected 30–50% of ultramarathon finishers, reflecting the
216 inadequate or incorrect hydration strategies of many runners (Glance et al. 2002; Hoffman et al.
217 2012; Lebus et al. 2010). Thus, better sodium intake during ultramarathons could contribute to
218 better performances by stimulating thirst, increasing voluntary fluid intake, enhancing intestinal
219 glucose and water absorption, optimizing extracellular and intracellular fluid balance, and
220 potentially mitigating the occurrence of clinically significant episodes of hyponatremia (Maughan
221 1991; Shirreffs and Sawka 2011; Speedy et al. 1999; Vrijens and Rehrer 1999).

222

2.2 Hydration policy and implementation

In a triathlon field study, oral salt supplementation improved half-Ironman performance through faster cycling ($p < 0.05$) and showed a similar trend in the running leg ($p = 0.06$), with reduced sweat rate and limited electrolyte deficit (Del Coso et al. 2016). The consensus recommendation is thus to ingest 0.5 to 0.7 g.L⁻¹.h⁻¹ for long endurance races (Von Duvillard et al. 2004) and up to 1.5 g.L⁻¹.h⁻¹ for athletes prone to develop muscle cramping (Bergeron 2003). However, adding salt to the consumed water is not sufficient to avoid hyponatremia if athletes over-drink during a race (73% of severe symptomatic hyponatremia found after an Ironman (Speedy et al. 1999)).

(Hoffman and Stuenkel 2014) observed no advantage to sodium-enriched beverages during a 161-km ultramarathon performed by all levels of athletes in heat (38°C). This result may have been due to a variation in sodium intake from solid food and/or to the tolerance of faster runners to hypohydration and hyperthermia (Baillot et al. 2014; Hue et al. 2014).

236

2.3 Cooling

Cooling strategies such as cold drink ingestion or cold-water immersion to reduce the thermal load may enhance performance in the heat (see chapter 7). This effect has been manifested in cyclists (Figure 4; (Burdon et al. 2013; Riera et al. 2016; Tran Trong et al. 2015) and runners as a longer time to exhaustion (Siegel et al. 2010; Siegel et al. 2012; Yeo et al. 2012). Wearing cooling garments prior to or during exercise has proven to be performance-effective in hot and humid climates (Arngrimsson et al. 2004; Hasegawa et al. 2005; Luomala et al. 2012), but they are difficult, impractical and potentially not allowed to be used during official long-distance events.

Other studies have focused on cooling strategies, but the exercise duration has been relatively short compared with ultra-endurance events. Nonetheless, they show interesting results that may be

247 extrapolated for long to ultra-long events. Spraying or pouring water over the face and/or body can
248 improve performance in tropical conditions. For example, pouring cold water over the skin will
249 reduce skin temperature before dripping off the body, and transiently improve thermal comfort.
250 Cooling the head in this manner resulted in a 51% increase in cycling time to exhaustion at 75%
251 $\text{VO}_{2\text{max}}$ (Ansley et al. 2008), with similar effects recently observed in running (Stevens et al.
252 2017b). Neck cooling during a 90-min running trial in a hot environment (30.4°C and 53% RH)
253 increased the distance covered by 7.4% with no change in rectal temperature (Tyler and Sunderland
254 2011). A similar effect was found with menthol ingestion, as it activates cold dermal sensors
255 (Stevens and Best 2017). Mixed into a cold beverage, menthol did not lower core body temperature
256 but had a positive effect on thermal sensation and running/cycling performances over various
257 distances in tropical climate (Stevens et al. 2016; Tran Trong et al. 2015). Indeed, Stevens et al.
258 2016 showed that a menthol mouth rinse every kilometre (25 mL at a concentration of 0.01%)
259 during a running time trial in the heat significantly improved 5-km performance time by 3%. A
260 cumulative effect of menthol and ice slurry or cold water was observed on performances during a
261 20-km cycling trial in hot environment (30.7±0.8°C and 78±0.03% RH) (Tran Trong et al. 2015).
262 Ultimately, menthol mouth rinsing and ingesting influence thermal perception and thermal comfort,
263 which in turn might contribute to enhance performance in hot climates (Stevens et al. 2017a;
264 Stevens et al. 2016).

265

266 2.4 Cooling policy and implementation

267 According to the literature, pre-cooling by ice or cold water has been successfully studied
268 during short duration exercise (<60 min) that is too brief to be a key factor of performance during
269 ultra-endurance trials (Siegel et al. 2012; Stevens et al. 2017b). Conversely, per-cooling by cold
270 water or ice slurry ingestion during exercise seems to be a very interesting intervention to improve
271 performances in a hot environment and possibly increase fluid intake, particularly for well-trained

athletes (see chapter 7). Indeed, most high-performing runners show the biggest increase in internal temperature, probably due to higher thermogenesis consecutive to higher workloads (Baillot et al. 2014; Hue et al. 2014). Faster trail runners (27 km; Baillot et al. 2014) and multi-stages trail runners (127 km on 6 days; Hue et al. 2014) also present greater post-race dehydration, which could be reduced by cold water due to its effect on voluntarily increasing water absorption during exercise. The higher temperatures and greater dehydration observed in faster trail runners also seem to indicate that performance in heat is associated with a better tolerance to hyperthermia in elite athletes (McKeag and Moeller 2007).

280

2.5 Heat acclimation

Heat acclimation (HA) before a race is an efficient way to increase performance in a hot environment (Armstrong and Maresh 1991; Nielsen 1994) (see chapter 8 for further details). The physiological adaptations of HA lead to better cardiovascular function (output, stroke volume, heart rate) and a decreased core temperature at rest and during exercise, in part due to an enhanced sweat rate and expanded plasma volume (Armstrong and Maresh 1991). HA is also known to reduce sodium loss by sweating (Kirby and Convertino 1986). However, HA does not fully restore long to ultra-long endurance performance to the level reached by athletes in temperate conditions (Voltaire et al. 2002).

If early arrival at the race location is not possible, HA in an environmental chamber is a strategy that may help preserve ultramarathon performance. (Costa et al. 2014) examined the effects of six 2-h sessions of running at 60% of VO_{2max} on a motorized treadmill in an environmental heat chamber at 30°C or 35°C. From the third session onward, mean heart rate was lower at 30°C, whereas mean heart rate and thermal comfort were lower at 35°C. The authors concluded that two bouts of running at 60% of VO_{2max} in a 30°C air temperature conditions were sufficient to induce heat acclimation in ultra-runners, which may enhance their performance in such environments.

297

298

299 3 Conclusions

300 The maintenance of triathlon or ultramarathon performance in a hot environment is a
301 considerable challenge which requires meticulous preparation and management during the race. To
302 limit the performance decrements induced by these harsh environmental conditions, several
303 countermeasures have been proposed, such as cold-water or ice slurry ingestion, external pre- and
304 per- cooling and/or menthol use. However, the limited duration of exercise performed in these
305 studies does not allow for direct application of the results to an ultra-endurance or ultra-triathlon
306 context. Thus, further studies are required to validate these approaches in long to triathlon and ultra-
307 endurance events.

308 Currently, one of the most relevant strategies to employ when performing in the heat is to
309 consume adequate fluids. Maintaining water availability along different race courses is thus crucial
310 due to the detrimental effects dehydration has on fluid balance and consequently heat loss via
311 sweating. Individual sweat rates are highly dependent on the athlete's morphology, the intensity of
312 exercise and the environment temperature and humidity. Sweat losses are accompanied by
313 electrolyte losses, particularly sodium, which may contribute to hyponatremia if not compensated.
314 To limit this risk, exogenous intake by enriched sodium beverages is the most effective strategy. In
315 addition to the traditional compounds found in most sport drinks (e.g. glucose, magnesium,), it
316 seems that ultramarathon beverages should contain about $0.7\text{--}1.2\text{ g.L}^{-1}$ of salt when conducted in
317 the heat. However, sodium-enriched beverages are insufficient if athletes overdrink by fear of
318 dehydration during the race, placing management of water intake as a key determinant of
319 performance.

320 Heat acclimation is also a relevant strategy that may enhance ultra-endurance and triathlon
321 performance in hot environments. Most of the acclimation benefits occur in the first 14 days of
322 exposure, but a short-term protocol of 5 days will induce significant early adaptations.

323

Figure 1. Individual and mean values ($\pm$ SD) of core temperature at each stage of the Guadeloupe half-Ironman held in tropical climate. T1: just before the race; T2: after the swim phase; T3: after the cycle phase; T4: at the end of the run phase (Baillot and Hue 2015). Temperatures were obtained from telemetric intestinal temperature devices ingested at least 6 hours before the race.

327

328 **Figure 2.** Nomogram representing the impact of WBGT on the relation between marathon finishing times
329 according to quartiles and relative performance decrement in comparison with WBGT at 5°C (Ely et al.
330 2007).

331

Figure 3. Proportion of runners abandoning the Western States Endurance Run (161 km; 6000m D+) race at each checkpoint along the course (A), and relationship between finish times for the 2006 (7-38°C) and 2007 (2-30°C) edition (B). (Parise and Hoffman 2011).

Figure 4. Trial times for 5 successive blocks (4 km cycling + 1.5 km running) with the ingestion of Neutral water (orange), Cold water (blue) and Ice-slurry (green).

^a Significantly different from Neutral water Cycling and Running Performance Using Ice-Slurry/Menthol (P<0.05).

^b Significantly different from Cold (P<0.05). Mean values and SD are shown.

β,† denote that block performance was affected by Time Period (P<0.007) and the Time Period x Drink Temperature interaction (P<0.004), respectively (Tran Trong et al. 2015).

337

338

339 4 Bibliography

340 Ansley L, Marvin G, Sharma A, Kendall MJ, Jones DA, Bridge MW (2008). The effects of head
341 cooling on endurance and neuroendocrine responses to exercise in warm conditions. *Physiol*
342 *Res.* 57:863-872

343 Armstrong LE, Maresh CM (1991). The induction and decay of heat acclimatisation in trained
344 athletes. *Sports Med.* 12:302-312

345 Arngrimsson SA, Petitt DS, Stueck MG, Jorgensen DK, Cureton KJ (2004). Cooling vest worn
346 during active warm-up improves 5-km run performance in the heat. *J Appl Physiol* (1985).
347 96:1867-1874

348 Baillot M, Hue O (2015). Hydration and thermoregulation during a half-ironman performed in
349 tropical climate. *J Sports Sci Med.* 14:263-268

350 Baillot M, Le Bris S, Hue O (2014). Fluid replacement strategy during a 27-Km trail run in hot and
351 humid conditions. *Int J Sports Med.* 35:147-152

352 Bergeron MF (2003). Heat cramps: fluid and electrolyte challenges during tennis in the heat. *J Sci*
353 *Med Sport.* 6:19-27

354 Britt R (2011) North American Ironman DNF Rates: Finishers and DNF by Race.
355 <http://www.runtri.com/2010/09/north-american-ironman-dnf-rates.html>.

356 Brotherhood JR (2008). Heat stress and strain in exercise and sport. *J Sci Med Sport.* 11:6-19

357 Brown JS, Connolly D (2015). Food and Fluid Intake During Extreme Heat: Experiences From The
358 Badwater Ultramarathon. *Wilderness & Environmental Medicine.* 26:e4

359 Burdon CA, Hoon MW, Johnson NA, Chapman PG, O'Connor HT (2013). The effect of ice slushy
360 ingestion and mouthwash on thermoregulation and endurance performance in the heat. *Int J*
361 *Sport Nutr Exerc Metab.* 23:458-469

362 Costa RJ, Crockford MJ, Moore JP, Walsh NP (2014). Heat acclimation responses of an ultra-
363 endurance running group preparing for hot desert-based competition. *Eur J Sport Sci.* 14
364 Suppl 1:S131-141

365 Del Coso J et al. (2016). Effects of oral salt supplementation on physical performance during a half-
366 ironman: A randomized controlled trial. *Scand J Med Sci Sports.* 26:156-164

367 Del Coso J et al. (2014). Relationship between physiological parameters and performance during a
368 half-ironman triathlon in the heat. *J Sports Sci.* 32:1680-1687

369 Ely MR, Cheuvront SN, Roberts WO, Montain SJ (2007). Impact of weather on marathon-running
370 performance. *Med Sci Sports Exerc.* 39:487-493

371 Gagnon D, Jay O, Kenny GP (2013). The evaporative requirement for heat balance determines
372 whole-body sweat rate during exercise under conditions permitting full evaporation. *J*
373 *Physiol.* 591:2925-2935

374 Galloway SD, Maughan RJ (1997). Effects of ambient temperature on the capacity to perform
375 prolonged cycle exercise in man. *Med Sci Sports Exerc.* 29:1240-1249

376 Giambelluca T et al. (2014). Evapotranspiration of Hawai'i: final report to the US Army Corps of
377 Engineers—Honolulu District and the Commission on Water Resource Management, State
378 of Hawai'i, 178 p.

379 Glace BW, Murphy CA, McHugh MP (2002). Food intake and electrolyte status of
380 ultramarathoners competing in extreme heat. *J Am Coll Nutr.* 21:553-559

381 Harris KM, Henry JT, Rohman E, Haas TS, Maron BJ (2010). Sudden death during the triathlon.
382 *JAMA.* 303:1255-1257

383 Hasegawa H, Takatori T, Komura T, Yamasaki M (2005). Wearing a cooling jacket during exercise
384 reduces thermal strain and improves endurance exercise performance in a warm
385 environment. *J Strength Cond Res.* 19:122-128

386 Hew-Butler T et al. (2008). Statement of the Second International Exercise-Associated
387 Hyponatremia Consensus Development Conference, New Zealand, 2007. Clin J Sport Med :
388 official journal of the Canadian Academy of Sport Medicine. 18:111-121

389 Hoffman MD, Stuenkel KJ (2014). Hydration strategies, weight change and performance in a 161
390 km ultramarathon. Res Sports Med. 22:213-225

391 Hoffman MD, Stuenkel KJ, Rogers IR, Weschler LB, Hew-Butler T (2012). Hyponatremia in the
392 2009 161-km Western States Endurance Run. Int J Sports Physiol Perform. 7:6-10

393 Hue O, Henri S, Baillot M, Sinnapah S, Uzel AP (2014). Thermoregulation, hydration and
394 performance over 6 days of trail running in the tropics. Int J Sports Med. 35:906-911

395 Hue O, Voltaire B, Hertogh C, Blonc S (2006). Heart rate, thermoregulatory and humoral responses
396 during a 9-day cycle race in a hot and humid climate. Int J Sports Med. 27:690-696

397 Ironman (2018) Rules and Regulations.
398 [http://m.eu.ironman.com/triathlon/events/emea/ironman/tallinn/athletes/rules-and-](http://m.eu.ironman.com/triathlon/events/emea/ironman/tallinn/athletes/rules-and-regulations.aspx)
399 [regulations.aspx](http://m.eu.ironman.com/triathlon/events/emea/ironman/tallinn/athletes/rules-and-regulations.aspx).

400 Kenefick RW (2018). Drinking Strategies: Planned Drinking Versus Drinking to Thirst. Sports
401 Med. 48:31-37

402 Kenefick RW, Cheuvront SN, Sawka MN (2007). Thermoregulatory function during the marathon.
403 Sports Med. 37:312-315

404 Kerr CG, Trappe TA, Starling RD, Trappe SW (1998). Hyperthermia during Olympic triathlon:
405 influence of body heat storage during the swimming stage. Med Sci Sports Exerc. 30:99-104

406 Kirby CR, Convertino VA (1986). Plasma aldosterone and sweat sodium concentrations after
407 exercise and heat acclimation. J Appl Physiol (1985). 61:967-970

408 Knechtle B, Knechtle P, Rosemann T (2011). Low prevalence of exercise-associated hyponatremia
409 in male 100 km ultra-marathon runners in Switzerland. Eur J Appl Physiol. 111:1007-1016

410 Latzka WA, Sawka MN (2000). Hyperhydration and glycerol: thermoregulatory effects during
411 exercise in hot climates. Can J Appl Physiol. 25:536-545

412 Latzka WA, Sawka MN, Montain SJ, Skrinar GS, Fielding RA, Matott RP, Pandolf KB (1997).
413 Hyperhydration: thermoregulatory effects during compensable exercise-heat stress. J Appl
414 Physiol (1985). 83:860-866

415 Laursen PB et al. (2006). Core temperature and hydration status during an Ironman triathlon. Br J
416 Sports Med. 40:320-325; discussion 325

417 Laursen PB, Watson G, Abbiss CR, Wall BA, Nosaka K (2009). Hyperthermic fatigue precedes a
418 rapid reduction in serum sodium in an ironman triathlete: a case report. Int J Sports Physiol
419 Perform. 4:533-537

420 Lebus DK, Casazza GA, Hoffman MD, Van Loan MD (2010). Can changes in body mass and total
421 body water accurately predict hyponatremia after a 161-km running race? Clin J Sport Med:
422 official journal of the Canadian Academy of Sport Medicine. 20:193-199

423 Luomala MJ, Oksa J, Salmi JA, Linnamo V, Holmér I, Smolander J, Dugue B (2012). Adding a
424 cooling vest during cycling improves performance in warm and humid conditions. Journal of
425 Thermal Biology. 37:47-55

426 Marino FE, Mbambo Z, Kortekaas E, Wilson G, Lambert MI, Noakes TD, Dennis SC (2000).
427 Advantages of smaller body mass during distance running in warm, humid environments.
428 Pflugers Arch. : European journal of physiology. 441:359-367

429 Maughan RJ (1991). Fluid and electrolyte loss and replacement in exercise. J Sports Sci. 9 Spec
430 No:117-142

431 Maughan RJ (2010). Distance running in hot environments: a thermal challenge to the elite runner.
432 Scand J Med Sci Sports. 20 Suppl 3:95-102

433 Maughan RJ, Otani H, Watson P (2012). Influence of relative humidity on prolonged exercise
434 capacity in a warm environment. Eur J Appl Physiol. 112:2313-2321

435 McKeag D, Moeller JL (2007). ACSM's primary care sports medicine. Wolters Kluwer/Lippincott
436 Williams & Wilkins,

437 Nielsen B (1994). Heat stress and acclimation. *Ergonomics*. 37:49-58
 438 Nielsen B (1996). Olympics in Atlanta: a fight against physics. *Med Sci Sports Exerc*. 28:665-668
 439 Noakes TD (2007). Drinking guidelines for exercise: what evidence is there that athletes should
 440 drink "as much as tolerable", "to replace the weight lost during exercise" or "ad libitum"? *J*
 441 *Sports Sci*. 25:781-796
 442 Noakes TD, Goodwin N, Rayner BL, Branken T, Taylor RK (2005). Water intoxication: a possible
 443 complication during endurance exercise, 1985. *Wilderness Environ Med*. 16:221-227
 444 Nybo L (2010). Cycling in the heat: performance perspectives and cerebral challenges. *Scand J Med*
 445 *Sci Sports*. 20 Suppl 3:71-79
 446 Parise CA, Hoffman MD (2011). Influence of temperature and performance level on pacing a 161
 447 km trail ultramarathon. *Int J Sports Physiol Perform*. 6:243-251
 448 Riera F, Trong T, Rinaldi K, Hue O (2016). Precooling does not enhance the effect on performance
 449 of midcooling with ice-slush/menthol. *Int J Sports Med*. 37:1025-1031
 450 Ross ML, Jeacocke NA, Laursen PB, Martin DT, Abbiss CR, Burke LM (2012). Effects of lowering
 451 body temperature via hyperhydration, with and without glycerol ingestion and practical
 452 precooling on cycling time trial performance in hot and humid conditions. *J Int Soc Sports*
 453 *Nutr*. 9:55
 454 Saat M, Sirisinghe RG, Singh R, Tochihara Y (2005a). Effects of short-term exercise in the heat on
 455 thermoregulation, blood parameters, sweat secretion and sweat composition of tropic-
 456 dwelling subjects. *J Physiol Anthropol Appl Human Sci*. 24:541-549
 457 Saat M, Tochihara Y, Hashiguchi N, Sirisinghe RG, Fujita M, Chou CM (2005b). Effects of
 458 exercise in the heat on thermoregulation of Japanese and Malaysian males. *J Physiol*
 459 *Anthropol Appl Human Sci*. 24:267-275
 460 Sawka MN, Francesconi RP, Young AJ, Pandolf KB (1984). Influence of hydration level and body
 461 fluids on exercise performance in the heat. *JAMA*. 252:1165-1169
 462 Sawka MN, Montain SJ, Latzka WA (2001). Hydration effects on thermoregulation and
 463 performance in the heat. *Comp Biochem Physiol A Mol Integr Physiol*. 128:679-690
 464 Sawka MN, Noakes TD (2007). Does dehydration impair exercise performance? *Med Sci Sports*
 465 *Exerc*. 39:1209
 466 Sheadler CM (2009) Glycerol hyperhydration and endurance running performance in the heat. The
 467 Ohio State University.
 468 Shirreffs SM, Sawka MN (2011). Fluid and electrolyte needs for training, competition, and
 469 recovery. *J Sports Sci*. 29 Suppl 1:S39-46
 470 Siegel R, Mate J, Brearley MB, Watson G, Nosaka K, Laursen PB (2010). Ice slurry ingestion
 471 increases core temperature capacity and running time in the heat. *Med Sci Sports Exerc*.
 472 42:717-725
 473 Siegel R, Mate J, Watson G, Nosaka K, Laursen PB (2012). Pre-cooling with ice slurry ingestion
 474 leads to similar run times to exhaustion in the heat as cold water immersion. *J Sports Sci*.
 475 30:155-165
 476 Speedy DB et al. (2001). Fluid balance during and after an ironman triathlon. *Clin J Sport Med*.
 477 11:44-50
 478 Speedy DB et al. (1999). Hyponatremia in ultradistance triathletes. *Med Sci Sports Exerc*. 31:809-
 479 815
 480 Stevens CJ, Bennett KJ, Sculley DV, Callister R, Taylor L, Dascombe BJ (2017a). A Comparison
 481 of Mixed-Method Cooling Interventions on Preloaded Running Performance in the Heat. *J*
 482 *Strength Cond Res*. 31:620-629
 483 Stevens CJ, Best R (2017). Menthol: A Fresh Ergogenic Aid for Athletic Performance. *Sports Med*.
 484 47:1035-1042
 485 Stevens CJ, Kittel A, Sculley DV, Callister R, Taylor L, Dascombe BJ (2017b). Running
 486 performance in the heat is improved by similar magnitude with pre-exercise cold-water
 487 immersion and mid-exercise facial water spray. *J Sports Sci*. 35:798-805

488 Stevens CJ, Thoseby B, Sculley DV, Callister R, Taylor L, Dascombe BJ (2016). Running
489 performance and thermal sensation in the heat are improved with menthol mouth rinse but
490 not ice slurry ingestion. *Scand J Med Sci Sports*. 26:1209-1216

491 Stiefel M, Knechtle B, Rüst CA, Rosemann T (2013). Analysis of performances at the'Ironman
492 Hawaii triathlon'and its qualifier events with respect to nationality. *Journal of Science and
493 Cycling*. 2:27

494 Stuempfle KJ, Hoffman MD, Weschler LB, Rogers IR, Hew-Butler T (2011). Race diet of finishers
495 and non-finishers in a 100 mile (161 km) mountain footrace. *J Am Coll Nutr*. 30:529-535

496 Tam N, Hew-Butler T, Papadopoulou E, Nolte H, Noakes TD (2009). Fluid intake and changes in
497 blood biochemistry, running speed and body mass during an 80 km mountain trail race. *Med
498 Sport*. 13:108-115

499 Taylor NA (2006). Challenges to temperature regulation when working in hot environments.
500 *Industrial health*. 44:331-344

501 Teunissen LP, Grabowski A, Kram R (2007). Effects of independently altering body weight and
502 body mass on the metabolic cost of running. *J Exp Biol*. 210:4418-4427

503 Tran Trong T, Riera F, Rinaldi K, Briki W, Hue O (2015). Ingestion of a cold temperature/menthol
504 beverage increases outdoor exercise performance in a hot, humid environment. *PloS one*.
505 10:e0123815

506 Trappe T, Pease D, Trappe S, Troup J, Burke E (1996). Physiological responses to swimming while
507 wearing a wet suit. *Int J Sports Med*. 17:111-114

508 Tyler CJ, Sunderland C (2011). Neck cooling and running performance in the heat: single versus
509 repeated application. *Med Sci Sports Exerc*. 43:2388-2395

510 Voltaire B, Berthouze-Aranda S, Hue O (2003). Influence of a hot/wet environment on exercise
511 performance in natives to tropical climate. *J Sports Med Phys Fitness*. 43:306-311

512 Voltaire B et al. (2002). Effect of fourteen days of acclimatization on athletic performance in
513 tropical climate. *Can J Appl Physiol*. 27:551-562

514 Von Duvillard SP, Braun WA, Markofski M, Beneke R, Leithauser R (2004). Fluids and hydration
515 in prolonged endurance performance. *Nutrition*. 20:651-656

516 Vrijens DM, Rehrer NJ (1999). Sodium-free fluid ingestion decreases plasma sodium during
517 exercise in the heat. *J Appl Physiol* (1985). 86:1847-1851

518 Wu SS, Peiffer JJ, Brisswalter J, Nosaka K, Lau WY, Abbiss CR (2015). Pacing strategies during
519 the swim, cycle and run disciplines of sprint, Olympic and half-Ironman triathlons. *Eur J
520 Appl Physiol*. 115:1147-1154

521 Yeo ZW, Fan PW, Nio AQ, Byrne C, Lee JK (2012). Ice slurry on outdoor running performance in
522 heat. *Int J Sports Med*. 33:859-866

523 Zaryski C, Smith DJ (2005). Training principles and issues for ultra-endurance athletes. *Curr Sports
524 Med Rep*. 4:165-170