Nature of Cp^*MoO_2 in Water and Intramolecular Proton-Transfer Mechanism by Stopped-Flow Kinetics and Density Functional Theory Calculations

Joo-Eun Jee, Aleix Comas-Vives, Chiara Dinoi, Gregori Ujaque, Rudi van Eldik, Agusti Lledos, Rinaldo Poli

► To cite this version:

Joo-Eun Jee, Aleix Comas-Vives, Chiara Dinoi, Gregori Ujaque, Rudi van Eldik, et al.. Nature of Cp^*MoO_2 in Water and Intramolecular Proton-Transfer Mechanism by Stopped-Flow Kinetics and Density Functional Theory Calculations. *Inorganic Chemistry*, 2007, 46 (10), pp.4103-4113. 10.1021/ic062409g . hal-03194455

HAL Id: hal-03194455

<https://hal.science/hal-03194455>

Submitted on 9 Apr 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Nature of $\text{Cp}^*\text{MoO}_2^+$ in water and intramolecular proton-transfer mechanism by stopped-flow kinetics and Density Functional Theory calculations

Joo-Eun Jee,^a Aleix Comas-Vives,^b Chiara Dinoi,^c Gregori Ujaque,^b Rudi van Eldik,^{*a} Agustí Lledós,^{*b} and Rinaldo Poli^{*c}

^a*Institute for Inorganic Chemistry, University of Erlangen-Nürnberg, Egerlandstr. 1, 91058 Erlangen, Germany*

^b*Unitat de Química Física, Departament de Química, Edifici Cn, Universitat Autònoma de Barcelona, 08193 Bellaterra, Catalonia, Spain*

^c*Laboratoire de Chimie de Coordination, UPR CNRS 8241 liée par convention à l'Université Paul Sabatier et à l'Institut National Polytechnique de Toulouse, 205 Route de Narbonne, 31077 Toulouse Cedex, France*

Proofs to :
Rinaldo Poli
Tel +33-561333173
Fax +33-561553003
E-mail poli@lcc-toulouse.fr

Abstract

A stopped-flow study of the $\text{Cp}^*\text{MoO}_3^-$ protonation at low pH (down to zero) in a mixed H_2O -MeOH (80:20) solvent at 25°C allows the simultaneous determination of the first acid dissociation constant of the oxo dihydroxo complex, $[\text{Cp}^*\text{MoO}(\text{OH})_2]^+$ ($\text{p}K_{\text{a}1} = -0.56$), and the rate constant of its isomerization to the more stable dioxo aqua complex $[\text{Cp}^*\text{MoO}_2(\text{H}_2\text{O})]^+$ ($k_{-2} = 28 \text{ s}^{-1}$). Variable temperature (5 - 25°C) and variable pressure (10 – 130 MPa) kinetics studies have yielded the activation parameters for the combined protonation/isomerization process ($k_{-2}/K_{\text{a}1}$) from $\text{Cp}^*\text{MoO}_2(\text{OH})$ to $[\text{Cp}^*\text{MoO}_2(\text{H}_2\text{O})]^+$, viz. $\Delta H^\ddagger = 5.1 \pm 0.1 \text{ kcal mol}^{-1}$, $\Delta S^\ddagger = -37 \pm 1 \text{ cal mol}^{-1} \text{ K}^{-1}$ and $\Delta V^\ddagger = -9.1 \pm 0.2 \text{ cm}^3 \text{ mol}^{-1}$. Computational analysis of the two isomers, as well as the $[\text{Cp}^*\text{MoO}_2]^+$ complex resulting from the dissociation of water, reveal a crucial solvent effect on both the isomerization and the water dissociation energetics. Introducing a solvent model by CPCM and especially by explicitly including up to three water molecules in the calculations led to the stabilization of the dioxo aqua species relative to the oxo dihydroxo isomer, and to the substantial decrease of the energy cost for the water dissociation process. The presence of a water dissociation equilibrium is invoked to account for the unusually low effective acidity ($\text{p}K_{\text{a}1}' = 4.19$) of the $[\text{Cp}^*\text{MoO}_2(\text{H}_2\text{O})]^+$ ion. In addition, the computational study reveals the positive role of external water molecules as simultaneous proton donors and acceptors, having the effect of dramatically lowering the isomerization energy barrier.

Introduction

The precise structure of aqua complexes and ions in water solution is rarely known. For instance, molybdic acid (H_2MoO_4) is commonly represented as $\text{Mo}(\text{OH})_6$, $\text{MoO}_2(\text{OH})_2(\text{H}_2\text{O})_2$ and $\text{MoO}_3(\text{H}_2\text{O})_3$,¹⁻⁶ while its first protonation product has been described as $\text{Mo}(\text{OH})_5(\text{H}_2\text{O})^+$ or $\text{MoO}_2(\text{OH})(\text{H}_2\text{O})_3^+$,^{1, 4, 5} although the isomeric formulation $\text{MoO}(\text{OH})_3(\text{H}_2\text{O})_2^+$ could also be envisaged. To the best of our knowledge, the factors regulating the relative energy of a transition metal dihydroxo complex and its isomeric aqua-oxo form (Scheme 1) are not well understood. In terms of the rate and mechanism of water exchange reactions, the correct assignment of such species is of utmost importance since it is well known that water exchange reactions follow the reactivity order $\text{M}-\text{OH}_2 \gg \text{M}-\text{OH} \gg \text{M}=\text{O}$.⁷

Scheme 1

In the organometallic area, the complex $\text{Cp}^*_2\text{Zr}(\text{OH})_2$ is a stable compound,⁸ whereas Cp^*_2ZrO is a reactive intermediate.⁹ On the other hand, Cp^*_2WO and $\text{Cp}^*_2\text{Ta}(\text{O})\text{H}$ are stable compounds that exchange water through $\text{Cp}^*_2\text{W}(\text{OH})_2$ and $\text{Cp}^*_2\text{Ta}(\text{OH})_2\text{H}$ intermediates.¹⁰ Finally, the closely related oxo and dihydroxo complexes $[\text{CpMoO}(\text{PMe}_3)_2]^+$ and $[(\eta^5\text{-C}_5\text{Et}_5)\text{Mo}(\text{OH})_2(\text{dppe})]^+$ have been isolated and structurally characterized.^{11, 12} In these systems, Cp, Cp* and related chelates are expected to have a drastic influence of the rate and mechanism of the water exchange process on the basis of data reported in the literature for water exchange on $[\text{Cp}^*\text{Rh}(\text{H}_2\text{O})_3]^{2+}$ and $[\text{Cp}^*\text{Ir}(\text{H}_2\text{O})_3]^{2+}$, which is 14 orders of magnitude faster than for the corresponding hexaaqua complexes and proceeds according to a more dissociative substitution mode.^{13, 14}

Some of us have initiated an investigation of the aqueous chemistry of high oxidation state organometallic compounds, focusing initially on Cp*Mo derivatives in a variety of oxidation states (VI, V, IV, and mixed-valence clusters).¹⁵ High oxidation state oxomolybdenum compounds are employed in a variety of catalytic reactions, such as olefin epoxidation, the selective oxidation of alcohols to aldehydes, the dehydrogenation and isomerization of alkenes, and even reductive processes such as the hydrosilylation of carbonyl compounds.¹⁶ Organometallic versions of these systems have shown high activities, notably in olefin epoxidation.¹⁷⁻²⁰ It would therefore be of interest to adapt these catalytic processes to an aqueous environment. To this end, knowledge of the nature of the Cp*Mo^{VI} aqua ion under different pH conditions is very useful.

A previous investigation of the aqueous speciation of Cp*Mo^{VI} has given the results summarized in Figure 1.²¹ In particular, the rate of conversion of compound Cp*MoO₂(OH) (generated quantitatively by the rapid protonation of Cp*MoO₃⁻ at pH ≤ 2) to the final cationic product was found to be first order in [H⁺]. The interpretation of this result leaves two possibilities as detailed in Scheme 2, both involving a proton addition pre-equilibrium to yield an intermediate cationic dihydroxo complex. The first possibility (path *a*) involves a rapid equilibrium rearrangement to an aqua-oxo isomer, followed by rate-determining loss of a water ligand, whereas the second one (path *b*) involves the intramolecular isomerization process as a rate-determining step. The [Cp*MoO(OH)₂]⁺ ion is likely a strong acid in water (pK_{a1} < 0), since most multiprotic inorganic oxo acids (e.g. H₂CO₃, H₂SO₃, H₃PO₄, ...) are characterized by a ΔpK_a of ca. 4-5 and the pK_{a2} (acid dissociation of Cp*MoO₂(OH)) is 3.65. Thus, the aquation state of the final cationic product is uncertain, [Cp*MoO₂(H₂O)_n]⁺ with either n = 0 or 1. The two possibilities cannot be distinguished on the basis of the kinetics experiment, nor by use of ¹H NMR spectroscopy because only the large solvent resonance was observed under all pH conditions, providing only negative evidence for water coordination (the resonance of the coordinated water molecule could either be overshadowed by the

much stronger solvent resonance, of the ligand exchange is in the fast regime leading to a single averaged resonance). IR spectroscopy is also useless for such dilute solutions in a strongly absorbing medium.

Figure 1. Thermodynamic and kinetic parameters related to the $\text{Cp}^*\text{Mo}^{\text{VI}}$ system in a H_2O - MeOH (80:20) solution.²¹

Scheme 2

Subsequent attempts to crystallize a salt of the $[\text{Cp}^*\text{MoO}_2(\text{H}_2\text{O})_n]^+$ ion and to determine its aquation state by X-ray crystallography have not been successful, always yielding instead the neutral dinuclear compound $\text{Cp}^*_2\text{Mo}_2\text{O}_5$.²² In fact, the low solubility product of this compound drives its precipitation by combination with the thermodynamically unfavorable (at low pH, see Figure 1) anionic hydrolysis product, $\text{Cp}^*\text{MoO}_3^-$. Another point that has remained unexplained from the previous study²¹ concerns the greater thermodynamic stability of the dioxo cation, $\text{Cp}^*\text{MoO}_2^+$ (or its water adduct) relative to the oxo dihydroxo isomer, $\text{Cp}^*\text{MoO}(\text{OH})_2^+$.

In this contribution, we present a more detailed stopped-flow kinetics investigation of the generation of the $[\text{Cp}^*\text{MoO}_2(\text{H}_2\text{O})_n]^+$ ion in the low pH regime, including variable temperature and variable pressure experiments, as well as a DFT computational study aimed at establishing (i) the aquation state of this ion and (ii) the mechanism of the slow process which leads to its generation. The results of this theoretical investigation are potentially of more general interest, as they illustrate the effect of the medium on the relative stability of oxo and dihydroxo transition metal species.

Results and Discussion

(a) Aquation state of $[\text{Cp}^*\text{MoO}_2(\text{H}_2\text{O})_n]^+$

The reaction shown in Equation 1 was studied computationally by DFT methods on the full systems (*i.e.* no ligand simplification was adopted). Both complexes were optimized in the gas phase. These two minima are shown in Figure 2. The unsolvated species give rise to a bent geometry, with the Mo-Cp* ring centroid axis forming an angle of 136.9° from the MoO_2 plane, which is only slightly smaller than the same angle in the water adduct (141.9°). Relative to the ideal planar structure (optimized independently under constrained C_s symmetry), this configuration is more stable by only $2.9 \text{ kcal mol}^{-1}$.

Equation 1

Figure 2. Optimized structures of the $[\text{Cp}^*\text{MoO}_2]^+$ and $[\text{Cp}^*\text{MoO}_2(\text{H}_2\text{O})]^+$ (**2a**) complexes.

The electronic structure of two-legged piano stool complexes has been analyzed before,²³⁻²⁵ including the effect of the coordination environment^{26, 27} and spin state^{28, 29} on the relative stability of planar and bent geometries. All previous studies, however, are restricted to electronically unsaturated (16-electron) d^6 systems, typically containing π -acidic (e.g. CO), neutral (e.g. H), or weakly donating (e.g. Cl) ligands. On the other hand, complex $[\text{Cp}^*\text{MoO}_2]^+$ features two strong double-sided π -donors on a d^0 metal center. Since the Cp^* ligand is electronically isolobal with the O^{2-} ligand (both are potentially 6-electron, $\sigma+2\pi$ donors), the compound can also be considered as isolobally related to MoO_3 , which is a d^0 MX_3 systems with strong double-sided π donor ligands. The geometric preference for d^0 MX_3 systems has been recently analyzed by Eisenstein *et al.*³⁰ for M = Group 3, Group 4 and lanthanide elements, and X = alkyl, halide and amido groups. It was found that the pyramidal structure is always preferred because of the d orbital participation in M-X σ bonding, whereas the ionic component should favor the planar structure. The contribution from π donation provides a driving force towards flattening the structure. Therefore, it seems that the Mo- Cp^* and Mo-O σ interactions dominates the geometric preference for the $[\text{Cp}^*\text{MoO}_2]^+$ ion. We have also calculated the isolobal MoO_3 system and found that the pyramidal

structure ($\text{O-Mo-O} = 109.7^\circ$) is again favored, in this case by $9.5 \text{ kcal mol}^{-1}$ with respect to the D_{3h} structure. The Mo-ligand interactions nevertheless have a significant contribution from ionicity. This is suggested by the values of the Mulliken charges, for instance the charges in the $[\text{Cp}^*\text{MoO}_2]^+$ ion are $+1.38$ for Mo and -0.52 (average) for the two O atoms, whereas in the MoO_3 molecule they are $+1.88$ and -0.63 , respectively, showing that the metal is electron-richer in $[\text{Cp}^*\text{MoO}_2]^+$ than in MoO_3 .

As expected, the addition of a water molecule results in an energetic stabilization of the $[\text{Cp}^*\text{MoO}_2]^+$ complex, corresponding to a potential energy gain of 39.7 kcal/mol in the gas phase. In addition to providing additional covalent bonding stabilization, this process is also favorable from an electrostatic point of view, since a positively charged Mo atom (Mulliken charge 1.37) is directly linked to the oxygen atom of a water molecule which has a marked anionic character (Mulliken charge -0.71). Although this addition process is entropically disfavored, the free energy difference of the reaction is still exergonic by 28.1 kcal/mol . The inclusion of solvent effects by means of the CPCM method changes significantly the results. The energetic difference remains in favor of the water adduct but the gain is reduced to 10.1 kcal/mol . Having the $\text{Cp}^*\text{MoO}_2^+$ and the H_2O molecules separated in the gas phase is a very energy costing process because a strong cation-dipole interaction is broken. Adding solvent effects stabilize the separated reactants by interacting with the dielectric continuum of the polar medium, leading to a considerable decrease in the energy difference between the isolated species and the aqua complex. The conclusion of this investigation is to establish the nature of the acidic form of $\text{Cp}^*\text{Mo}^{\text{VI}}$ in water as the aqua complex $[\text{Cp}^*\text{MoO}_2(\text{H}_2\text{O})]^+$, although the dissociation of the water ligand to afford the $[\text{Cp}^*\text{MoO}_2]^+$ intermediate may be a facile process. As mentioned in the Introduction, from a kinetics point of view, the Cp^* ligand is known to exert a strong *trans* effect on ligand dissociation processes. The introduction of the Cp^* ligand on the very inert hexa-aqua complexes of Ir(III) and Rh(III), *i.e.* on going from the $[\text{M}(\text{H}_2\text{O})_6]^{3+}$ to the $[\text{Cp}^*\text{M}(\text{H}_2\text{O})_3]^{2+}$ complexes ($\text{M} = \text{Rh}, \text{Ir}$), increases the water exchange rate

by 12-14 orders of magnitude.¹³ A ^{17}O NMR investigation of the water exchange process for the $\text{Cp}^*\text{Mo}^{\text{VI}}_{\text{aq}}$ system as a function of pH is currently ongoing in our laboratories and will be reported in detail in a separate contribution, but we are already able to state that this appears to be an extremely fast process. Therefore, the mechanism of the low-pH transformation leading from $\text{Cp}^*\text{MoO}_2(\text{OH})$ to $[\text{Cp}^*\text{MoO}_2(\text{H}_2\text{O})]^+$ may be formulated as path *b* in Scheme 2, the slow step being the intramolecular proton transfer process, leading from the intermediate dihydroxo complex $[\text{Cp}^*\text{MoO}(\text{OH})_2]^+$ to the final product.

(b) Protonation kinetics in the low pH regime

In an effort to determine the first acid dissociation constant of $[\text{Cp}^*\text{MoO}(\text{OH})_2]^+$ ($\text{p}K_{\text{a1}}$), the acidification kinetics of $[\text{Cp}^*\text{MoO}_3]^-$ was studied at low pH (down to zero), where the complex is present in the $[\text{Cp}^*\text{MoO}_2(\text{OH})]$ form ($\text{p}K_{\text{a2}} = 3.65$, see Figure 1). Consistent with the previous study,²¹ acidification kinetics showed a linear $[\text{H}^+]$ dependence up to 0.35 M ($\text{pH} = 0.45$), but a slight saturation effect was visible at concentrations up to 1 M (pH down to 0) as shown in Figure 3. This behavior is fully consistent with the previously established scheme (Figure 1), because the rate law for the acidification reaction can be expressed as in Equation 2 (the equilibrium K_{a1} is rapidly maintained on the time scale of k_{-1} and k_{-2}).²¹ At low $[\text{H}^+]$ ($K_{\text{a1}}^{-1}[\text{H}^+] \ll 1$), the rate law further simplifies to Equation 3, which accounts for the linear dependence observed under such conditions. The small intercept could be ascribed to a minor contribution of the parallel k_{-1} path. It is useful to remind the reader here that at low pH the k_{-1} pathway, corresponding to direct loss of OH^- from $\text{Cp}^*\text{MoO}_2(\text{OH})$, should be negligible ($k_{-1} = 6.32 \cdot 10^{-7} \text{ s}^{-1}$) compared to the k_{-2} pathway, corresponding to protonation and isomerization of the

$[\text{Cp}^*\text{MoO}(\text{OH})_2]^+$ intermediate, *e.g.* $k_{-2}K_{a1}^{-1}[\text{H}^+] = 1.73 \text{ s}^{-1}$ at pH 1. Alternatively, the intercept can also be due to a contribution of the back reaction. i.e. the k_2 path.

Figure 3. Plots of k_{obs} versus $[\text{H}^+]$ for the acidification of $[\text{Cp}^*\text{MoO}_3]^-$ with HNO_3 in the range 0.01 to 0.95 M. Experimental conditions; $[\text{Cp}^*\text{Mo}_2\text{O}_5] = 4 \times 10^{-4} \text{ M}$, $\lambda_{\text{det}} = 390 \text{ nm}$, 20% MeOH – H_2O , temp = 25°C , $\mu_{\text{tot}} = 1 \text{ M}$ (adjusted with NaNO_3).

$$k_{\text{obs}} = \{k_{-1} + k_{-2}K_{a1}^{-1}[\text{H}^+]\} / \{1 + K_{a1}^{-1}[\text{H}^+]\}$$

Equation 2

$$k_{\text{obs}} \sim k_{-1} + k_{-2}K_{a1}^{-1}[\text{H}^+] \quad (\text{if } [\text{H}^+] \text{ small})$$

Equation 3

The fit of the data in Figure 3 with the expression of Equation 4, for which the small contribution of the intercept was ignored, resulted in $k_{-2} = 28 \pm 1 \text{ s}^{-1}$ ($\log k_{-2} = 1.44$) and $K_{a1} = 3.6 \pm 0.6 \text{ M}$, from which

$pK_{a1} = -0.56$. It follows that k_{-2} is almost 8 orders of magnitude larger than k_{-1} and that ΔpK_a (*i.e.* $pK_{a2} - pK_{a1}$) = 4.1. These values are consistent with expectations, since as mentioned above the ΔpK_a of a typical inorganic diprotic oxoacid is ≥ 4 . Thus, the new results described here fully confirm the validity of the previously proposed kinetic scheme (Figure 1), where a rapid pre-equilibrium protonation precedes a slow step, which is now known to be the intramolecular proton transfer, see next sections. It is worth to mention that the proton transfer to the oxo ligands in compound $\text{Re}(\text{O})\text{I}(\text{2,7-nonadiyne})$ by $\text{CF}_3\text{SO}_3\text{H}$ in MeCN was shown to be relatively slow ($11.9 \text{ M}^{-1} \text{ s}^{-1}$ at -40°C).³¹

$$k_{\text{obs}} \sim k_{-2}K_{a1}^{-1}[\text{H}^+]/\{1 + K_{a1}^{-1}[\text{H}^+]\} \quad (\text{if } k_{-1} \text{ negligible})$$

Equation 4

It should be noted that the individual values of K_{a1} and k_{-2} obtained by the above analysis give rise to a ratio $k_{-2}/K_{a1} = 7.8$, in relatively good agreement with the previously determined value (17.2).²¹ The discrepancy may be attributed to the difference in ionic strength used in these studies. However, one point remains to be discussed on the thermodynamics of the system, before we can turn to the kinetics. The combination of the individual rate constants for the tautomerization of the cationic system in both directions (k_2 and k_{-2} as shown in Scheme 3) yields the tautomerization equilibrium constant, $K_2 = k_2/k_{-2} = 1.8 \cdot 10^{-5}$, which corresponds to a free energy difference of $6.5 \text{ kcal mol}^{-1}$ in favor of the oxo aqua isomer. The combination of this value with the experimentally determined pK_{a1} value (-0.56) according to the thermodynamic cycle shown in Scheme 3 (the deprotonation of the two isomeric cations affords the same neutral hydroxo complex) yields the thermodynamic proton dissociation constant of the dioxo aqua species, $K_{a1}' = 6.5 \cdot 10^{-5} \text{ M}$ ($pK_{a1}' = 4.19$). This compares well with the value calculated from the k_{-2}/K_{a1} determined in the previous study ($K_{a1}' = 2.9 \cdot 10^{-5} \text{ M}$; $pK_{a1}' = 4.54$). Note that this value is also given by

the expression $(k_{+1}/k_{-1})K_s$, from the thermodynamic cycle in Figure 1. At first sight, it may appear unreasonable that the first acid dissociation constant of this aqua complex is weaker than the second one. However, this observation can be rationalized if the water ligand is extensively dissociated due to the labilization caused by the Cp* chelate as referred to above. In this case, the calculated pK_{a1} ' value must be treated as an apparent pK_a value since it includes the equilibrium constant for the dissociation of water, which is expected to be large in order to offset the observed pK_a value. Thus the effective thermodynamic acidity of the cationic complex is a weighted average of the coordinated water molecule acidity, which is expected to be intrinsically very high, and the acidity of the free water molecule ($pK_a = 15.6$). The hypothesis of an extensive water dissociation from the cationic complex is not inconsistent with the computational study, since the water adduct is calculated as only 10.1 kcal/mole more stable than the dissociated species according to the CPCM, but this calculation does not consider explicit interactions of the adduct and the separate fragments with additional water molecules. For instance, the stabilization of a water molecule by hydrogen bonding to additional water molecules is expected to contribute significantly to the equilibrium energetics (the solvation free energy of a water molecule in water has recently been estimated at -6.3 kcal/mol).³² It is also interesting to recall that an electrospray mass spectrometric investigation of a H₂O-MeOH solution of Cp*Mo^{VI} at low pH (ca. 4 and 1) revealed the presence of both Cp*MoO₂⁺ and Cp*MoO₂(H₂O)⁺ species, as well as the related Cp*MoO₂(MeOH)⁺ adduct.^{33, 34} Whether the rapid equilibrium highlighted at the left hand side of Scheme 3 involves only species Cp*MoO₂⁺ and Cp*MoO₂(H₂O)⁺, or whether also a methanol solvate is present, does not affect the measured values of all rate and equilibrium constants, nor their significance.

Scheme 3

In general it is expected that ground state labilization caused by metal-carbon bonds in the *trans* position to coordinated water molecules will not only drastically accelerate the water exchange process, but will also significantly increase the $\text{p}K_a$ value of the coordinated water as found in the present case. Bond weakening of the M-OH₂ bond will cause a decrease in the acidity of the coordinated water molecule.^{7, 14} In addition, the bond strength will follow the order M=O >> M-OH >> M-OH₂ as mentioned before. Thus, the combination of bond labilization and pH will control the lability of coordinated water and lead to apparent $\text{p}K_a$ values that seem to be ‘abnormal’.

It is also interesting to analyze the $\text{Cp}^*\text{Mo}^{\text{VI}}$ system in terms of the isolobal analogy between the O^{2-} and Cp^{*-} ligands. Thus, the $\text{Cp}^*\text{MoO}_2^+$ system is isolobally related to MoO_3 , $\text{Cp}^*\text{MoO}_2(\text{OH})$ to HMoO_4^- , and $\text{Cp}^*\text{MoO}_3^-$ to MoO_4^{2-} . Interestingly, while the MoO_4^{2-} and HMoO_4^- are always described as tetrahedral ions, aqueous MoO_3 is described as having coordination number 6 (either as $\text{MoO}_3(\text{H}_2\text{O})_3$ or $\text{MoO}_2(\text{OH})_2(\text{H}_2\text{O})_2$), possibly because an octahedral environment is found for MoO_3 ,³⁵ as well as for its water adduct $\text{MoO}_3 \cdot 2\text{H}_2\text{O}$,³⁶ in the solid state. However, equilibria with 5- or 4-coordinate species may in fact exist. A comparison between the $\text{p}K_a$ of HMoO_4^- (3.48)⁴ and $\text{Cp}^*\text{MoO}_2(\text{OH})$ (3.65) suggests that the Cp^{*-} ligand is a marginally better electron donor than the isolobal O^{2-} ligand for this system in an aqueous solvent. This means that the Mo center is electron-richer in $\text{Cp}^*\text{MoO}_2^+$ than in MoO_3 as supported by the

Mulliken charges in Mo atom (+1.38 vs. +1.88). In combination with the greater steric bulk of the Cp* ligand vs. the oxo ligand, this may lead to a weaker interaction between $[\text{Cp}^*\text{MoO}_2]^+$ and water.

(c) Activation parameters for the low-pH transformation

The temperature and pressure dependence of the acidification process for the protonation reaction in the low acidity range have also been investigated. The data were collected only in the pH range where the rate depends linearly on $[\text{H}^+]$, see Figure 4. The obtained activation parameters are $\Delta H^\ddagger_{\text{slope}} = 5.1 \pm 0.1$ kcal mol⁻¹ and $\Delta S^\ddagger_{\text{slope}} = -37 \pm 1$ cal mol⁻¹ K⁻¹. The double logarithmic plots (log k vs. pH) are linear with slopes very close to unity, demonstrating that the intercept plays a minor role in fitting the data.

Figure 4. (a) Plots of k_{obs} versus $[\text{H}^+]$ for the acidification of $[\text{Cp}^*\text{MoO}_3]^-$ with HNO_3 in the temperature range 5 – 25 °C measured by stopped-flow. (b) Corresponding Eyring plot for the slope in (a). Experimental conditions; $[\text{Cp}^*\text{MoO}_3] = 4 \times 10^{-4}$ M, $\lambda_{\text{det}} = 390$ nm, 20% MeOH – H₂O, $\mu_{\text{tot}} = 1$ M (adjusted with NaNO_3).

The pressure dependence of the reaction was studied at 0.4 M acid for which the data are shown in Figure 5. The obtained activation parameter, $\Delta V_{\text{slope}}^\ddagger$ is $-9.1 \pm 0.2 \text{ cm}^3 \text{ mol}^{-1}$. The activation parameters for the slope of the plots in Figure 4 and Figure 5 represent those for k_2/K_{a1} . Since nothing is presently known about the temperature dependence of K_{a1} , it is difficult to speculate about the meaning of the reported activation parameters. It is reasonable to speculate that K_{a1} should not show a significant pressure dependence since it involves no changes in electrostriction and at the most can be slightly positive. The overall significantly negative activation entropy and volume values favor a process in which significant bond formation or charge creation occurs. According to the slow reaction for path (b) of Scheme 2, formation of the aqua complex involves bond formation between OH and H, as well as shortening of the Mo-O bond during conversion of Mo-OH to Mo=O. These processes are suggested to account for the observed volume collapse and decrease in entropy on going to the transition state.

Figure 5. Plot of $\ln(k_{\text{obs}})$ versus pressure for the acidification of $[\text{Cp}^*\text{MoO}_3]^-$ with HNO_3 in the range 10 – 130 MPa. Experimental conditions; $[\text{Cp}^*_2\text{Mo}_2\text{O}_5] = 4 \times 10^{-4} \text{ M}$, $[\text{H}^+] = 0.4 \text{ M}$, $\lambda_{\text{det}} = 390 \text{ nm}$, 20% MeOH – H_2O , temp = 25 °C, $\mu_{\text{tot}} = 1 \text{ M}$ (adjusted with NaNO_3).

(d) Computational study of the intramolecular proton transfer process

In order to validate the intramolecular proton transfer mechanism and find the possible origin of the negative activation parameters (entropy and volumes), we carried out calculations on the initial (**1a**) and end product (**2a**) of Equation 5, as well as on the transition state (**ts1a**), in the absence and in the presence of additional water molecules. The gas phase and water solution energy profiles obtained in the absence of additional water molecules are shown in Figure 6, whereas the optimized **1a** and **ts1a** species are shown in Figure 7. The barrier height is quite high, having a value of 43.1 kcal/mol in the gas phase, whereas the reaction is endoergic by 10.6 kcal/mol. Both results disagree with the experimental evidence, since the isomerization process is rather facile and the oxo-aqua complex is the thermodynamically more stable product. When the non-specific solvent effects are accounted for by use of the CPCM method, the barrier is nearly the same (43.9 kcal/mol), whereas the reaction is endoergic by 7.2 kcal/mol, so the energetic difference has decreased by 3.4 kcal/mol by the inclusion of the solvent dielectric effects. Concerning the free Gibbs energy values (in the gas phase), they are very similar to the potential energy ones, for example, the difference between **1a** and **2a** is of 10.6 kcal/mol in potential energy whereas the free energy difference is 9.9 kcal/mol. The high activation barrier is probably related to the distortion of the OMoO fragment as reflected in the large decrease in the O-Mo-O angle on going from **1a** (106.8°) to the transition state structure **ts1a** (71.3°). The transition state is also characterised by distances of 1.221 Å and 1.276 Å for the O-H bonds being broken and formed, respectively, when going from **1a** to **2a**. Thus, the old O-H bond is already almost fully broken, whereas the new one has not yet formed to a significant extent, and the two Mo-O bond distances have intermediate values between those of the corresponding distances in the starting and final compounds. In conclusion, the intramolecular proton transfer implies a remarkable change in the geometry and this is associated with a high energetic cost.

Equation 5

Figure 6. Relative energy profile (in kcal mol⁻¹) in gas phase and water solution for the starting complex(**1a**), transition state (**ts1a**) and product (**2a**) of Equation 5.

Figure 7. Optimized structures of the dihydroxo complex (**1a**) and the proton transfer transition state (**ts1a**) of Equation 5 (distances in Å; angles in degrees).

In order to further probe the mechanistic details of this reaction, we considered that the explicit inclusion of water molecules could afford an easier proton transfer pathway, as the system contains both donor and acceptor sites for the establishment of hydrogen bonds with water molecules. Many theoretical works have already reported the active participation of water chains in tautomerization³⁷⁻³⁹ and proton

exchange processes.^{40, 41} The mechanism is described as a water-assisted reaction in which one or more water molecules act as bifunctional catalyst. Recent theoretical studies of the tautomerization process between the hydrated oxide, $[\text{MO}(\text{H}_2\text{O})]^+$, and the dihydroxide $[\text{M}(\text{OH})_2]^+$ cations ($\text{M} = \text{V}, \text{Nb}$ and Ta) also point out that the participation of water acting as proton donor and acceptor can effectively lower the barrier height for the isomerization process.⁴² The same water-catalysis has been found for the isomerization of $\text{UO}_2(\text{OH})_2$.⁴³ The resulting energy profiles with one additional water molecule are shown in Figure 8. In Figure 9 the main geometric characteristics of the optimized structures are depicted. First of all, the interaction of the additional water molecule with systems **1a** and **2a** gives rise to a significant energetic stabilization (15.5 and 8.1 kcal mol⁻¹ in the gas phase and in water solution, respectively, for the dihydroxo complex **1b**; 21.0 and 11.3 kcal mol⁻¹ under the same conditions for the oxo aqua complex **2b**). The stabilization of **2b** is slightly greater than that of **1b**, which renders the isomerization process less endothermic relative to the situation of the isolated system ($\Delta E = +5.1$ and $+4.0$ kcal mol⁻¹ in the gas phase and in water solution). The corresponding free energy difference in the gas phase is $+3.4$ kcal mol⁻¹. This change seems related to the stronger hydrogen bonding of water with the proton of the aqua ligand in **2b**, relative to the proton of one of the two hydroxo ligands in **1b** (as measured by the greater O-H bond elongation, 1.025 vs. 0.997 Å, and by the shorter O...H distance, 1.511 vs. 1.655 Å). The most dramatic effect relative to the water free system, however, is observed at the relative barrier height, which is now only 12.7 kcal mol⁻¹ (vs. 43.1 without water) in the gas phase and only 10.3 kcal mol⁻¹ (vs. 43.9 without water) in solution above the **1b** species. This effect is related to the ability of the additional water molecule to act, at the same time, as a proton acceptor for the donating O-H ligand and as a proton donor for the receiving O-H ligand, thereby mediating the proton transfer. Both hydrogen-bonding interactions are established at the level of the transition state (**ts1b**) giving a six-membered transition state, as shown in Figure 9. The most likely cause of such dramatic decrease in activation barrier is the smaller distortion of

the OMoO fragment in the transition states, as evidenced by the wider O-Mo-O angle in **ts1b** (92.5°) in comparison with the 71.3° value optimized for the **ts1a** structure. The optimized O-Mo-O angles in **1b** and **2b**, on the other hand, are essentially unchanged relative to those of **1a** and **2a**. Thus, the structural rearrangement in **ts1b** is not as marked as in **ts1a**, with respect to the relative reactants. It is worth to mention that a very slow rate of intramolecular proton transfer was reported for complex $\text{Re}(\text{O})(^{18}\text{OH})(\text{MeC}\equiv\text{CMe})_2$ in benzene solution.⁴⁴ Although the rate of transformation was shown not to strongly depend on the presence or absence of stoichiometric amounts of ethanol, trace amounts of $\text{CF}_3\text{SO}_3\text{H}$ strongly accelerate the reaction. It is possible that the transformation could also be accelerated by excess water. Another slow intramolecular proton transfer process was reported for *trans*- $[\text{Ru}^{\text{IV}}(\text{tpy})(\text{O})(\text{O}=\text{P}(\text{Ph})_2\text{CH}_2\text{CH}_2\text{PPh}_2)(\text{H}_2\text{O})]^{2+}$ in MeCN solution.⁴⁵

Another noteworthy characteristic of **ts1b** is that the proton originating from the donating O-H group has already migrated onto the water molecule (1.088 Å), whereas the proton that eventually ends up in the aqua ligand is still bonded to the added water molecule (1.026 Å). These distances suggest that the transition state involves the interaction between a hydronium ion (H_3O^+) and the neutral $\text{Cp}^*\text{MoO}_2(\text{OH})$ molecule, through the establishment of two hydrogen bonds (1.382 Å to the oxo ligand and 1.591 Å to the hydroxo ligand). We have reported a similar transition-state in hydrogenation performed by gold catalysts for the heterolytic cleavage of H_2 assisted by a solvent molecule (ethanol).⁴⁶ In conclusion, the explicit consideration of a water molecule confirms the feasibility of the rearrangement of **1b** to **2b** by solvent assisted intramolecular proton transfer.

Equation 6

Figure 8. Relative energy profiles (in kcal mol⁻¹) in gas phase and water solution for the starting complex (**1b**), transition state (**ts1b**) and product (**2b**) of Equation 6.

Figure 9. Optimized structures of the starting complex (**1b**, above), transition state (**ts1b**, middle) and product (**2b**, below) of Equation 6 (distances in Å; angles in degrees).

On the basis of the above results, we have imagined that the addition of a second water molecule could further lower the energy of the transition state. An eight-membered transition-state (**ts1c**) having two water molecules acting as proton and donor acceptors concurrently between the two complexes **1c** and **2c** (Equation 7) may further reduce the necessary geometric reorganization to reach the transition state. The resulting energy profiles in the gas phase and water solution on this system are shown in Figure 10 and the optimized structures for the related species are drawn in Figure 11. As already seen on going from the water-free to the H₂O adduct, going from the mono to the bis(water) adduct further stabilizes the oxo-aqua isomer relative to the dihydroxo isomer. The oxo-aqua complex is now only 1.4 kcal mol⁻¹ less stable than the dihydro isomer in the gas phase (*cf.* 5.1 and 10.6 kcal mol⁻¹ for the mono(water) and water-

free system, respectively), and actually *more stable* by a small margin (0.4 kcal mol⁻¹) in water solution. The gas phase free energy is still in favor of the dihydroxo isomer, but only by 2.4 kcal mol⁻¹. Undoubtedly, the correct relative energy can be obtained at this level of theory, provided the effect of the water solvation is considered by the explicit inclusion of a sufficient number of water molecules in the calculation. Although the relative free energy computed by including two water molecules is still a bit far from the experimental value of 6.5 kcal mol⁻¹ in favor of the oxo aqua isomer, the explicit inclusion of water molecules in the calculations moves the energetic balance in the right direction. In conclusion, our calculations indicate an intrinsic preference of the system for the dihydroxide form (gas phase). The non-specific interactions (continuum calculations) are not sufficient to reverse this stability order, whereas a reversal, in agreement with the experiment, is observed upon inclusion of the specific interactions (hydrogen bonds) with the water solvent. Undoubtedly, the correct relative energy can be obtained at this level of theory, provided the effect of the water solvation is considered by the explicit inclusion of a sufficient number of water molecules in the calculation.

Equation 7

Figure 10. Relative energy profiles (in kcal mol⁻¹) in gas phase and water solution for the starting complex (**1c**) transition state (**ts1c**) and product (**2c**) of Equation 7.

Figure 11. Optimized structures of the starting complex (**1c**, above), transition state (**ts1c**, middle) and product (**2c**, below) of Equation 7 (distances in Å; angles in degrees).

Concerning the energetic profile, the proton transfer barrier is now reduced to a very low value (1.3 kcal/mol and 3.5 kcal/mol in potential and free energy, respectively). In CPCM single point calculations the energy of the transition-state is even lower than the corresponding reactant energy. We consider that this difference is within the error of the CPCM method. Compared to **ts1b** and **ts1a**, the structure of **ts1c** is characterized by an even wider O-Mo-O angle (102.0°). Now the Mo-O bonds barely have to bend in order to reach the suitable configuration for the proton transfer. The explicit inclusion of the two water molecules also has the effect of further weakening the O-H bonds for the hydroxo ligands (1.049 Å in **1c**, vs. 0.997 Å in **1b**) and the aqua ligand (1.132 Å in **2c**, vs. 1.025 Å in **2b**). Thus, the proton becomes effectively more transferred to the medium. As was the case for **ts1b**, the structure of **ts1c** may also be viewed as resulting from the hydrogen bonding interaction of a hydronium ion (in this case H₅O₂⁺) with neutral Cp*MoO₂(OH), according to the observed pattern of O-H and O...H distances.

For the sake of completion, we have pursued the study by adding a third water molecule to the system. In this case, however, the optimization of both the hydroxo and the oxo aqua isomers **1d** and **2d** resulted in an optimized structure where the water proton is completely transferred to the water cluster,

which thereby becomes a H_7O_3^+ hydronium ion and interacts via hydrogen bonding with the neutral $\text{Cp}^*\text{MoO}_2(\text{OH})$ complex, see Figure 12. These results confirm the trend reported in **2b** and **2c**, where the O-H bond became more elongated as more water molecules were added in the system. This trend reflects the acidity of the cationic species, thus a more water-rich environment favors the proton transfer to the medium. The two structures **1d** and **2d** are isomers in terms of the arrangement of the hydrogen-bonded water molecules. Structure **2d** is more stable than **1d** by $1.8 \text{ kcal mol}^{-1}$ in water solution. It is interesting to note that the protonated water molecule (H_3O^+) interacts with the OH group in structure **2d**, whereas it prefers an oxo ligand in structure **1d**. Although this difference is fairly small and the final point depends on the starting point in the optimization, this difference may be attributed to the fact that in **2d** there is an additional hydrogen bond. Moreover, the O(OH) interacting with H_3O^+ in **2d** presents a more negative Mulliken charge than the O interacting with H_3O^+ in **1d**: -1.11 vs. -0.89.

Figure 12. Optimized structure of **1d** and **2d**, featuring the $\text{Cp}^*\text{MoO}_2(\text{OH})$ molecule interacting with the $[(\text{H}_3\text{O})(\text{H}_2\text{O})_2]^+$ cluster (distances in Å; angles in degrees).

A comparison of all isomeric pairs shows that the aqua O-H bond in the oxo aqua isomer systematically experiences a greater lengthening effect than the hydroxo O-H bond in the dihydroxo

isomer (1.025 Å in **2b** vs. 0.997 Å in **1b**, 1.132 Å in **2c** vs. 1.049 Å in **1c**; these distances are essentially identical in **1a** and **2a**). This suggests that $[\text{Cp}^*\text{MoO}_2(\text{H}_2\text{O})]^+$ is intrinsically more acidic than its dihydroxo isomer, which may seem in contradiction with the experimental evidence (oxo aqua isomer: $\text{p}K_{\text{a}} = 4.19$; dihydroxo isomer: $\text{p}K_{\text{a}} = -0.56$). However, as argued above, the high effective $\text{p}K_{\text{a}}$ value of $[\text{Cp}^*\text{MoO}_2(\text{H}_2\text{O})]^+$ is proposed to result from an equilibrium with $[\text{Cp}^*\text{MoO}_2]^+$ and free H_2O . Thus, these computational results indirectly validate the water dissociation hypothesis.

The calculations agree with the experimental evidence that the cationic systems, $[\text{Cp}^*\text{MoO}(\text{OH})_2]^+$ and $[\text{Cp}^*\text{MoO}_2(\text{H}_2\text{O})]^+$, can be deprotonated in a slightly acidic medium. However, the predominant species in very acidic media is a cationic complex. As a very simple way of modeling the effect of lowering the pH, we have added an extra proton to the system, i.e. using the $[\text{H}_3\text{O}(\text{H}_2\text{O})_2]^+$ cluster as a model of the medium. The resulting system is dipositive. This is a rough simulation of a low pH aqueous solution but can give some insights of what occurs when increasing the acidity of the medium. Several theoretical studies have addressed the question of the nature of the hydronium species in solution and its solvation shell.⁴⁷⁻⁵⁰ The $[\text{H}_3\text{O}(\text{H}_2\text{O})_2]^+$ cluster has been used as a model of the hydronium species in solution in other reactions involving transition metal complexes.^{51, 52} Different structures were optimized, exploring several starting points, and the various optimized species are shown in Figure 13. For species **1e** and **2e**, the hydronium ion (H_3O^+) is located in the second coordination sphere, whereas species **1e2**, **1e3** and **2e2** feature this ion in the first coordination sphere, directly interacting with the molybdenum complex. Obviously, these are not necessarily all the possible minima, but probably the most favored ones.

Figure 13. Optimised structures of the $[\text{Cp}^*\text{MoO}(\text{OH})_2]^+$ and $[\text{Cp}^*\text{MoO}_2(\text{H}_2\text{O})]^+$ complexes interacting with the $[(\text{H}_3\text{O})(\text{H}_2\text{O})_2]^+$ cluster (relative energies in kcal mol⁻¹ in parentheses).

In no case was the proton of the cationic complex $[\text{Cp}^*\text{MoO}_2(\text{H}_2\text{O})]^+$ or $[\text{Cp}^*\text{MoO}(\text{OH})_2]^+$ transferred to the medium, conversely of what occurred for the monopositive system with three water molecules. This is not a surprising result, the medium (modeled here as $[\text{H}_3\text{O}(\text{H}_2\text{O})_2]^+$) being too acidic in the gas phase to accept an additional proton, because it has an excess of positive charge. Nevertheless, this approach gives a nice representation of what occurs when lowering the pH value because the cationic species become now favorable and can be obtained as real minima in the potential energy surface for both isomers.

Concerning the relative energies, placing the hydronium ion in the first coordination sphere carries an energy cost of ca. 2 kcal/mol, relative to having it in the second sphere (e.g. going from **1e** to **1e2** and **1e3** for the dihydroxo complex, and from **2e** to **2e2** for the oxo aqua complex). On the other hand, the

dihydroxo isomer is more stable than the oxo aqua species by 5.3 kcal mol⁻¹ (**1e** vs. **2e** structure). As discussed before, we believe that a larger number of water molecules (at least the complete first solvation sphere) should be included in order to reproduce the experimental relative stability.

The experimental data also show that the dominant species at pH > 5 is the anionic complex [Cp*MoO₃]⁻.²¹ We suggest that when increasing the hydroxide concentration, the proton of the [Cp*MoO₂(OH)] complex can be easily transferred to hydroxide in solution, analogously to what occurs to one proton of the aqua complex when water molecules are added to the system. Nevertheless, although theoretical calculations reported here provide some relevant energy trends, geometries and activation barriers, a complete theoretical analysis of the species present in solution as a function of pH is beyond the scope of this study, mainly because of the intrinsic difficulty of simulating an aqueous solution at a given pH. More information could be extracted from dynamics simulations.

Conclusions

A combination of kinetic and computational investigations on the Cp*Mo^{VI} system in acidic aqueous medium has greatly improved our understanding of what factors regulate the properties of the oxo dihydroxo ([Cp*MoO(OH)₂]⁺) and dioxo aqua ([Cp*MoO₂(H₂O)]⁺) isomers in solution. The computational study shows that the dihydroxo isomer is favored for the isolated system in the gas phase, but the subsequent introduction of the solvent model (by CPCM) and especially the explicit introduction of water molecules in the calculations attenuate and even reverse this stability trend. The study further suggests that the medium strongly affects the water dissociation energy cost from the dioxo aqua isomer (10.1 kcal mol⁻¹ in water solution vs. 39.7 kcal mol⁻¹ in the gas phase). A generalization of these trends suggests that the dihydroxo species for any given system have a better chance to be stable in apolar

organic solvents, whereas a rearrangement to oxo complexes accompanied by water dissociation may be favored in stronger dielectric solvents.

The above trend also rationalizes the unusually low acidities that are sometimes found for high oxidation state cationic complexes. While the water ligand in the oxo aqua isomer, $[\text{Cp}^*\text{MoO}_2(\text{H}_2\text{O})]^+$, is an intrinsically stronger proton donor than the hydroxo ligand in the dihydroxo isomer, $[\text{Cp}^*\text{MoO}(\text{OH})_2]^+$, the much lower observed acidity for the former ($\text{p}K_{\text{a}} = 4.19$; vs. -0.56 for the latter) is related to a water dissociation equilibrium. In turn, this is related to the coordination sphere and particularly to how the ligands are able to satisfy the electron deficiency at the metal center upon loss of the water lone pair donation. In the present system the extremely high lability of the coordinated water molecule can be ascribed to the *trans* labilization effect of the Cp^* chelate, which in turn favors the formation of the dioxo complex and is responsible for the apparent high $\text{p}K_{\text{a}}$ value of the oxo aqua isomer. We are not aware of any other analogous study in the literature.

Concerning the isomerization mechanism leading from the dihydroxo to the oxo aqua species, the direct intramolecular proton transfer from the dihydroxo to the aqua complex can be ruled out due to the high calculated barrier for the direct proton transfer, viz. $43.9 \text{ kcal mol}^{-1}$ including solvent effects. The assistance of either one or two water molecules renders proton transfer feasible by dramatically lowering the barrier height. The mechanism involving assistance by two water molecules even yields a lower energy for the transition state than for the reactant and product when including the CPCM, suggesting that this is indeed a very facile process. The assistance by amphiphilic external molecules, such as water, is increasingly found to be crucial to lower activation barriers of intramolecular proton transfer rearrangements.

Acknowledgement. We gratefully acknowledge the European Commission for funding this work through the AQUACHEM Research Training Network (Contract n° MRTN-CT-2003-503864). Financial support from the Spanish MEC through Project CTQ2005-09000-C02-01 and from Generalitat de Catalunya through grant 2005SGR00896 is gratefully acknowledged. A.C.-V. acknowledges the Spanish MEC for a FPU fellowship.

Supporting information available. Complete references 19, 51 and 56, and absolute energies and cartesian coordinates of the optimized structures. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Experimental Section

Solution Preparation and Measurements. All chemicals used in this study were of analytical grade. The solution pH was controlled by using HNO₃. The total ionic strength (μ) was kept constant at 1 M with NaNO₃. pH measurements were carried out on a Metrohm 623 pH meter equipped with a Sigma glass electrode. UV-vis spectra were recorded in gastight cuvettes on a Shimadzu UV-2100 spectrophotometer equipped with a thermostated cell compartment CDS-260.

Stopped-flow kinetic measurements on the acidification [Cp*MoO₃][−] with HNO₃ in the pH range from 0.025 to 0.35 were carried out using an Applied Photophysics SX-18MV stopped-flow spectrophotometer. 20% MeOH–H₂O solutions of [Cp*MoO₃][−] were rapidly mixed with varying pH solution in a gas-tight syringe. The kinetics of the reaction was monitored at 390 nm where the change in absorbance is a maximum for the acidified molybdenum product. The rate constant for acidification was determined from the slope of linear plots of k_{obs} versus [H⁺], as described in more detail under Results and Discussion. All kinetic experiments were performed under pseudo-first order conditions, i.e., with at least

10-fold excess of $[H^+]$ over the $Cp^*_2Mo_2O_5$ complex. Reported rate constants are the mean values of at least five kinetics runs, and the quoted uncertainties are based on the standard deviation. High-pressure stopped-flow studies were performed on a custom-built instrument (from 10 to 130 MPa).⁵³ Kinetic traces were recorded on an IBM-compatible computer and analyzed with the use of OLIS KINFIT (Bogart, GA) set of programs.

Computational details. Calculations were carried out using the Gaussian 03 package⁵⁴ at the DFT level by means of the B3LYP functional.⁵⁵⁻⁵⁷ For the Mo atom, the LANL2DZ pseudopotential⁵⁸ was used with the addition of f polarisation functions.⁵⁹ The 6-31G(d) basis set was used for C atoms whether for O atoms additional diffuse functions were added due to their anionic character (6-31+G(d)). For the hydrogens, the 6-31G(d,p) basis set was employed. IRC calculations were made in order to get the two minima linked by every transition-state.⁶⁰⁻⁶²

Solvent effects were included by means of CPCM single point calculations.^{63, 64} Additional spheres were included for all the hydrogens except for the Cp^* hydrogens by means of the SPHEREONH option. Frequency calculations were made in order to get the free Gibbs energy. The temperature used was 298.15 K.

References

- ¹ Cruywagen, J. J.; Heyns, J. B. B.; Rohwer, E. F. C. H., *J. Inorg. Nucl. Chem.* **1976**, 38, 2033-2036.
- ² Cruywagen, J. J.; Draaijer, A. G.; Rypstra, T., *South African Journal of Chemistry* **1988**, 41, 89-96.
- ³ Castro-Garcia, S.; Pecquenard, B.; Bender, A.; Livage, J.; Julien, C., *Ionics* **1997**, 3, 104-109.
- ⁴ Cruywagen, J. J.; Draaijer, A. G.; Heyns, J. B. B.; Rohwer, E. A., *Inorg. Chim. Acta* **2002**, 331, 322-329.
- ⁵ Paffett, M. T.; Anson, F. C., *Inorg. Chem.* **1981**, 20, 3967-72.
- ⁶ Cruywagen, J. J., *Adv. Inorg. Chem.* **2000**, 49, 127-182.
- ⁷ Richens, D. T., *Chem. Rev.* **2005**, 105, 1961-2002.
- ⁸ Bortolin, R.; Patel, V.; Munday, I.; Taulor, N. J.; Carty, A. J., *J. Chem. Soc., Chem. Commun.* **1985**, 456-458.
- ⁹ Carney, M. J.; Walsh, P. J.; Bergman, R. G., *J. Am. Chem. Soc.* **1990**, 112, 6426-6428.
- ¹⁰ Parkin, G.; Bercaw, J. E., *Polyhedron* **1988**, 7, 2053-2082.

Fettinger, J. C.; Kraatz, H.-B.; Poli, R.; Quadrelli, E. A., *J. Chem. Soc., Dalton Trans.* **1999**, 497-508.

Morales, D.; Pleune, B.; Poli, R.; Richard, P., *J. Organometal. Chem.* **2000**, 596, 64-69.

Dadci, L.; Elias, H.; Frey, U.; Hoernig, A.; Koelle, U.; Merbach, A. E.; Paulus, H.; Schneider, J. S., *Inorg. Chem.* **1995**, 34, 306-15.

Helm, L.; Merbach, A. E., *Chem. Rev.* **2005**, 105, 1923-1959.

Poli, R., *Chem. Eur. J.* **2004**, 10, 332-341.

Reis, P. M.; Romão, C. C.; Royo, B., *J. Chem. Soc., Dalton Trans.* **2006**, 1842-1846.

Abrantes, M.; Santos, A.; Mink, J.; Kühn, F.; Romão, C., *Organometallics* **2003**, 22, 2112-2118.

Zhao, J.; Santos, A. M.; Herdtweck, E.; Kühn, F. E., *J. Mol. Catal. A* **2004**, 222, 265-271.

Martins, A. M., *et al.*, *Organometallics* **2005**, 24, 2582-2589.

Zhao, J.; Herdtweck, E.; Kühn, F. E., *J. Organomet. Chem.* **2006**, 691, 2199-2206.

Collange, E.; Garcia, J.; Poli, R., *New J. Chem.* **2002**, 26, 1249-1256.

Collange, E.; Metteau, L.; Richard, P.; Poli, R., *Polyhedron* **2004**, 23, 2605-2610.

Hofmann, P., *Angew. Chem., Int. Ed. Engl.* **1977**, 16, 536-537.

Schilling, B. E. R.; Hoffmann, R.; Lichtenberger, D. L., *J. Am. Chem. Soc.* **1979**, 101, 585-591.

Schilling, B. E. R.; Hoffmann, R.; Faller, J. W., *J. Am. Chem. Soc.* **1979**, 101, 592-598.

Johnson, T. J.; Folting, K.; Streib, W. E.; Martin, J. D.; Huffman, J. C.; Jackson, S. A.; Eisenstein, O.; Caulton, K. G., *Inorg. Chem.* **1995**, 34, 488-499.

Ward, T. R.; Schafer, O.; Daul, C.; Hofmann, P., *Organometallics* **1997**, 16, 3207-3215.

Smith, K. M.; Poli, R.; Legzdins, P., *Chem. Commun.* **1998**, 1903-1904.

Smith, K. M.; Poli, R.; Legzdins, P., *Chem. Eur. J.* **1999**, 5, 1598-1608.

Perrin, L.; Maron, L.; Eisenstein, O., *Faraday Disc.* **2003**, 124, 25-39.

Han, Y.; Harlan, C. J.; Stoessel, P.; Frost, B. J.; Norton, J. R.; Miller, S.; Bridgewater, B.; Xu, Q., *Inorg. Chem.* **2001**, 40, 2942-2952.

Kelly, C. P.; Cramer, C. J.; Truhlar, D. G., *Journal of Physical Chemistry B* **2006**, 110, 16066-16081.

Gun, J.; Modestov, A.; Lev, O.; Saurenz, D.; Vorotyntsev, M. A.; Poli, R., *Eur. J. Inorg. Chem.* **2003**, 482-492.

Gun, J.; Modestov, A.; Lev, O.; Poli, R., *Eur. J. Inorg. Chem.* **2003**, 2264-2272.

Wooster, N., *Nature (London, United Kingdom)* **1931**, 127, 93.

Krebs, B., *Acta Crystallogr., Sect. C: Struct. Crystallogr. Cryst. Chem.* **1972**, 28, 2222-31.

Lledós, A.; Bertrán, J., *Tetrahedron Lett.* **1981**, 22, 775-8.

Ventura, O. N.; Lledos, A.; Bonaccorsi, R.; Bertran, J.; Tomasi, J., *Theor. Chim. Acta* **1987**, 72, 175-95.

Lima, M. C. P.; Coutinho, K.; Canuto, S.; Rocha, W. R., *J. Phys. Chem. A* **2006**, 110, 7253-7261.

Bergquist, C.; Bridgewater, B. M.; Harlan, C. J.; Norton, J. R.; Friesner, R. A.; Parkin, G., *J. Am. Chem. Soc.* **2000**, 122, 10581-10590.

Prabhakar, R.; Blomberg, M. R. A.; Siegbahn, P. E. M., *Theoretical Chemistry Accounts* **2000**, 104, 461-470.

Sambrano, J. R.; Andres, J.; Gracia, L.; Safont, V. S.; Beltran, A., *Chem. Phys. Lett.* **2004**, 384, 56-62.

Hratchian, H. P.; Sonnenberg, J. L.; Hay, P. J.; Martin, R. L.; Bursten, B. E.; Schlegel, H. B., *J. Phys. Chem. A* **2005**, 109, 8579-8586.

Erikson, T. K. G.; Mayer, J. M., *Angew. Chem., Int. Ed. Engl.* **1988**, 27, 1527-1529.

Dovletoglou, A.; Meyer, T. J., *J. Am. Chem. Soc.* **1994**, 116, 215-23.

46 Comas-Vives, A.; Gonzalez-Arellano, C.; Corma, A.; Iglesias, M.; Sanchez, F.; Ujaque, G., *J. Am.*
Chem. Soc. **2006**, 128, 4756-4765.
 47 Tunon, I.; Silla, E.; Bertran, J., *J. Phys. Chem.* **1993**, 97, 5547-52.
 48 Wei, D.; Salahub, D. R., *J. Chem. Phys.* **1994**, 101, 7633-42.
 49 Marx, D.; Tuckerman, M. E.; Hutter, J.; Parrinello, M., *Nature (London)* **1999**, 397, 601-604.
 50 Schmitt, U. W.; Voth, G. A., *J. Chem. Phys.* **1999**, 111, 9361-9381.
 51 Kovács, G.; Ujaque, G.; Lledós, A.; Joó, F., *Organometallics* **2006**, 25, 862-872.
 52 Kovács, G.; Schubert, G.; Joó, F.; Papai, I., *Organometallics* **2005**, 24, 3059-3065.
 53 Van Eldik, R.; Gaede, W.; Wieland, S.; Kraft, J.; Spitzer, M.; Palmer, D. A., *Review of Scientific*
Instruments **1993**, 64, 1355-7.
 54 Frisch, M. J., *et al.*, *Gaussian 03, Revision C.02*. ed.; Gaussian, Inc.: Wallingford CT, 2004.
 55 Becke, A. D., *J. Chem. Phys.* **1993**, 98, 5648-5652.
 56 Lee, C. T.; Yang, W. T.; Parr, R. G., *Phys. Rev. B* **1988**, 37, 785-789.
 57 Stephens, P.; Devlin, F.; Chabalowski, C.; Frisch, M., *J. Phys. Chem.* **1994**, 98, 11623-11627.
 58 Hay, P. J.; Wadt, W. R., *J. Chem. Phys.* **1985**, 82, 270-283.
 59 Ehlers, A. W., *et al.*, *Chem. Phys. Lett.* **1993**, 208, 111-14.
 60 Fukui, K., *Acc. Chem. Res.* **1981**, 14, 363-8.
 61 Gonzalez, C.; Schlegel, H. B., *J. Chem. Phys.* **1989**, 90, 2154-61.
 62 Gonzalez, C.; Schlegel, H. B., *J. Phys. Chem.* **1990**, 94, 5523-7.
 63 Barone, V.; Cossi, M., *J. Phys. Chem. A* **1998**, 102, 1995-2001.
 64 Cossi, M.; Rega, N.; Scalmani, G.; Barone, V., *J. Comput. Chem.* **2003**, 24, 669-681.

Synopsis

Combined protonation kinetics and DFT studies bring to light the regulating factors for the relative energy of dihydroxo vs. oxo aqua isomers, the water dissociation equilibrium, and abnormal acid dissociation constants, for the organometallic “Cp*MoO₂(OH) + H⁺” system.

Table of Contents graphics