



**HAL**  
open science

## Role of the North Atlantic Oscillation in decadal temperature trends

Carley Iles, Gabriele Hegerl

► **To cite this version:**

Carley Iles, Gabriele Hegerl. Role of the North Atlantic Oscillation in decadal temperature trends. *Environmental Research Letters*, 2017, 12 (11), pp.114010. 10.1088/1748-9326/aa9152 . hal-03226698

**HAL Id: hal-03226698**

**<https://hal.science/hal-03226698>**

Submitted on 16 May 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

LETTER • OPEN ACCESS

## Role of the North Atlantic Oscillation in decadal temperature trends

To cite this article: Carley Iles and Gabriele Hegerl 2017 *Environ. Res. Lett.* **12** 114010

View the [article online](#) for updates and enhancements.

### Related content

- [Forcing of the wintertime atmospheric circulation by the multidecadal fluctuations of the North Atlantic ocean](#)  
Yannick Peings and Gudrun Magnusdottir
- [Winter Eurasian cooling linked with the Atlantic Multidecadal Oscillation](#)  
Dehai Luo, Yanan Chen, Aiguo Dai et al.
- [Tropical origin for the impacts of the Atlantic Multidecadal Variability on the Euro-Atlantic climate](#)  
Paolo Davini, Jost von Hardenberg and Susanna Corti

### Recent citations

- [Observed changes and trends in heatwave characteristics in Turkey since 1950](#)  
Ecmel Erlat *et al*
- [Evolution of observed and modelled temperatures in Finland in 1901–2018 and potential dynamical reasons for the differences](#)  
Kimmo Ruosteenoja and Jouni Räisänen
- [The Spatial Transformation Process and Critical Time Node Detection in Global Extreme High Temperature Clusters](#)  
Ting Zhang *et al*

# Environmental Research Letters



## LETTER

# Role of the North Atlantic Oscillation in decadal temperature trends

### OPEN ACCESS

RECEIVED  
22 June 2017

REVISED  
27 September 2017

ACCEPTED FOR PUBLICATION  
5 October 2017

PUBLISHED  
2 November 2017

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Carley Iles<sup>1,2,3</sup>  and Gabriele Hegerl<sup>1</sup>

<sup>1</sup> School of Geosciences, University of Edinburgh, Alexander Crum Brown Road, Edinburgh EH9 3FF, United Kingdom

<sup>2</sup> LSCE, IPSL, Université Paris-Saclay, Gif sur Yvette, France

<sup>3</sup> Author to whom any correspondence should be addressed.

E-mail: [carley.iles@lsce.ipsl.fr](mailto:carley.iles@lsce.ipsl.fr)

**Keywords:** North Atlantic Oscillation, climate variability, CMIP5, atmospheric circulation, decadal variability, temperature trends

Supplementary material for this article is available [online](#)

## Abstract


Global temperatures have undergone periods of enhanced warming and pauses over the last century, with greater variations at local scales due to internal variability of the climate system. Here we investigate the role of the North Atlantic Oscillation (NAO) in decadal temperature trends in the Northern Hemisphere for periods with large decadal NAO trends. Using a regression based technique we find a best estimate that trends in the NAO more than halved (reduced by 57%, 5%–95%: 47%–63%) the winter warming over the Northern Hemisphere extratropics (NH; 30N–90N) from 1920–1971 and account for 45% ( $\pm 14\%$ ) of the warming there from 1963–1995, with larger impacts on regional scales. Over the period leading into the so-called warming hiatus, 1989–2013, the NAO reduced NH winter warming to around one quarter (24%; 19%–31%) of what it would have been, and caused large negative regional trends, for example, in Northern Eurasia. Warming is more spatially uniform across the Northern Hemisphere after removing the NAO influence in winter, and agreement with multi-model mean simulated trends improves. The impact of the summer NAO is much weaker, but still discernible over Europe, North America and Greenland, with the downward trend in the summer NAO from 1988–2012 reducing warming by about a third in Northern Europe and a half in North America. A composite analysis using CMIP5 control runs suggests that the ocean response to prolonged NAO trends may increase the influence of decadal NAO trends compared to estimates based on interannual regressions, particularly in the Arctic. Results imply that the long-term NAO trends over the 20th century alternately masked or enhanced anthropogenic warming, and will continue to temporarily offset or enhance its effects in the future.

## 1. Introduction

Global temperatures have not warmed uniformly over the last century, instead undergoing periods of enhanced warming and pauses, with greater variability at regional scales. Such variations are expected due to internal variability of the climate system, for example through modes of atmospheric and ocean variability, including the El Niño Southern Oscillation, the Atlantic Multidecadal Oscillation, the Arctic Oscillation and atmospheric blocking (Hasselmann 1976, Hawkins and Sutton 2009, see Medhaug *et al* 2017). Atmospheric dynamics accounted for around 40% of the observed winter warming over the Northern Hemisphere

extratropical continents from 1965–2000 (Wallace *et al* 2012), and one third of the winter warming over North America over the last 50 years, with larger or even opposite contributions for some sub-regions (Deser *et al* 2016b). Variability of the annular modes is the main source of spread in climate projections up to 2060 in the mid and high latitudes in a large model ensemble (Deser *et al* 2012). Removing the effect of circulation reduces ensemble spread and improves agreement with observations (Wallace *et al* 2012, Deser *et al* 2016b).

The North Atlantic Oscillation (NAO) is the leading mode of atmospheric circulation variability over the North Atlantic sector, associated with the sea level pressure gradient between Iceland and the Azores. It



is related to the position of the jet stream and storm tracks, affecting temperature and precipitation patterns over Eurasia, North American and Greenland, particularly in boreal winter, with impacts on both ecosystems and human activities (Hurrell *et al* 2003 and references therein). The NAO varies on a wide range of timescales from days to multiple decades (e.g. Hurrell 1995, Hurrell *et al* 2003) and is closely related to the Northern Annular Mode (NAM) or Arctic Oscillation (AO), the leading mode of the entire Northern Hemispheric (NH) cold season circulation throughout the atmosphere (e.g. Thompson and Wallace 1998). The winter NAO underwent a multidecadal downward trend from around 1920–1970, an upward trend until the early 1990s and downward until 2013 (figure 1(a)). Spatial patterns of NH temperature trends during these periods resemble the appropriate phase of the NAO (Ostermeier and Wallace 2003 and Nagato and Tanaka 2012), but also show some resemblance with the pattern expected from global warming during the latter two periods. The upward trend in the NAM to the 1990s accounted for 30% of the winter warming over the NH extratropics, 50% of the warming over Eurasia and much of the cooling over eastern-Canada to Greenland from 1968–1997 (Thompson *et al* 2000, Hurrell 1996). Over the recent period of reduced warming, a substantial portion of the winter cooling over the NH continents (Cohen *et al* 2012, Saffioti *et al* 2015) and Europe (Saffioti *et al* 2016) can be accounted for by the NAO or AO, with improved agreement with CMIP5 simulated trends over Europe following dynamical adjustment (Saffioti *et al* 2016). Over the next 50 years, it is expected that multi-decadal NAO variations will cause a large spread in projections of temperature and precipitation over Europe, at least in the CESM large ensemble (Deser *et al* 2016a).

Here we systematically quantify the contribution of trends in the winter and summer NAO to decadal temperature trends in the Northern Hemisphere, following the method of Thompson *et al* (2000), but extended over the entire 20th century. We further remove the NAO contribution to trends, deriving a pattern that is closer to the direct thermodynamic effect of warming. We focus on individual periods of the NAO's maximum decadal trends and compare residual regional temperature changes to CMIP5 simulations of the last century. We then investigate whether the effect of long-term NAO trends may be enhanced via ocean processes (Deser *et al* 2016a, Delworth *et al* 2016, Delworth and Zeng 2016, Li *et al* 2013).

## 2. Data and methods

### 2.1. Observational data

The NAO is generally characterised either using the difference in normalised station based pressure measurements between Iceland and Portugal or the Azores, or as the first Empirical Orthogonal Function (EOF) of sea level pressure over the North Atlantic sector. We use the EOF based index since it captures the full spatial NAO pattern and the seasonal migration of its pressure centres better than the station based index and is less affected by noise induced by small-scale weather features (Hurrell *et al* 2003). The EOF based index for 1899 onwards was downloaded from the NCAR climate data guide (CDG) website for the extended winter season (DJFM), when the NAO is strongest, and for contrast the summer (JJA) when the NAO is weaker and shifted to the North-East, and European climate is less dominated by the large scale circulation (Bladé *et al* 2012, Hurrell *et al* 2003). EOFs are calculated for the region

20 to 80°N and 90°W to 40°E across the whole time period. Some studies use the high summer index (JA) instead of JJA since the July and August NAO are more closely related to each other than that in June (Folland *et al* 2009, Bladé *et al* 2012); but we found little difference when basing our analysis on a calculated JA EOF-based index instead (not shown). We also tested sensitivity to using station-based indices, downloaded from NCAR CGD. These are based on stations from Iceland and Portugal for DJFM, and Iceland and the Azores for JJA. In winter results were very similar to those using the EOF based indices. While there were some differences in summer (figure 1), the effect of the summer NAO (SNAO) on temperature trends remained small.

We used the HadCRUT4.4 observed surface temperature dataset, which contains monthly temperature anomalies from 1850 to present relative to 1961–1990 climatology on a 5° latitude-longitude grid (Morice *et al* 2012). These are based on station measurements of surface air temperature (SAT) over land and ship and buoy measurements of sea surface temperatures over ocean, with gridboxes not containing data labelled as missing. To calculate seasonal data, we required all months in a given season to be present. Observed sea level pressure (SLP) data is taken from HadSLP2, which contains monthly values on a 5° latitude-longitude grid from 1850–2004. Data are spatially interpolated to give globally complete coverage (Allan and Ansell 2006).

## 2.2. Climate model data

Observational results are compared to climate model simulations from the CMIP5 (Coupled Model Inter-comparison Project Phase 5) archive (Taylor *et al* 2012). SAT from historical ‘ALL’ forcing runs is used for the period 1850–2005, extended to the present by appending the first few years of the appropriate Representative Concentration Pathway 4.5 (RCP 4.5) simulation, which reaches a radiative forcing of 4.5  $\text{Wm}^{-2}$  by 2100 relative to preindustrial conditions (results should not be sensitive to scenario over this short timeframe considered; Kirtman *et al* 2013). The historical ‘ALL’ runs are forced with observed records of greenhouse gases, anthropogenic aerosols, volcanic aerosols, solar variability, land-use and ozone. Note that over the most recent period, the assumed forcing may not realistically capture low-level volcanic and solar forcing (e.g. Kaufmann *et al* 2011, Santer *et al* 2014, 2015) and hence slightly overestimate recent trends. We select all available model runs that have data for both the historical and RCP4.5 periods (38 models, 108 runs; see table S1). Model data are regridded to the grid of HadCRUT4.4, with missing values in observations masked in simulations to replicate the same coverage.

In order to investigate longer-term impacts of NAO trends on temperature (see methods and S1) we use SAT and SLP data from CMIP5 control runs, which use forcings fixed at preindustrial levels (table S1). Data are regridded to a common 2.5° grid.


## 2.3. Methods

To calculate the influence of the NAO on SAT we regress the standardised time series of the NAO index for a given season on the temperature time series for the same season for each grid cell for the period 1900–2013, using detrended data. This yields a regression coefficient for each grid cell showing the change in temperature per unit change in the NAO (figures 2(c) and (d)), and its significance levels assuming white noise. A requirement of at least 30 years of data per grid cell was used. The same procedure was applied to sea-level pressure, but based on the period 1900–2004, when the dataset based on consistent methods ends (figures 2(a) and (b)).

In order to examine the impact of the NAO on decadal temperature trends, we first select periods of the strongest observed NAO trends of at least 20 years for both seasons. These are shown in figures 1(a) and (b) and figure S1 and are discussed below. For each year the NAO contribution to temperature is calculated for each grid cell by multiplying the NAO index value by the regression coefficients calculated above. The NAO contribution is then subtracted from the raw temperature anomalies to calculate residual anomalies for each year. Temperature trends are calculated for each grid cell for the raw temperature, residual temperature, and NAO related temperature, and are used if 80% of data are available over a given time period.

NH extratropical (30–90 N) and regional temperature time series with and without NAO effects are calculated, using the regions shown in figures 2(c) and (d) and table S2 (see also figure S5). The regions are selected to illustrate the maximum possible impact of the NAO on temperature trends based on the interannual regressions above. The contribution of the NAO is calculated based on a regression of the detrended regional mean temperature time series against the detrended NAO index. Calculating trends and NAO contributions on a grid cell basis first and then regional averages gives very similar results (figure S9). Our method assumes a symmetrical temperature response between positive and negative NAO states. Whilst not strictly true (Hurrell *et al* 2003, Heape *et al* 2013), figure S8, which is based on a composite of NAO trend periods of either sign in CMIP5 simulations, suggests any asymmetry is small and unlikely to substantially affect our findings.

Calculating the effect of decadal NAO trends on temperature based on interannual regressions may miss any further response caused by the ocean integrating short-term atmospheric variations, e.g. in temperature and wind forcing (Hasselmann 1976, Thompson 2015), which can enhance the NAO response (Deser *et al* 2016a). Therefore, we also examine the effect of decadal NAO trends in CMIP5 model control runs. We compare CMIP5 results obtained by applying the method detailed above to results obtained by compositing temperature trends over multiple NAO trend periods and multiple model



runs, which should capture longer-term aspects of the response (see supplementary methods available at [stacks.iop.org/ERL/12/114010/mmedia](http://stacks.iop.org/ERL/12/114010/mmedia)). We examine NAO trends of varying lengths (11, 16, 21, 25, 31 and 51 years) that cover the range of periods examined in the first part of our analysis. Significance of the difference in results using the two techniques is evaluated using a two sample student t-test that does not assume equal variances.

#### 4. Results

Figures 1(a) and (b) show the time series of the winter and summer NAO, for both the EOF and station-based indices. In winter, the two indices are very similar, but in summer there is a noticeable difference, including in the overall trend in some decades, particularly the 1960s to 1980s. This difference may arise because the stations used do not coincide well with the pressure centres of the SNAO, which is shifted to the north-west of its winter counterpart. Therefore the following analysis focuses on the EOF-based index. The periods of decadal NAO trends analysed are shown as trend lines. These are 1920–1971, 1963–1995 and 1989–2013 for winter,

during which periods the NAO index shows a trend of  $-1.3$ ,  $+2.4$  and  $-2.3$  standard deviations respectively, and 1902–1942, 1936–1960, 1957–1984 and 1988–2012 for summer, with a change of  $+1.9$ ,  $-1.2$ ,  $+1.4$  and  $-2.5$  standard deviations over time respectively. These are individual periods of maximum NAO trends that overlap slightly. Results for non-overlapping periods are similar, although NAO trends are weaker for the most recent period (see figure S3 for DJFM).

Figure 2 shows the interannual regression of the NAO indices on sea level pressure and temperature for both seasons. The effect of the NAO on seasonal temperatures and temperature trends arises due to the cumulative effects of its synoptic time scale variations. A positive phase of the NAO in winter is associated with anomalously low pressure over Iceland and the North Pole and high pressure over the Azores. This leads to a strengthened storm track angled towards the North-East which brings warm maritime air from the Atlantic over Northern Eurasia, causing a strong warming signal there (Hurrell *et al* 2003, Thompson and Wallace 2000). Over Eastern Canada and Greenland cold anomalies are associated with stronger than average northerly winds. A weak warming signal can also be seen over the United States and cooling over the Mediterranean and Middle





Figure 3. The contribution of the NAO to winter (DJFM) decadal temperature trends. Top row: raw temperature trends, middle row: the contribution of the NAO, bottom row: the residual temperature trend once the NAO contribution has been subtracted. Left hand column 1920–1971, middle column 1963–1995, right hand column 1989–2013. Stipples are the same those in figure 2 and indicate grid cells with a significant *interannual* relationship between the NAO and temperature. Units are °C per decade.

East, associated with stronger clockwise flow around the subtropical Atlantic high. These patterns, including the effects over Asia, agree with those found by previous studies (see Hurrell 1995, Hurrell *et al* 2003, Thompson and Wallace 1998, 2000, Thompson *et al* 2000, Deser *et al* 2016a). A negative NAO is associated with opposite pressure anomalies, leading to a more wavy jet stream and a weaker, more zonal storm track. This allows north-easterly anomalous winds to occur over Eurasia bringing cold temperatures to northern Europe. The NAO's effects can extend beyond the Atlantic region through its association with the Arctic Oscillation, and through interactions with other synoptic scale phenomena. For example, interactions with Ural Blocking, the Siberian high and the East Asian Monsoon are thought to extend its influence over Asia (Luo *et al* 2016, He *et al* 2017 and references therein), whilst interactions with European blocking allow its influence to reach the Middle East (Yao *et al* 2016). The SNAO is weaker, with north-eastward shifted pressure centres. It leads to weak temperature anomalies reminiscent of the winter NAO, with significant impacts largely limited to Northern Europe,

North America, and the North Atlantic (see Folland *et al* 2009, Bladé *et al* 2012).

For the time periods identified above we calculate the contribution of the NAO trends to decadal trends in temperature across the Northern Hemisphere (see methods). Figure 3 shows the spatial pattern of winter temperature trends for each period, the contribution from the NAO and the residual trends once the NAO has been subtracted. From 1920–1971 there was a cooling trend over northern Eurasia and the eastern United States, with warming elsewhere. The NAO downtrend explains the Eurasian cooling, and part of the North American cooling, with a more uniform warming signal following its removal. From 1963–1995, the upward NAO trend accounts for a large part of the strong warming trend over northern Eurasia and part of the cooling trend over Eastern Canada–Greenland (see also Thompson *et al* 2000). Finally, during the most recent period (1989–2013), most of the large cooling signal over Northern Eurasia and part of the warming signal over Eastern Canada–Greenland is accounted for by the concurrent NAO downtrend, with a more amorphous warming signal following NAO removal. In summer



**Figure 4.** Temperature trend (yellow), NAO contribution (blue) and residual trend once the NAO influence has been subtracted (red) for winter (DJFM) (a)–(f) and summer (JJA) (g)–(i) averaged across the regions shown in figure 2, for the time periods shown in figures 1(a) and (b). Also shown is the CMIP5 multi-model mean trend for each region (grey). 5%–95% confidence intervals for the NAO and residual reflect uncertainty in the regression of the NAO on regional mean temperature.


the contribution of NAO trends to temperature trends is much weaker, with only subtle changes in spatial patterns following NAO removal (figure S4).

The impact of NAO-induced temperature trends is strongest at regional scales. Figure 4 shows the contribution of the NAO to temperature trends for the regions shown in figure 2(c and d) (see methods). In winter, the NAO has a large influence on temperature trends in all of the regions examined. For most regions the downward trend of the NAO has a cooling effect from 1920–1971, contributing to a negative or small positive temperature trend overall, except for Eastern Canada-Greenland and the Mediterranean. From 1963–1995 the NAO makes a substantial contribution to the warming trend seen particularly over Northern Eurasia (contributing 82% of the warming ( $\pm 15\%$  based on 5%–95% uncertainty range of the NAO vs SAT regression coefficients) or  $0.65 (\pm 0.11) ^\circ\text{C decade}^{-1}$ ), but also over the USA and the Arctic ( $65 (\pm 23)\%$  and  $55 (\pm 28)\%$  respectively) (see also Thompson *et al* 2000). Over Eastern Canada-Greenland and the Mediterranean, the NAO uptrend causes a cooling trend over this period, with a positive residual trend once the NAO influence is removed. During the most recent period (1989–2013) the negative NAO trend caused a large cooling over northern Eurasia ( $-0.70 ^\circ\text{C decade}^{-1}$ ), with a weak residual warming trend of  $0.12 (\pm 0.15) ^\circ\text{C decade}^{-1}$  following its removal. The NAO reduced the warming trend in the USA and the Arctic over this same

period, whilst in Eastern Canada-Greenland and the Mediterranean it greatly enhanced the warming trend, doubling it in the former case, and tripling it in the latter. Averaged over the NH extratropics, the NAO reduced the winter warming trend to 43% (37%–53%) of what it would have been from 1920–1970, accounts for  $45 (\pm 14)\%$  of the warming signal from 1963–1995, consistent with Thompson *et al* (2000) and reduces it to 24% (19%–31%) of what might have occurred from 1989–2013. Overall, removing the NAO influence yields residual trends that are mostly positive and tend to increase over time (see figure 4). Comparing with the CMIP5 multi-model mean temperature trends over the same regions and time periods, removing the NAO influence from the observations improves agreement between the observed and model simulated trends in 14 out of 18 cases (6 regions over 3 time periods).

In summer, the contribution of NAO trends to temperature trends is smaller than in winter, and not noticeable if averaged over the NH extratropics (not shown). However, its effects are noticeable over North America and Europe, with opposite sign over Eastern Canada-Greenland (figures 4(g)–(i)). For example, the downward trend in the SNAO from 1988–2012 reduced warming by 32% (13%–44%) in Northern Europe and 48% (31%–58%) in North America, and enhanced it in Eastern Canada-Greenland by 58% (32%–97%). Removing the NAO influence improves agreement






of observed temperature trends with CMIP5 multi-model mean simulated trends in 9 of 12 cases. Unlike in winter, the residual trends are not consistently positive or growing over time, suggesting that other factors are at play, including possibly aerosol forcing (Boucher *et al* 2013)

The analysis is designed to explore the maximum influence that the NAO has had over selected periods of the last century. Therefore, the NAO contributions are smaller if examining other periods. Figure 5, left column shows the contribution of the NAO to the recent warming hiatus period (1998–2013). The hiatus was most pronounced over northern extratropical regions in winter and figure 5 shows that the NAO accounts for part of the winter cooling trend in these regions. Since the NAO was preferentially in a low phase during this period, the trend starting in 1989

captures the NAO induced winter cooling into the hiatus period best (figure 3). Figure S6 shows that during the hiatus itself, the remaining NAO trend accounts for 81 ( $\pm 14$ )% of the cooling over northern Eurasia, 77 ( $\pm 16$ )% of the warming over Canada-Greenland and 87 ( $\pm 27$ )% of the cooling over the NH extratropics. Residual trends are still negative across northern Eurasia and the NH extratropics, in contrast to the multi-model mean, which shows a strong warming trend. Other factors contributing to the cooling in NH boreal winter may include positive pressure anomalies over Eurasia (Deser *et al* 2017), including an increased frequency of Ural Blocking (Luo *et al* 2016), with global temperatures influenced by increased Pacific Ocean heat uptake, AMO variability, the recent solar minimum and low level volcanism (e.g. Flato *et al* 2013 and references therein).



Many detection and attribution studies are based on the period 1950–2005 (Bindoff *et al* 2013). While the global contribution of the NAO to winter temperature trends over this period is negligible (4.8 ( $\pm 7$ )% of the observed warming; figure S6), its effect is noticeable on regional scales (figure 5, right column, S6), contributing 25 ( $\pm 8$ )% of the warming over the NH Extratropics and 54 ( $\pm 10$ )% of the warming over Northern Eurasia. This suggests that the best estimate of the human contribution from detection and attribution over those regions may be overestimated in winter, but as detection and attribution methods account for climate variability (Hegerl *et al* 1997, 2007, Tett *et al* 1999), uncertainty ranges should include the true value. In Eastern Canada–Greenland, the NAO has masked a positive trend (figure S6). Removing the NAO influence brings observed trends closer to the multi-model mean trend (figure S6). Extending the analysis to 2013 gives similar results although both raw temperature trends and the contribution of the NAO are weaker (not shown).

### 5. Longer term temperature responses

The analysis so far has been based on the NAO–temperature relationship derived from interannual regressions. However, this approach does not capture

the effect of long-term NAO trends on the ocean. For instance, Delworth and Zeng (2016) found using the GFDL climate model that a positive NAO trend of longer than 10 years can strengthen the Atlantic Meridional Overturning Circulation (AMOC). Cold, strong westerlies extract heat from the subpolar gyre, increasing deep water formation and speeding up gyre circulation. The resulting increased AMOC circulation warms the extratropics with a delay of about 10 years from the initial NAO forcing, amplified by melting sea ice and snow-related albedo feedbacks, and increased heat flux from ice-free oceans. These effects increase for longer NAO trends, and are opposite for a negative NAO trend.

We use CMIP5 control runs to test whether the long-term NAO trends analysed here might be amplified by the ocean response (see also Deser *et al* 2016a). We examine the temperature change for a one standard deviation NAO trend, comparing the interannual regression technique to results from a composite technique based on control simulations (see methods and SI). The difference between the two techniques is shown in figure 6 for 16, 31 and 51 year trend lengths (for other periods see figure S7), and the absolute values are shown in figure S8.

The composite method results in a larger warming trend, particularly over North-East Russia and the Barents Sea compared to the interannual regression

method for positive NAO trends (cooling for downwards NAO trends), which gets stronger, increases in area and becomes more significant as the trend length increases. For 51 year trends the signal extends over much of the Northern Hemisphere, but is strongest in the Barents Sea. A classic tripole pattern response to the NAO can be seen over the North Atlantic for most periods (e.g. Visbeck *et al* 2003) (note also a small response in the Pacific). The Atlantic response is consistent with the findings of Delworth and Zeng (2016) and Delworth *et al* (2016), and the strengthened warming over the Eurasian high latitudes with Deser *et al* (2016a). These results imply that the contribution of the NAO to twentieth century temperature trends is likely to be somewhat larger than that calculated through the interannual regression technique, particularly in the Barents Sea and high latitudes of Eurasia.

Using the composite technique enhances the temperature response particularly in the Arctic (by around 60%–90%), but also in Northern Eurasia (14%–25%), Eastern Canada–Greenland (5%–17%) and the NH extratropics as a whole (by 26%–46%) (figure S9; table S3). However, the limited observational coverage in the Arctic does not capture such an effect well (compare figures 3, 6). When explicitly removing the NAO effect using the response of control run-based trends, this tends to overcompensate rather than improve agreement with multi-model mean CMIP5 simulated historical temperature trends in around half of cases (not shown); although results are difficult to interpret given the limited data coverage for the ocean effect.

## 6. Conclusions

Decadal trends in the NAO have had a large effect on NH extratropical winter temperature trends over various periods since 1900. The NAO reduced warming over the NH extratropics by 57% (47%–63%) from 1920–1971, accounted for 45 ( $\pm 14$ )% of the warming there from 1963–1995 and reduced warming to 24% (19%–31%) of what it would have been from 1989–2013. The largest effects occur in Northern Eurasia, with opposite effects in Eastern Canada to Greenland. After removing trends explained by the NAO, the residual shows a more amorphous pattern of warming in all periods considered, with observed trends in better agreement with CMIP5 multi-model mean trends in most cases. In summer, the NAO influence is weaker, but made a notable contribution to temperature trends in northern North America, Europe and Eastern Canada–Greenland. The NAO also explains some of the boreal winter cooling trend over northern Eurasia and the NH extratropics over the ‘hiatus’ period itself, although observed residual trends still disagree with the warming simulated by CMIP5 models during this period.

Our method of removing the NAO is based on the temperature response to interannual NAO variations.

A composite method analysing the effect of multi-decadal NAO trends in CMIP5 control runs yields a stronger temperature response, particularly over the Barents Sea and high latitude Eurasia, and more so for longer NAO trend periods. This is in agreement with previous studies that suggest an influence of the NAO on temperature via changes in the AMOC and sea-ice and snow feedbacks. However, taking into account these longer-term effects does not consistently improve agreement with multi-model mean CMIP5 simulated temperature trends, either due to further variability, missing data in key regions, or errors in the model response to NAO trends.

In conclusion, long-term NAO trends have had a strong effect on winter temperature trends during the 20th century, particularly the most recent ones, and modest effects on some regional summer trends. These results highlight the need to better understand and predict circulation responses, such as that of the NAO, to forcing. Even due to variability alone, our findings highlight long-term uncertainty in future temperature trends.

## Acknowledgments

The authors thank David Thompson and Vikki Thompson nee Frith for discussions. We acknowledge the World Climate Research Programme’s Working Group on Coupled Modelling, which is responsible for CMIP, and we thank the climate modelling groups for producing and making available their model output. For CMIP the US Department of Energy’s Program for Climate Model Diagnosis and Intercomparison provides coordinating support and led development of software infrastructure in partnership with the Global Organization for Earth System Science. We would also like to thank two anonymous reviewers for their helpful comments and suggestions. CI and GH are supported by the ERC funded project TITAN (EC-320691). GH was supported by NCAS and the Wolfson Foundation and the Royal Society as a Royal Society Wolfson Research Merit Award (WM130060) holder.

## ORCID iDS

Carley Iles  <https://orcid.org/0000-0003-3504-2107>

## References

- Allan R and Ansell T 2006 A new globally complete monthly historical gridded mean sea level pressure dataset (HadSLP2): 1850–2004 *J. Clim.* **19** 5816–42
- Bladé I *et al* 2012 Observed and simulated impacts of the summer NAO in Europe: implications for projected drying in the mediterranean region *Clim. Dyn.* **39** 709–27
- Bindoff N *et al* 2013 Detection and attribution: from global to regional *Climate Change, 2013. Contribution of Working Group 1 to the Fifth Assessment report of the Intergovernmental Panel on Climate Change* ed T F Stocker *et al* (Cambridge: Cambridge University Press)

- Boucher O *et al* 2013 Clouds and aerosols *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed T F Stocker *et al* (Cambridge: Cambridge University Press)
- Cohen J L *et al* 2012 Arctic warming, increasing snow cover and widespread boreal winter cooling *Environ. Res. Lett.* **7** 14007
- Delworth T L *et al* 2016 The North Atlantic Oscillation as a driver of rapid climate change in the Northern hemisphere *Nat. Geosci.* **9** 509–12
- Delworth T L and Zeng F 2016 The impact of the North Atlantic Oscillation on climate through its influence on the Atlantic meridional overturning circulation *J. Clim.* **29** 941–62
- Deser C *et al* 2012 Uncertainty in climate change projections: the role of internal variability *Clim. Dyn.* **38** 527–46
- Deser C, Hurrell J W and Phillips A S 2016a The role of the North Atlantic Oscillation in European climate projections *Clim. Dyn.* (<https://doi.org/10.1007/s00382-016-3502-z>)
- Deser C, Terray L and Phillips A S 2016b Forced and internal components of winter air temperature trends over North America during the past 50 years: mechanisms and implications *J. Clim.* **29** 2237–58
- Deser C, Guo R and Lehner F 2017 The relative contributions of tropical pacific sea surface temperatures and atmospheric internal variability to the recent global warming hiatus *Geophys. Res. Lett.* **44** 7945–54
- Flato G *et al* 2013 Evaluation of climate models *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed T F Stocker *et al* (Cambridge: Cambridge University Press) pp 741–882
- Folland C K *et al* 2009 The summer North Atlantic Oscillation: past, present, and future *J. Clim.* **22** 1082–103
- Hasselmann K 1976 Stochastic climate models *Theory Tellus* **28** 473–85
- Hawkins E and Sutton R 2009 The potential to narrow uncertainty in regional climate predictions *Bull. Am. Meteorol. Soc.* **90** 1095–107
- He S *et al* 2017 Impact of Arctic Oscillation on the East Asian climate: a review *Earth Sci. Rev.* **164** 48–62
- Heape R, Hirschi J and Sinha B 2013 Asymmetric response of European pressure and temperature anomalies to NAO positive and NAO negative winters *Weather* **68** 73–80
- Hegerl G C *et al* 1997 Multi-fingerprint direction and attribution analysis of greenhouse gas, greenhouse gas-plus-aerosol and solar forced climate change *Clim. Dyn.* **14** 71–2
- Hegerl G C *et al* 2007 Understanding and attributing climate change *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* ed T F Stocker *et al* (Cambridge: Cambridge University Press)
- Hurrell J W *et al* 2003 An overview of the North Atlantic Oscillation *The North Atlantic Oscillation, Climatic Significance and Environmental Impact* ed J W Hurrell *et al* (Washington, DC: AGU Geophysical Monograph) pp 1–35
- Hurrell J W 1995 Decadal trends in the North Atlantic Oscillation: regional temperatures and precipitation *Science* **269** 676–9
- Hurrell J W 1996 Influence of variations in extratropical wintertime teleconnections on Northern hemisphere temperature *Geophys. Res. Lett.* **23** 665–8
- Kaufmann R K *et al* 2011 Reconciling anthropogenic climate change with observed temperature 1998–2008 *Proc. Natl Acad. Sci. USA* **108** 11790–3
- Kirtman B *et al* 2013 Near-term climate change: projections and predictability *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* T F Stocker *et al* (Cambridge: Cambridge University Press)
- Li J, Sun C and Jin F F 2013 NAO implicated as a predictor of Northern Hemisphere mean temperature multidecadal variability *Geophys. Res. Lett.* **40** 5497–502
- Luo D *et al* 2016 Impact of ural blocking on winter warm Arctic–cold Eurasian anomalies. Part I: blocking-induced amplification *J. Clim.* **29** 3925–47
- Medhaug A, Stolpe M B, Fischer E and Knutti R 2017 Reconciling controversies about the global warming hiatus *Nature* **545** 41–47
- Morice C P *et al* 2012 Quantifying uncertainties in global and regional temperature change using an ensemble of observational estimates: the HadCRUT4 data set *J. Geophys. Res. Atmos.* **117** 1–22
- Nagato Y and Tanaka H L 2012 Global warming trend without the contributions from decadal variability of the Arctic Oscillation *Polar Sci.* **6** 15–22
- Ostermeier G M and Wallace J M 2003 Trends in the North Atlantic Oscillation—Northern hemisphere annular mode during the twentieth century *J. Clim.* **16** 336–41
- Saffioti C *et al* 2016 Reconciling observed and modeled temperature and precipitation trends over Europe by adjusting for circulation variability *Geophys. Res. Lett.* **43** 8189–98
- Saffioti C, Fischer E M and Knutti R 2015 The contribution of atmospheric circulation and coverage bias to the warming hiatus *Geophys. Res. Lett.* **42** 2385–91
- Santer B D *et al* 2015 Observed multivariable signals of late 20th and early 21st century volcanic activity *Geophys. Res. Lett.* **42** 500–9
- Santer B D *et al* 2014 Volcanic contribution to decadal changes in tropospheric temperature *Nat. Geosci.* **7** 185–9
- Taylor K E, Stouffer R J and Meehl G A 2012 An Overview of CMIP5 and the experiment design *Bull. Am. Meteorol. Soc.* **93** 485–98
- Thompson D W J and Wallace J M 1998 The Arctic Oscillation signature in the wintertime geopotential height and temperature fields *Geophys. Res. Lett.* **25** 1297
- Thompson D W J and Wallace J M 2000 Annular modes in the extratropical circulation. Part I: month-to-month variability *J. Clim.* **13** 1000–1016
- Thompson D W J, Wallace J M and Hegerl G C 2000 Annular modes in the extratropical circulation. Part II: trends *J. Clim.* **13** 1018–1036
- Thompson V 2015 The early twentieth century warming *PhD Thesis* The University of Reading
- Tett S *et al* 1999 Causes of twentieth-century temperature change near the Earth's surface *Nature* **399** 569–72
- Visbeck M *et al* 2003 The ocean's response to North Atlantic Oscillation variability *The North Atlantic Oscillation, Climatic Significance and Environmental Impact* ed J W Hurrell *et al* (Washington, DC: AGU Geophysical Monograph) pp 113–45
- Wallace J M *et al* 2012 Simulated versus observed patterns of warming over the extratropical Northern hemisphere continents during the cold season *Proc. Natl Acad. Sci. USA* **109** 14337–42
- Yao Y, Luo D, Dai A and Feldstein S B 2016 The positive North Atlantic Oscillation with downstream blocking and middle East snowstorms impacts of the North Atlantic jet *J. Clim.* **29** 1853–76