Access to 12-Membered Cyclic ortho , meta -Diarylheptanoids: Total Synthesis of Actinidione via Isomyricanone

Paul Massé, Sabine Choppin, Lucia Chiumminto, Françoise Colobert, Gilles
Hanquet

► To cite this version:

Paul Massé, Sabine Choppin, Lucia Chiumminto, Françoise Colobert, Gilles Hanquet. Access to 12-Membered Cyclic ortho , meta -Diarylheptanoids: Total Synthesis of Actinidione via Isomyricanone. Journal of Organic Chemistry, 2021, 86 (3), pp.3033-3040. 10.1021/acs.joc.0c02489 . hal-03228806

HAL Id: hal-03228806

<https://hal.science/hal-03228806>

Submitted on 18 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Access to 12-Membered Cyclic *ortho,meta*-Diarylheptanoids: Total Synthesis of Actinidione *via* Isomyricanone

Paul Massé,[†] Sabine Choppin,^{*†} Lucia Chiumminto,[‡] Françoise Colobert,[†] Gilles Hanquet^{*†}

[†] Laboratoire d'Innovation Moléculaire et Applications (UMR CNRS 7042), Université de Strasbourg | Université de Haute Alsace, ECPM, 25 Rue Becquerel, 67087 Strasbourg, France

[‡] Department of Science, University of Basilicata, Via dell'Ateneo lucano, 10, 85100 Potenza, Italy

Supporting Information Placeholder

ABSTRACT: We describe herein the first access to 12-membered cyclic [7,0]*ortho,meta*-diarylheptanoids. The key features of the synthesis include both a Suzuki-Miyaura coupling and a ring closing metathesis. Actinidione, a promising natural product, along with a bioactive tetracyclic derivative were obtained in 14 steps for the first time from cheap commercially available substrates with an overall yield of 18-21%. Our *modus operandi* complies with the principles of the synthesis ideality by using notably strategic reactions.

Since the discovery of curcumin in 1815 by Vogel and Pelletier,¹ diarylheptanoids have been thoroughly investigated due to their unique architectural features as well as their broad range of biological activities.² Indeed, many approaches have been explored for the synthesis of the appealing family of cyclic 13-membered *meta,meta*-bridged biphenyls.³ However, to the best of our knowledge, no route scouting has been described to prepare 12-membered *ortho,meta*-bridged biphenyls.

The tetracyclic diarylheptanoid **1** represents an interesting compound of this family. A structure-activity relationship study revealed that **1**, synthesized by Dickey *et al.* and resulting from an acid-catalyzed cyclodehydration of myricanol **2** (Scheme 1A), exhibits a robust protein tau reduction activity, hence an anti-Alzheimer's disease effect, on a par with (-)-aS,11R-myricanol (EC₅₀ = 35 μM).⁴ This rearranged molecule is unexpected due to its isomyricanone-type biaryl axis but also by the presence of the tetralin moiety. The two enantiomers have been studied separately and have commensurate activity.

Additionally, isomyricanone **3** was the first 12-membered [7,0]*ortho,meta*-cyclophane, prepared by Whiting *et al.* by isomerization of natural myricanone **4** with boron trifluoride etherate.⁵ **3** unveils a potent inhibitory activity (IC₅₀ = 350 mol

ratio/TPA) against the Epstein-Barr virus early antigens, which induces lymphomas and carcinomas.⁶ Initially, only a migration of the alkyl chain on the aromatic cycle from *meta* to *ortho* position to release the ring strain was inferred (Scheme 1A, structure A). Nonetheless, Nagai *et al.* reported in 1991 the revision of the structure based on further NMR studies (see Supporting Information for the proposed mechanism).⁷

Furthermore, actinidione **5** was isolated in 2006 by Zhang *et al.* from leaves and twigs of *Actinidiaceae*, endemic to China (Scheme 1C).⁸ These species are employed as traditional medicine for hernia, hepatitis, hematemesis and rheumatic diseases.⁹ As far as we are aware, it is the first natural cyclic diarylheptanoid bearing both a [7,0]*ortho,meta*-cyclophane scaffold and a 1,4-benzoquinone. Its cytotoxicity has been shown to be promising against human breast cancer (Bre-04, GI₅₀ = 26.67 μM), lung cancer (Lu-06, GI₅₀ = 31.82 μM) and neuroma (Neu-04, GI₅₀ = 15.02 μM) cell lines. This compound was then isolated from the bark of *Myrica nana*, *Myrica Rubra* and *Myrica adenophora*.¹⁰ As might be expected from the 1,4-benzoquinone motif, actinidione demonstrates a strong antioxidant activity (IC₅₀ = 7.93 μM) against superoxide dismutase as well.^{10b}

This paper details a versatile access to a large range of *ortho,meta*-bridged diarylheptanoids via a key intermediate,

isomyricanone **3**, and exemplifies the preparation of **1** and **5**. This total synthesis was performed keeping in mind the outline of Scheme 1. Previous work in the literature and our retrosynthetic approach

lines of the ideality to prepare a natural product.¹¹ The latter has been

numerically expressed by Baran *et al.*^{11a} and graphically crafted by Christmann *et al.*^{11d} Hence, we envisaged a new retrosynthetic approach of **3** involving two evident disconnections and therefore two key reactions: a Suzuki- Miyaura coupling (SM coupling) along with a ring closing metathesis (RCM) between a type I and type II olefin for the macrocyclization step (Scheme 1C). Indeed, in the literature, only two examples for the formation of 12-membered macrocycles featuring a (hetero)biaryl scaffold have been described. The first one was illustrated by Kündig *et al.* who synthesized vertine, an alkaloid commonly known as cryogenine, by means of a RCM.¹² Macrocyclisation was enabled by the Hoveyda-Grubbs second-generation catalyst on the pre-conformationally prone to cyclize seco-precursor, which allowed to isolate vertine in 37% yield (Scheme 1B). In the second instance, Bristol-Myers Squibb has described the synthesis of a 12-membered macrocyclic compound containing a heterocycle via RCM using Grubbs second-generation catalyst in 71% yield (Scheme 1B).¹³ It is important to highlight that no cyclic diarylheptanoid has ever been prepared by RCM.

Our synthesis of isomyricanone **3**, the key precursor, begins with the preparation of the first coupling partner **7**, as depicted in Scheme 2. Thereupon, cheap 3,4,5-trimethoxybromobenzene **10** undergoes a Rieche formylation. Subsequently, successive AlCl_3 -promoted regioselective

demethylation in *ortho* position of the aldehyde,¹⁴ addition of the allyl Grignard reagent to build the alkyl chain and reduction of the resulting carbinol by triethylsilane were performed to afford **11** smoothly. Since a lithium-bromine exchange followed by insertion of a boron-containing electrophile ($\text{B}(\text{OMe})_3$ or $\text{B}(\text{O}i\text{Pr})_3$) did not allow us to afford the required boronic acid as pure product, we ended up using a one-pot procedure to isolate trifluoroborate salt **7** on exposure to potassium hydrogen bifluoride, in 81% yield.

Concurrently, the second building block **8** was quantitatively obtained in one step through iodination of the commercially available **9** with I_2 and silver sulfate. Notwithstanding, during scale up, we observed predominantly the formation of a by-product resulting from the rearrangement of the benzylic ether into the *ortho*-benzyl alkylated phenol.¹⁵ In the aftermath of this displacement, we eventually had to tweak the conditions by using NIS in acidic medium, that afforded **8** in quantitative yield, regardless of the scale.

With both coupling partners **7** and **8** in hand, obtained without column chromatography on multigram scale, the SM coupling proceeded in 35% yield in the presence of $\text{Pd}(\text{PPh}_3)_4$ and K_2CO_3 in water. We observed the formation of the corresponding carboxylic acid due to *in situ* saponification. To overcome this problem, we tuned the solvent ratio to dioxane/ H_2O (8/2)

in order to get the coupled product in 67% yield.¹⁶ Adding a ligand such as Xphos or SPhos led to the isomerization of the terminal olefin.¹⁷ The preparation of the Weinreb amide at 0 °C ensued, pending treatment with vinyl Grignard to give rise

to the terminal α,β -unsaturated ketone **6**, the seco-precursor of the RCM. RCM of **6** was performed using Grubbs second-generation

Scheme 2. Synthesis of isomyricanone **3** and isomyricanol **13**^a with its Molecular Structure Showing 50% Ellipsoids

^a Procedures: (a) Dichloromethyl methyl ether, TiCl_4 , CH_2Cl_2 , 0 °C to r.t. ; (b) AlCl_3 , CH_2Cl_2 , 15 °C ; (c) allylmagnesium bromide, THF, 0 °C ; (d) Et_3SiH , TFAA, CH_2Cl_2 , 0 °C then TBAF, THF ; (e) BnBr , NaH, NaI, DMF, 0 °C ; (f) $t\text{-BuLi}$, THF, -78 °C then $i\text{-PrOBpin}$, -78 °C to 40 °C then KHF_2 , H_2O , MeOH, r.t. ; (g) NIS, TFA, MeCN, r.t. ; (h) $\text{Pd(PPh}_3)_4$, K_2CO_3 , dioxane/ H_2O (8/2), 100 °C ; (i) $\text{CH}_3\text{NHOCH}_3\cdot\text{HCl}$, AlMe_3 , PhMe, 0 °C to r.t. ; (j) vinylmagnesium bromide, THF, 0 °C ; (k) Grubbs II, CH_2Cl_2 , 40 °C ; (l) H_2 (55 psi), Pd/C, AcOEt, r.t. ; (m) NaBH_4 , MeOH, r.t.

Scheme 3. Synthesis of **1** and **5** from **13**.

^a Procedures: (a) PTSA, PhCH_3 , 90 °C ; (b) salcomin, O_2 , DMF, r.t.

catalyst at reflux in DCM in high-dilution conditions. Pleasingly, the conversion of starting material was complete and only one addition of catalyst was required to reach expected macrocyclized synthon **12** in 85% yield. This implies that the ring strain is fully released in an *ortho,meta*-diarylheptanoid compared to a *meta,meta*-diarylheptanoid.¹⁸ Subsequently, an hydrogenation was achieved in order to saturate the heptylene chain and to deprotect the benzyl groups, giving prominence to pressure to ensure reproducibility. Finally, isomyricanone **3**, the 12-membered key intermediate was furnished in 12 steps with an overall yield of 33%.

At this stage, we envisaged the post-derivation of **3** to access other derivatives of interest. The first one to be supplied was

isomyricanol **13**, obtained as a pair of enantiomers, by reduction of **3** using sodium borohydride.¹⁰ The latter was recrystallized out of a mixture of ethyl acetate and chloroform at room temperature to afford a single crystal, giving rise to an X-ray structure.

Stemming from **13**, challenging tetracyclic diarylheptanoid **1** was obtained by acid-dehydration on the heptylene chain followed by electrophilic aromatic substitution as a pair of enantiomers (presumably *aR,R* and *aS,S*)⁴ in 79% yield after purification via semi-preparative HPLC (Scheme 3).

Scheme 4. Flowchart representation of the synthesis of **5**

The last envisaged post-derivation delivered the natural bioactive actinidione **5**. Indeed, this *ortho,meta*-diarylheptanoid was obtained by a regioselective phenol oxidation of **13** in a satisfying yield of 68%, by engaging salcomin under oxygen atmosphere in DMF (Scheme 3).¹⁹ To the best of our knowledge, no precedent synthesis was described in the literature. By using Frémy's catalyst,²⁰ we observed several products including one stemming from a dehydrogenation followed by an *ipso*-hydroxylation in *ortho* of the phenol on the southern aromatic ring.

In the attempt to optimize our synthesis of **3**, we tackled the step-economy by avoiding protecting groups at most, in order to attain a sufficient ideality, still having in mind the complexity of our targeted molecule. Our tactic focused likewise on optimizing each step to get a high overall yield, even on scalability. Furthermore, we set out to develop a modular approach to gain access to diverse *ortho,meta*-diarylheptanoids through a common intermediate, **3**. To that extent, we decided to follow the model of Christmann *et al.*, who recently promoted a color-coded flowchart to apprehend the ideality of a synthesis (Scheme 4).^{11d} Concerning our synthesis of **5**, out of 14 steps and an overall yield of 18%, it is unequivocal that most of the steps are green, to wit strategic. We exclusively account for three functional group interconversions (iodination, boronation, Weinreb amide preparation) and one required benzyl-protecting step. Accordingly, we have a high degree of ideality of 71%, which is calculated by the strategic steps out of the total number of steps.^{11a}

In conclusion, we undertook a novel, rapid, efficient and modular approach to the appealing family of 12-membered *ortho,meta*-diarylheptanoids via two key metal-mediated reactions (RCM and SM coupling). Attempts to prepare cyclic diarylheptanoids by RCM beforehand had failed.²¹ It is worth

noting that the key precursor **3** was obtained in 12 steps in 33% overall yield and late-stage diversification thereof gives access to promising compounds **1**, **5** and **13**. As an outcome, we have reached a high degree of ideality combined with a decent overall yield of 18% for the first synthesis of natural actinidione **5**.

EXPERIMENTAL SECTION

General Information All reactions were conducted under argon atmosphere unless otherwise noted and THF was distilled by standard technique prior to use. Commercially available reagents were used without additional purification, unless otherwise stated. **9** was bought from Fluorochem and **10** from TCI Chemicals. Hydrogenation reactions were performed using a QianCapTM Q-Tube-Purging-12-SS system pressure reactor. NMR analysis were recorded in CDCl₃ or CD₃CN. ¹H and proton-decoupled ¹³C NMR spectra were either recorded on Bruker AV 400 or Bruker AV 500. Chemical shifts are reported in parts per million (ppm) and are referenced to the residual solvent resonance as the internal standard (CDCl₃: δ [¹H] = 7.267 and accordingly δ [¹³C] = 77.16 ppm). Coupling constants (*J*) are given in Hz. IR spectra were recorded on Perkin Elmer Spectrum UATR two equipped with a diamond detector and an ATR unit. HRMS (Q-TOF) analysis were performed by the analytical facility at the University of Strasbourg. Crystal X-ray diffraction analyses were carried out by the Radiocrystallography Service of the University of Strasbourg by using a Bruker PHOTON III pixel detector diffractometer equipped with two micro-sources IμS Mo and IμS Diamond Cu. Chiral HPLC measurements were performed on a Shimadzu system with a quaternary low-pressure LC-20AD pump, an automatic SIL-20A HT injector, a CTO-10 AS oven and a SPD-M20 A diode array detector (DAD). The injection volume was 1 μL, the temperature of the oven set to 25°C and the concentration of the sample 1 g/L. Semi-preparative HPLC was performed with a Varian Prostar 210PDA apparatus equipped with a semi-preparative Daicel column (Chiralpak IA 20 mm x 250 mm, 5 μm) at 215 nm.

Preparation of 3-(benzyloxy)-1-bromo-2-(but-3-en-1-yl)-4,5-dimethoxybenzene (11) starting from (10).

6-bromo-2,3,4-trimethoxybenzaldehyde (14) A magnetically stirred solution of dichloromethyl methyl ether (2.05 equiv., 15 mL, 165.75 mmol) and TiCl₄ (2.25 equiv., 20 mL, 182.4 mmol) in CH₂Cl₂ (300 mL) was treated with a solution of commercially available **10** (1 equiv., 20 g, 80.94 mmol) in anhydrous CH₂Cl₂ (100 mL) at 0 °C. The red mixture was stirred at room temperature for 5 h. HCl (10%, 40 mL) was added at 0 °C, the stirring was continued for 15 min. The organic layer was then separated, and the aqueous phase extracted with CH₂Cl₂ (3x 200 mL). The combined extracts were washed with a saturated solution of NaHCO₃ (50 mL), brine, dried over MgSO₄, filtered off and concentrated under reduced pressure to afford **14** as a yellow solid. (22 g, 80.13 mmol, 99% yield) without further purification. Analytical data are consistent with those reported in the literature.²²

6-bromo-2-hydroxy-3,4-dimethoxybenzaldehyde (15) To a solution of **14** (1 equiv., 10 g, 36.35 mmol) in dry CH₂Cl₂ (300 mL) was added AlCl₃ (1 equiv., 4.85 g, 36.35 mmol) portionwise at 0 °C under inert atmosphere. After 1 h, another equivalent of AlCl₃ was added. The mixture was stirred for another 3 h below 15 °C. The flask was dived in an ice bath at 0 °C. The reaction mixture was then poured carefully into a mixture of ice and HCl (10%) over a few minutes, also dived in an ice bath. The mixture was stirred for 1 h. Aqueous phase was extracted with CH₂Cl₂ (3x 200 mL). The combined organic layer were washed successively with a saturated solution of NaHCO₃ (100 mL), brine, dried over anhydrous Na₂SO₄, filtered off through a pad of silica gel and concentrated under reduced pressure to afford **15** as a yellow oil (8.26 g, 31.62 mmol, 87% yield) without further purification. Analytical data are consistent with those reported in the literature.²³

3-bromo-2-(1-hydroxybut-3-en-1-yl)-5,6-dimethoxyphenol

(16) Allylmagnesium bromide (2.09 equiv., 1 M in Et₂O, 72 mL, 72 mmol) was added dropwise over 5 minutes to a stirred solution of **15** (1 equiv., 9 g, 34.47 mmol) in THF (170 mL) at 0 °C. The mixture was stirred at 0 °C for 30 minutes. The reaction mixture was quenched at 0 °C by addition of a saturated solution of NH₄Cl (150 mL). The aqueous phase was extracted with EtOAc (3x 200 mL). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered off and evaporated under reduced pressure. Crude was filtered through a pad of silica/celite/activated charcoal to afford **16** as a yellow oil (10.44 g, 34.45 mmol, 99% yield) without further purification; R_f 0.41 (SiO₂, ⁶Hex/EtOAc 7:3); ¹H NMR (500 MHz, CDCl₃) δ 8.33 (s, 1H), 6.58 (s, 1H), 5.88 (ddt, J = 17.2, 10.2, 7.1 Hz, 1H), 5.19 – 5.06 (m, 3H), 3.84 (s, 3H), 3.81 (s, 3H), 2.62 – 2.50 (m, 2H); ¹³C{¹H} NMR (125 MHz, CDCl₃) δ 152.3, 149.9, 136.0, 134.3, 120.3, 118.3, 115.8, 108.0, 74.7, 60.9, 56.1, 40.7; IR (cm⁻¹) ν 3389, 3078, 2935, 2841, 1603, 1571, 1493, 1444, 1417, 1301, 1244, 1123, 1045, 919, 811; HRMS (ESI+) m/z: [M + Na]⁺ calcd for C₁₂H₁₅⁷⁹BrO₄Na 325.0046; found 325.0068.

3-bromo-2-(but-3-en-1-yl)-5,6-dimethoxyphenol (17) Under inert atmosphere, **16** (1 equiv., 5.3 g, 17.52 mmol) was dissolved in CH₂Cl₂ (85 mL) and the mixture was cooled to 0 °C. Triethylsilane (1.13 equiv., 3.2 mL, 19.81 mmol) and trifluoroacetic anhydride (1.31 equiv., 3.2 mL, 23.0 mmol) were added. The reaction was stirred for 1.5 h at 0 °C. The reaction mixture was quenched with H₂O (10 mL) at 0 °C and the aqueous phase was extracted with CH₂Cl₂ (3x 100 mL). The combined organic layers were dried over Na₂SO₄, filtered off and evaporated under reduced pressure. The residue was dissolved in THF (105 mL) and TBAF (1.02 equiv., 1M in THF, 18 mL, 18 mmol) was added. The mixture was stirred for 30 minutes at room temperature. HCl (10%, 50 mL) was added at 0 °C. Et₂O was added and the organic phase was washed with HCl (3x 80 mL). Organic layer was washed with brine (50 mL), dried over Na₂SO₄, filtered off and the solvent was concentrated under reduced pressure to afford **17** as a yellow oil (4.63 g, 16.12 mmol, 92% yield) without any further purification. R_f 0.36 (SiO₂, ⁶Hex/EtOAc 9:1). ¹H NMR (500 MHz, CDCl₃) δ 6.69 (s, 1H), 5.97 – 5.84 (m, 2H), 5.11 – 4.91 (m, 2H), 3.89 (s, 3H), 3.84 (s, 3H), 2.88 – 2.77 (m, 2H), 2.32 – 2.27 (m, 2H). ¹³C{¹H} NMR (125 MHz, CDCl₃) δ

150.6, 148.0, 138.4, 134.7, 121.2, 118.8, 114.7, 108.0, 61.1, 56.1, 33.0, 29.3. IR (cm⁻¹) ν 3495, 3071, 2998, 2935, 2853, 1639, 1576, 1607, 1493, 1454, 1421, 1315, 1237, 1197, 1123, 1043, 911, 797. HRMS (ESI-) m/z: [M - H]⁺ calcd for C₁₂H₁₄⁷⁹BrO₃ 285.0132; found 285.0130.

3-(benzyloxy)-1-bromo-2-(but-3-en-1-yl)-4,5-

dimethoxybenzene (11) To a solution of **17** (1 equiv., 4.25 g, 14.8 mmol) in DMF (60 mL) was added NaH (1.2 equiv., 710 mg, 17.82 mmol) at 0 °C. After 10 minutes of stirring, benzyl bromide (1.01 equiv., 1.77 mL, 14.8 mmol) and NaI (9 mol%, 200 mg, 1.32 mmol) were added. The reaction mixture was stirred at room temperature for 1.5 h. A saturated solution of NH₄Cl (25 mL) was added slowly to quench the reaction at 0 °C. The aqueous phase was extracted with Et₂O (3x 100 mL). Combined organic layers were washed with cold water (3x 250 mL), dried over Na₂SO₄, filtered off and concentrated under reduced pressure. Crude was purified by flash chromatography on silica gel (gradient was performed from 100/0, V/V cyclohexane/ethyl acetate to 8/2, V/V) to afford **11** (5.02 g, 13.32 mmol, 90% yield) as a yellow oil. R_f 0.56 (SiO₂, ⁶Hex/EtOAc 9:1). ¹H NMR (400 MHz, CDCl₃) δ 7.49 – 7.44 (m, 2H), 7.43 – 7.32 (m, 3H), 6.90 (s, 1H), 5.87 (ddt, J = 16.9, 10.2, 6.7 Hz, 1H), 5.07 (s, 2H), 5.05 – 4.93 (m, 2H), 3.86 (s, 6H), 2.82 – 2.73 (m, 2H), 2.28 – 2.19 (m, 2H); ¹³C{¹H} NMR (125 MHz, CDCl₃) δ 152.2, 151.5, 142.1, 138.3, 137.7, 128.6, 128.4, 128.1, 128.1, 118.3, 114.8, 112.1, 75.5, 61.0, 56.3, 33.7, 29.9; IR (cm⁻¹) ν 3065, 3031, 2964, 2934, 2841, 1639, 1589, 1481, 1444, 1429, 1412, 1367, 1284, 1231, 1212, 1120, 1091, 1045, 996, 909, 795, 734, 696; HRMS (ESI+) m/z: [M + K]⁺ calcd for C₁₉H₂₁⁷⁹BrO₃K 415.0306; found 415.0305.

Potassium

(3-(benzyloxy)-2-(but-3-en-1-yl)-4,5-dimethoxyphenyl)trifluoroborate (7) To a solution of **11** (1 equiv., 4.42 g, 11.72 mmol) in THF (200 mL) was added *t*-BuLi (2.02 equiv., 1.7 M in pentane, 13.7 mL, 23.37 mmol) dropwise at -78 °C under argon. It was stirred for 30 minutes at -78 °C. Then, distilled 2-Isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (5 equiv., 0.25 mL, 1.23 mmol) was added quickly at -78 °C. Dry ice bath was removed, and the reaction medium was stirred at room temperature for 4 h. Solvent was evaporated under reduced pressure. The residue was taken up in CH₃OH (120 mL) and a sonicated aqueous solution of potassium hydrogen fluoride (5.57 equiv., 5 g KHF₂, 4.5 M in H₂O, 14.5 mL H₂O, 65.25 mmol) was added at room temperature. The reaction was stirred for 1.5 h minutes at room temperature. PhCH₃ (3x 50 mL) was added, and the mixture was concentrated under reduced pressure to eliminate water and to afford a yellowish solid. Acetone was added and the mixture was filtered off to eliminate inorganic salts. The filtrate was evaporated to afford a yellowish solid. Hexane (3x 20 mL) was then added and supernatant was removed. White powder was washed thrice with Et₂O to afford **7** as a white solid (3.84 g, 9.49 mmol, 81% yield) without any further purification. ¹H NMR (500 MHz, CD₃CN) δ 7.52 (d, J = 6.9 Hz, 2H), 7.43 – 7.30 (m, 3H), 6.95 (s, 1H), 5.91 (ddt, J = 16.9, 10.2, 6.5 Hz, 1H), 4.98 (s, 2H), 4.98 – 4.93 (m, 1H, 20), 4.88 (ddt, J = 10.2, 2.4, 1.3 Hz, 1H), 3.80 (s, 3H), 3.78 (s, 3H), 2.81 – 2.76 (m, 2H), 2.26 – 2.19 (m, 2H); ¹³C{¹H} NMR (125 MHz, CD₃CN) δ 151.2, 151.0, 141.4, 139.9, 132.0, 129.3, 128.7, 128.5, 113.7, 112.8, 75.4, 60.9, 56.3, 36.8, 29.4; ¹¹B

NMR (160 MHz, CD₃CN) δ 3.31; **¹⁹F NMR (470 MHz, CD₃CN)** δ -138.35; **IR (cm⁻¹)** ν 3622, 3564, 3034, 3001, 2967, 2938, 2878, 2838, 1648, 1413, 1372, 1318, 1121, 1069, 980, 920, 859, 796, 752, 697; **HRMS (ESI-)** m/z : [M - K]⁺ calcd for C₁₉H₂₁BF₃O₃ 365.1545; found 365.1530.

Methyl 3-(4-(benzyloxy)-3-iodophenyl)propanoate (8) To a solution of **9** (1 equiv., 10 g, 36.99 mmol) in CH₃CN (150 mL) and TFA (0.3 equiv., 0.82 mL, 11.10 mmol) was added NIS (1.5 equiv., 12.5 g, 55.56 mmol) at 0 °C. The mixture was stirred at room temperature for 5 h. EtOAc (300 mL) was added and the organic phase was washed with a saturated aqueous solution of Na₂S₂O₃ (70 mL) then NaOH (6N, 200 mL), dried over Na₂SO₄, filtered off and concentrated under reduced pressure to afford **8** as a yellow oil (14.7 g, 36.97 mmol, 99% yield) without any further purification. **R_f** 0.33 (SiO₂, ^CHex/EtOAc 9:1). The analytical data are consistent with those reported in the literature.³⁶

Preparation of 5-(3',6-bis(benzyloxy)-2'-(but-3-en-1-yl)-4',5'-dimethoxy-[1,1'-biphenyl]-3-yl)pent-1-en-3-one (6) starting from (7) and (8).

methyl 3-(3',6-bis(benzyloxy)-2'-(but-3-en-1-yl)-4',5'-dimethoxy-[1,1'-biphenyl]-3-yl)propanoate (18) To a round-bottomed flask was added **7** (1.5 equiv., 1.91 g, 4.71 mmol) and **8** (1 equiv., 1.27 g, 3.21 mmol). Then, H₂O (10 mL) and dioxane (40 mL) were added, followed by tetrakis(triphenylphosphine)palladium(0) (11 mol%, 414 mg, 0.358 mmol) and dry K₂CO₃ (5 equiv., 2.22 g, 16.06 mmol). The reaction mixture was stirred at 100 °C using an aluminum heating block for 16 h. Once the reaction mixture was back at room temperature, H₂O (30 mL) was added. The aqueous phase was extracted with CH₂Cl₂ (3x 100 mL). The combined organic layers were dried over Na₂SO₄, filtered off and evaporated under reduced pressure. Crude was purified by flash chromatography on silica gel (gradient was performed from 100/0, V/V cyclohexane/ethyl acetate to 8/2, V/V) to afford **18** (1.2 g, 2.15 mmol, 67% yield) as a colorless oil. **R_f** 0.23 (SiO₂, ^CHex/EtOAc 9:1); **¹H NMR (400 MHz, CDCl₃)** δ 7.56 – 7.45 (m, 2H), 7.41 – 7.27 (m, 3H), 7.27 – 7.17 (m, 5H), 7.12 (dd, J = 8.4, 2.4 Hz, 1H), 7.03 (d, J = 2.3 Hz, 1H, 16), 6.92 (d, J = 8.4 Hz, 1H), 6.55 (s, 1H, 3), 5.59 (ddt, J = 17.0, 10.4, 6.7 Hz, 1H), 5.09 (AB, J = 10.8 Hz, $\Delta\nu_{AB}$ = 36.5 Hz, 2H), 5.00 (s, 2H), 4.79 – 4.71 (m, 2H), 3.94 (s, 3H), 3.82 (s, 3H), 3.66 (s, 3H), 2.92 (t, J = 7.8, 7.8 Hz, 2H), 2.69 – 2.56 (m, 3H), 2.43 – 2.30 (m, 1H), 2.16 – 1.91 (m, 2H); **¹³C{¹H} NMR (125 MHz, CDCl₃)** δ 173.4, 154.2, 151.0, 150.9, 141.6, 138.9, 138.2, 137.5, 134.1, 132.8, 131.4, 131.2, 128.5, 128.4, 128.3, 128.0, 127.8, 127.6, 127.3, 126.6, 114.1, 113.2, 109.9, 75.2, 70.3, 61.0, 56.0, 51.7, 36.0, 34.7, 30.2, 27.6; **IR (cm⁻¹)** ν 3065, 3031, 2929, 2856, 1737, 1636, 1599, 1487, 1453, 1414, 1341, 1251, 1126, 1045, 1024, 736, 697; **HRMS (ESI+)** m/z : [M + Na]⁺ calcd for C₃₆H₃₈O₆Na 589.2561; found 589.2533.

3-(3',6-bis(benzyloxy)-2'-(but-3-en-1-yl)-4',5'-dimethoxy-[1,1'-biphenyl]-3-yl)-N-methoxy-N-methylpropanamide (19) To a slurry mixture of *N,O*-dimethylhydroxylamine hydrochloride (5 equiv., 1.04 g, 10.66 mmol) in dry PhCH₃ (6 mL) was added dropwise trimethylaluminum (5 equiv., 2 M in PhCH₃, 5.3 mL, 10.6 mmol) over 10 minutes at 0 °C under argon

atmosphere (vigorous bubbling was observed). Then, the solution was allowed to warm to room temperature for 1 h. The mixture was re-cooled at 0 °C and a sonicated solution of **18** (1 equiv., 1.2 g, 2.12 mmol) in PhCH₃ (6 mL) was added. The yellow mixture was vigorously stirred at room temperature for 4 h. The reaction was hydrolysed slowly at 0 °C with HCl (10%, 20 mL, vigorous bubbling was observed). An aqueous saturated solution of Rochelle's salt (potassium sodium tartrate, 30 mL) was added and it was stirred for 10 minutes. The aqueous layer was extracted with EtOAc (3x 150 mL). The combined organic layers were washed with an aqueous saturated solution of NaHCO₃ (100 mL), dried over Na₂SO₄, filtered off and evaporated under reduced pressure to afford **19** (1.25 g, 2.10 mmol, 99%) without any further purification as a colorless oil. **R_f** 0.22 (SiO₂, ^CHex/EtOAc 7:3); **¹H NMR (500 MHz, CDCl₃)** δ 7.55 – 7.46 (m, 2H), 7.43 – 7.29 (m, 3H), 7.29 – 7.19 (m, 5H), 7.17 (dd, J = 8.3, 2.4 Hz, 1H), 7.07 (d, J = 2.4 Hz, 1H), 6.93 (d, J = 8.4 Hz, 1H), 6.57 (s, 1H), 5.61 (ddt, J = 17.0, 10.4, 6.7 Hz, 1H), 5.10 (AB, J = 12.0 Hz, $\Delta\nu_{AB}$ = 46.0 Hz, 2H), 5.00 (s, 2H), 4.83 – 4.65 (m, 2H), 3.94 (s, 3H), 3.83 (s, 3H), 3.61 (s, 2H), 3.18 (s, 3H), 2.97 – 2.91 (m, 2H), 2.77 – 2.71 (m, 2H), 2.66 – 2.57 (m, 1H), 2.44 – 2.33 (m, 1H), 2.20 – 1.96 (m, 2H); **¹³C{¹H} NMR (125 MHz, CDCl₃)** δ 173.8, 154.1, 151.0, 150.9, 141.6, 139.0, 138.2, 137.5, 134.2, 133.6, 131.3, 131.3, 128.5, 128.4, 128.0, 127.8, 127.6, 127.3, 126.7, 114.0, 113.2, 109.9, 75.2, 70.3, 61.3, 61.0, 56.0, 34.7, 34.0, 32.3, 29.9, 27.6; **IR (cm⁻¹)** ν 3065, 3028, 2933, 2853, 1664, 1596, 1571, 1487, 1453, 1414, 1374, 1341, 1251, 1125, 1024, 908, 736, 697; **HRMS (ESI+)** m/z : [M + H]⁺ calcd for C₃₇H₄₂NO₆ 596.3007; found 596.2994.

5-(3',6-bis(benzyloxy)-2'-(but-3-en-1-yl)-4',5'-dimethoxy-[1,1'-biphenyl]-3-yl)pent-1-en-3-one (6) To a sonicated solution of **19** (1 equiv., 1.45 g, 2.10 mmol) in dry THF (6 mL) was added vinylmagnesium bromide (2.4 equiv., 1 M in THF, 6 mL, 5.0 mmol) at 0 °C. The yellow mixture was then stirred at 0 °C for 1 h. The reaction mixture was quickly quenched at 0 °C with HCl (10%, 40 mL). The aqueous phase was extracted with EtOAc (3x 150 mL) and the combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, filtered off and concentrated under reduced pressure to afford **6** (1.37 g, 2.10 mmol, 99%) without any further purification as an orange oil. **R_f** 0.21 (SiO₂, ^CHex/EtOAc 9:1); **¹H NMR (500 MHz, CDCl₃)** δ 7.52 – 7.47 (m, 2H), 7.40 – 7.29 (m, 3H), 7.28 – 7.17 (m, 5H), 7.13 (dd, J = 8.4, 2.4 Hz, 1H), 7.03 (d, J = 2.3 Hz, 1H), 6.92 (d, J = 8.4 Hz, 1H), 6.56 (s, 1H), 6.35 (X(AMX), J = 17.7, 10.5 Hz, 1H), 6.02 (AM(AMX), J = 17.7, 10.5, 0.9 Hz, $\Delta\nu_{AM}$ = 156.5 Hz, 2H), 5.60 (ddt, J = 17.0, 10.4, 6.7 Hz, 1H), 5.14 (AB, J = 10.7 Hz, $\Delta\nu_{AB}$ = 36.3 Hz, 1H), 5.00 (s, 2H), 4.79 – 4.72 (m, 2H), 3.94 (s, 3H), 3.82 (s, 3H), 2.95 – 2.87 (m, 4H), 2.66 – 2.56 (m, 1H), 2.43 – 2.32 (m, 1H), 2.19 – 1.94 (m, 2H); **¹³C{¹H} NMR (125 MHz, CDCl₃)** δ 199.9, 154.2, 151.1, 150.9, 141.6, 139.0, 138.2, 137.5, 136.6, 134.1, 133.3, 131.4, 131.3, 128.5, 128.4, 128.3, 128.0, 127.9, 127.6, 127.3, 126.7, 114.0, 113.2, 109.9, 75.2, 70.3, 61.0, 56.1, 41.5, 34.7, 29.1, 27.6; **IR (cm⁻¹)** ν 3062, 3034, 2930, 2859, 1702, 1679, 1638, 1596, 1571, 1486, 1453, 1412, 1372, 1340, 1284, 1250, 1123, 1044, 1022, 908, 842, 811, 735, 696; **HRMS (ESI+)** m/z : [M + H]⁺ calcd for C₃₇H₃₉O₅ 563.2792; found 563.2804.

(*E*)-13,26-bis(benzyloxy)-14,15-dimethoxy-1(1,2),2(1,3)-dibenzenacyclononaphan-6-en-5-one (**12**) A solution of **6** (1 equiv., 97 mg, 0.170 mmol) in CH₂Cl₂ (165 mL) was degassed for 30 minutes under a flow of argon. The reaction mixture was heated at reflux by using a glycerol bath. Then, Grubbs-II catalyst (20 mol%, 30 mg, 0.035 mmol) was added in one portion. The reaction mixture was stirred at reflux for 16 h. The reaction mixture was filtered through a pad of both activated charcoal and silica gel and solvent was evaporated under reduced pressure to **12** (78 mg, 0.15 mmol, 85%) without any further purification as a colorless oil. **R_f** 0.21 (SiO₂, ^CHex/EtOAc 9:1); ¹H NMR (400 MHz, CDCl₃) δ 7.44 – 7.36 (m, 2H), 7.32 – 7.21 (m, 3H), 7.21 – 7.09 (m, 5H), 6.95 (dd, *J* = 8.2, 2.3 Hz, 1H), 6.78 (d, *J* = 8.3 Hz, 1H), 6.65 (d, *J* = 2.3 Hz, 1H), 6.53 (s, 1H), 5.83 (dt, *J* = 15.8, 7.8 Hz, 1H), 5.56 (d, *J* = 16.0 Hz, 1H), 5.07 (AB, *J* = 11.3 Hz, Δ*v*_{AB} = 63.9 Hz, 2H), 4.93 (AB, *J* = 12.2 Hz, Δ*v*_{AB} = 16.5 Hz, 2H), 3.86 (s, 3H), 3.78 (s, 3H), 2.95 – 2.83 (m, 2H), 2.83 – 2.72 (m, 1H), 2.69 – 2.58 (m, 1H), 2.51 – 2.41 (m, 1H), 2.29 – 2.08 (m, 2H), 1.88 – 1.76 (m, 1H); ¹³C{¹H} NMR (100 MHz, CDCl₃) δ 203.8, 154.3, 151.8, 151.2, 151.1, 141.4, 138.1, 137.4, 135.7, 134.9, 133.9, 131.9, 130.3, 128.6, 128.5, 128.4, 127.9, 127.9, 127.7, 126.7, 125.7, 112.8, 107.9, 75.1, 70.5, 61.0, 56.0, 40.8, 33.8, 33.5, 27.6; **IR** (cm⁻¹) *v* 3031, 2924, 2854, 1685, 1654, 1596, 1486, 1415, 1374, 1344, 1249, 1126, 1104, 1023, 736, 697; **HRMS** (**ESI**⁺) *m/z*: [*M* + *K*]⁺ calcd for C₃₅H₃₄O₅K 573.2038; found: 573.2048.

1,2-dihydroxy-14,15-dimethoxy-1(1,2),2(1,3)-dibenzenacyclononaphan-5-one (isomyricanone, **3**) To a solution of **12** (1 equiv., 50 mg, 0.094 mmol) in EtOAc (1 mL) was added palladium on activated charcoal (1 equiv., 10 mg, 0.103 mmol) and stirred for 16 h under a H₂ atmosphere (55 psi). The reaction mixture was filtered through a pad of celite and solvent was evaporated under reduced pressure to afford isomyricanone **3** (33 mg, 0.093 mmol, 99% yield) without any further purification as a colorless oil. **R_f** 0.15 (SiO₂, ^CHex/EtOAc 7:3); Analytical data are consistent with those reported in the literature.⁴

1,1-dimethoxy-1(1,2),2(1,3)-dibenzenacyclononaphane-1,2,5-triol, (isomyricanol, **13**) To a solution of isomyricanone **3** (1 equiv., 200 mg, 0.56 mmol) in CH₃OH (7 mL) was added sodium borohydride (6 equiv., 126 mg, 3.34 mmol) at 0 °C under inert atmosphere and was stirred for 1 h at room temperature. H₂O (5 mL) and EtOAc (25 mL) were added. The aqueous phase was extracted with EtOAc (3x 30 mL). Combined organic layers were dried over Na₂SO₄, filtered off and concentrated under reduced pressure to afford isomyricanol **13** (168 mg, 0.47 mmol, 84%) without any further purification as a white solid. Analytical data are consistent with those reported in the literature.⁴ This compound is present as a 50:50 ratio of two enantiomers and chiral HPLC are available in the SI.

1⁶,7-dihydroxy-2⁴,2⁵-dimethoxy-1(1,3)-benzena-2(1,2)-cyclohexanacyclononaphane-2⁴,2⁴-diene-2³,2⁶-dione (actinidione, **5**)⁸ Isomyricanol **13** (1 equiv., 50 mg, 0.14 mmol) and salcomine (0.4 equiv., 18 mg, 0.054 mmol) were charged and DMF (2.5 mL) was added. Oxygen gas was continuously bubbled through the stirring mixture for 5 h. Afterwards, an additional 18 mg of catalyst was added. The reaction

was complete after 16 h under oxygen atmosphere. EtOAc (30 mL) was added. The organic phase was washed with cold H₂O (5x 10 mL), dried over anhydrous MgSO₄, filtered off and concentrated under reduced pressure to afford actinidione **5** (35 mg, 0.093 mmol, 68%) without any further purification as an orange oil. **R_f** 0.40 (SiO₂, ^CHex/EtOAc 3:7); ¹H NMR (500 MHz, CDCl₃) δ 7.07 (dd, *J* = 8.2, 2.2 Hz, 1H), 6.88 (d, *J* = 8.3 Hz, 1H), 6.81 (d, *J* = 2.2 Hz, 1H), 5.36 (s, 1H), 4.06 (s, 3H), 4.02 (s, 3H), 3.73 (t, *J* = 9.3, 9.3 Hz, 1H), 2.83 (dt, *J* = 13.8, 4.2, 4.2 Hz, 1H), 2.67 (td, *J* = 12.9, 12.3, 3.7 Hz, 1H), 2.41 (ddd, *J* = 12.1, 10.4, 7.6 Hz, 1H), 1.96 (dt, *J* = 13.7, 4.3, 4.3 Hz, 1H), 1.93 – 1.84 (m, 0H), 1.73 – 1.55 (m, 1H), 1.25 – 1.12 (m, 1H), 1.06 – 0.96 (m, 1H), 0.79 (t, *J* = 14.5 Hz, 1H). ¹³C{¹H} NMR (125 MHz, CDCl₃) δ 184.3, 184.2, 151.0, 146.6, 145.0, 144.5, 141.1, 133.7, 131.2, 130.6, 120.8, 117.6, 72.2, 61.4, 61.3, 39.9, 34.8, 33.7, 26.9, 26.4, 24.2; **IR** (cm⁻¹) *v* 3391, 3013, 2925, 2854, 1651, 1613, 1599, 1507, 1454, 1287, 1202, 1144, 1130, 1070, 1016, 822, 758; **HRMS** (**ESI**⁺) *m/z*: [*M* + *Na*]⁺ calcd for C₂₁H₂₄O₆Na 395.1465; found 395.1450.

(*S*)-8,9-dimethoxy-2,3,3a,4,5,6-hexahydro-1*H*-benzo[4,5]cycloocta[1,2,3-*de*]naphthalene-7,11-diol (**1**)

To a solution of isomyricanol **13** (1 equiv., 50 mg, 0.139 mmol) in PhCH₃ (10 mL) was added *para*-toluenesulfonic acid (3.3 equiv., 79 mg, 0.459 mmol). The solution was heated at 90 °C with an aluminum heating block for 5 h under air atmosphere. After cooling, Et₂O (15 mL) was added. The organic phase was washed with a saturated solution of NaHCO₃ (5 mL), dried over MgSO₄, filtered off and evaporated under reduced pressure. Crude was purified by semi-preparative HPLC to afford **1** (37.4 mg, 0.110 mmol) as a pair of enantiomers. **R_f** 0.69 (SiO₂, ^CHex/EtOAc 6:4). Analytical data are consistent with those reported in the literature.⁴ This compound is present as a 50:50 ratio of two enantiomers and chiral HPLC are available in the SI.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Proposed mechanism, detailed synthetic schemes, ¹H, ¹³C{¹H}, ¹⁹F and ¹¹B NMR spectra of all compounds and X-ray molecular structure for compound **13** CCDC 2012419 (PDF).

AUTHOR INFORMATION

Corresponding Author

* Sabine Choppin, sabine.choppin@unistra.fr ORCID NUMBER 0000-0002-9642-7396

* Gilles Hanquet, ghanquet@unistra.fr ORCID NUMBER 0000-0001-8829-1225

Author Contributions

All authors have given approval to the final version of the manuscript.

ACKNOWLEDGMENTS

We would like to thank Matthieu Chessé for the purification of tetracyclic **1** via semi-preparative HPLC and the French government for Paul Massé PhD fellowship.

REFERENCES

- (1) Vogel, H.; Pelletier, J. Examen Chimique de La Racine de Curcuma. *J. Pharm. des Sci. Accessoires* **1815**, 289–300.
- (2) (a) Zhu, J.; Islas-Gonzalez, G.; Bois-Choussy, M. Recent Progress in Isolation, Bioactivity Evaluation and Total Synthesis of Diarylheptanoids. *Org. Prep. Proced. Int.* **2000**, 32, 505–546. (b) Alberti, Á.; Riethmüller, E.; Béni, S. Characterization of Diarylheptanoids: An Emerging Class of Bioactive Natural Products. *J. Pharm. Biomed. Anal.* **2018**, 147, 13–34. (c) Sun, D. J.; Zhu, L. J.; Zhao, Y. Q.; Zhen, Y. Q.; Zhang, L.; Lin, C. C.; Chen, L. X. Diarylheptanoid: A Privileged Structure in Drug Discovery. *Fitoterapia* **2020**, 142.
- (3) (a) Jahng, Y.; Park, J. G. Recent Studies on Cyclic 1,7-Diarlylheptanoids: Their Isolation, Structures, Biological Activities, and Chemical Synthesis. *Molecules* **2018**, 23, 3107–3149. (b) Bochicchio, A.; Schiavo, L.; Chiumminto, L.; Lupattelli, P.; Funicello, M.; Hanquet, G.; Choppin, S.; Colobert, F. Convergent Total Synthesis of (±)-Myricanol, a Cyclic Natural Diarylheptanoid. *Org. Biomol. Chem.* **2018**, 16, 8859–8869.
- (4) Martin, M. D.; Calcul, L.; Smith, C.; Jinwal, U. K.; Fontaine, S. N.; Darling, A.; Seeley, K.; Wojtas, L.; Narayan, M.; Gestwicki, J. E.; Smith, G. R.; Reitz, A. B.; Baker, B. J.; Dickey, C. A. Synthesis, Stereochemical Analysis, and Derivatization of Myricanol Provide New Probes That Promote Autophagic Tau Clearance. *ACS Chem. Biol.* **2015**, 10, 1099–1109.
- (5) Begley, M. J.; Campbell, R. V. M.; Crombie, L.; Tuck, B.; Whiting, D. A. Constitution and Absolute Configuration of Meta,Meta-Bridged, Strained Biphenyls from Myrica Nagi; X-Ray Analysis of 16-Bromyricanol. *J. Chem. Soc.* **1971**, 3634–3642.
- (6) Ishida, J.; Kozuka, M.; Tokuda, H.; Nishino, H.; Nagumo, S.; Lee, K. H.; Nagai, M. Chemopreventive Potential of Cyclic Diarylheptanoids. *Bioorganic Med. Chem.* **2002**, 10, 3361–3365.
- (7) Sakurai, N.; Yaguchi, Y.; Hirakawa, T.; Nagai, M.; Inoue, T. Two Myricanol Glycosides from Myrica Rubra and Revision of the Structure of Isomyricanone. *Phytochemistry* **1991**, 30, 3077–3079.
- (8) Guo, H.; Li, B. G.; Wu, Z. J.; Zhang, G.-L. Lupane Triterpenoids and a Diaryl-Heptanoid from Clematoclethra Actinidioides. *Planta Med.* **2006**, 72, 180–183.
- (9) Zhang, H.; Zhang, Z. *Handbook of Chinese Traditional Medicine Resources*; Beijing: Academic Press, 1994.
- (10) (a) Wang, J.; Dong, S.; Wang, Y.; Lu, Q.; Zhong, H.; Du, G.; Zhang, L.; Cheng, Y. Cyclic Diarylheptanoids from Myrica Nana Inhibiting Nitric Oxide Release. *Bioorganic Med. Chem.* **2008**, 16, 8510–8515. (b) Yoshimura, M.; Yamakami, S.; Amakura, Y.; Yoshida, T. Diarylheptanoid Sulfates and Related Compounds from Myrica Rubra Bark. *J. Nat. Prod.* **2012**, 75, 1798–1802. (c) Ting, Y. C.; Ko, H. H.; Wang, H. C.; Peng, C. F.; Chang, H. S.; Hsieh, P. C.; Chen, I. S. Biological Evaluation of Secondary Metabolites from the Roots of Myrica Adenophora. *Phytochemistry* **2014**, 103, 89–98.
- (11) (a) Gaich, T.; Baran, P. S. Aiming for the Ideal Synthesis. *J. Org. Chem.* **2010**, 75, 4657–4673. (b) Newhouse, T.; Baran, P. S.; Hoffmann, R. W. The Economies of Synthesis. *Chem. Soc. Rev.* **2009**, 38, 3010–3021. (c) Burns, N. Z.; Baran, P. S.; Hoffmann, R. W. Redox Economy in Organic Synthesis. *Angew. Chem. Int. Ed.* **2009**, 48, 2854–2867. (d) Schwan, J.; Christmann, M. Enabling Strategies for Step Efficient Syntheses. *Chem. Soc. Rev.* **2018**, 47, 7985–7995.
- (12) Chausset-Boissarie, L.; Árvai, R.; Cumming, G. R.; Besnard, C.; Kündig, E. P. Total Synthesis of (±)-Vertine with Z-Selective RCM as a Key Step. *Chem. Commun.* **2010**, 46, 6264–6266.
- (13) Corte, J. R.; Pinto, D. J. P.; Fang, T.; Osuna, H.; Yang, W.; Wang, Y.; Lai, A.; Clark, C. G.; Sun, J. H.; Rampulla, R.; Mathur, A.; Kaspady, M.; Neithnadka, P. R.; Li, Y. X. C.; Rossi, K. A.; Myers, J. E.; Sheriff, S.; Lou, Z.; Harper, T. W.; Huang, C.; Zheng, J. J.; Bozarth, J. M.; Wu, Y.; Wong, P. C.; Crain, E. J.; Seiffert, D. A.; Luetgen, J. M.; Lam, P. Y. S.; Wexler, R. R.; Ewing, W. R. Potent, Orally Bioavailable, and Efficacious Macrocyclic Inhibitors of Factor XIa. Discovery of Pyridine-Based Macrocycles Possessing Phenylazole Carboxamide P1 Groups. *J. Med. Chem.* **2020**, 63, 784–803.
- (14) Du, Z.; Lu, J.; Yu, H.; Xu, Y.; Li, A. A Facile Demethylation of *Ortho* Substituted Aryl Methyl Ethers Promoted by AlCl₃. *J. Chem. Res.* **2010**, 34, 222–227.
- (15) (a) Cullen, A.; Muller, A. J.; Williams, D. B. G. Protecting Group-Free Use of Alcohols as Carbon Electrophiles in Atom Efficient Aluminium Triflate-Catalysed Dehydrative Nucleophilic Displacement Reactions. *RSC Adv.* **2017**, 7, 42168–42171. (b) Bering, L.; Jeyakumar, K.; Antonchick, A. P. Metal-Free C–O Bond Functionalization: Catalytic Intramolecular and Intermolecular Benzoylation of Arenes. *Org. Lett.* **2018**, 20, 3911–3914.
- (16) Hoffmann, I.; Blumenröder, B.; Onodi née Thumann, S.; Dommer, S.; Schatz, J. Suzuki Cross-Coupling in Aqueous Media. *Green Chem.* **2015**, 17, 3844–3857.
- (17) Ahn, H.; Son, I.; Lee, J.; Lim, H. Palladium-Hydride-Catalyzed Regiodivergent Isomerization of 1-Alkenes. *Asian J. Org. Chem.* **2017**, 6, 335–341.
- (18) Massé, P.; Choppin, S.; Chiumminto, L.; Hanquet, G.; Colobert, F. Unintended Formation of a 26-Membered Cycle in the Course of a Novel Approach to Myricanol, a Strained [7,0]-Metacyclopentane. *Synlett* **2020**, 31, 559–564.
- (19) (a) Lipshutz, B. H.; Lower, A.; Berl, V.; Schein, K.; Wetterich, F. An Improved Synthesis of the “Miracle Nutrient” Coenzyme Q 10. *Org. Lett.* **2005**, 7, 4095–4097. (b) Greene, L. E.; Godin, R.; Cosa, G. Fluorogenic Ubiquinone Analogue for Monitoring Chemical and Biological Redox Processes. *J. Am. Chem. Soc.* **2016**, 138, 11327–11334.
- (20) Goto, G.; Okamoto, K.; Okutani, T.; Imada, I. A Facile Synthesis of 1,4-Benzoquinones Having a Hydroxy Alkyl Side Chain. *Chem. Pharm. Bull.* **1985**, 33, 4422–4431.
- (21) Kishore Kumar, G. D.; Natarajan, A. Total synthesis of ovalifoliolatin B, acerogenins A and C. *Tetrahedron Lett.* **2008**, 49, 2103–2105.
- (22) Yang, W.-R.; Choi, Y.-S.; Jeong, J.-H. Efficient synthesis of polymethoxyselenoflavones via regioselective direct C–H arylation of selenochromones. *Org. Biomol. Chem.* **2017**, 15, 3074–3083.
- (23) Peschko, C.; Winklhofer, C.; Terpin, A.; Steglich, W. Biomimetic Syntheses of Lamellarin and Lukianol-Type Alkaloids. *Synthesis* **2006**, 18, 3048–3057.