Manganese – new prominent actor in transfer hydrogenation catalysis

Karim Azouzi, Dmitry A. Valyaev, Stéphanie Bastin, Jean-Baptiste Sortais

► To cite this version:

Karim Azouzi, Dmitry A. Valyaev, Stéphanie Bastin, Jean-Baptiste Sortais. Manganese – new prominent actor in transfer hydrogenation catalysis. *Current opinion in green and sustainable chemistry*, 2021, 31, pp.100511. 10.1016/j.cogsc.2021.100511 . hal-03230650

HAL Id: hal-03230650

<https://hal.science/hal-03230650>

Submitted on 4 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Manganese – new prominent actor in transfer hydrogenation catalysis

Karim Azouzi^a, Dmitry A. Valyaev^a, Stéphanie Bastin^{a*}, Jean-Baptiste Sortais^{a,b*}

^a LCC-CNRS, Université de Toulouse, CNRS, UPS, 205 route de Narbonne, 31077 Toulouse Cedex 4, France

^b Institut Universitaire de France, 1 rue Descartes, 75231 Paris Cedex 5, France

Corresponding authors: stephanie.bastin@lcc-toulouse.fr, jean-baptiste.sortais@lcc-toulouse.fr

Abstract: Transfer hydrogenation processes catalyzed by manganese organometallic complexes are comprehensively reviewed. The reaction scope includes the reduction of aldehydes, ketones, imines, polyaromatic nitrogen heterocycles, esters, nitriles and semi-hydrogenation of internal alkynes. Special attention is devoted to ligand design for Mn-catalyzed asymmetric transfer hydrogenation of prochiral ketones leading to enantiomerically-enriched alcohols.

Keywords: Transfer hydrogenation, Manganese, Bidentate ligands, Pincer ligands, Asymmetric catalysis

1. Introduction.

Transfer hydrogenation (TH) is a reduction process that involves hydrogen donors (H₂-donor) such as primary and secondary alcohols (typically isopropanol), diols (such as ethyleneglycol), formic acid, and aminoboranes to cite a few, instead of molecular H₂. For this reason, TH is considered as a safer alternative to direct hydrogenation but also easier to operate since it does not require any use of high-pressure equipment such as autoclaves or Fischer porter vessels.

The first example of late-transition metal-catalyzed TH reaction was reported in 1964 by Henbest and Mitchell with an iridium complex in the presence of isopropanol as the hydrogen donor for the reduction of cyclohexanone in cyclohexanol [1,2]. Later on, in 1973, Blum and Sasson reported the TH of diones with RuCl₂(PPh₃)₃ [3,4]. Since then, many examples have flourished in literature with noble metals [5-7].

If noble metals arguably proved to be very efficient and selective, they are nevertheless quite toxic and very expensive due to their rarity and limited availability. Therefore, interest in non-noble metals has been revived, and several catalytic systems based on late first row transition metals such as iron [8,9] or cobalt [9], have emerged as powerful TH promoters. Contrary to the former 3d transition metals, manganese was disregarded at first, most certainly, because it has long been considered as a metal promoting mainly oxidation reactions. It was not until 2017 and seminal Beller's work that the first catalysts for the TH of ketones to alcohols appeared in literature [10].

This breakthrough was the starting point for enormous progress in this field in parallel with the extensive development of Mn-catalyzed hydrogenation processes,[11-15] whether through the design of new efficient ligands and their combination with different hydrogen donors, the development of a chiral version of this reaction (abbreviated ATH for asymmetric transfer hydrogenation), or the broadening of the application scope to a wider range of substrates. All these aspects will be highlighted in this review.

2.1. Reduction of ketones, aldehydes, imines, nitroarenes and N-heterocycles

In 2017, Beller reported the first example of Mn-based pre-catalyst for TH supported by a tridentate NNN ligand, namely dipicolylamine (Figure 1) [10]. This catalytic system was highly efficient for the reduction of (hetero)aryl and alkyl ketones affording the corresponding alcohols in excellent yields (up to 99%), under mild conditions (70 °C, 2 mol% of base) using 1 mol% catalyst loading. This Mn-based catalytic system showed an excellent chemoselectivity towards ketones over other reducible functions (esters, nitriles) and no deactivation for the substrates containing potentially coordinating O-, S- and N-groups. The same year, our group demonstrated that tridentate ligands were not mandatory to stabilize the catalyst [16]. High TON up to 2000 and TOF up to 3600 h⁻¹ were recorded at 80 °C in the presence of 0.5 mol% of Mn-based catalyst supported by very simple and inexpensive bidentate 2-(aminomethyl)pyridine ligand and the reaction can proceed even at 30 °C albeit more slowly. It should be noted that this Mn-NN catalyst allowed reduction of aldehydes unlike Beller's system with negligible amounts of aldol-type condensation products and preserved conjugated C=C bonds. The ligand's system was even simplified further as catalyst generated *in situ* from Mn(CO)₅Br and ethylenediamine proved to efficiently promote TH of ketones with the same catalyst charge [17].

Figure 1. Mn-based catalytic systems for transfer hydrogenation of carbonylated substrates.

These promising results paved the way to further investigations performed by the groups of Leitner,[18] Kundu,[19] Khusnutdinova,[20] Pidko [21] and Chen [22]. They contributed to the field by expanding the library of catalytic systems for the reduction of ketones and aldehydes. Most of these systems relies on bidentate tricarbonyl Mn-NN complexes and are selective for C=O bond reduction. However, TH of chalcones catalyzed by PN complex developed by Lacy [23] proceeds selectively at the conjugated C=C bond, being consistent with the results of hydrogenation of these substrates using related Mn-PN complex [24]. Although demonstrating reasonable activities, there was still a gap to be bridged to approach the activity recorded with catalytic systems involving noble metals. Pidko and coll.[21] demonstrated that the replacement of one N-donor moiety in NN ligand by a N-heterocyclic carbene (NHC) resulted in a significant improvement in Mn-catalyzed TH of ketones. Indeed, such a Mn-NC system could operate at 75 ppm of catalyst loading at 60 °C with TON up to *ca.* 17000.

A further step has been taken in the field when the groups of Kundu,[19] Khusnutdinova [20] and Sortais [25] demonstrated that their catalytic systems were also able to reduce imines,

which are more reluctant substrates towards TH. The catalyst developed by Khusnutdinova was particularly active for the reduction of aldimines, the corresponding amines were produced in good to excellent yields up to 96% in the presence of 0.5 mol% of catalyst at 80 °C. Its applicability could be extended to the TH of polyaromatic nitrogen heterocycles. Although the scope remains rather limited (4 examples, yields up to 94%), this protocol represents the first example in the field. Finally, selective TH of 2-nitropyridine to form the corresponding amine was recently reported by Morrill using 5 mol% of $[(i\text{Pr}_2\text{PNHPyNHPiPr}_2)\text{Mn}(\text{CO})_3]\text{Br}$ in MeOH at 110-130 °C [26]. Nitro compounds ArNO_2 with more electron-donating aryl groups under these conditions afforded the corresponding ArNHMe products probably as a sequence of TH and aniline alkylation with methanol [27].

2.2. Reduction of carboxylic acid derivatives

Unlike carbonyl derivatives, carboxylic acid derivatives are more difficult to hydrogenate. In this respect, the results reported by Garcia's group in transfer hydrogenation of nitriles to amines constitute an important breakthrough in the field (Figure 2) [28]. Though TH of benzonitrile with 2-butanol catalyzed with Mn-diphosphine complex **Mn1** initially leads to a mixture of target BnNH_2 and its condensation products with 2-butanone $\text{BnN}=\text{C}(\text{Me})\text{Et}$ and benzaldehyde $\text{BnN}=\text{C}(\text{H})\text{Ph}$, they could all be converted to the corresponding ammonium salt upon acidic treatment. The choice of the H_2 -donor appeared to be crucial for the selectivity of this transformation, since the use of more classical isopropanol led to undesired hydration and aldol condensation processes. Under optimized conditions, excellent selectivity was reached at 3 mol% of catalyst loading at 100°C for 3h in the presence of 10 mol% of *t*BuOK affording $[\text{BnNH}_3]\text{Cl}$ in 96% isolated yield. While the presence of electron-donating groups such as *p*-Me and *p*-OMe was well tolerated, *p*- $\text{CF}_3\text{C}_6\text{H}_4\text{CN}$ bearing electron-withdrawing group was more reluctant to reduction and only 39% product yield was obtained despite an increase in of the catalytic charge to 5 mol%. Aliphatic nitriles can also undergo TH using this system, but higher catalyst loading (5 to 10 mol%) was required to achieve good conversions.

TH reduction of nitriles

Garcia, 2019

Mn1

Adhikari / Maji, 2021

Mn2

TH reduction of esters

Clarke, 2020

Figure 2. Mn-based catalytic systems for TH of carboxylic acid derivatives.

Using ammonia-borane as H₂-donor, Adhikari and Maji developed very recently a chemo-divergent reduction of nitriles to primary amines or secondary amines by using n-hexane or isopropanol as solvent, respectively, using 3 mol% of complex **Mn2** supported by hemilabile tridentate SNN ligands (Figure 2) [29]. Unsymmetrical products ArCH₂NHR were even readily obtained by reduction of nitriles ArCN in the presence of external primary amines RNH₂.

In 2020, Clarke and co-workers reported the first example of TH of ester catalyzed by a ferrocene-based Mn(I) complex using ethanol as the H₂-donor (Figure 2).[30] The catalytic system displayed high activity and the catalyst loading could be reduced to 1 mol% at 100 °C in the presence of *t*BuOK (20 mol%). Good tolerance to halogen and heterocycles was observed, whereas the presence of cyano or amino group was detrimental to the activity. The scope of the reaction was successfully extended to aliphatic esters as well as to esters derived from ethanol, methanol and isopropanol. It is worth noting that (*S*)-Naproxen was successfully reduced using this protocol in 82% yield, however with the chiral center epimerization.

2.3. Stereoselective reduction of alkynes to alkenes via TH

Another milestone in the field was reached thanks to the discovery of Mn-based stereoselective transfer semi-hydrogenation of alkynes into alkenes. In 2018, two catalytic systems were reported both involving the use of ammonia borane as H₂-donor and Mn(II) pre-catalysts, but leading to complementary stereoselectivity (Figure 3). [31,32] The groups of El-Sepelgy and Azofra performed the stereoselective transfer semi-hydrogenation of aromatic and aliphatic alkynes to (*Z*)-alkenes using Mn(II) pre-catalyst (1 mol%) supported by tridentate PNP ligands in the presence of 1 equiv. of H-donor in methanol at 60°C. [31] High yields (up to 98%) and stereoselectivities (up to >99:1 *Z/E* ratio) were obtained on a broad scope of internal alkynes. On the other side, Driess and coll. succeeded to reverse the stereoselectivity of the reaction by using N-heterocyclic silylene Mn(II) pre-catalysts (1 mol%) and (*E*)-alkenes were produced in high conversions and stereoselectivities at 55°C.[32] Although the latter catalytic system tolerates the presence of sensitive functional groups, limitations appeared regarding the reduction of dialkyl substrates. Indeed, 6-dodecyne was reduced with lower conversion (67%) and without any stereoselectivity and no reduction was observed with disilylated alkyne derivatives.

Figure 3. Mn-based catalytic systems for transfer semi-hydrogenation of alkynes.

Later Rueping and coll. reported that methanol could also act as an hydrogen donor for Z-selective transfer semi-hydrogenation of alkynes using 2 mol% of Mn(I) pre-catalyst bearing PNP ligand.[33] In this case, the addition of 2 equivalents of weak base Cs_2CO_3 was required and the reaction proceeds under rather harsh conditions at 130-150 °C. However, a variety of aromatic and aliphatic (Z)-alkenes was obtained by this procedure including substrates containing various nitrogen and sulfur heterocycles. Deuterium labelling experiments performed in the presence of CD_3OD evidenced the *cis*-selective hydrogen incorporation onto the alkyne giving important information regarding a plausible catalytic cycle.

2.4. Asymmetric transfer hydrogenation of ketones

Due to the importance of chiral alcohols in industry, and the low price and toxicity of manganese, it became evident to investigate the development of chiral catalysts for asymmetric hydrogenation of ketones. The first report was made by Zirakzadeh and Kirchner in 2017, who investigated the implementation of a Mn(I) catalyst supported by a tridentate PNP ferrocenyl-based chiral ligand. [34] The catalytic system displayed high activity and selectivity, acetophenone was reduced in 95% conversion in 85% *ee* (room temperature, 1 mol% catalyst loading, 4 mol% of *t*BuOK). While *para*-substitution of the phenyl group of aryl ketones did not affect the enantioselectivity to great extent, stereoselectivity dropped to 20 and 46% with acetyl ferrocene and tetralone, respectively. ^1H NMR spectroscopy experiments and theoretical calculations demonstrated that the catalytic cycle involves the formation of a monohydride dicarbonyl Mn(I) catalytic intermediate and suggest that an outer sphere transfer of hydrogen occurs without direct ligand participation. In the same vein, Clarke and coll.[35] investigated the implementation of a Mn(I) catalyst derived from a closely related ferrocenyl tridentate PNN ligand in ATH of ketones. The enantiomeric induction recorded was modest under ATH conditions, whereas hydrogen pressure profoundly improved the catalysts efficiency.

Figure 4. Mn-catalyzed ATH of prochiral ketones

When developing a catalytic system, different aspects have to be considered before envisaging any industrial application. First, the number of additives should be limited. Large-amount of base is often used to activate the pre-catalyst, but reduces functional group tolerance. It is in this context that the catalytic system developed by Morris reveals all its importance since it constitutes the first example of a catalytic system operating in the absence of base.[36] During the course of a study aiming at evaluating the catalytic efficiency of manganese complexes

supported by chiral tri- and tetradentate ligands in ATH, they have isolated a borohydride complex capable of reducing ketones under base-free conditions for the first time. Another point to take into account is the total cost of catalysts. If the replacement of noble metals by base metals lower the price of hydrogenation catalysts, it should be noted that the price of the chiral ligand is non-negligible. With this in mind, the Beller's group has focused its research on the application of inexpensive commercial ligands in ATH. They have tested C2-symmetric potentially tetradentate ligands derived from cheap chiral aminoalcohols.[37] While catalyst systems based on these ligands in combination with $\text{Mn}(\text{CO})_5\text{Br}$ have led to poor enantiomeric excesses (36% *ee* for the best) in the hydrogenation of aromatic ketones, these systems have proven to be particularly effective in the TH of dialkyl ketones. The reduction of the cyclohexyl methyl ketone led to the corresponding alcohol with 88% *ee*. While catalytic system generated *in situ* from $\text{Mn}(\text{CO})_5\text{Br}$ and ONNO-ligand was efficient for the reduction of different (hetero)cyclic alkyl ketones with *ee* up to 93%, the replacement of manganese with iron and ruthenium did not reach the same efficiency. Indeed, iron salts associated with chiral oxamide ligands were unable to reduce cyclohexyl methyl ketone and use of ruthenium precursors did not allow to surpass the enantioselectivity obtained with Mn-based system. Though the catalytic systems described above rely on the use on tri- and tetradentate chiral ligands, our group showed that the combination of simpler bidentate diamine ligands with $\text{Mn}(\text{CO})_5\text{Br}$ can also conduct to efficient catalytic systems with *ee* values up to 90% obtained using commercially available (1*R*,2*R*)-*N,N'*-dimethyl-1,2-diphenylethane-1,2-diamine.[17] This work was complemented by Pidko and coll. in 2019, who performed a detailed experimental and theoretical mechanistic study with well-defined Mn catalysts based on (1*R*,2*R*)-*N,N'*-dimethyl-1,2-diaminocyclohexane scaffold.[38] As part of our research on bidentate ligands in reduction, we have investigated the implementation of chiral aminophosphines in ATH.[39] It emerges from our study that a bulky binepine moiety on the phosphorus atom is necessary to get excellent enantioselectivities (up to 99%) for hindered carbonyl derivatives.

Conclusion and perspectives

Since the discovery of the efficiency of Mn-based catalysts in transfer hydrogenation reactions in 2017, the field has been in constant evolution. A large number of catalytic systems has been developed by playing on the ligand design, the choice of the hydrogen donors and the optimization of the reaction conditions, which resulted in significant progress in terms of catalytic activity, (enantio)selectivity and application scope. These promising results shows a great potential for future development of Mn-catalyzed TH processes and open the way for industrial applications. The biocompatibility of manganese, its low cost and its abundance in the Earth's crust compared to noble metals are all elements in favor of the future application of Mn-based catalytic systems in flavor and fragrance industries as well as for the synthesis of pharmaceutical and agrochemical products.

Acknowledgements.

We thank the Centre National de la Recherche Scientifique (CNRS) and the Institut Universitaire de France (IUF) for a general support of this project. K.A. is grateful to the "Région Occitanie" for a PhD fellowship.

References:

Papers of particular interest, published within the period of review, have been highlighted as:

* of special interest

** of outstanding interest

[1] Haddad YMY, Henbest HB, Husbands J, Mitchell TRB: **Reduction of cyclohexanones to axial alcohols via iridium containing catalysts.** *Proc Chem Soc* 1964: 361.

- [2] Trocha-Grimshaw J, Henbest HB: **Catalysis of the transfer of hydrogen from propan-2-ol to α,β -unsaturated ketones by organo-iridium compounds. A carbon-iridium compound containing a chelate keto-group.** *Chem Commun* 1967: 544.
- [3] Sasson Y, Blum J, Dunkelblum E: **$\text{RuCl}_2(\text{PPh}_3)_3$ -Catalyzed transfer hydrogenation of cyclohexane-1,3-diones.** *Tetrahedron Lett* 1973, 3199–3202.
- [4] Sasson Y, Blum J: **Dichlorotris(triphenylphosphine)ruthenium-catalyzed hydrogen transfer from alcohols to saturated and α,β -unsaturated ketones.** *J Org Chem* 1975, 40: 1887–1896.
- [5] Wang D, Astruc, D: **The golden age of transfer hydrogenation.** *Chem Rev* 2015, 115: 6621–6686.
- [6] Ito J-I, Nishiyama H: **Recent topics of transfer hydrogenation.** *Tetrahedron Lett* 2014, 55: 3133–3146.
- [7] Noyori R, Hashiguchi S: **Asymmetric transfer hydrogenation catalyzed by chiral ruthenium complexes.** *Acc Chem Res* 1997, 30: 97–102.
- [8] Morris RH: **Iron group hydrides in Noyori bifunctional catalysis.** *Chem Rec* 2016, 16: 2644–2658.
- [9] Li YY, Yu SL, Shen WY, Gao JX: **Iron-, cobalt-, and nickel-catalyzed asymmetric transfer hydrogenation and asymmetric hydrogenation of ketones.** *Acc Chem Res* 2015, 48: 2587–2598.
- [10] Perez M, Elangovan S, Spannenberg A, Junge K, Beller M: **Molecularly defined manganese pincer complexes for selective transfer hydrogenation of ketones.** *ChemSusChem* 2017, 10: 83–86.
- [11] Filonenko GA, van Putten R, Hensen EJ, Pidko EA: **Catalytic (de)hydrogenation promoted by non-precious metals – Co, Fe and Mn: recent advances in an emerging field.** *Chem Soc Rev* 2018, 47: 1459–1483.
- [12] Kallmeier F, Kempe R: **Manganese complexes for (de)hydrogenation catalysis: a comparison to cobalt and iron catalysts.** *Angew Chem Int Ed* 2018, 57: 46–60.
- [13] Mukherjee A, Milstein D: **Homogeneous catalysis by cobalt and manganese pincer complexes.** *ACS Catal* 2018, 8: 11435–11469.
- [14] Wang Y, Wang M, Li Y, Liu Q: **Homogeneous manganese-catalyzed hydrogenation and dehydrogenation reactions.** *Chem* 2020 DOI: 10.1016/j.chempr.2020.11.013.
- [15] Sortais JB, Buhaibeh R, Canac Y: **Manganese-catalyzed hydrogenation and hydrogen transfer reactions**, in: Sortais JB (Ed.), *Manganese catalysis in organic synthesis*, Wiley-VCH, 2021, *in press*.
- [16] Bruneau-Voisine A, Wang D, Dorcet V, Roisnel T, Darcel C, Sortais J-B: **Transfer hydrogenation of carbonyl derivatives catalyzed by an inexpensive phosphine-free manganese precatalyst.** *Org Lett* 2017, 19: 3656–3659.
- [17] Wang D, Bruneau-Voisine A, Sortais J-B: **Practical (asymmetric) transfer hydrogenation of ketones catalyzed by manganese with (chiral) diamines ligands.** *Catal Commun* 2018, 105: 31–36.
- [18] Martínez-Ferraté O, Werlé C, Franciò G, Leitner W: **Aminotriazole Mn(I) complexes as effective catalysts for transfer hydrogenation of ketones.** *ChemCatChem* 2018, 10: 4514–4518.

[19] Ganguli K, Shee S, Panja D, Kundu S: **Cooperative Mn(i)-complex catalyzed transfer hydrogenation of ketones and imines.** *Dalton Trans* 2019, **48**: 7358–7366.

[20] Dubey A, Rahaman SMW, Fayzullin RR, Khusnutdinova JR: **Transfer hydrogenation of carbonyl groups, imines and N-heterocycles catalyzed by simple, bipyridine-based Mn(I) complexes.** *ChemCatChem* 2019, **11**: 3844–3852.

[21] van Putten R, Benschop J, de Munck VJ, Weber M, Müller C, Filonenko GA, Pidko EA: **Efficient and practical transfer hydrogenation of ketones catalyzed by a simple bidentate Mn-NHC complex.** *ChemCatChem* 2019, **11**: 5232–5235.

*Highly active catalytic system operating at very low catalyst loading (75 ppm).

[22] Zhang C, Hu B, Chen D, Xia H: **Manganese(I)-catalyzed transfer hydrogenation and acceptorless dehydrogenative condensation: promotional influence of the uncoordinated N-heterocycle.** *Organometallics* 2019, **38**: 3218–3226.

[23] Vigneswaran V, MacMillan SN, Lacy DC: **β -Amino phosphine Mn catalysts for 1,4-transfer hydrogenation of chalcones and allylic alcohol isomerization.** *Organometallics* 2019, **38**: 4387–4391.

[24] Wei D, Bruneau-Voisine A, Chauvin T, Dorcet V, Roisnel T, Valyaev DA, Lugan N, Sortais J-B: **Hydrogenation of carbonyl derivatives catalysed by manganese complexes bearing bidentate pyridinyl-phosphine ligands.** *Adv Synth Catal* 2018, **360**: 676–681.

[25] Wei D, Bruneau-Voisine A, Dubois M, Bastin S, Sortais J-B: **Manganese-catalyzed transfer hydrogenation of aldimines.** *ChemCatChem* 2019, **11**: 5256–5259.

[26] Reed-Berendt BG, Mast N, Morrill LC: **Manganese-catalyzed one-pot conversion of nitroarenes into N-methylarylamines using methanol.** *Eur J Org Chem* 2020, 1136–1140.

[27] Bruneau-Voisine A, Wang D., Dorcet V, Roisnel T, Darcel C, Sortais J-B: **Mono-N-methylation of anilines with methanol catalyzed by a manganese pincer-complex.** *J Catal* 2017, **347**: 57–62.

[28] Garduño JA, Flores-Alamo M, García JJ: **Manganese-catalyzed transfer hydrogenation of nitriles with 2-butanol as the hydrogen donor.** *ChemCatChem* 2019, **11**: 5330–5338.

*First example of transfer hydrogenation of nitriles, the choice of 2-butanol as H₂-donor and solvent proved to be crucial for the reaction selectivity.

[29] Sarkar K, Das K, Kundu A, Adhikari D, Maji B: **Phosphine-free manganese catalyst enables selective transfer hydrogenation of nitriles to primary and secondary amines using ammonia-borane.** *ACS Catal* 2021, **11**: 2786–2794.

** Solvent-dependent chemo-divergent synthesis of primary amines or secondary amines by reduction of nitriles using n-hexane or isopropanol as solvent, respectively.

[30] Oates CL, Widegren MB, Clarke ML: **Manganese-catalysed transfer hydrogenation of esters.** *Chem Commun* 2020, **56**: 8635–8638.

** The only Mn-based catalytic system efficient in transfer of hydrogenation of esters.

[31] Brzozowska A, Azofra LM, Zubar V, Atodiresei I, Cavallo L, Rueping M, El-Sepelgy O: **Highly chemo- and stereoselective transfer semihydrogenation of alkynes catalyzed by a stable, well-defined manganese(II) complex.** *ACS Catal* 2018, **8**: 4103–4109.

[32] Zhou YP, Mo Z, Luecke MP, Driess M: **Stereoselective transfer semi-hydrogenation of alkynes to E-olefins with N-heterocyclic silylene-manganese catalysts.** *Chem Eur J* 2018, **24**: 4780–4784.

[33] Sklyaruk J, Zubar V, Borghs JC, Rueping M: **Methanol as the hydrogen donor in the selective transfer hydrogenation of alkynes enabled by a manganese pincer complex.** *Org Lett* 2020, **22**: 6067–6071.

* First example of Mn-catalyzed (Z)-selective transfer semi-hydrogenation of alkynes using MeOH as H₂-donor.

[34] Zirakzadeh A, de Aguiar SRMM, Stöger B, Widhalm M, Kirchner K: **Enantioselective transfer hydrogenation of ketones catalyzed by a manganese complex containing an unsymmetrical chiral PNP' tridentate ligand.** *ChemCatChem* 2017, **9**: 1744–1748.

[35] Widegren MB, Clarke ML: **Towards practical earth abundant reduction catalysis: design of improved catalysts for manganese catalysed hydrogenation.** *Catal Sci Technol* 2019, **9**: 6047–6058.

[36] Demmans KZ, Olson ME, Morris RH: **Asymmetric transfer hydrogenation of ketones with well-defined manganese(I) PNN and PNNP complexes.** *Organometallics* 2018, **37**: 4608–4618.

[37] Schneekönig J, Junge K, Beller M, **Manganese catalyzed asymmetric transfer hydrogenation of ketones using chiral oxamide ligands.** *Synlett* 2019, **30**: 503–507.

[38] van Putten R, Filonenko GA, Gonzalez de Castro A, Liu C, Weber M, Müller C, Lefort L, Pidko EA: **Mechanistic complexity of asymmetric transfer hydrogenation with simple Mn-diamine catalysts.** *Organometallics* 2019, **38**: 3187–3196.

[39] Azouzi K, Bruneau-Voisine A, Vendier L, Sortais J-B, Bastin S: **Asymmetric transfer hydrogenation of ketones promoted by manganese(I) pre-catalysts supported by bidentate aminophosphines.** *Catal Commun* 2020, **142**: 106040.

Highlights:

Mn complexes can catalyze transfer hydrogenation of C=O, C=N, C=C, C≡N and C≡C bonds

Mn-catalyzed transfer hydrogenations often display high chemo- and stereoselectivity

Chiral Mn complexes have good potential for enantioselective reduction processes

TOC graphic