Changes in foraging mode caused by a decline in prey size have major bioenergetic consequences for a small pelagic fish

Elisa Thorald, Quentin Queiros, Damien Roussel, Gilbert Dutto, Eric Gasset, David J Mckenzie, Caroline Romestaing, Jean-marc Fromentin, Claire Saraux, Loïc Teulier

► To cite this version:

Elisa Thorald, Quentin Queiros, Damien Roussel, Gilbert Dutto, Eric Gasset, et al.. Changes in foraging mode caused by a decline in prey size have major bioenergetic consequences for a small pelagic fish. Journal of Animal Ecology, 2021, 90, pp.2289-2301. 10.1111/1365-2656.13535 . hal-03234605

HAL Id: hal-03234605

<https://hal.science/hal-03234605>

Submitted on 25 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution| 4.0 International License

Changes in foraging mode caused by a decline in prey size have major bioenergetic consequences for a small pelagic fish

Elisa Thorat¹, Quentin Queiros², Damien Roussel¹, Gilbert Dutto³, Eric Gasset⁴, David J. McKenzie⁵,
Caroline Romestaing¹, Jean-Marc Fromentin², Claire Saraux^{2,6} and Loïc Teulier¹.

¹Univ Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, UMR 5023 LEHNA, F-69622, Villeurbanne, France

²MARBEC, Univ Montpellier, CNRS, IFREMER, IRD, Sète, France

³Ifremer (Institut Français de Recherche pour l'Exploitation de la MER), Laboratoire SEA, Chemin de Maguelonne, 34250 Palavas-Les-Flots, France

⁴MARBEC, Univ Montpellier, CNRS, IFREMER, IRD, Palavas-Les-Flots, France

⁵MARBEC, Univ Montpellier, CNRS, IFREMER, IRD, Montpellier, France

⁶IPHC, UMR7178, Université de Strasbourg, CNRS, 67000 Strasbourg, France

Corresponding author: Loïc TEULIER, Université de Lyon, UMR 5023 Ecologie des Hydrosystèmes Naturels et Anthropisés, Université Lyon 1, ENTPE, CNRS, F-69622 Villeurbanne, France, loic.teulier@univ-lyon1.fr

Authors' contributions: QQ, J-MF, GD, EG, DMK, DR, CS and LT conceived and designed the study. ET, QQ, GD, EG, DR, CR, DMK, CS and LT carried out the study. GD & EG took care of the fish and participated to the experimental protocol. ET, QQ, DR, DMK, CS and LT analyzed the data and drafted the manuscript. All authors reviewed and revised the article critically and approved the final version.

Acknowledgements: This study was funded through the MONALISA project, financed by the European Union and the French Ministry of Agriculture through the European Maritime and Fisheries Fund (EMFF/FEAMP) as well as France Filière Pêche. Experimental procedures were approved by the ethics committee for animal experimentation n° 036 of the French Ministère de l'Enseignement Supérieur, de la Recherche et de l'Innovation, with reference number APAFIS n° 7097-2016093008412692. We warmly thank the reviewers for their constructive remarks and comments on the manuscript.

Data availability statement: Data available from the Dryad Digital Repository:
<https://doi.org/10.5061/dryad.xpvnvx0kfp> (Thoral *et al.* 2021a)

Abstract

1. Global warming is causing profound modifications of aquatic ecosystems and one major outcome appears to be a decline in adult size of many fish species. Over the last decade, sardine populations in the Gulf of Lions (NW Mediterranean Sea) have shown severe declines in body size and condition as well as disappearance of the oldest individuals, which could not be related to overfishing, predation pressure or epizootic diseases.
2. In this study, we investigated whether this situation reflects a bottom-up phenomenon caused by reduced size and availability of prey that could lead to energetic constraints. We fed captive sardines with food items of two different sizes eliciting a change in feeding mode (filter-feeding on small items and directly capturing larger ones) at two different rations for several months, and then assessed their muscle bioenergetics to test for changes in cellular function.
3. Feeding on smaller items was associated with a decline in body condition, even at high ration, and almost completely inhibited growth by comparison to sardines fed large items at high ration.
4. Sardines fed on small items presented specific mitochondrial adjustments for energy sparing, indicating a major bioenergetic challenge. Moreover, mitochondria from sardines in poor condition had low basal oxidative activity but high efficiency of ATP production. Notably, when body condition was below a threshold value of 1.07, close to the mean observed in the wild, it was directly correlated with basal mitochondrial activity in muscle.
5. The results show a link between whole-animal condition and cellular bioenergetics in

the sardine, and reveal physiological consequences of a shift in feeding mode. They demonstrate that filter-feeding on small prey leads to poor growth, even under abundant food and an increase in the efficiency of ATP production. These findings may partially explain the declines in sardine size and condition observed in the wild.

Keywords: Bioenergetics, red muscle, mitochondria, fish shrinking, sardine, foraging behaviour, food restriction

65 **Résumé**

66 Le changement global entraîne de profondes modifications des écosystèmes aquatiques, l'une
67 des principales étant le déclin de la taille des adultes chez de nombreuses espèces de poissons.
68 Au cours de la dernière décennie, les populations de sardines du Golfe du Lion (Nord-Ouest de
69 la Méditerranée) ont montré une importante diminution de leur taille et de leur condition
70 corporelle ainsi qu'une disparition des individus les plus âgés, qui n'ont pas pu être liées à la
71 surpêche, à la pression de prédation ou aux épizooties.

72 Dans cette étude, nous avons cherché à savoir si cette situation reflète un phénomène ascendant
73 causé par la réduction de la taille et de la disponibilité des proies qui pourrait entraîner des
74 contraintes énergétiques chez la sardine. Nous avons ainsi nourri des sardines captives avec des
75 granulés de deux tailles différentes provoquant un changement de mode d'alimentation
76 (filtration des petits granulés et capture directe des plus gros) et à deux rations différentes
77 pendant plusieurs mois, puis nous avons évalué leur bioénergétique musculaire pour tester les
78 changements au niveau de leur fonction cellulaire.

79 L'alimentation à base de petits granulés a été associée à un déclin de la condition corporelle,
80 même à une ration élevée, et à une croissance quasiment inhibée par rapport aux sardines
81 nourries avec des plus gros granulés à une ration élevée.

82 Les sardines nourries avec des petits granulés ont également présenté des ajustements
83 mitochondriaux spécifiques pour économiser de l'énergie, indiquant un défi bioénergétique
84 majeur. De plus, les mitochondries des sardines en mauvaise condition présentaient une faible
85 activité oxydative basale, mais une efficacité élevée de production d'ATP. Notamment, lorsque
86 la condition corporelle était inférieure à une valeur seuil de 1,07, proche de la moyenne observée
87 dans la nature, elle était directement corrélée à l'activité mitochondriale basale dans le muscle.

88 Ces résultats montrent un lien entre la condition de l'animal entier et la bioénergétique cellulaire
89 chez la sardine, et révèlent les conséquences physiologiques d'un changement de mode
90 d'alimentation. Ils démontrent que le nourrissage via la filtration de petites proies entraîne une
91 faible croissance, même en cas de nourriture abondante, et une augmentation de l'efficacité de
92 la production d'ATP. Ces résultats peuvent expliquer en partie le déclin de la taille et de la
93 condition des sardines observé dans la nature.

1. Introduction

Marine ecosystems face strong pressures from a combination of climate forcing, anthropogenic disturbance (pollution, navigation, recreational activities, by-catch, climate change) and anthropogenic competition for resources (e.g. fisheries) (Halpern *et al.*, 2008), all of which can impact organismal phenotypes. While water warming and a relaxation of density-dependence phenomena through fishing should promote an increase in growth rates of ectothermic organisms (Morrongiello, Sweetman, & Thresher, 2019), there is evidence that the final adult size of many aquatic ectotherms is declining globally (Daufresne, Lengfellner, & Sommer, 2009; Forster, Hirst, & Atkinson, 2012). Whereas recently debated (Audzijonyte *et al.*, 2020), this size reduction occurs within and amongst species, throughout the food web, from top predators (Sibert *et al.*, 2006) to primary producers (Sommer, Paul, & Moustaka-Gouni, 2015). The decrease in size could result from fishery-induced evolution (as fisheries targeting mostly larger organisms should favour the selection of slow growth and smaller phenotypes, Conover & Munch, 2002; Jørgensen *et al.*, 2007) and/or from within-generation plastic physiological responses. It could also result from direct effects of warming on physiology (e.g. the temperature-size rule; Atkinson, 1994) or global-change-derived modifications of primary production (Bopp *et al.*, 2013).

There has been a profound decline in size and body condition of sardines *Sardina pilchardus* and anchovies *Engraulis encrasicolus* in the Gulf of Lions (northwestern Mediterranean Sea), due to lower growth and disappearance of larger and older individuals (Brosset *et al.*, 2017; Van Beveren *et al.*, 2014). Extensive research has established that this is not due to top-down effects, such as predation or fishing pressure, nor to pathogens (reviewed in Saraux *et al.*, 2019). Rather, a bottom-up process appears to be at play, linked to a shift in plankton communities that the fishes feed upon, towards smaller zooplankton species (Brosset *et al.*, 2016a; Queiros *et al.*, 2019). When maintained in captivity on artificial feeds, wild

sardines exhibited high growth rates and condition, indicating that the decline in size and condition was a plastic rather than an evolutionary response (Saraux *et al.*, 2019). While it is obvious that food quantity can have a significant impact on growth and body condition in fishes (Lee *et al.*, 2018; Sun & Chen, 2009), the potential implications of prey size are less well known. They deserve to be investigated because of climate-change related regime shifts in food webs (Garzke, Ismar, & Sommer, 2015). A reduction in prey size is often accompanied by lower energy content (Zarubin *et al.*, 2014), which may present a major energetic challenge to predators, even for sardines that are opportunistic planktivores (Palomera *et al.*, 2007; Rumolo *et al.*, 2016).

It has been demonstrated that, for rations of similar energy content, sardines fed small prey have lower growth and condition than those fed larger prey (Queiros *et al.*, 2019). The sardine modifies its feeding behaviour with prey size, switching from filter feeding on small prey to direct capture of larger prey, regardless of prey availability (Costalago, Garrido, & Palomera, 2015; Garrido *et al.*, 2007). Foraging duration was much higher for sardines fed on small food items than those fed on larger ones (Queiros *et al.*, 2019). Thus, a possible mechanism for the poor growth and condition may relate to high energy requirements for extended swimming activity during filter-feeding, compared to shorter bouts of swimming to capture large items (Costalago & Palomera, 2014; Crowder, 1985).

Foraging behaviour in fishes is sustained by skeletal muscle activity, with the muscles being structurally separated into discrete slow-twitch oxidative ‘red’ muscle, used for steady aerobic swimming, and fast-twitch glycolytic ‘white’ muscle, used for short bursts of high-speed swimming (Bone, 1978; Webb, 1998). Muscular contraction is powered by ATP but, its production depends on muscle type: while red muscle generates this aerobically by mitochondrial oxidative phosphorylation using substrates and oxygen delivered in the bloodstream, white muscle generates ATP mostly through anaerobic glycolysis of intracellular

144 substrates (Fig. 1, Bone, 1978; Webb, 1998). In the sardine, swimming muscle represents up to
145 70% of body mass and, although the red muscle mass only represents ~10% of the body mass,
146 its mitochondrial activity accounts for 70% of muscle metabolic activity (Teulier *et al.*, 2019).
147 Mitochondrial oxidative phosphorylation is a series of reactions that couple substrate oxidation
148 (catabolism of carbohydrates, lipids or proteins) and oxygen consumption with the generation
149 of a proton motive force, which is then used by ATP synthase to synthesize ATP (Fig. 1). A
150 significant proportion of mitochondrial oxygen consumption is not, however, coupled to ATP
151 synthesis but is used to compensate for proton leakage across the inner mitochondrial
152 membrane. Mitochondrial proton leak is a ubiquitous waste of potential energy, which reduces
153 the efficiency of ATP production (*i.e.* the ATP/O ratio) (Brand, 2005; Fig. 1). The ATP/O ratio
154 defines how much oxygen and nutrients are needed to meet cell ATP demands and, ultimately,
155 sustain animal performance (Brand, 2005). For instance, mitochondrial ATP/O has been
156 positively related to muscle performance (Conley *et al.*, 2013; Distefano *et al.*, 2018) and
157 growth efficiency (Salin *et al.*, 2019; Toyomizu *et al.*, 2011). Mitochondrial efficiency is also
158 flexible, varying among individuals over time, notably in response to environmental and
159 physiological constraints such as temperature, oxygen or food availability (Roussel & Voituron,
160 2020; Salin *et al.*, 2015; Salin *et al.*, 2012; Thorat *et al.*, 2021b). For example, mitochondrial
161 efficiency can increase to enhance energy conservation during food deprivation (Bourguignon
162 *et al.*, 2017; Monternier *et al.*, 2017) or decrease to promote heat generation, as in brown
163 adipose tissue of cold-adapted mammals (Nedergaard & Cannon, 2018). The maintenance of a
164 high ATP/O ratio and related efficiency may carry significant costs, especially increased
165 production of reactive oxygen species (Brand, 2000), and/or a decreased carbon flux from
166 mitochondria that can compromise aspects of cellular biosynthesis (Rolfe & Brand, 1997).
167 Overall, there is strong evidence that mitochondrial efficiency (ATP/O ratio) can be adjusted

168 plastically, in response to metabolic constraints and energy demands, leading to a dynamic
169 interplay with the biological condition of an organism (Koch *et al.*, 2021).

170 In this study we investigated whether aerobic muscle energy metabolism and, in
171 particular, mitochondrial function, might represent a mechanistic nexus for how sardine
172 foraging mode impacts upon their physical condition (Fig. 1). We thus investigated the
173 oxidative activity of red muscle fibres and the coupling efficiency of mitochondrial oxidative
174 phosphorylation in sardines fed for 7 months with either small or large food items (0.1mm and
175 1.2mm diameter commercial pellets), at either low or high ration (0.3% and 0.6% of fish mass
176 per day). We predicted that fish fed on small items would show specific mitochondrial
177 adjustments for energy sparing, with decreased oxygen consumption but enhanced efficiency
178 (ATP/O ratio), which would indicate that increased metabolic costs of foraging mode
179 represented an energetic challenge.

181 2. Material and methods

182 2.1 Experimental conditions

183 All details of the experiment can be found in Queiros *et al.*, 2019. Briefly, wild sardines
184 were captured off Sète (South of France) in October 2016 and brought back to the Ifremer
185 Palavas-les-Flots research station. The mean age of sardines was 1.24 years. After acclimation
186 and weaning onto commercial pellets, sardines were weighed, measured and marked
187 individually using RFID tags (Biolog-id, Bernay, France). Thereafter, they were distributed
188 among 8 tanks (vol. 300L, 56-57 sardines per tank) with similar mean ($\pm$ SD) of mass and length
189 in each (mass: 14.1 ± 3.2 g; length: 11.9 ± 0.7 cm). A 7-month feeding experiment was
190 performed, from 14/11/2016 to 15/6/2017, with natural photoperiod and water temperature
191 (water was pumped directly from the sea). Animals were fed four times a day with commercial
192 pellets (except on weekends where one daily meal was provided). The experiment included the

reproductive season, which occurs once a year between December and March. As detailed in Queiros *et al.*, 2019, the daily weight gain of sardines *in natura* was 0.2% with good condition and 0.1% with bad condition. A preliminary study found that these growth rates were achieved with daily feeding rates of 0.6% and 0.3% of fish mass. Thus, the combination of two pellet sizes (0.1mm and 1.2mm, within the natural range of sardine prey size) (Le Bourg *et al.*, 2015) and two rations (0.3% and 0.6% of fish biomass per day) resulted in four treatments: (1) small items in small quantity (SI-SQ), (2) small items in large quantity (SI-LQ), (3) large items in small quantity (LI-SQ), (4) large items in large quantity (LI-LQ). The two sizes elicited two distinct foraging modes, filtration *vs.* particulate capture (Queiros *et al.*, 2019). Each treatment was randomly assigned to two tanks for replicates. Biometries were performed each month to correct food quantity against increasing biomass, adjusted each week based on linear extrapolation of biomass across successive biometries.

2.2 Sardine sacrifices and morphometric measurements

At the end of the experiment, 1 or 2 sardines were randomly removed from each tank every day over a period of 1 week, *i.e.* 11 to 12 fish per group, and killed by an overdose of benzocaine (1000 ppm). Sardines were fasted for 24h before euthanasia. Mass and size of each sardine were recorded. Intramuscular fat was estimated by fatmeter (Distell Fish Fatmeter MFM-992, as described in Brosset *et al.*, 2015a) and sex was determined visually. Experiments were performed in June when sardines were out of the breeding season (Brosset *et al.*, 2016b). Cumulative growth was estimated based on the increase in size divided by the experiment duration (mm/day), and the relative growth (in %) was calculated as follows:

$$Relative\ growth = \frac{Final\ size - Initial\ size}{Initial\ size}$$

Where initial and final sizes are expressed in mm.

Body condition was calculated following the Le Cren index K_n as estimated by Brosset *et al.*, 2015b (based on more than 24,000 wild sardines sampled over the last 50 years):

$$K_n = \frac{BM}{0.00607 \times TL^{3.057}}$$

where TL is the total length (cm) and BM is the wet body mass (g). This index compares the actual wet mass of an individual to the mass predicted by the relationship of mass to length for the entire wild population. As such, $K_n < 1$ (or $K_n > 1$) means that body condition is lower (or higher) than the long-term average observed in the wild population

2.3 Mitochondrial respiration

Oxygen consumption measurements were performed in red muscle at two levels: in muscle fibres and in isolated mitochondria. At the mitochondrial level, oxygen consumption was expressed in nmol O.min⁻¹.mg⁻¹ of protein, and it was expressed in nmol O.min⁻¹.mg⁻¹ of red muscle at the fibre level. The oxygen consumption was then estimated at the muscle level by multiplying the oxygen consumption obtained at the fibre level by the red muscle mass (μmol O.min⁻¹) or by the relative red muscle mass (nmol O.min⁻¹.g⁻¹ of fish) (see Table 2).

2.3.a Red muscle fibre preparation and oxidative activity

After dissection, the total mass of red muscle was measured and a ~5 mg sample of fibres was withdrawn (Fig. 1). Fibre respiration was monitored at 20°C using high resolution respirometers (Oxygraph-2K, Oroboros® Instruments – WGT Austria) in air-saturated respiration buffer (Mir05: 110 mM sucrose, 0.5 mM EGTA, 3 mM MgCl₂, 60 mM K-lactobionate, 20 mM taurine, 10 mM KH₂PO₄, 1 g/L fatty acid-free bovine serum albumin and 20 mM Hepes, pH 7.1) using different combination of respiratory substrates (5 mM glutamate/2.5 mM malate; 40 μM palmitoyl-carnitine/2.5 mM malate; 5 mM pyruvate/2.5 mM malate/5 mM succinate) with a protocol adapted from Teulier *et al.*, 2019. These different

combinations of substrates enabled us to assay mitochondrial respiration using substrates derived from amino acid (glutamate), carbohydrate (pyruvate) and lipid (palmitoyl-carnitine) metabolic pathways. In addition, the mixture of NADH- and FADH₂-linked substrates (pyruvate/malate and succinate) allowed a full activation of mitochondrial oxidative activity by providing electrons to complex I and complex II of the electron transport system (ETS). Phosphorylating respiration was then initiated by adding 1mM ADP, and the integrity of mitochondria within muscle fibres was subsequently tested by adding 10µM cytochrome c. Finally, a sequential addition of 1 µM p-trifluoromethoxy-carbonyl-cyanide-phenyl hydrazine (FCCP) was performed to measure maximal respiratory activity of the fibres.

2.3.b Mitochondrial isolation, coupling efficiency, respiratory capacity and content

Mitochondria were isolated from red skeletal muscle in an ice-cold isolation buffer containing (100 mM sucrose, 50 mM KCl, 5 mM EDTA, 50 mM Tris-base, pH 7.4 at 4°C), according to previously described protocol for small tissue samples (Boël *et al.*, 2019), involving potter homogenizer, partial protease digestion, and differential centrifugation with mitochondria pelleted at 9000 × g, all steps at 4°C. Protein content of the preparation was assayed in duplicate at 540nm using the biuret method, with bovine serum albumin as standard (Gornall *et al.*, 1949). Absorbance of the same volume of mitochondria was also assayed at 540nm in a solution containing 0.6% Na-K-tartrate and 3% NaOH and subtracted, to correct for any contamination with pigments absorbing at 540nm.

Mitochondrial oxidative phosphorylation efficiency was assessed at 20°C by measuring the rates of ATP synthesis and oxygen consumption in respiratory buffer (120 mM KCl, 5 mM KH₂PO₄, 1 mM EGTA, 2 mM MgCl₂, 0.3% fatty acid-free bovine serum albumin (w/v), and 3mM HEPES, pH 7.4) supplemented with 20 mM glucose, 1.5 U/mL hexokinase (Colinet, Renault, & Roussel, 2017; Teulier *et al.*, 2010). Mitochondria were energized with a mixture

267 of 5 mM pyruvate, 2.5 mM malate, 5 mM succinate to fully activate the ETS. ADP (500 μ M)
268 was added to initiate ATP synthesis. After recording the phosphorylating respiration rate using
269 a Clark electrode (Rank Brother Ltd, Cambridge, UK), four 100 μ L samples of mitochondrial
270 suspension were withdrawn from the respiratory chamber every 2 min and immediately
271 quenched in 100 μ L of ice-cold perchloric acid solution (10% HClO₄ and 25mM EDTA).
272 Production of ATP was determined from the slope of the linear accumulation of glucose-6-
273 phosphate content over the sampling time interval (6 min) as described previously (Colinet *et*
274 *al.*, 2017; Teulier *et al.*, 2010). To make sure that ATP synthesis rate was specific to
275 mitochondrial ATP synthase activity, we determined oxygen consumption and ATP synthesis
276 rates in the presence of oligomycin (1 μ g/mL) that inhibits mitochondrial ATP synthase. These
277 values were taken into account (*i.e.* subtracted from the raw ATP synthesis) to calculate the rate
278 of mitochondrial ATP synthesis (Teulier *et al.*, 2010). Of note, the basal non-phosphorylating
279 oxygen consumption recorded in the presence of oligomycin is a measure of the maximal rate
280 of proton leak across the inner membrane of mitochondria. Due to the linear relationship
281 between ATP synthesis and oxygen consumption in isolated mitochondria from different tissues
282 and organisms (Beavis & Lehninger, 1986; Fontaine *et al.*, 1996; Nogueira *et al.*, 2002; Roussel
283 *et al.*, 2015; Teulier *et al.*, 2010), we presented these data on a single graph by drawing a linear
284 relationship between the basal non-phosphorylating oxygen consumption activity and the
285 maximal oxidative phosphorylation activity, which is related to maximal rates of oxygen
286 consumption and corresponding ATP synthesis (Fig. 1B).

287 Maximal activity of the electron transport system (*i.e.* the respiratory capacity) was
288 measured at 20°C by adding 2 μ M FCCP in the respiratory buffer supplemented with 5mM
289 pyruvate, 2.5 mM malate, 5 mM succinate, and 1 μ g/mL oligomycin. Mitochondrial content of
290 skeletal muscle was estimated from the ratio between the maximal oxygen consumption rates

of muscle fibres (expressed per gram of muscle) and mitochondria (expressed per milligram of protein).

2.5. Statistical analyses

Linear mixed-models (LMM) were used to evaluate the effects of food size and quantity on phenotypic traits, red muscle mitochondrial respiration and efficiency. These models all followed a similar structure, with food size and quantity as well as their interaction as fixed effects, while sex was included as a random effect to remove any potential bias associated to differences between males and females. When an interaction between size and quantity of items was significant ($P < 0.05$) or tended to be significant ($P < 0.07$), post-hoc tests comparing all four treatments (*lsmeans*) were performed. When the interaction was not significant, it was removed from the model and the p- values obtained in the simplified model were reported in the results section.

The relationship between individual body condition and basal mitochondrial respiration was analysed by piecewise linear regression (Fig. 3). Once the breakpoint was estimated, linear regressions were evaluated over each segment of condition (*i.e.* before and after the breakpoint).

Data are presented as means $\pm$ SE. Sample sizes varied among models because some data were missing for some fish, or at the fibre level, data were removed from analyses if addition of cytochrome c caused oxygen consumption to increase by 30% or more (see 2.3.a). Statistical analyses were performed in *R* v. 3.6.1. with the packages *lme4* and *lsmeans*.

3. Results

3.1. Effect of food size and quantity on body condition and growth

For the effect of food size and quantity on body condition, fat content, growth and final mass, the interaction between size and quantity was always significant or close to significance

(LMMs: $P = 0.025$ - 0.061). LI-LQ sardines showed the highest body condition, fat content, body mass, cumulative and relative growth compared to the other three groups (post-hoc lsmeans: $P < 0.010$), while LI-SQ, SI-LQ and SI-SQ did not differ significantly (Table 1). By contrast, there was no interaction between food size and quantity for effects on red muscle mass, whether in absolute terms ($F = 2.359$; $P = 0.133$, $n = 9$) or relative to body mass ($F = 0.002$; $P = 0.962$, $n = 9$). Still, LI sardines had greater absolute red muscle mass than SI sardines ($F = 15.152$; $P < 0.001$), as did LQ compared to SQ sardines ($F = 4.551$; $P = 0.039$). Finally, the size of items also had an effect on relative red muscle mass ($F = 16.515$; $P < 0.001$) with LI sardines being higher than SI (Table 1).

3.2. Oxidative metabolism at different levels of organisation

The phosphorylating respiration state of skeletal muscle fibres obtained with different substrates is presented in Supplementary Material (Figure S1). Apart from glutamate/malate, there was no effect of food size, quantity or their interaction on the phosphorylating state of fibres (all $P > 0.05$). Regarding glutamate/malate, the interaction between food size and quantity tended to be significant ($F = 3.614$, $P = 0.066$), and muscle phosphorylating respiration was affected by food size ($F = 4.220$, $P = 0.048$, post-hoc tests: SI-LQ < LI-LQ, $P = 0.040$).

Table 2 reports the maximal respiration state following addition of FCCP, as measured in isolated mitochondria and muscle fibres, and further estimated in whole red muscle and fish. Maximal respiration of isolated mitochondria was only affected by food size ($F = 9.334$; $P = 0.004$), and was significantly higher in LI than in SI sardines (Table 2). Similarly, mitochondrial content was only affected by food size ($F = 4.604$; $P = 0.040$) but in an opposite direction, being significantly lower in LI groups. The interaction between food size and quantity was close to significance ($F = 3.228$; $P = 0.082$), suggesting that the negative effect of food size on mitochondrial content was mostly observed in sardines fed in small quantity. At the fibre level,

while the interaction between food size and quantity was significant for the maximal respiration state measured at this level ($F = 6.697$; $P = 0.014$), no differences in fibre respiration were highlighted among groups. Finally, the food treatments had no effect on maximal respiration rates at the red muscle or whole fish level ($P > 0.05$ for size, quantity, and its interaction).

3.3. Mitochondrial oxidative phosphorylation efficiency

At the mitochondrial level, there was no interaction between food size and quantity on oxidative phosphorylation activity (*i.e.* the rates of oxygen consumption and associated ATP synthesis). However, activity was higher in sardines fed large food items (oxygen consumption: $F = 9.807$; $P = 0.003$; ATP production: $F = 10.638$; $P = 0.002$), regardless of quantity ($P > 0.05$, Fig. 2A). Basal non-phosphorylating respiration state measured in the presence of oligomycin (*i.e.* the intercept with the x-axis) was also significantly higher in LI sardines ($F = 11.746$; $P = 0.002$, see box in Fig. 2A). There was no significant interaction of size and quantity, and no effect of size or quantity on the slopes of the relationships between rates of ATP synthesis and oxygen consumption (Size*Quantity: $F = 0.014$; $P = 0.906$; Size: $F = 0.784$; $P = 0.382$; Quantity: $F = 2.017$; $P = 0.164$). Thus, these slopes were similar between SI and LI sardines (Slope_{SI}: 3.19 ± 0.15 ; Slope_{LI}: 3.03 ± 0.10). Altogether, these results indicate that the regression line relating ATP synthesis to oxygen consumption was shifted to the right in LI compared to SI groups, meaning that mitochondria from LI sardines had to consume more oxygen to produce the same amount of ATP, due to a higher basal non-phosphorylating respiration or “leak” respiration (Fig. 2A).

When mitochondrial function was quantified for the entire red muscle, oxidative phosphorylation activity did not differ among treatments except for LI-LQ sardines, which exhibited higher rates of ATP synthesis and corresponding phosphorylating respiration, and higher basal non-phosphorylating respiration than the other groups ($P < 0.05$, Fig. 2B).

However, as the slopes were similar between LI-LQ and the other groups (Slope_{LI-LQ}: 3.17 ± 0.19 ; Slope_{other groups}: 2.96 ± 0.18 ; $P > 0.05$), red muscles of LI-LQ sardines needed to consume more oxygen to produce the same amount of ATP compared to the sardines from the other three groups (Fig. 2B).

3.4. Basal respiration in relation to body condition

The rate of basal non-phosphorylating respiration at the mitochondrial level was significantly related to Le Cren body condition (Fig. 3). The segmented regression analysis showed a significant increase in basal non-phosphorylating respiration rate with body condition, until a condition threshold of 1.07 ($R^2 = 0.45$; $P < 0.001$) beyond which no significant relation was found ($P > 0.05$, Fig. 3).

4. Discussion

The results demonstrate clear links between sardine red muscle energetics and foraging modes, indicating effects of both food size and ration. Reducing the size and/or quantity of food led to a series of mitochondrial energy sparing mechanisms (decrease in maximum oxygen consumption and ATP production, enhanced ATP production efficiency and reduced waste of oxygen linked to proton leak), presumably to maintain energy homeostasis. Such energy saving processes suggest that sardines feeding on small food items or small food quantities faced an energetic challenge. These metabolic processes were linked across different levels, from macroscopic integrative traits such as body mass and body condition to intracellular mechanisms such as mitochondrial efficiency. The effects of food ration were stronger at the whole animal level than at the tissue or mitochondrial level, whereas food size also induced marked mitochondrial adjustments beyond the effects at tissue or whole animal scale (Fig. 4). This shows that energy saving processes were not adequate to counteract detrimental effects of

foraging on small particles. Indeed, despite an enhanced mitochondrial efficiency to produce ATP, SI sardines still exhibited lower body condition, body mass and growth performance, regardless of the ration provided (Queiros *et al.*, 2019). Our results therefore support the hypothesis that filter-feeding behaviour is more costly than particulate-feeding in Mediterranean sardines (Queiros *et al.*, 2019), unlike in other small pelagic species (Gibson & Ezzi, 1992; Sanderson & Cech, 1992; Van Der Lingen, 1995), and that this induces plastic responses in cellular energetics.

4.1. Food size induces subcellular bioenergetic adjustments.

Why would the size of food items affect traits like body condition, body mass or growth performance? Part of the answer may reside in the most striking result of our study, which is that small food items triggered bioenergetic adjustments in sardine muscle mitochondria. The effects of food size overwhelmed any potential effect of quantity. Small food items may have induced an increased utilisation of lipids, as intramuscular lipids were ~50% lower in SI-SQ compared to LI-LQ sardines. Moreover, a reduction in glutamate oxidation is a known protein-sparing mechanism, which suggests that protein catabolism may have been occurring. This, together, may explain why SI sardine exhibited lower red muscle mass both in absolute and relative terms.

Feeding on small food items was also associated with decreased oxidative capacity and reduced energy (ATP) outflow, from mitochondria up to whole muscle. This involved a severe loss of mitochondrial power, *i.e.* a ~50% decrease in both maximal rates of ATP synthesis and oxygen consumption. These reduced maximal rates were balanced by a higher coupling efficiency; mitochondria of SI sardines consumed less oxygen than LI sardines to produce a given amount of ATP. The lower maximal respiratory capacity of skeletal muscle and economic management of cellular resources are energy saving mechanisms, decreasing the quantity of

substrates required to produce ATP. Similar adjustments have been reported in skeletal muscle of birds at a critical stage of fasting, when endogenous lipid stores are depleted (Bourguignon *et al.*, 2017; Monternier *et al.*, 2017), and also in skeletal muscle of mammals fed on protein-deficient diets (Zangarelli *et al.*, 2006). Overall, these results indicate that mitochondrial efficiency improves when energy balance is severely threatened in the sardine. Consequently, our results indicate that sardines were unable to cope with the high energy expenditure imposed by filter-feeding on small items, even when these were provided at high ration.

4.2. Food quantity mainly influenced the whole organism.

It is well established that food shortage triggers adaptive behavioural, physiological and cellular responses to lower costs of living, reducing metabolic rates, spontaneous activity and somatic growth (Metcalf, Van Leeuwen, & Killen, 2016). For example, the standard metabolic rate of juvenile brown trout *Salmo trutta* changes as a function of food availability, decreasing or increasing when fish are fed on a low or high ration, respectively (Auer *et al.*, 2016), which was closely related to their growth rate (Auer *et al.*, 2015). In our study, a 50% reduction in ration negatively impacted body condition (-26%), body mass (-33%) and dramatically slowed growth rate of LI sardines (-64%, similarly to what Queiros *et al.*, 2019 found). While a similar trend was observed for growth rate in SI sardines when comparing between rations (-68%) the difference was not significant, presumably because the energetic challenge of foraging on small items transcended any effect of ration (Table 1).

Energy conservation during food shortage is also enacted at a cellular level, by reductions in energy demanding processes and oxidative activity of metabolically active tissues, such as liver and skeletal muscles (McCue, 2010). For instance, the oxidative activity of brown trout muscle is positively correlated with food intake and maximum metabolic rate (Salin *et al.*, 2016a; 2016b). In the sardines, bioenergetics of whole red muscle in the LI group was positively

correlated with food availability, which is likely due to muscle mass. Because of lower body mass and red muscle mass in food-restricted fish, maximal oxygen consumption and ATP synthesis rates of the muscle are depressed. Down regulation of aerobic metabolic pathways has been reported in the red muscle of food-restricted gilthead sea bream *Sparus aurata* (Bermejo-Nogales *et al.*, 2011).

The lowered respiratory capacity of whole muscle in LI-SQ sardines appears mainly to be due to decreased mitochondrial content rather than to changes in mitochondrial activity or skeletal muscle protein content. Changes in mitochondrial content could be caused by dysregulation of autophagy and biogenesis. The former could be induced by caloric restriction (reviewed in Mehrabani *et al.*, 2020) while the latter, which is extremely costly, is probably inhibited to conserve energy. It is also possible that the decrease in mitochondrial content and associated loss of metabolic power in the sardines is linked to decreased use of slow-twitch aerobic red muscle for a particulate capture feeding mode (Hood, 2009; Novak *et al.*, 2005), as prey capture tends to use fast-twitch anaerobic white muscle (Bone, 1978).

4.3. Body condition index and skeletal muscle mitochondria.

The energetic constraints associated with the filtering mode led to very low body condition in the sardines. It is fascinating that, below a threshold value of 1.07, declines in body condition index were strongly correlated with declines in basal non-phosphorylating respiration rate of muscle mitochondria. This threshold is surprisingly close to the mean body condition of sardines observed in the wild, around 1 (Brosset *et al.*, 2015b), which may imply that wild populations are also at a threshold of bioenergetic disequilibrium. Basal non-phosphorylating respiration represents maximal proton leakage across the inner mitochondrial membrane. Limiting proton leak lowers costs of basal maintenance (Boutilier & St-Pierre, 2002) but may carry an oxidative risk, such as an increased reactive oxygen species production (Salin *et al.*,

2018). A reduction of maintenance costs could also be achieved by decreasing mitochondrial content (Guderley, 2004), as observed in LI-SQ sardines (Fig 2b). That is, a decrease in mitochondrial content seems less harmful than directly reducing basal respiration rate itself, as LI-SQ body condition stayed closer to 1 than the SI groups.

Hence, the decline in mitochondrial energy wastage in sardines with poor condition suggests that modifying mitochondrial function provides an economical phenotype when metabolic costs of foraging are high compared to energy acquired.

4.4. Conclusions

Within our experimental conditions, we provide clear evidence that, in sardines, feeding on small prey can present a greater energetic challenge than reduced availability of large prey. Feeding sardines a restricted ration of large food items reduced their body condition, growth rate and body mass, revealing an unbalanced energy budget, but did not stimulate plastic metabolic responses within mitochondria. By contrast, feeding small items, whatever the quantity, induced energy-saving adjustments in mitochondria that are typical of a severe energetic challenge (Bourguignon *et al.*, 2017; Costalago & Palomera, 2014). The food size effect seems to be mediated by feeding mode in the sardines, with the switch from prey capture to filtering. Therefore, impacts of prey size on population condition could be profoundly non-linear. Prey size might only have minor influence on condition as long as sardines use a capture feeding mode, but when prey species are small enough to trigger a change to filtering mode, there are dramatic consequences for body condition.

While this study focuses on sardines in the north-western Mediterranean Sea, it may be indicative of a widespread phenomenon linked to climate change. That is, if global warming causes systematic reductions in prey size (Daufresne *et al.*, 2009), the extra time and energy required to forage could cause energy challenges at multiple levels of the food web. Coupled

with increased metabolic demands of ectotherms in a warmer environment, this could lead to reduced growth rates and be a major contributor to declining final adult sizes in fishes (Gardner *et al.*, 2011).

References

- Atkinson, D. (1994). Temperature and Organism Size-A Law for Ectotherms? *Advances in Ecological Research*, 25, 1–58.
- Audzijonyte, A., Richards, S. A., Stuart-Smith, R. D., Pecl, G., Edgar, G. J., Barrett, N. S., ... Blanchard, J. L. (2020). Fish body sizes change with temperature but not all species shrink with warming. *Nature Ecology and Evolution*, 4(6), 809–814. doi: 10.1038/s41559-020-1171-0
- Auer, S. K., Salin, K., Rudolf, A. M., Anderson, G. J., & Metcalfe, N. B. (2015). Flexibility in metabolic rate confers a growth advantage under changing food availability. *Journal of Animal Ecology*, 84(5), 1405–1411. doi: 10.1111/1365-2656.12384
- Auer, S. K., Salin, K., Rudolf, A. M., Anderson, G. J., & Metcalfe, N. B. (2016). Differential effects of food availability on minimum and maximum rates of metabolism. *Biology Letters*, 12(10), 0–4. doi: 10.1098/rsbl.2016.0586
- Beavis, A. D., & Lehninger, A. L. (1986). The upper and lower limits of the mechanistic stoichiometry of mitochondrial oxidative phosphorylation: Stoichiometry of oxidative phosphorylation. *European Journal of Biochemistry*, 158(2), 315–322. doi: 10.1111/j.1432-1033.1986.tb09753.x
- Bermejo-Nogales, A., Benedito-Palos, L., Caldach-Giner, J. A., & Pérez-Sánchez, J. (2011). Feed restriction up-regulates uncoupling protein 3 (UCP3) gene expression in heart and red muscle tissues of gilthead sea bream (*Sparus aurata* L.). New insights in substrate oxidation and energy expenditure. *Comparative Biochemistry and Physiology - A Molecular and Integrative Physiology*, 159(3), 296–302. doi: 10.1016/j.cbpa.2011.03.024
- Boël, M., Romestaing, C., Voituron, Y., & Roussel, D. (2019). Allometry of mitochondrial efficiency is set by metabolic intensity. *Proceedings of the Royal Society B: Biological Sciences*, 286(1911). doi: 10.1098/rspb.2019.1693
- Bone, Q. (1978). *Locomotor Muscle* (Vol. 7). Academic Press. doi: 10.1007/BF00007710
- Bopp, L., Resplandy, L., Orr, J. C., Doney, S. C., Dunne, J. P., Gehlen, M., ... Vichi, M. (2013). Multiple stressors of ocean ecosystems in the 21st century: Projections with CMIP5 models. *Biogeosciences*, 10(10), 6225–6245. doi: 10.5194/bg-10-6225-2013
- Bourguignon, A., Rameau, A., Toullec, G., Romestaing, C., & Roussel, D. (2017). Increased mitochondrial energy efficiency in skeletal muscle after long-term fasting: Its relevance to animal performance. *Journal of Experimental Biology*, 220(13), 2445–2451. doi: 10.1242/jeb.159087
- Boutilier, R. G., & St-Pierre, J. (2002). Adaptive plasticity of skeletal muscle energetics in

hibernating frogs: Mitochondrial proton leak during metabolic depression. *Journal of Experimental Biology*, 205(15), 2287–2296.

Brand, M. (2005). The efficiency and plasticity of mitochondrial energy transduction. *Biochemical Society Transactions*, 33(5), 897–904. doi: 10.1042/BST20050897

Brand, M. D. (2000). Uncoupling to survive? The role of mitochondrial inefficiency in ageing. *Experimental Gerontology*, 35(6), 811–820. doi: [https://doi.org/10.1016/S0531-5565\(00\)00135-2](https://doi.org/10.1016/S0531-5565(00)00135-2)

Brosset, P., Fromentin, J. M., Ménard, F., Pernet, F., Bourdeix, J. H., Bigot, J. L., ... Saraux, C. (2015a). Measurement and analysis of small pelagic fish condition: A suitable method for rapid evaluation in the field. *Journal of Experimental Marine Biology and Ecology*, 462, 90–97. doi: 10.1016/j.jembe.2014.10.016

Brosset, P., Fromentin, J. M., Van Beveren, E., Lloret, J., Marques, V., Basilone, G., ... Saraux, C. (2017). Spatio-temporal patterns and environmental controls of small pelagic fish body condition from contrasted Mediterranean areas. *Progress in Oceanography*, 151, 149–162. doi: 10.1016/j.pocean.2016.12.002

Brosset, P., Le Bourg, B., Costalago, D., Bănar, D., Van Beveren, E., Bourdeix, J. H., ... Saraux, C. (2016a). Linking small pelagic dietary shifts with ecosystem changes in the Gulf of Lions. *Marine Ecology Progress Series*, 554, 157–171. doi: 10.3354/meps11796

Brosset, P., Lloret, J., Muñoz, M., Fauvel, C., Van Beveren, E., Marques, V., ... Saraux, C. (2016b). Body reserves mediate trade-offs between life-history traits: New insights from small pelagic fish reproduction. *Royal Society Open Science*, 3(10). doi: 10.1098/rsos.160202

Brosset, P., Ménard, F., Fromentin, J. M., Bonhommeau, S., Ulses, C., Bourdeix, J. H., ... Saraux, C. (2015b). Influence of environmental variability and age on the body condition of small pelagic fish in the Gulf of Lions. *Marine Ecology Progress Series*, 529, 219–231. doi: 10.3354/meps11275

Colinet, H., Renault, D., & Roussel, D. (2017). Cold acclimation allows *Drosophila* flies to maintain mitochondrial functioning under cold stress. *Insect Biochemistry and Molecular Biology*, 80, 52–60. doi: 10.1016/j.ibmb.2016.11.007

Conley, K. E., Jubrias, S. A., Cress, M. E., & Esselman, P. (2013). Exercise efficiency is reduced by mitochondrial uncoupling in the elderly. *Experimental Physiology*, 98(3), 768–777. doi: 10.1113/expphysiol.2012.067314

Conover, D. O., & Munch, S. B. (2002). Sustaining fisheries yields over evolutionary time scales. *Science*, 297(5578), 94–96. doi: 10.1126/science.1074085

Costalago, D., Garrido, S., & Palomera, I. (2015). Comparison of the feeding apparatus and diet of European sardines *Sardina pilchardus* of Atlantic and Mediterranean waters: Ecological implications. *Journal of Fish Biology*, 86(4), 1348–1362. doi: 10.1111/jfb.12645

Costalago, D., & Palomera, I. (2014). Feeding of European pilchard (*Sardina pilchardus*) in the northwestern Mediterranean: from late larvae to adults. *Scientia Marina*, 78(1), 41–54. doi: 10.3989/scimar.03898.06d

Crowder, L. B. (1985). Optimal foraging and feeding mode shifts in fishes. *Environmental Biology of Fishes*, 12(1), 57–62. doi: 10.1007/BF00007710

573 Daufresne, M., Lengfellner, K., & Sommer, U. (2009). Global warming benefits the small in
574 aquatic ecosystems. *Proceedings of the National Academy of Sciences of the United*
575 *States of America*, 106(31), 12788–12793. doi: 10.1073/pnas.0902080106

576 Distefano, G., Standley, R. A., Zhang, X., Carnero, E. A., Yi, F., Cornnell, H. H., & Coen, P.
577 M. (2018). Physical activity unveils the relationship between mitochondrial energetics,
578 muscle quality, and physical function in older adults. *Journal of Cachexia, Sarcopenia*
579 *and Muscle*, 9(2), 279–294. doi: 10.1002/jcsm.12272

580 Fontaine, E. M., Moussa, M., Devin, A., Garcia, J., Ghisolfi, J., Rigoulet, M., & Leverve, X.
581 M. (1996). Effect of polyunsaturated fatty acids deficiency on oxidative phosphorylation
582 in rat liver mitochondria. *Biochimica et Biophysica Acta - Bioenergetics*, 1276(3), 181–
583 187. doi: 10.1016/0005-2728(96)00075-8

584 Forster, J., Hirst, A. G., & Atkinson, D. (2012). Warming-induced reductions in body size are
585 greater in aquatic than terrestrial species. *Proceedings of the National Academy of*
586 *Sciences of the United States of America*, 109(47), 19310–19314. doi:
587 10.1073/pnas.1210460109

588 Gardner, J. L., Peters, A., Kearney, M. R., Joseph, L., & Heinsohn, R. (2011). Declining body
589 size: A third universal response to warming? *Trends in Ecology and Evolution*, 26(6),
590 285–291. doi: 10.1016/j.tree.2011.03.005

591 Garrido, S., Marçalo, A., Zwolinski, J., & Van Der Lingen, C. D. (2007). Laboratory
592 investigations on the effect of prey size and concentration on the feeding behaviour of
593 *Sardina pilchardus*. *Marine Ecology Progress Series*, 330, 189–199. doi:
594 10.3354/meps330189

595 Garzke, J., Ismar, S. M. H., & Sommer, U. (2015). Climate change affects low trophic level
596 marine consumers: warming decreases copepod size and abundance. *Oecologia*, 177(3),
597 849–860. doi: 10.1007/s00442-014-3130-4

598 Gibson, R. N., & Ezzi, I. A. (1992). The relative profitability of particulate- and filter-feeding
599 in the herring, *Clupea harengus* L. *Journal of Fish Biology*, 40(4), 577–590. doi:
600 10.1111/j.1095-8649.1992.tb02607.x

601 Gornall, A. G., Bardawill, C. J., & David, M. M. (1949). Determination of serum proteins by
602 means of the biuret reaction. *The Journal of Biological Chemistry*, 177(2), 751–766.

603 Guderley, H. (2004). Metabolic responses to low temperature in fish muscle. *Biological*
604 *Reviews of the Cambridge Philosophical Society*, 79(2), 409–427. doi:
605 10.1017/S1464793103006328

606 Halpern, B. S., Walbridge, S., Selkoe, K. A., Kappel, C. V., Micheli, F., D'Agrosa, C., ...
607 Watson, R. (2008). A Global Map of Human Impact on cadherin-mediated cell adhesion
608 in metazoans may reflect the co-option of a class of proteins whose earliest function was
609 to interpret and re- Marine Ecosystems. *Science*, 319(February), 948–953.

610 Hood, D. A. (2009). Mechanisms of exercise-induced mitochondrial biogenesis in skeletal
611 muscle. *Applied Physiology, Nutrition and Metabolism*, 34(3), 465–472. doi:
612 10.1139/H09-045

613 Jørgensen, C., Enberg, K., Dunlop, E. S., Arlinghaus, R., Boukal, D. S., Brander, K., ...
614 Rijnsdorp, A. D. (2007). Ecology: Managing evolving fish stocks. *Science*, 318(5854),
615 1247–1248. doi: 10.1126/science.1148089

616 Koch, R. E., Buchanan, K. L., Casagrande, S., Crino, O., Dowling, D. K., Hill, G. E., ... Stier,
617 A. (2021). Integrating Mitochondrial Aerobic Metabolism into Ecology and Evolution.
618 *Trends in Ecology and Evolution*, 1–12. doi: 10.1016/j.tree.2020.12.006

619 Le Bourg, B., Bănar, D., Saraux, C., Nowaczyk, A., Le Luherne, E., Jadaud, A., ... Richard,
620 P. (2015). Trophic niche overlap of sprat and commercial small pelagic teleosts in the
621 Gulf of Lions (NW Mediterranean Sea). *Journal of Sea Research*, 103(September), 138–
622 146. doi: 10.1016/j.seares.2015.06.011

623 Lee, S., Lee, Y. M., Kim, K. H., Kim, H. C., Park, C. J., Park, J. W., ... Hwang, H. K. (2018).
624 Effects of food availability on growth performance and immune-related gene expression
625 of juvenile olive flounder (*Paralichthys olivaceus*). *Fish and Shellfish Immunology*,
626 80(May), 348–356. doi: 10.1016/j.fsi.2018.06.021

627 McCue, M. D. (2010). Starvation physiology: Reviewing the different strategies animals use
628 to survive a common challenge. *Comparative Biochemistry and Physiology - A*
629 *Molecular and Integrative Physiology*, 156(1), 1–18. doi: 10.1016/j.cbpa.2010.01.002

630 Mehrabani, S., Bagherniya, M., Askari, G., Read, M. I., & Sahebkar, A. (2020). The effect of
631 fasting or calorie restriction on mitophagy induction: a literature review. *Journal of*
632 *Cachexia, Sarcopenia and Muscle*, 11(6), 1447–1458. doi: 10.1002/jcsm.12611

633 Metcalfe, N. B., Van Leeuwen, T. E., & Killen, S. S. (2016). Does individual variation in
634 metabolic phenotype predict fish behaviour and performance? *Journal of Fish Biology*,
635 88(1), 298–321. doi: 10.1111/jfb.12699

636 Monternier, P. A., Teulier, L., Drai, J., Bourguignon, A., Collin-Chavagnac, D., Hervant, F.,
637 ... Roussel, D. (2017). Mitochondrial oxidative phosphorylation efficiency is
638 upregulated during fasting in two major oxidative tissues of ducklings. *Comparative*
639 *Biochemistry and Physiology -Part A : Molecular and Integrative Physiology*, 212(May),
640 1–8. doi: 10.1016/j.cbpa.2017.06.018

641 Morrongiello, J. R., Sweetman, P. C., & Thresher, R. E. (2019). Fishing constrains
642 phenotypic responses of marine fish to climate variability. *Journal of Animal Ecology*,
643 88(11), 1645–1656. doi: 10.1111/1365-2656.12999

644 Nedergaard, J., & Cannon, B. (2018). Brown adipose tissue as a heat-producing
645 thermoeffector. In *Handbook of Clinical Neurology* (1st ed., Vol. 156). Elsevier B.V.
646 doi: 10.1016/B978-0-444-63912-7.00009-6

647 Nogueira, V., Walter, L., Avéret, N., Fontaine, E., Rigoulet, M., & Leverve, X. M. (2002).
648 Thyroid status is a key regulator of both flux and efficiency of oxidative phosphorylation
649 in rat hepatocytes. *Journal of Bioenergetics and Biomembranes*, 34(1), 55–66. doi:
650 10.1023/A:1013822820840

651 Novak, C. M., Jiang, X., Wang, C., Teske, J. A., Kotz, C. M., & Levine, J. A. (2005). Caloric
652 restriction and physical activity in zebrafish (*Danio rerio*). *Neuroscience Letters*, 383(1–
653 2), 99–104. doi: 10.1016/j.neulet.2005.03.048

654 Palomera, I., Olivar, M. P., Salat, J., Sabatés, A., Coll, M., García, A., & Morales-Nin, B.
655 (2007). Small pelagic fish in the NW Mediterranean Sea: An ecological review. *Progress*
656 *in Oceanography*, 74(2–3), 377–396. doi: 10.1016/j.pocean.2007.04.012

657 Queiros, Q., Fromentin, J. M., Gasset, E., Dutto, G., Huiban, C., Metral, L., ... Saraux, C.
658 (2019). Food in the sea: Size also matters for pelagic fish. *Frontiers in Marine Science*,

659 6(JUL), 1–13. doi: 10.3389/fmars.2019.00385

660 Rolfe, D. F. S., & Brand, M. D. (1997). The physiological significance of mitochondrial
661 proton leak in animal cells and tissues. *Bioscience Reports*, 17(1), 9–16. doi:
662 10.1023/A:1027327015957

663 Roussel, D., Salin, K., Dumet, A., Romestaing, C., Rey, B., & Voituron, Y. (2015). Oxidative
664 phosphorylation efficiency, proton conductance and reactive oxygen species production
665 of liver mitochondria correlates with body mass in frogs. *Journal of Experimental*
666 *Biology*, 218(20), 3222–3228. doi: 10.1242/jeb.126086

667 Roussel, D., & Voituron, Y. (2020). Mitochondrial Costs of Being Hot: Effects of Acute
668 Thermal Change on Liver Bioenergetics in Toads (*Bufo bufo*). *Frontiers in Physiology*,
669 11(March), 1–10. doi: 10.3389/fphys.2020.00153

670 Rumolo, P., Bonanno, A., Barra, M., Fanelli, E., Calabrò, M., Genovese, S., ... Basilone, G.
671 (2016). Spatial variations in feeding habits and trophic levels of two small pelagic fish
672 species in the central Mediterranean Sea. *Marine Environmental Research*, 115, 65–77.
673 doi: 10.1016/j.marenvres.2016.02.004

674 Salin, K., Auer, S. K., Anderson, G. J., Selman, C., & Metcalfe, N. B. (2016a). Inadequate
675 food intake at high temperatures is related to depressed mitochondrial respiratory
676 capacity. *Journal of Experimental Biology*, 219(9), 1356–1362. doi: 10.1242/jeb.133025

677 Salin, K., Auer, S. K., Rey, B., Selman, C., & Metcalfe, N. B. (2015). Variation in the link
678 between oxygen consumption and ATP production, and its relevance for animal
679 performance. *Proceedings of the Royal Society B: Biological Sciences*, 282(1812). doi:
680 10.1098/rspb.2015.1028

681 Salin, K., Auer, S. K., Rudolf, A. M., Anderson, G. J., Selman, C., & Metcalfe, N. B. (2016b).
682 Variation in metabolic rate among individuals is related to tissue-specific differences in
683 mitochondrial leak respiration. *Physiological and Biochemical Zoology*, 89(6), 511–523.
684 doi: 10.1086/688769

685 Salin, K., Roussel, D., Rey, B., & Voituron, Y. (2012). David and Goliath: A Mitochondrial
686 Coupling Problem? *Journal of Experimental Zoology Part A: Ecological Genetics and*
687 *Physiology*, 317(5), 283–293. doi: 10.1002/jez.1722

688 Salin, K., Villasevil, E. M., Anderson, G. J., Auer, S. K., Selman, C., Hartley, R. C., ...
689 Metcalfe, N. B. (2018). Decreased mitochondrial metabolic requirements in fasting
690 animals carry an oxidative cost. *Functional Ecology*, 32(9), 2149–2157. doi:
691 10.1111/1365-2435.13125

692 Salin, K., Villasevil, E. M., Anderson, G. J., Lamarre, S. G., Melanson, C. A., McCarthy, I.,
693 ... Metcalfe, N. B. (2019). Differences in mitochondrial efficiency explain individual
694 variation in growth performance. *Proceedings of the Royal Society B: Biological*
695 *Sciences*, 286(1909). doi: 10.1098/rspb.2019.1466

696 Sanderson, S. L., & Cech, J. J. (1992). Energetic Cost of Suspension Feeding versus
697 Particulate Feeding by Juvenile Sacramento Blackfish. *Transactions of the American*
698 *Fisheries Society*, 121(2), 149–157. doi: 10.1577/1548-
699 8659(1992)121<0149:ecosfv>2.3.co;2

700 Saraux, C., Van Beveren, E., Brosset, P., Queiros, Q., Bourdeix, J. H., Dutto, G., ...
701 Fromentin, J. M. (2019). Small pelagic fish dynamics: A review of mechanisms in the

702 Gulf of Lions. *Deep-Sea Research Part II: Topical Studies in Oceanography*, 159(XXXX),
703 52–61. doi: 10.1016/j.dsr2.2018.02.010

704 Sibert, J., Hampton, J., Kleiber, P., & Maunder, M. (2006). Biomass, Size, and Trophic Status
705 of Top Predators in the Pacific Ocean. *Science*, 314(March), 1773–1777. doi:
706 10.1126/science.1135347

707 Sommer, U., Paul, C., & Moustaka-Gouni, M. (2015). Warming and ocean acidification
708 effects on phytoplankton - From species shifts to size shifts within species in a
709 mesocosm experiment. *PLoS ONE*, 10(5), 1–17. doi: 10.1371/journal.pone.0125239

710 Sun, L., & Chen, H. (2009). Effects of ration and temperature on growth, fecal production,
711 nitrogenous excretion and energy budget of juvenile cobia (*Rachycentron canadum*).
712 *Aquaculture*, 292(3–4), 197–206. doi: 10.1016/j.aquaculture.2009.04.041

713 Teulier, L., Rouanet, J. L., Letexier, D., Romestaing, C., Belouze, M., Rey, B., ... Roussel, D.
714 (2010). Cold-acclimation-induced non-shivering thermogenesis in birds is associated
715 with upregulation of avian UCP but not with innate uncoupling or altered ATP
716 efficiency. *Journal of Experimental Biology*, 213(14), 2476–2482. doi:
717 10.1242/jeb.043489

718 Teulier, L., Thorat, E., Queiros, Q., McKenzie, D. J., Roussel, D., Dutto, G., ... Saraux, C.
719 (2019). Muscle bioenergetics of two emblematic Mediterranean fish species: *Sardina*
720 *pilchardus* and *Sparus aurata*. *Comparative Biochemistry and Physiology -Part A :*
721 *Molecular and Integrative Physiology*, 235(June), 174–179. doi:
722 10.1016/j.cbpa.2019.06.008

723 Thorat, E., Queiros, Q., Roussel, D., Dutto, G., Gasset, E., McKenzie, D. J., ... Teulier, L.
724 (2021a). Changes in foraging mode caused by a decline in prey size have major
725 bioenergetic consequences for a small pelagic fish, Dryad, Dataset,
726 <https://doi.org/10.5061/dryad.xpnvx0kfp>

727 Thorat, E., Roussel, D., Chinopoulos, C., Teulier, L., & Salin, K. (2021b). Low oxygen levels
728 can help to prevent the detrimental effect of acute warming on mitochondrial efficiency
729 in fish. *Biology Letters*, 17.

730 Toyomizu, M., Kikusato, M., Kawabata, Y., Azad, M. A. K., Inui, E., & Amo, T. (2011).
731 Meat-type chickens have a higher efficiency of mitochondrial oxidative phosphorylation
732 than laying-type chickens. *Comparative Biochemistry and Physiology - A Molecular and*
733 *Integrative Physiology*, 159(1), 75–81. doi: 10.1016/j.cbpa.2011.01.020

734 Van Beveren, E., Bonhommeau, S., Fromentin, J. M., Bigot, J. L., Bourdeix, J. H., Brosset,
735 P., ... Saraux, C. (2014). Rapid changes in growth, condition, size and age of small
736 pelagic fish in the Mediterranean. *Marine Biology*, 161(8), 1809–1822. doi:
737 10.1007/s00227-014-2463-1

738 Van Der Lingen, C. D. (1995). Respiration rate of adult pilchard *Sardinops sagax* in relation
739 to temperature, voluntary swimming speed and feeding behaviour. *Marine Ecology*
740 *Progress Series*, 129(1–3), 41–54. doi: 10.3354/meps129041

741 Webb, P. W. (1998). Entrainment by river chub *Nocomis micropogon* and smallmouth bass
742 *Micropterus dolomieu* on cylinders. *Journal of Experimental Biology*, 201(16), 2403–
743 2412.

744 Zangarelli, A., Chanseau, E., Morio, B., Brugère, C., Mosoni, L., Rousset, P., ... Walrand,

- 745 S. (2006). Synergistic effects of caloric restriction with maintained protein intake on
746 skeletal muscle performance in 21-month-old rats: a mitochondria-mediated pathway.
747 *The FASEB Journal*, 20(14), 2439–2450. doi: 10.1096/fj.05-4544com
- 748 Zarubin, M., Farstey, V., Wold, A., Falk-Petersen, S., & Genin, A. (2014). Intraspecific
749 differences in lipid content of calanoid copepods across fine-scale depth ranges within
750 the photic layer. *PLoS ONE*, 9(3), 1–10. doi: 10.1371/journal.pone.0092935

FIGURES

Figure 1. Illustration of the integrative levels of the red muscle bioenergetics in sardine, from isolated mitochondria, to the whole tissue.

Substrates are oxidized by the complexes of the electron transport system (ETS) and ATP is produced through a phosphorylation process by the ATP synthase. The ETS and ATP synthase are enclosed in the inner membrane of the mitochondria (panel A). Mitochondrial respiration rates associated to either basal non-phosphorylating state (BASAL) and/or phosphorylating state (PHOSPHO) were assessed at different levels of integration. 1) Mitochondrial oxygen consumption was coupled to the amount of ATP produced, to obtain the mitochondrial efficiency which is the slope of the linear regression between oxygen consumption and ATP production, called P/O ratio (panel B). 2) The muscle fibre respiration rates were induced by addition of different substrates to discriminate the metabolic pathways fuelling the red muscle (panel C). 3) The whole red muscle oxidation phosphorylation activity was calculated to estimate the relative importance of the whole muscle bioenergetics. The red muscle is positioned along the lateral line and is clearly separated from the white muscle (panel D). ETS: Electron transport system; ATP: Adenosine triphosphate; ADP: adenosine diphosphate; Pi: Phosphate inorganic; IMS: intermembrane space of the mitochondrion; M: matrix of the mitochondrion. See the “Material and Methods” section for further details.

Figure 2. Mitochondrial efficiency of sardine that received different quantities and sizes of food items, at the mitochondrial level (Panel A; LI-LQ: large food items in large quantities, n = 11; LI-SQ: large food items in small quantities, n = 12; SI-LQ: small food items in large quantities, n = 11; SI-SQ: small food items in small quantities, n = 9) **and red muscle level** (Panel B, LI-LQ: n = 11, LI-SQ: n = 11; SI-LQ: n = 10; SI-SQ: n = 9). In panel A, the two trend curves drawn are those of the “large food items” (LI: $y = 3.03x - 44.40$) and “small food items” (SI: $y = 3.19x - 37.96$) groups, and in panel B, the trend curves are those of LI-LQ group ($y = 3.17x - 4.52$) and of the other groups together ($y = 2.97x - 1.77$). These data were analysed using a linear-mixed effects model with size and quantity of items as fixed-effects and sex as random effect. The values shown are means $\pm$ s.e.m. (the error bars of the basal oxygen consumption are small enough that they are hidden by the points, see zoom boxes). An asterisk represents a significant difference in maximal oxygen consumption, ATP production and basal oxygen consumption (see zoom boxes) between diets composed of different size food items (Panel A, difference between LI and SI) and between LI-LQ sardines and other groups (Panel B; $*P < 0.05$; $**P < 0.01$).

787 **Figure 3. Basal oxygen consumption at the mitochondrial level as a function of sardine body condition index.** Fish of each treatment are
788 represented by a coloured circle. Sardines fed with large food items are represented in blue for large quantity (LI-LQ, n = 11) and in green for small
789 quantity (LI-SQ, n = 12). Yellow and red circles represent sardines fed with small food items, in large (SI-LQ, n = 10) and small (SI-SQ, n = 11)
790 quantities, respectively. Basal mitochondrial respiration in $\text{nmol O} \cdot \text{min}^{-1} \cdot \text{mg protein}^{-1}$ is positively correlated to the body condition index when the
791 BCI is less than 1.07 ($P < 0.001$). Above this threshold, there is no longer a regression between these two parameters.

809 **Figure 4. Whole-organism, muscle, and subcellular consequences of the quantity and/or size of food.** In order to investigate the effect of size
810 and/or quantity of food, a fully factorial protocol has been set up. Size and quantity impacted the whole-organism and muscle parameters, whereas
811 subcellular bioenergetics have been modified only by size of food items. Significant effects are illustrated by an asterisk. As foraging mode depends
812 on food size (Costalago *et al.*, 2015; Garrido *et al.*, 2007), the effect described by size is called “foraging mode” (FM). Lowering the quantity of
813 food by 50% induces a severe “caloric restriction” (CR). [1] Queiros *et al.*, 2019

814
815
816

817

818 **Table 1. Characteristics of sardines used to measure the respiration of red muscle fibres.** These data correspond to the measurements made
819 on the day sardines were euthanised. Values are mean $\pm$ s.e.m. N corresponds to the number of individuals randomly split into 4 treatments, sorted
820 in three sex categories (male, female and indeterminate). Cumulative growth represents the total length gain in mm, over the entire experiment.
821 RM/BM ratio corresponds to the relative red muscle mass, which is the proportion of the red muscle mass over the body mass. Data were analysed
822 with a linear mixed-effect model with size and quantity as fixed-effects and sex as random effect. Data with different superscripts are significantly
823 different at $P < 0.05$. The boxes with the asterisks indicated a global effect of item size (** $P < 0.001$).

	LI-LQ	LI-SQ	SI-LQ	SI-SQ	Statistical analysis		
N	8-11	10-12	9-11	10-12			
(M/F/indeterminate)	(6/3/2)	(5/6/1)	(3/5/2)	(3/8/1)			
Final BCI	1.17 ± 0.04 ^a	0.98 ± 0.02 ^b	0.93 ± 0.05 ^b	0.87 ± 0.02 ^b	Size: <i>P</i> < 0.001	824	
					Quantity: <i>P</i> = 0.001	825	
					Size*Quantity: <i>P</i> = 0.061	826	
Fat content (%)	13.46 ± 1.40 ^a	8.86 ± 0.76 ^b	8.12 ± 0.81 ^b	6.93 ± 0.37 ^b	Size: <i>P</i> < 0.001	827	
					Quantity: <i>P</i> = 0.002	828	
					Size*Quantity: <i>P</i> = 0.055	829	
Cumulative growth (mm)	14.29 ± 1.68 ^a	5.05 ± 1.07 ^b	5.27 ± 1.42 ^b	1.64 ± 0.46 ^b	Size: <i>P</i> < 0.001	830	
					Quantity: <i>P</i> < 0.001	831	
					Size*Quantity: <i>P</i> = 0.025	832	
Relative growth (%)	12.55 ± 1.56 ^a	4.35 ± 0.91 ^b	4.60 ± 1.29 ^b	1.34 ± 0.36 ^b	Size: <i>P</i> < 0.001	833	
					Quantity: <i>P</i> < 0.001	834	
					Size*Quantity: <i>P</i> = 0.029	835	
Body mass (g)	19.23 ± 1.38 ^a	13.98 ± 0.62 ^b	14.22 ± 1.19 ^b	12.81 ± 0.73 ^b	Size: <i>P</i> = 0.005	836	
					Quantity: <i>P</i> = 0.001	837	
					Size*Quantity: <i>P</i> = 0.058	838	
Red muscle mass (g)	1.59 ± 0.17 ^a	1.18 ± 0.06 ^{a,b}	0.98 ± 0.13 ^b	0.91 ± 0.07 ^b	Size: <i>P</i> < 0.001	839	
					Quantity: <i>P</i> = 0.039	840	
RM/BM (%)	8.11 ± 0.39	8.46 ± 0.20	**	6.70 ± 0.45	7.09 ± 0.30	Size: <i>P</i> < 0.001	841

842

843

844

845

846

847 **Table 2. Maximal respiration ETS activity and mitochondrial content of sardine red muscle for each feeding treatment. The maximal**
848 **respiration rate of ETS induced by FCCP is represented at three different levels of organisation: at the subcellular level (nmol O.min⁻¹.mg**
849 **proteins⁻¹) obtained through respiration measurements of isolated mitochondria; at the red muscle fibre level (pmol O₂.s⁻¹.mg muscle⁻¹) obtained**
850 **through respiration measurements of a red muscle sample, and at the red muscle level estimated by multiplying maximal respiration rate at the**
851 **fibre level by red muscle mass or relative red muscle mass (total: pmol O₂.s⁻¹; specific: pmol O₂.s⁻¹.g fish⁻¹). Values are mean ± s.e.m. The**
852 **mitochondrial content of red muscle is expressed in milligram of mitochondrial proteins per gram of red muscle and was estimated as the ratio**
853 **between respiration rates at the fibre level and mitochondrial level. Data were analysed with a linear mixed-effects model with size and quantity**
854 **as fixed-effects and sex as random effect. Different letters indicate a significant difference among groups at P < 0.05. The boxes with the asterisks**
855 **on the measurements indicates a global effect of item size (*P < 0.05; **P < 0.01).**
856

	LI-LQ	LI-SQ	SI-LQ	SI-SQ	Statistical analysis
N	9-11	11-12	9-11	9-12	
Maximal respiration ETS activity					
Mitochondrial level (nmol O.min ⁻¹ .mg protein ⁻¹)	90 ± 9	96 ± 7	59 ± 12	65 ± 12	Size: P = 0.004
Red muscle fibre level (nmol O.min ⁻¹ .mg muscle ⁻¹)	5 ± 0.8 ^a	3.5 ± 0.4 ^a	3.3 ± 0.4 ^a	5.3 ± 0.9 ^a	Size*Quantity: P = 0.014
Red muscle level	Total (μmol O.min ⁻¹)	6.3 ± 0.9 ^a	4.4 ± 0.7 ^a	3.6 ± 0.7 ^a	4.1 ± 0.8 ^a n.s.
	Specific (nmol O.min ⁻¹ .g fish ⁻¹)	350 ± 44 ^a	306 ± 38 ^a	245 ± 34 ^a	307 ± 58 ^a n.s.
Mitochondrial content (mg protein. g muscle ⁻¹) ^x	50.9 ± 7.7	34.8 ± 5.4	52.6 ± 8.2	62.2 ± 7.6	Size: P = 0.040

857