One-Pot Synthesis of 2,5-Disubstituted Furans through In Situ Formation of Allenes and Enolization Cascade

Yann Bernhard, Joachim Gilbert, Till Bousquet, Audrey Favrelle-Huret,
Philippe Zinck, Sylvain Pellegrini, Lydie Pelinski

► To cite this version:

Yann Bernhard, Joachim Gilbert, Till Bousquet, Audrey Favrelle-Huret, Philippe Zinck, et al.. One-Pot Synthesis of 2,5-Disubstituted Furans through In Situ Formation of Allenes and Enolization Cascade. European Journal of Organic Chemistry, 2019, European Journal of Organic Chemistry, 2019 (48), pp.7870-7873. 10.1002/ejoc.201901669 . hal-03263313

HAL Id: hal-03263313

<https://hal.science/hal-03263313>

Submitted on 17 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Article

Title: One-pot synthesis of 2,5-disubstituted furans through in-situ formation of allenes and enolization cascade

Authors: Lydie Pelinski, Yann Bernhard, Till Bousquet, Audrey Favrelle-Huret, Philippe Zinck, Joachim Gilbert, and Sylvain Pellegrini

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Eur. J. Org. Chem.* 10.1002/ejoc.201901669

Link to VoR: <http://dx.doi.org/10.1002/ejoc.201901669>

COMMUNICATION

One-pot synthesis of 2,5-disubstituted furans through in-situ formation of allenes and enolization cascade

Yann Bernhard, Joachim Gilbert, Till Bousquet, Audrey Favrelle-Huret, Philippe Zinck, Sylvain Pellegrini, and Lydie Pelinski*[a]

Dedication ((optional))

Abstract: A one-pot synthesis of 2,5-disubstituted furans from γ -ketoacids is reported. In situ formation of allenates by action of chloroformate on carboxylic acid following by enolization of ketone affords to furan derivatives by cyclization. The reaction was extended on a wide scope of ketoacids and phosphonium salts. This methodology was applied on levulinic acid and derivatives, one of the biosourced platform chemicals.

Substituted furans represent an important class of heterocycles, often found in bioactive natural products and pharmacologically active compounds.^[1] As such the attractiveness of furan-containing structures lead to a continuous development of new efficient methods for the synthesis of this heterocycle.^[2] Among the many approaches explored towards their synthesis, homogeneous metal-catalyzed reactions are attractive due to their efficiency and functional group compatibility.^[3] Particularly, a strategy involving cycloisomerization of alkyne using transition-metal catalysts such as rhodium, copper, gold, and palladium complexes is the most studied.^[4] In contrast to metal-mediated processes, organocatalytic approaches to the construction of substituted furan are also described.^[5] Recently, a convenient scandium(III) triflate-catalyzed synthesis of 2,5-disubstituted furans was developed by employing oxiranes with γ -hydroxyketones.^[6]

Allenes are highly valuable synthetic precursors in preparative organic chemistry due to their high reactivity and rich chemistry associated with the two orthogonal carbon-carbon double bonds.^[7] Among the transformations investigated, the transition-metal-catalyzed intramolecular hydrofunctionalization of allenes with heteroatom nucleophiles displays an elegant approach for the synthesis of heterocycles.^[8] Particularly, allenyl ketones or alcohols (Scheme 1, pathway a)^[9] and allenyl amides (Scheme 1, pathway b)^[10] are common starting materials for transition-metal catalyzed furan synthesis.^[11] More recently, multisubstituted furans were obtained by reaction of an allenate with α -halocarbonyl ketones or dicarbonyl compounds (Scheme 1, pathway c).^[12]

Previous work

This work

Scheme 1. Previous studies and our contribution on furan synthesis

Despite these processes, there is still a requirement to develop new and efficient strategies particularly based on easily accessible starting materials. To this end, we developed a one-pot synthesis of 2,5-disubstituted furans involving an unprecedented formation of allenes as key intermediates (Scheme 1). It is worth noting that the most useful synthesis of allenes consists of a Wittig reaction by reacting a phosphorane or phosphonium salt with activated carboxylic acids such as acid chlorides.^[13] In our study, we propose the possibility to use chloroformates as an alternative to thionyl chloride and oxalyl chloride as activating reagent of carboxylic acid.

We initiated our study with 3-benzoyl propionic acid **1a** as a model substrate. In a typical procedure, the ketoacid **1a** (1 equiv) was first activated by the chloroformate **2** (1 equiv) in the presence of triethylamine (2 equiv) in dichloromethane at 0 °C. After 10 min, the phosphonium salt **3** (0.5 equiv) was added and the mixture was refluxed for 10 h (Table 1). After screening various chloroformates (entries 1-4), we discovered that the desired furan product **4a** was formed in 82% yield in the presence of ethyl chloroformate (entry 2). When the reaction was performed with an equimolar quantity of ketoacid **1a** and phosphonium salt, the yield

[a] Dr. Y. Bernhard, J. Gilbert, Dr. T. Bousquet, Dr A. Favrelle-Huret, Prof. P. Zinck, Dr. S. Pellegrini, Prof. Dr. L. Pelinski
Univ. Lille, ENSCL, Centrale Lille, Univ. Artois, UMR 8181 –UCCS–
Unité de Catalyse et Chimie du Solide, F-59000 Lille, France
E-mail: lydie.pelinski@univ-lille.fr
<http://uccs.univ-lille1.fr/index.php/fr/catalyse-chimie-moleculaire/caseco>

Supporting information for this article is given via a link at the end of the document. ((Please delete this text if not appropriate))

COMMUNICATION

dropped from 82 to 62% (Entry 5). Likewise, the reaction did not proceed to completion at room temperature (entry 6). With the exception of replacing the dichloromethane by dimethylformamide (entry 12), the solvents were found to decrease the yields of the transformation of ketoacid (entries 7–11).

The mechanism suggested for this transformation is outlined in Scheme 2. In a first step, the ketoacid reacts with ethyl chloroformate in the presence of triethylamine to form the mixed anhydride carbonate. The latter by action of the phosphonium salt leads to the allene which readily undergoes intramolecular cyclization to furnish the furan derivative. It is worth noting that the assumption of allene formation was confirmed when an additional reaction was set from 3-phenylpropionic acid. In fact, this substrate, which is unable to perform the last cyclization step, allows to isolate the allene in a 60% yield.

Scheme 2. Proposed mechanism

Table 1. Optimisation of the reaction conditions^[a]

$\text{Ph-C(=O)-CH}_2\text{-C(=O)OH} \xrightarrow[2) \text{ BrP(Ph)}_3^+\text{CH}_2\text{CO}_2\text{Et } \mathbf{3} \text{ (0.5 equiv), 45 }^\circ\text{C, 10 h}]{1) \text{ ClCO}_2\text{R } \mathbf{2} \text{ (1 equiv), NEt}_3 \text{ (2 equiv), solvent, 0 }^\circ\text{C, 10 min}}$			
1a			4a
Entry	2	Solvent	Yield [%] ^[b]
1	ClCO ₂ Me	CH ₂ Cl ₂	68
2	ClCO ₂ Et	CH ₂ Cl ₂	82
3	ClCO ₂ <i>i</i> -Bu	CH ₂ Cl ₂	68
4	ClCO ₂ <i>p</i> -NO ₂ Ph	CH ₂ Cl ₂	0
5 ^[c]	ClCO ₂ Et	CH ₂ Cl ₂	62
6 ^[d]	ClCO ₂ Et	CH ₂ Cl ₂	38
7	ClCO ₂ Et	CHCl ₃	62
8	ClCO ₂ Et	CH ₃ CN	76
9	ClCO ₂ Et	Dioxane	38
10	ClCO ₂ Et	1,2-DCE	70
11	ClCO ₂ Et	Toluene	23
12	ClCO ₂ Et	DMF	79

[a] Reaction conditions: **1a** (1.28 mmol), **2** (1.28 mmol), NEt₃ (300 μL), **3** (0.64 mmol), solvent (10 mL). [b] Isolated yield. [c] In presence of 1.28 mmol of **3**. [d] RT.

With the optimized conditions in hand, we began to investigate the scope of the method. At first, different kinds of substituted benzoylpropionic acids, easily prepared by a Friedel-Crafts reaction between substituted benzene derivatives and succinic anhydride, were evaluated. As shown in Table 2, a broad range of benzoylpropionic acids bearing electron-donating and -withdrawing substituents reacted smoothly with the phosphonium salt **3** in moderate to good yields. The best result of 93% yield was obtained from the meta-nitro substituted substrate. Additionally, electron donating groups such as methoxy, alkyl substituents are compatible with the reaction conditions. However, the lower 56% yield observed for the synthesis of **4d** was likely due to the steric hindrance of the methoxy moiety in the ortho position. The presence of halogens on the benzene ring did not affect the yield (73% for **4g** and 70% for **4h**). The ferrocenyl furan **4j** was obtained in only 35% yield due to heat decomposition.

The scope of the reaction was then extended to various phosphonium salts containing other functions such as ketones or nitrile (Table 3). Apart from furan **6a** (56% yield), the condensation of phosphonium salt **5** led to furan derivatives in good 65–86% yields.

Table 2. Scope of the ketoacid substrates

COMMUNICATION

[a] Reaction conditions: **1** (1.28 mmol), **2** (140 μ L), NEt₃ (300 μ L), **3** (0.64 mmol), solvent (10 mL). Yields of the isolated product are given.

Table 3. Scope of the phosphonium salt

[a] Reaction conditions: **1a** (1.28 mmol), **2** (140 μ L), NEt₃ (300 μ L), **5** (0.64 mmol), solvent (10 mL). Yields of the isolated product are given.

Finally, this new synthetic method was performed from the levulinic acid **7**, a promising biosourced platform chemical (Scheme 3).^[14] The furan **4l** was then isolated in a good yield of 65%. Additionally, the ketoacid **10**, obtained by reaction of ethyl levulinate **9** on benzaldehyde following by hydrolysis of ester, led the the furan **4m** with a disappointing yield of 34%. A degradation of the mixture was observed during the synthesis. The reaction of levulinic acid-based phosphonium salt **11** on benzoylpropionic acid **1** led to the furanyl ketoester **6g** in a modest yield of 46%.

Scheme 3. Furan synthesis starting from levulinic acid derivatives

In conclusion, we have developed an efficient method for the synthesis of 2,5-disubstituted furans from γ -ketoacids. The reaction is operationally simple and presents a wide substrate scope. This methodology was further extended to levulinic acid and derivatives in order to valorize a promising biobased platform chemical.

Experimental Section

Typical procedure for the one-pot synthesis of 2,5-disubstituted furans: to ketoacid (1.28 mmol) in CH₂Cl₂ (10 mL) was added triethylamine (2.15 mmol) then ethylchloroformate (1.47 mmol) at 0 °C. After stirring for 10 min, the phosphonium salt (0.64 mmol) was added and the mixture was heated under reflux for 10 h. The mixture was neutralized by aqueous HCl solution 1 M and extracted with CH₂Cl₂ (3 x 15 mL). The combined organic layers were dried over MgSO₄ and concentrated in vacuo. The product was purified by column chromatography over silica gel (eluent: petroleum ether/diethylether 9/1 then petroleum ether/diethylether 7/3) to give the furan derivatives.

Full experimental details and characterization data are given in the Supporting Information.

Acknowledgements

This work was financially supported by the FWV ALPO Interreg Grant. Chevreul institute (FR 2638), Ministère de l'Enseignement Supérieur et de la Recherche, Région Nord – Pas-de-Calais and FEDER are acknowledged for supporting and funding this work. We also would like to thank Céline Delabre for her technical support.

Keywords: Furan derivatives • phosphonium salt • allene intermediate • levulinic acid

[1] a) B. A. Keay, J. M. Hopkins, P. W. Dibble, In *Comprehensive Heterocyclic Chemistry III*, Vol. 3 (Eds.: A. R. Katritzky, C. A. Ramsden,

COMMUNICATION

- E. F. V. Scriven, R. J. K. Taylor), Elsevier, Amsterdam, **2008**, pp 571; b) K.-S. Yeung, X.-S. Peng, J. Wu, R. Fan, X.-L. Hou, *Prog. Heterocycl. Chem.* **2013**, 25, 183; c) Y. Wang, Y.-C. Luo, X.-Q. Hu, P.-F. Xu, *Org. Lett.* **2011**, 13, 5346; d) C.-C. Wu, W.-Y. Hung, T.-L. Liu, L.-Z. Zhang, T.-Y. Luh, *J. Appl. Phys.* **2003**, 93, 5465; e) M. D. Mullican, R. J. Sorenson, D. T. Connor, D. O. Thueson, J. A. Kennedy, M. C. Conroy, *J. Med. Chem.* **1991**, 34, 2186; f) S. N. Abramson, J. A. Trischman, D. M. Tapiolas, E. E. Harold, W. Fenical, P. Taylor, *J. Med. Chem.* **1991**, 34, 1798; g) B. Tang, C. D. Bray, G. Pattenden, J. Rogers, *Tetrahedron* **2010**, 66, 2492; h) S. M. Rahmathullah, J. E. Hall, B. C. Bender, D. R. McCurdy, R. R. Tidwell, D. W. Boykin, *J. Med. Chem.* **1999**, 42, 3994; i) D. S. Mortensen, A. L. Rodriguez, K. E. Carlson, J. Sun, B. S. Katzenellenbogen, J. Katzenellenbogen, *J. Med. Chem.* **2001**, 44, 3838.
- [2] a) J. Shen, J. Zhao, B. Hu, Y. Chen, L. Wu, Q. You, L. Zhao, *Green Chem.* **2018**, 20, 600; b) S. F. Kirsch, *Org. Biomol. Chem.* **2006**, 4, 2076; c) T. J. Donohoe, J. F. Bower, *Proc. Natl. Acad. Sci. U. S. A.* **2010**, 107, 3373; d) Keay, B. A. *Chem. Soc. Rev.* **1999**, 28, 209; e) Y. Xiao, J. Zhang, *Chem. Commun.* **2009**, 3594.
- [3] a) A. S. Karpov, E. Merkul, T. Oeser, T. J. J. Müller, *Chem. Commun.* **2005**, 2581; b) A. V. Gulevich, A. S. Dudnik, N. Chernyak, V. Gevorgyan, *Chem. Rev.* **2013**, 113, 3084; c) S. P. Morcillo, D. Leboeuf, C. Bour, V. Gandon, *Chem. Eur. J.* **2016**, 22, 16974.
- [4] a) M. Zhang, H.-F. Jiang, H. Neumann, M. Beller, P. H. Dixneuf, *Angew. Chem., Int. Ed.* **2009**, 48, 1681; b) A. V. Kellin, V. Gevorgyan, *J. Org. Chem.* **2002**, 67, 95; c) J. T. Kim, A. V. Kellin, V. Gevorgyan, *Angew. Chem., Int. Ed.* **2003**, 42, 98; d) A. W. Sromek, A. V. Kellin, V. Gevorgyan, *Angew. Chem., Int. Ed.* **2004**, 43, 2280; e) A. Gille, J. Rehbein, M. Hiersemann, *Org. Lett.* **2011**, 13, 2122; f) M. H. Suhre, M. Reif, S. F. Kirsch, *Org. Lett.* **2005**, 7, 3925; g) H. Cao, H. Jiang, R. Mai, S. Zhu, C. Qia, *Adv. Synth. Catal.* **2010**, 352, 143; h) C. Arroniz, G. Chaubet, E. A. Anderson, *ACS Catal.* **2018**, 8, 8290.
- [5] a) C.-K. Jung, J.-C. Wang, M. J. Krische, *J. Am. Chem. Soc.* **2004**, 126, 4118; b) L. Albrecht, L. K. Ransborg, B. Gschwend, K. A. Jørgensen, *J. Am. Chem. Soc.* **2010**, 132, 17886; c) H. Kuroda, E. Hanaki, H. Izawa, M. Kano, H. Itahashi, *Tetrahedron* **2004**, 60, 1913; d) J. S. Clark, A. Boyer, A. Aimon, P. E. Garcia, D. M. Lindsay, A. D. F. Symington, Y. Danoy, *Angew. Chem. Int. Ed.* **2012**, 51, 12128.
- [6] K. Mondal, S. C. Pan, *J. Org. Chem.* **2017**, 82, 4415.
- [7] a) S. Ma, *Chem. Rev.* **2005**, 105, 2829; b) S. Yu, S. Ma, *Angew. Chem. Int. Ed.* **2012**, 51, 3074; c) B. Alcaide, P. Almendros, C. Aragoncillo, *Chem. Soc. Rev.* **2014**, 43, 3106; d) R. Zimmer, H.-U. Reissig, *Chem. Soc. Rev.* **2014**, 43, 2888; e) K. M. Brummond, J. E. DeForrest, *Synthesis* **2007**, 6, 795.
- [8] a) O. El-Sepelgy, A. Brzozowska, J. Sklyaruk, Y. K. Jang, V. Zubar, M. Rueping, *Org. Lett.* **2018**, 20, 696; b) S. Ma, *Acc. Chem. Res.* **2003**, 36, 701; c) S. Pashikanti, J. A. Calderone, M. K. Nguyen, C. D. Sibley, W. L. Santos, *Org. Lett.* **2016**, 18, 2443; d) Z. Zhang, C. Liu, R. E. Kinder, X. Han, H. Qian, R. A. Widenhoefer, *J. Am. Chem. Soc.* **2006**, 128, 9066; e) Z. Zhang, R. A. Widenhoefer, *Angew. Chem. Int. Ed.* **2007**, 46, 283.
- [9] a) N. Krause, C. Winter, *Chem. Rev.* **2011**, 111, 1994; b) A. W. Sromek, M. Rubina, V. Gevorgyan, *J. Am. Chem. Soc.* **2005**, 127, 10500; c) S. Ma, J. Zhang, L. Lu, *Chem.-Eur. J.* **2003**, 9, 2447; d) J. A. Marshall, X.-J. Wang, *J. Org. Chem.* **1991**, 56, 960; e) J. A. Marshall, E. D. Robinson, *J. Org. Chem.* **1990**, 55, 3450; f) A. S. K. Hashmi, L. Schwarz, J.-H. Choi, T. M. Frost, *Angew. Chem. Int. Ed.* **2000**, 39, 2285; g) Y. Li, J. P. Brand, J. Waser, *Angew. Chem. Int. Ed.* **2013**, 52, 6743; h) A. S. K. Hashmi, *Angew. Chem. Int. Ed.* **1995**, 34, 1581; i) Y. Xia, A. S. Dudnik, V. Gevorgyan, Y. Li, *J. Am. Chem. Soc.* **2008**, 130, 6940; j) Z. Zhang, C. Liu, R. E. Kinder, X. Han, H. Qian, R. A. Widenhoefer, *J. Am. Chem. Soc.* **2006**, 128, 9066; k) Z. Zhang, R. A. Widenhoefer, *Angew. Chem. Int. Ed.* **2007**, 46, 283.
- [10] a) J. Zhang, M. Wu, W. Lu, S. Wang, Y. Zhang, C. Cheng, G. Zhu, *J. Org. Chem.* **2017**, 82, 11134; b) Y. Wan, J. Zhang, Y. Chen, L. Kong, F. Luo, G. Zhu, *Org. Biomol. Chem.* **2017**, 15, 7204; c) A. Kondoh, S. Ishikawa, T. Aoki, M. Terada, *Chem. Commun.* **2016**, 52, 12513.
- [11] a) A. S. K. Hashmi, in *Modern Allene Chemistry*, Vol. 2 (Eds.: Norbert Krause and A. Stephen K. Hashmi), Wiley-VCH, Weinheim, **2004**, pp 889; b) R. C. D. Brown, *Angew. Chem. Int. Ed.* **2005**, 44, 850.
- [12] D. Bakshi, A. Singh, *Org. Biomol. Chem.* **2017**, 15, 3175.
- [13] a) R. W. Lang, H.-J. Hansen, *Helv. Chim. Acta* **1980**, 63, 438; b) H.-J. Bestmann, H. Hartung, *Chem. Ber.* **1966**, 99, 1198; c) E. Kohl-Mines, H.-J. Hansen, *Helv. Chim. Acta* **1985**, 68, 2244; d) R. W. Lang, H.-J. Hansen, *Org. Synth.* **1984**, 62, 202; e) K. Kumar, S. Kaur, M. P. S. Ishar, *Synlett* **1999**, 1237.
- [14] a) U. Omoruyi, S. Page, J. Hallett, P. W. Miller, *ChemSusChem* **2016**, 9, 2037; b) B. V. Timokhin, V. A. Baransky, G. D. Eliseeva, *Russ. Chem. Rev.* **1999**, 68, 73; c) Z. Xue, Q. Liu, J. Wang, T. Mu, *Green Chem.* **2018**, 20, 4391; d) L. Yan, Q. Yao, Y. Fu, *Green Chem.*, **2017**, 19, 5527-5547; e) L. Hu, J. Xu, S. Zhou, A. He, X. Tang, L. Lin, J. Xu, Y. Zhao, *ACS Catal.* **2018**, 8, 2959.

COMMUNICATION

Entry for the Table of Contents (Please choose one layout)

Layout 1:

COMMUNICATION

Text for Table of Contents

Page No. – Page No.

Title

((Insert TOC Graphic here))

Layout 2:

COMMUNICATION

Y. Bernhard, J. Gilbert, T. Bousquet, A. Favrelle-Huret, P. Zinck, S. Pellegrini, L. Pelinski *

Page No. – Page No.

Title

Furan synthesis: A one-pot synthesis of 2,5-disubstituted furans from γ -ketoacids is reported. In situ formation of allenoates by action of chloroformate on carboxylic acid following by enolization of ketone affords to furan derivatives by cyclization.