1 **The *foraging* gene as a modulator of division of labour in social insects**

2 Christophe Lucas^a and Yehuda Ben-Shahar^b

3 *^aInstitut de Recherche sur la Biologie de l’Insecte (UMR7261), CNRS – University of Tours,*
4 *Tours, France; ^bDepartment of Biology, Washington University in St. Louis, St. Louis MO*
5 *63130, USA*

6 [^achristophe.lucas@univ-tours.fr](mailto:christophe.lucas@univ-tours.fr)

7 [^bbenshahary@wustl.edu](mailto:benshahary@wustl.edu)

8 **The *foraging* gene as a modulator of division of labour in social insects**

9 **Abstract**

10 The social ants, bees, wasps, and termites include some of the most ecologically-successful
11 groups of animal species. Their dominance in most terrestrial environments is attributed to their
12 social lifestyle, which enable their colonies to exploit environmental resources with remarkable
13 efficiency. One key attribute of social insect colonies is the division of labour that emerges
14 among the sterile workers, which represent the majority of colony members. Studies of the
15 mechanisms that drive division of labour systems across diverse social species have provided
16 fundamental insights into the developmental, physiological, molecular, and genomic processes
17 that regulate sociality, and the possible genetic routes that may have led to its evolution from a
18 solitary ancestor. Here we specifically discuss the conserved role of the *foraging* gene, which
19 encodes a cGMP-dependent protein kinase (PKG). Originally identified as a behaviourally
20 polymorphic gene that drives alternative foraging strategies in the fruit fly *Drosophila*
21 *melanogaster*, changes in *foraging* expression and kinase activity were later shown to play a key
22 role in the division of labour in diverse social insect species as well. In particular, *foraging*
23 appears to regulate worker transitions between behavioural tasks and specific behavioural traits
24 associated with morphological castes. Although the specific neuroethological role of *foraging* in
25 the insect brain remains mostly unknown, studies in genetically tractable insect species indicate
26 that PKG signalling plays a conserved role in the neuronal plasticity of sensory, cognitive and
27 motor functions, which underlie behaviours relevant to division of labour, including appetitive
28 learning, aggression, stress response, phototaxis, and the response to pheromones.

29

30 Keywords (5 to 6): *foraging* gene; sociality; insects; behaviour; division of
31 labour

32

33 Subject classification codes: review

Introduction

Animal social grouping is common because it increases the fitness of individuals by, for example, maximizing their access to resources and providing improved defences against predators (Wilson, 1987). Some of the most derived forms of sociality are exhibited by species of eusocial insects in the orders Hymenoptera (ants, bees, wasps) and Isoptera (termites), which together are estimated to represent more than 30% of the global animal biomass (Hölldobler & Wilson, 2009). All social insect colonies exhibit reproductive division of labour, in which majority of a colony is comprised of sterile female workers and just one or few reproductive individuals (the “queen” in social Hymenoptera and the “royal couple” in termites) (Oster & Wilson, 1978). The sterile workers, in turn, self-organize to become specialized in different tasks that are necessary for the growth and maintenance of the colony such as foraging, brood care and colony defence (Corona et al., 2016; Gordon, 2016; Leitner & Dornhaus, 2019). Although division of labour among workers is common across many different social insect taxa, different species show varying levels of individual specializations. At one extreme, workers might exhibit almost no task specialization, while at the other extreme workers might differentiate into developmentally-determined morphological castes (Oster & Wilson, 1978) (Figure 1).

Despite the overwhelming complexity associated with insect sociality and the regulation of division of labour, the evolutionary emergence of division of labour systems across different social insect species most likely depended on the co-option of existing sensory pathways, and associated modulatory neural circuits, to generate differential behavioural syndromes across individuals that share a common social and physical environment. Subsequently, although sensory stimuli for all tasks are simultaneously present in the colony, division of labour systems maintain differential behavioural response thresholds to task specific cues in different individuals (Figure 1). The highly-tuned network of semi-autonomous individuals provides a robust, systems-level modulation of task allocation across

time and space based on the overall needs of the colony (Buttstedt et al., 2016; Haydak, 1970; Krista K Ingram, 2002; Kamakura, 2011; Romain Libbrecht & Keller, 2013; Nygaard et al., 2011; Patel et al., 1960; Schwander & Keller, 2008; Schwander et al., 2010; Wheeler, 1991). Phylogenetic studies indicate that division of labour systems likely evolved multiple times across, and within, different lineages of the social hymenopterans (Gordon, 2016; Rueffler et al., 2012). Therefore, division of labour in colonies of social insects has emerged as an excellent model for understanding how animal social systems have evolved and are regulated at the physiological, molecular, and cellular levels (Kapheim et al., 2015; Robinson et al., 2005; Smith et al., 2008; Toth & Robinson, 2007; A. L. Toth et al., 2010; Whitfield et al., 2006). Transcriptomic analyses of gene expression in the brains of workers that are engaged in specific tasks, for example, have indicated that division of labour depends on broad and complex changes in many different neuronal signalling pathways across different parts of the brain and other tissues (Figure 1). This appears to occur via differential promoter activity, regulatory non-coding miRNAs, and epigenetic processes (Cash et al., 2005; Greenberg et al., 2012; K. M. Kapheim et al., 2020; Lattorff & Moritz, 2013; Simola et al., 2016; Sinha et al., 2006; Toth et al., 2007; Whitfield et al., 2006). To date, changes in the expression levels of several genes have been associated with division of labour across diverse social insect species. However, only a few have been studied in any depth. The highly conserved cGMP-dependent protein kinase gene *foraging* (*for*), in particular, has been well-studied for its role in regulating division of labour. Previous work has established a causal association between *for* and task specialization, which is conserved across multiple independently-evolved lineages. Here we review the role of *for*, first identified as a behaviourally polymorphic gene in *Drosophila* associated with larval and adult foraging decisions (J. S. de Belle et al., 1989; K. A. Osborne et al., 1997), as a conserved signalling pathway that has been co-opted for the regulation of division of labour across diverse social insect species.

The study of sociality in the era of molecular genetics

Over 50 years ago, the pioneering ethologist Niko Tinbergen published his seminal paper entitled ‘On aims and methods of ethology’ (Burkhardt, 2014; Tinbergen, 1963). With brilliant simplicity, Tinbergen argued that if biologists want to really understand “behaviour” then they need to ask the following four questions (rephrased): 1) What is the studied trait good for (its impact on fitness)? 2) How does it develop during the lifetime of an individual (development/ontogeny)? 3) How did it evolve over the history of the species (trait phylogeny)? 4) How does it work (mechanism/causation)? The emergence of modern neuroscience and neurogenetics followed in the footsteps of ethology by providing a mechanistic framework as a powerful approach to the design of behavioural studies and their interpretation in the context of brain functions (Tinbergen question #4).

However, modern neuroscience research is now largely framed in the context of causation and mechanism (Tinbergen question #4). Consequently, the diversity of animal species used for basic neurogenetic research has, historically, been reduced to just a few genetically tractable species that have sequenced genomes, including the roundworm *Caenorhabditis elegans* (Bargmann, 1998), the fruit fly *Drosophila melanogaster* (Bellen et al., 2010), the zebrafish *Danio rerio* (Stewart et al., 2014), and the laboratory mouse *Mus musculus* (Lehner, 2013). While there is no doubt that these models have been instrumental in the phenomenal progress made in cellular and molecular neuroscience over the past three decades, the decline in model diversity is clearly an unfortunate side effect, and has negatively impacted our ability to understand behaviour in ecologically and evolutionary relevant contexts (Fitzpatrick et al., 2005; Walton et al., 2020).

Yet, recent advances in genetic and genomic approaches have enabled investigators to adopt modern molecular approaches for understanding the behaviour of a wider assortment of animal species in the lab and in their natural environments. Some of the first non-drosophilid

insects for which genomic and molecular tools for studying behaviour have made a tremendous impact are the social insects, particularly in the several independent social clades within the order Hymenoptera. One of the best established model species for sociogenomic and genetic studies is the European honey bee *Apis mellifera*. As the most dominant domesticated insect, the long tradition of beekeeping has provided investigators with a rich knowledge of its behaviour and physiology, including well-established techniques for manipulating colony demography and colony genetics via artificial inseminations (Page et al., 2002; Vanengelsdorp & Meixner, 2010). Not surprisingly, the economic importance of the honey bee has led to the development of diverse molecular and genetic resources, including a relatively well-annotated genome, high-resolution population genetics, and successful genome transformations (Ament et al., 2010; Ben-Shahar, 2014; Ben-Shahar et al., 2004; Chen et al., 2021; Foret et al., 2009; Sinha et al., 2006; Y. Wang et al., 2006; Weinstock et al., 2006; Whitfield et al., 2006; Whitfield et al., 2003). Subsequently, the progress in analysing honey bee behaviour with molecular and genetic tools has paved the way for similar breakthroughs in many additional social insect species (Karen M. Kapheim et al., 2020; Kapheim et al., 2015; R. Libbrecht et al., 2013; Sadd et al., 2015; A. L. Toth et al., 2010). Nevertheless, the genetic mechanisms by which specific genes and neuronal signalling pathways regulate the complex behaviour of social insects had remained mostly a mystery until the discovery of the gene *foraging* (*for*), a cGMP-dependent protein kinase (PKG), and its role in the regulation of behavioural plasticity in colonies of social insects (Figure 1D). Subsequent studies of this gene in the context of division of labour have provided some important mechanistic insights into animal sociality.

The role of *foraging* in regulating division of labour in social insect colonies

The *foraging* (*for*) gene encodes a cGMP-dependent protein kinase (K. Osborne et al., 1997), a highly conserved enzyme present in all animal genomes sequenced to date (Anreiter &

135 Sokolowski, 2019; Lucas, Kornfein, et al., 2010). The *for* gene was first identified as a
136 naturally polymorphic locus that regulates larval foraging behaviour in *Drosophila* (J.S. de
137 Belle & Sokolowski, 1987; Sokolowski, 1980), and was later identified as a gene that encodes
138 one of the two cGMP-dependent protein kinases (PKG) encoded by the *Drosophila* genome
139 (K. Osborne et al., 1997). The highly conserved PKG family is broadly expressed in both
140 neuronal and non-neuronal tissues, and has been implicated in numerous cellular and
141 physiological functions via its kinase activity (Dason et al., 2019; Douglas et al., 2005;
142 Francis et al., 2010; Lohmann et al., 1997; X. Wang & Robinson, 1997). In *Drosophila*, *for* is
143 expressed in diverse tissues, including the peripheral (PNS) and central (CNS) nervous
144 systems of both larvae and adults (Anreiter et al., 2017). Several genetic studies have revealed
145 that the natural polymorphism in *for* contributes to diverse sensory functions, including the
146 visual, olfaction, and gustatory pathways (Lin et al., 1992). In the CNS, *for* is expressed in the
147 Kenyon cells of the mushroom bodies, the central complex, and the antennal lobes. It has
148 been implicated in regulating neuronal and behavioural plasticity in diverse cognitive and
149 motor functions such as neuronal sensitization and habituation, learning and memory, and
150 sleep (Eddison et al., 2012; Engel et al., 2000; Scheiner et al., 2004). In addition to regulating
151 behaviour via neuronal functions, *for* also indirectly contributes to feeding and foraging
152 behaviours via its role in metabolic plasticity and the response to stress (Dason et al., 2020;
153 Kaun et al., 2008; Kaun & Sokolowski, 2009; Kent et al., 2009).

154 The role of the cGMP-PKG signalling axis in regulating feeding behaviours is conserved in
155 other animal species. For example, cGMP-signalling has been shown to regulate the response
156 threshold of insects to pheromones (Boekhoff et al., 1993; Ziegelberger et al., 1990), and to
157 influence olfactory communication in rats (Kroner et al., 1996; Moon et al., 1998). In the
158 worm *Caenorhabditis elegans*, the PKG-ortholog *egl-4* plays a role in foraging, olfaction,
159 sleep, body size and lifespan (Fujiwara et al., 2002; L'Etoile et al., 2002; Raizen et al., 2008).

Yet, in spite of its broad role in regulating different forms of behavioural plasticity in diverse animal species, the identities of the specific neuronal substrates modulated by *for* in the context of foraging behaviour and social insect division of labour remain mostly unknown. However, studies in genetically tractable species provide some mechanistic insights into how PKG signalling might be regulating neuronal processes associated with division labour. For example, studies in the mouse, indicate that PKG activity can have broad effects on long-term neuronal processes and behavioural plasticity via diverse downstream effectors. Neuronal PKG activity can modulate long-term potentiation (LTP) in the mammalian brain via a direct modulation of cAMP-phosphodiesterases, which regulate cAMP levels and synaptic plasticity (Bollen et al., 2014). In addition, studies of *Egl-4*, the ortholog of *for* in the worm *C. elegans*, has shown that sensory-induced neuronal plasticity within the chemosensory system depends on the transition of activated PKG to the nucleus, which leads to changes in neuronal gene expression via the modulation of heterochromatin binding proteins (Lee et al., 2010). Together, these examples illustrate the broad role *for* and related kinase genes play in regulating neuronal plasticity and organismal behaviour via diverse molecular and cellular mechanisms in the peripheral and central nervous systems of invertebrates and vertebrates.

The *foraging* gene in social bees

The realization that *for* plays a key role in regulating feeding-related behavioural plasticity in *Drosophila* suggested that the phylogenetically conserved PKG signalling pathway is also likely to be driving similar foraging-related decisions in other insect species, including in the regulation of foraging vs. brood care behaviour in social insect division of labour. Recently, *for* has also been implicated in the regulation of social networks in *Drosophila* (Alwash et al., 2021). By using transcriptomics and gene candidate approaches (Fitzpatrick et al., 2005), multiple studies have identified several key genes that play a role in regulating division of labour in honey bees including the acetylcholine esterase (*AchE*) gene (Shapira et

al., 2001), the *foraging* (*for*) gene (Ben-Shahar et al., 2003; Ben-Shahar et al., 2002), and the manganese transporter *malvolio* (Ben-Shahar et al., 2004). In terms of how *for* exerts its action in honey bees, follow-up studies have identified an increase in brain expression levels of *for* just prior to the initiation of foraging, which suggests that its activity may serve as a trigger for the transition from in-hive behaviours to foraging outside (Heylen et al., 2008). Furthermore, QTL analysis of foraging-related traits in honey bee populations has identified an association with QTLs in the *for* locus, suggesting that in addition to its role at the developmental and physiological timescales, *for* may play a role in shaping division of labour at the evolutionary timescale as well (Ruppell et al., 2004).

The discovery of *for* as a putative key regulator of division of labour in the honey bee was followed by studies investigating its role in regulating division of labour across the social hymenopteran phylogeny. It was shown that changes in *for* expression are also associated with division of labour in other species of social bees, including the Asian honey bee *A. cerana* (Ma et al., 2018), as well as different bumblebee species (Kodaira et al., 2009; Tobback et al., 2011). Interestingly, in contrast to the reported increased *for* expression in the brains of honey bee foragers, the expression of *for* in association with the size-dependent regulation of division of labour in bumblebees was variable, indicating higher *for* expression in *Bombus terrestris* foragers but lower in brains of *B. ignitus* foragers (Couvillon et al., 2010; Kodaira et al., 2009; Shpigler et al., 2013). Together, these studies suggest that the functional relationship between *for* activity in the brain and the behavioural states that define the probability of performing specific tasks by individual bees is complex and likely involve both central and peripheral neuronal pathways.

The *foraging* gene in ants

In ants, *for* has been shown to be involved in age-related division of labour, as in honey bees (Krista K Ingram et al., 2016; K. K. Ingram et al., 2005; Lucas et al., 2015; Manfredini et al., 2014; Oettler et al., 2015). However, in contrast to social bees, which primarily regulate division of labour on a physiological timescale, studies of *for* in ants revealed its role in regulating polyethisms in association with both age-dependent and developmentally-regulated morphological castes. For example, colonies of the ant *Pheidole pallidula* have two morphologically-distinct workers (Passera, 1985), each associated with specific behavioural repertoires. The majors have a large head with large mandibles and are mainly involved in the defence of the colony and processing of large food items, while the minors are smaller and are mainly involved in brood care and foraging. This morphological and behavioural division of labour was associated with changes in brain activity; minors had lower brain PKG activity than majors (Lucas & Sokolowski, 2009). In addition, although minors and majors typically perform distinct roles in the colony, when confronted with changing needs of the colony, majors may assist minors in foraging activities and minors may help with defence tasks (Aarab, 1991). Accordingly, experimental manipulations showed that majors that respond to foraging stimuli show minor-like, low brain PKG activity. Similarly, when participating in colony defence, minors showed a significant increase in brain PKG activity to major-like levels (Lucas & Sokolowski, 2009). Moreover, pharmacological treatment of minors with the PKG activator 8-Br-cGMP induced brain PKG activity, and led to an increase in defence and decrease in foraging behaviours (Lucas & Sokolowski, 2009), thus establishing a causal association between PKG activity and worker behavioural plasticity. The *foraging* gene in *P. pallidula* therefore appears to modulate both foraging behaviour and defence behaviour (Anreiter & Sokolowski, 2019; Lucas, Hughson, et al., 2010; Lucas & Sokolowski, 2009) and, under conditions that require plastic task reallocation, can drive neuronal plasticity even in species with a robust morphological division of labour

system (Anreiter & Sokolowski, 2019; Lucas, Hughson, et al., 2010). These findings suggest that *for* exerts at least two levels of control over worker behaviour in ants: (1) Differences in baseline brain PKG activity are associated with caste polyethism at the developmental timescale; and (2) Plasticity in PKG activity allows colony-level responses to environmental changes at the physiological timescale. This link between defence and PKG activity has also been demonstrated in a field study, where feeding the ant *Allomerus octoarticulatus* with a PKG activator, increased defence behaviour of the ant-plant *Cordia nodosa* against herbivores, suggesting a molecular basis for ant-plant mutualism (Malé et al., 2017).

In another ant species, *Solenopsis invicta*, *foraging* gene expression in workers was influenced by colony demography, in particular whether the colony had one queen (monogynous) or multiple queens (polygynous) (Lucas et al., 2015). Workers from monogynous colonies of *S. invicta*, which are very territorial and aggressive (Chirino et al., 2012; Tschinkel, 2006), had higher brain expression levels of *for* relative to workers from the more docile polygynous colonies (Lucas et al., 2015). Altogether, these results suggest that the *for* gene might be involved in regulating differential levels of aggression associated with territorial foraging or caste polyethisms. Recently, *for* has been shown to regulate aggression in flies as well, providing further support to the idea that the role of *for* in regulating behavioural plasticity across distant species is conserved not only for foraging behaviour but may be also for defence behaviour (S. Wang & Sokolowski, 2017).

Moreover in the ant *S. invicta*, *for* expression level was shown to be associated with reproductive state (future queen vs. workers) and the number of future queens (monogynous vs. polygynous) (Lucas et al., 2015), suggesting that in addition to its role in regulating behavioural plasticity in workers, *for* also contributes to reproductive division of labour in *S. invicta*. Interestingly, variations in colony demography (monogynous vs. polygynous) exist in

several ant species, and are thought to confer colony-level benefits by increasing genetic diversity, and therefore, amplifying worker behavioural differences that underlie division of labour (Cronin et al., 2013; Krista K Ingram, 2002; Manfredini et al., 2014; Pearcy et al., 2004; Ross & Keller, 1995).

The neuroethological context of the regulation of division of labour by *for*

Based on theoretical and empirical studies, one of the primary models for self-organized division of labour in social insect colonies is based on the idea that animals at different behavioural states exhibit different behavioural response thresholds to task-specific stimuli (Beshers & Fewell, 2001; Duarte et al., 2012). Consequently, although workers are exposed to many or all task-stimuli in the colony, the probability that an individual worker will respond to the stimulus for a particular task varies (Bonabeau et al., 1998; Detrain et al., 1999; Duarte et al., 2012; Leitner & Dornhaus, 2019). Subsequently, molecular, genetic, and neuroethological studies have suggested that the PKG activity of *for* regulates division of labour, at least in part, via the modulation of task-specific sensory thresholds (Ben-Shahar et al., 2003; George et al., 2020; Heylen et al., 2008; Thamm & Scheiner, 2014).

As a kinase, *for* is likely to have numerous protein targets in the insect brain (X. Wang & Robinson, 1997). To date, the specific protein targets that mediate its role in behavioural plasticity in general, and division of labour in particular, remain mostly elusive, however, studies in various species suggest that *for* may have targets within the sensory system. Studies in *Drosophila*, for example, have indicated that *for* contributes to sucrose responsiveness in adults (Scheiner et al., 2004) and the larval nociception threshold (Dason et al., 2020). Similar effects on sensory thresholds have also been identified in *C. elegans* and rodent models, further indicating a conserved role for cGMP-dependent signalling in modulating behavioural states via sensory functions (Gangadharan et al., 2017; Jaumann et al., 2012; Levy & Bargmann, 2020). Subsequently, several studies in the honey bee have indicated that *for*

activity plays a role in regulating the response threshold to light, indicating that increased PKG activity drives positive phototaxis in foragers (Ben-Shahar, 2005; Ben-Shahar et al., 2003). In addition, *for* has also been implicated in honey bee sucrose responsiveness and the response threshold to queen pheromones (Fussnecker et al., 2011; Thamm & Scheiner, 2014). Together, these studies suggest that *for* is likely to regulate division of labour via multimodal modulation of sensory functions.

In addition to its role in regulating sensory functions, *for* has been shown to contribute to both simple forms of neuronal plasticity, such as the habituation in the giant fiber escape circuit responsiveness to sugar (Engel et al., 2000; Scheiner et al., 2004), as well as more complex processes such as visual memories (Kuntz et al., 2017; Z. Wang et al., 2008) and associative learning (Kaun et al., 2007; Mery et al., 2007) in flies. Because division of labour is often associated with differential learning and memory capacities (Ben-Shahar & Robinson, 2001; Ben-Shahar et al., 2000; Roussel et al., 2009), it has been hypothesized that *for* is also likely to also modulate division of labour via its action in central neuronal circuits associated with higher level cognitive functions (Ben-Shahar, 2005; Menzel et al., 2006). Interestingly, in *Pheidole* ants, the expression pattern of *for* in the brains of majors is specific to a subregion of the mushroom bodies, while both minor and major workers share its expression in several brain regions including the lobula, which plays a role in the integration of visual cues (Lucas & Sokolowski, 2009). Although it is still unknown how the action of *for* in this specific brain region is contributing to the behaviours exhibited by minors and majors, the well-established anatomical and physiological role of the mushroom bodies in regulating division of labour in diverse social insects suggest that *for*-dependent signalling in this brain region is important for cognitive processes associated with division of labour. This is further supported by the recent findings that *for* activity is associated with interindividual differences in the dance intensity of returning foragers (George et al., 2020), and the impact of cGMP pharmacological treatment

on spatial learning in honey bee foragers (Tsvetkov et al., 2019).

The *foraging* gene in other social contexts

The complex role of *for* in regulating division of labour in bee and ant species indicates that this gene is highly pleiotropic, regulating not only foraging but also many other behavioural aspects of division of labour and other sociality-related traits. The study of *for* in other social species, such as wasps and termites, will yield important discoveries for understanding how *for* regulates social behaviours across diverse insect taxa. For example, a study on the comparisons of brain gene expression between the social wasp *Polistes metricus* and the honey bee *Apis mellifera* allowed to identify common molecular roots for division of labour, demonstrating the importance of *for* across lineages (Amy L Toth et al., 2010). In the termite *Reticulitermes flavipes*, a recent study showed differences in mRNA expression depending on caste, temperature and photoperiod (Merchant et al., 2019), suggesting the possibility that the differential expression of *for* may very well be important for this social insect group as well. Future comparisons between termites and social hymenopteran will likely shed even more light on how *for* regulates division of labour across phylogenetically distant social systems (Hymenoptera/Isoptera) with distinct developmental processes (e.g., holometabolism vs. hemimetabolism).

Although the precise origin of insect eusociality remains mostly unknown at the molecular level, it is clear that some key neuronal signalling pathways, such as *for*, contribute to social interactions between conspecifics in non-social insects as well. For example, flies with low *for* expression are more sensitive to social context in learning paradigms than those with high expression levels (Kohn et al., 2013). In addition, *for* has been shown to be associated with population density-dependent regulation of polyphenisms in some aphid species (Tarès et al., 2013), and between solitary and gregarious forms in the locust

Schistocerca gregaria (Lucas, Kornfein, et al., 2010; Rogers et al., 2003; Tobback et al., 2013).

Concluding remarks

The association between the PKG activity of the *foraging* gene and behavioural plasticity in response to changes in the social and biotic environments, especially in the context of feeding and foraging behaviours, is conserved across a broad range of animal species, from insects to humans (Anreiter & Sokolowski, 2019; Armstrong et al., 2010; Chardonnet et al., 2014; Kohlmeier et al., 2019; Lucas, Kornfein, et al., 2010; Malé et al., 2017; Merchant et al., 2019; Robertson & Sillar, 2009; Struk et al., 2019; Tarès et al., 2013; Tobback et al., 2013). Additionally, it remains unknown whether all or only some of the alternative transcripts transcribed by *for* in at least some social insect species play a role in regulating neuronal plasticity and behaviour in social insects (Lucas & Sokolowski, 2009; Merchant et al., 2019; NCBI, 2021). Furthermore, the specific molecular targets and neuronal signalling pathways modulated by *for* activity across the evolutionary, developmental, and physiological timescales remain a mystery. Nevertheless, recent developments in transposon-mediated transgenesis and *Cas9*/CRISPR-dependent genome editing in several social insect species (Ben-Shahar, 2014; Chen et al., 2021; Kohno et al., 2016; McAfee et al., 2019; Schulte et al., 2014; Tribble et al., 2017) will likely yield new insights into the molecular and cellular pathways that underlie the role of *for*, and other important signalling pathways, in the “social” brain.

Acknowledgements

We thank Jeffrey Dason, Ina Anreiter, and Chun-Fang Wu for their kind invitation to contribute to this special issue; Joël Meunier and Romain Libbrecht for the pictures of the ants; and Nicole Leitner for helpful comments on previous drafts of this manuscript.

Disclosure statement

The authors report no conflict of interests.

Funding

This work was supported by the European Union's Horizon 2020 research and innovation program under the grant agreement 773324, and APR-IR 2017-00117111 from Région Centre-Val de Loire to CL; and grants R21NS089834 from the NIH, and 1754264 and 1707221 from the NSF to YB.

ORCID

Christophe Lucas [0000-0003-2839-7583](https://orcid.org/0000-0003-2839-7583)

Yehuda Ben-Shahar [0000-0002-2956-2926](https://orcid.org/0000-0002-2956-2926)

References

- Aarab, A. (1991). *Polyéthisme, régulation sociale et éthogénèse chez les deux sous-castes morphologiques de la fourmi Pheidole pallidula*, Université Paris-Nord].
- Alwash, N., Allen, A.M., Sokolowski, M.B., Levine, J. (2021) The *Drosophila melanogaster* foraging gene affects social networks. *Journal of Neurogenetics*.
- Ament, S. A., Wang, Y., & Robinson, G. E. (2010). Nutritional regulation of division of labor in honey bees: toward a systems biology perspective. *Wiley Interdiscip Rev Syst Biol Med*, 2(5), 566-576. <https://doi.org/10.1002/wsbm.73>
- Anreiter, I., Kramer, J. M., & Sokolowski, M. B. (2017). Epigenetic mechanisms modulate differences in *Drosophila* foraging behavior. *Proceedings of the National Academy of Sciences*, 114(47), 12518-12523.
- Anreiter, I., & Sokolowski, M. B. (2019). The foraging gene and its behavioral effects: Pleiotropy and plasticity. *Annual Review of Genetics*, 53.
- Armstrong, G. A., López-Guerrero, J. J., Dawson-Scully, K., Peña, F., & Robertson, R. M. (2010). Inhibition of protein kinase G activity protects neonatal mouse respiratory network from hyperthermic and hypoxic stress. *Brain Research*, 1311, 64-72.
- Bargmann, C. I. (1998). Neurobiology of the *Caenorhabditis elegans* genome. *Science*, 282(5396), 2028-2033. <http://www.ncbi.nlm.nih.gov/pubmed/9851919>
- Bellen, H. J., Tong, C., & Tsuda, H. (2010). 100 years of *Drosophila* research and its impact on vertebrate neuroscience: a history lesson for the future [10.1038/nrn2839]. *Nat Rev Neurosci*, 11(7), 514-522. <http://dx.doi.org/10.1038/nrn2839>
- Ben-Shahar, Y. (2005). The foraging gene, behavioral plasticity, and honeybee division of labor. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol*, 191(11), 987-994. <https://doi.org/10.1007/s00359-005-0025-1>
- Ben-Shahar, Y. (2014). A piggyBac route to transgenic honeybees. *Proc Natl Acad Sci U S A*, 111(24), 8708-8709. <https://doi.org/10.1073/pnas.1407876111>
- Ben-Shahar, Y., Dudek, N. L., & Robinson, G. E. (2004). Phenotypic deconstruction reveals involvement of manganese transporter malvolio in honey bee division of labor. *J Exp Biol*, 207(Pt 19), 3281-3288. <https://doi.org/10.1242/jeb.01151>
- Ben-Shahar, Y., Leung, H. T., Pak, W. L., Sokolowski, M. B., & Robinson, G. E. (2003). cGMP-dependent changes in phototaxis: a possible role for the foraging gene in honey bee division of labor. *J Exp Biol*, 206(Pt 14), 2507-2515. <http://www.ncbi.nlm.nih.gov/pubmed/12796464>

401 Ben-Shahar, Y., Robichon, A., Sokolowski, M. B., & Robinson, G. E. (2002). Influence of
 402 gene action across different time scales on behavior. *Science*, 296(5568), 741-744.
 403 <https://doi.org/10.1126/science.1069911>

404 Ben-Shahar, Y., & Robinson, G. E. (2001). Satiation differentially affects performance in a
 405 learning assay by nurse and forager honey bees. *J Comp Physiol A Neuroethol Sens*
 406 *Neural Behav Physiol*, 187(11), 891-899.
 407 <http://www.ncbi.nlm.nih.gov/pubmed/11866187>

408 Ben-Shahar, Y., Thompson, C. K., Hartz, S. M., Smith, B. H., & Robinson, G. E. (2000).
 409 Differences in performance on a reversal learning test and division of labor in honey bee
 410 colonies. *Animal Cognition*, 3(3), 119-125. <https://doi.org/10.1007/s100710000068>

411 Beshers, S. N., & Fewell, J. H. (2001). Models of division of labor in social insects. *Annual*
 412 *Review of Entomology*, 46, 413-440. <Go to ISI>://000167877300015

413 Boekhoff, I., Seifert, E., Goggerle, S., Lindemann, M., Kruger, B. W., & Breer, H. (1993).
 414 Pheromone-induced second-messenger signalling in insect antennae. *Insect. Biochem.*
 415 *Molec. Biol.*, 23, 757-762.

416 Bollen, E., Puzzo, D., Rutten, K., Privitera, L., De Vry, J., Vanmierlo, T., Kenis, G., Palmeri,
 417 A., D'Hooge, R., Balschun, D., Steinbusch, H. M., Blokland, A., & Prickaerts, J. (2014).
 418 Improved long-term memory via enhancing cGMP-PKG signaling requires cAMP-PKA
 419 signaling. *Neuropsychopharmacology*, 39(11), 2497-2505.
 420 <https://doi.org/10.1038/npp.2014.106>

421 Bonabeau, E., Theraulaz, G., & Deneubourg, J.-L. (1998). Fixed response thresholds and the
 422 regulation of division of labor in insect societies. *Bulletin of Mathematical Biology*,
 423 60(4), 753-807. <https://doi.org/10.1006/bulm.1998.0041>

424 Burkhardt, R. W. (2014). Tribute to Tinbergen: Putting Niko Tinbergen's 'Four Questions' in
 425 Historical Context. *Ethology*, 120(3), 215-223. <https://doi.org/10.1111/eth.12200>

426 Buttstedt, A., Ihling, C. H., Pietzsch, M., & Moritz, R. F. (2016). Royalactin is not a royal
 427 making of a queen. *Nature*, 537(7621), E10.

428 Cash, A. C., Whitfield, C. W., Ismail, N., & Robinson, G. E. (2005). Behavior and the limits
 429 of genomic plasticity: power and replicability in microarray analysis of honeybee
 430 brains. *Genes Brain Behav*, 4(4), 267-271. [https://doi.org/10.1111/j.1601-](https://doi.org/10.1111/j.1601-183X.2005.00131.x)
 431 [183X.2005.00131.x](https://doi.org/10.1111/j.1601-183X.2005.00131.x)

432 Chardonnet, F., Capdevielle-Dulac, C., Chouquet, B., Joly, N., Harry, M., Le Ru, B., Silvain,
 433 J.-F., & Kaiser, L. (2014). Food searching behaviour of a Lepidoptera pest species is

- modulated by the foraging gene polymorphism. *Journal of Experimental Biology*, 217(19), 3465-3473.
- Chen, Z., Traniello, I. M., Rana, S., Cash-Ahmed, A. C., Sankey, A. L., Yang, C., & Robinson, G. E. (2021). Neurodevelopmental and transcriptomic effects of CRISPR/Cas9-induced somatic orco mutation in honey bees. *J Neurogenet*, 1-13. <https://doi.org/10.1080/01677063.2021.1887173>
- Chirino, M. G., Gilbert, L. E., & Folgarait, P. J. (2012). Behavioral discrimination between monogyne and polygyne red fire ants (Hymenoptera: Formicidae) in their native range. *Annals of the Entomological Society of America*, 105(5), 740-745.
- Corona, M., Libbrecht, R., & Wheeler, D. E. (2016). Molecular mechanisms of phenotypic plasticity in social insects. *Current Opinion in Insect Science*, 13, 55-60. <https://doi.org/https://doi.org/10.1016/j.cois.2015.12.003>
- Couvillon, M. J., Jandt, J. M., Duong, N., & Dornhaus, A. (2010). Ontogeny of worker body size distribution in bumble bee (*Bombus impatiens*) colonies. *Ecol Entomol*, 35(4), 424-435. <https://doi.org/10.1111/j.1365-2311.2010.01198.x>
- Cronin, A. L., Molet, M., Doums, C., Monnin, T., & Peeters, C. (2013). Recurrent evolution of dependent colony foundation across eusocial insects. *Annual Review of Entomology*, 58, 37-55.
- Dason, J. S., Allen, A. M., Vasquez, O. E., & Sokolowski, M. B. (2019). Distinct functions of a cGMP-dependent protein kinase in nerve terminal growth and synaptic vesicle cycling. *Journal of Cell Science*, 132(7), 1-12.
- Dason, J. S., Cheung, A., Anreiter, I., Montemurri, V. A., Allen, A. M., & Sokolowski, M. B. (2020). *Drosophila melanogaster* foraging regulates a nociceptive-like escape behavior through a developmentally plastic sensory circuit. *Proceedings of the National Academy of Sciences*, 117(38), 23286-23291.
- de Belle, J. S., Hilliker, A. J., & Sokolowski, M. B. (1989). Genetic localization of foraging (for): a major gene for larval behavior in *Drosophila melanogaster*. *Genetics*, 123(1), 157-163. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=2509284
- de Belle, J. S., & Sokolowski, M. B. (1987). Heredity of rover/sitter: alternative foraging strategies of *Drosophila melanogaster* larvae. *Heredity*, 59, 73-83.
- Detrain, C., Deneubourg, J. L., & Pasteels, J. M. (1999). *Information processing in social insects*. Berlin: Birkhäuser Verlag.

468 Douglas, S. J., Dawson-Scully, K., & Sokolowski, M. B. (2005). The neurogenetics and
469 evolution of food-related behaviour. *Trends in Neurosciences*, 28(12), 644-652.

470 Duarte, A., Pen, I., Keller, L., & Weissing, F. J. (2012). Evolution of self-organized division
471 of labor in a response threshold model. *Behavioral Ecology and Sociobiology*, 66(6),
472 947-957. <https://doi.org/10.1007/s00265-012-1343-2>

473 Eddison, M., Belay, A. T., Sokolowski, M. B., & Heberlein, U. (2012). A genetic screen for
474 olfactory habituation mutations in *Drosophila*: analysis of novel foraging alleles and an
475 underlying neural circuit. *PLoS ONE*, 7(12), e51684.
476 <https://doi.org/10.1371/journal.pone.0051684>

477 Engel, J. E., Xie, X. J., Sokolowski, M. B., & Wu, C. F. (2000). A cGMP-dependent protein
478 kinase gene, foraging, modifies habituation-like response decrement of the giant fiber
479 escape circuit in *Drosophila*. *Learn Mem*, 7(5), 341-352.
480 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cita](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11040266)
481 [tion&list_uids=11040266](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11040266)

482 Fitzpatrick, M. J., Ben-Shahar, Y., Smid, H. M., Vet, L. E., Robinson, G. E., & Sokolowski,
483 M. B. (2005). Candidate genes for behavioural ecology. *Trends Ecol Evol*, 20(2), 96-
484 104. <https://doi.org/10.1016/j.tree.2004.11.017>

485 Foret, S., Kucharski, R., Pittelkow, Y., Lockett, G. A., & Maleszka, R. (2009). Epigenetic
486 regulation of the honey bee transcriptome: unravelling the nature of methylated genes.
487 *BMC Genomics*, 10, 472. <https://doi.org/10.1186/1471-2164-10-472> [pii]
488 10.1186/1471-2164-10-472

489 Francis, S. H., Busch, J. L., Corbin, J. D., & Sibley, D. (2010). cGMP-dependent protein
490 kinases and cGMP phosphodiesterases in nitric oxide and cGMP action. *Pharmacol*
491 *Rev*, 62(3), 525-563. <https://doi.org/10.1124/pr.110.002907>

492 Fujiwara, M., Sengupta, P., & McIntire, S. L. (2002). Regulation of body size and behavioral
493 state of *C. elegans* by sensory perception and the EGL-4 cGMP-dependent protein
494 kinase. *Neuron*, 36(6), 1091-1102. <Go to ISI>://000180011600012

495 Fussnecker, B. L., McKenzie, A. M., & Grozinger, C. M. (2011). cGMP modulates responses
496 to queen mandibular pheromone in worker honey bees. *J Comp Physiol A Neuroethol*
497 *Sens Neural Behav Physiol*, 197(9), 939-948. [https://doi.org/10.1007/s00359-011-0654-](https://doi.org/10.1007/s00359-011-0654-5)
498 [5](https://doi.org/10.1007/s00359-011-0654-5)

499 Gangadharan, V., Wang, X., & Luo, C. (2017). Cyclic GMP-dependent protein kinase-I
500 localized in nociceptors modulates nociceptive cortical neuronal activity and pain

hypersensitivity. *Mol Pain*, 13, 1744806917701743.
<https://doi.org/10.1177/1744806917701743>

George, E. A., Bröger, A.-K., Thamm, M., Brockmann, A., & Scheiner, R. (2020). Inter-individual variation in honey bee dance intensity correlates with expression of the foraging gene. *Genes, Brain and Behavior*, 19(1), e12592.
<https://doi.org/10.1111/gbb.12592>

Gordon, D. M. (2016). From division of labor to the collective behavior of social insects. *Behav Ecol Sociobiol*, 70, 1101-1108. <https://doi.org/10.1007/s00265-015-2045-3>

Greenberg, J. K., Xia, J., Zhou, X., Thatcher, S. R., Gu, X., Ament, S. A., Newman, T. C., Green, P. J., Zhang, W., Robinson, G. E., & Ben-Shahar, Y. (2012). Behavioral plasticity in honey bees is associated with differences in brain microRNA transcriptome. *Genes Brain Behav*, 11(6), 660-670. <https://doi.org/10.1111/j.1601-183X.2012.00782.x>

Haydak, M. H. (1970). Honey bee nutrition. *Annual Review of Entomology*, 15(1), 143-156.

Heylen, K., Gobin, B., Billen, J., Hu, T. T., Arckens, L., & Huybrechts, R. (2008). Amfor expression in the honeybee brain: a trigger mechanism for nurse-forager transition. *J Insect Physiol*, 54(10-11), 1400-1403. [https://doi.org/S0022-1910\(08\)00154-6](https://doi.org/S0022-1910(08)00154-6) [pii] 10.1016/j.jinsphys.2008.07.015

Hölldobler, B., & Wilson, E. O. (2009). *The superorganism : the beauty, elegance, and strangeness of insect societies*. W.W. Norton & Company.

Ingram, K. K. (2002). Plasticity in queen number and social structure in the invasive Argentine ant (*Linepithema humile*). *Evolution*, 56(10), 2008-2016.

Ingram, K. K., Gordon, D. M., Friedman, D. A., Greene, M., Kahler, J., & Peteru, S. (2016). Context-dependent expression of the foraging gene in field colonies of ants: the interacting roles of age, environment and task. *Proceedings of the Royal Society B: Biological Sciences*, 283(1837), 20160841.

Ingram, K. K., Oefner, P., & Gordon, D. M. (2005). Task-specific expression of the *foraging* gene in harvester ants. *Molecular Ecology*, 14(3), 813-818. <Go to ISI>://000227132300013

Jaumann, M., Dettling, J., Gubelt, M., Zimmermann, U., Gerling, A., Paquet-Durand, F., Feil, S., Wolpert, S., Franz, C., Varakina, K., Xiong, H., Brandt, N., Kuhn, S., Geisler, H. S., Rohbock, K., Ruth, P., Schlossmann, J., Hutter, J., Sandner, P., Feil, R., Engel, J., Knipper, M., & Rüttiger, L. (2012). cGMP-Prkg1 signaling and Pde5 inhibition shelter cochlear hair cells and hearing function. *Nat Med*, 18(2), 252-259.
<https://doi.org/10.1038/nm.2634>

535 Kamakura, M. (2011). Royalactin induces queen differentiation in honeybees. *Nature*,
536 473(7348), 478-483.

537 Kapheim, K. M., Jones, B. M., Sovik, E., Stolle, E., Waterhouse, R. M., Bloch, G., & Ben-
538 Shahr, Y. (2020). Brain microRNAs among social and solitary bees. *R Soc Open Sci*,
539 7(7), 200517. <https://doi.org/10.1098/rsos.200517>

540 Kapheim, K. M., Pan, H., Li, C., Salzberg, S. L., Puiu, D., Magoc, T., Robertson, H. M.,
541 Hudson, M. E., Venkat, A., Fischman, B. J., Hernandez, A., Yandell, M., Ence, D.,
542 Holt, C., Yocum, G. D., Kemp, W. P., Bosch, J., Waterhouse, R. M., Zdobnov, E. M.,
543 Stolle, E., Kraus, F. B., Helbing, S., Moritz, R. F., Glastad, K. M., Hunt, B. G.,
544 Goodisman, M. A., Hauser, F., Grimmelikhuijzen, C. J., Pinheiro, D. G., Nunes, F. M.,
545 Soares, M. P., Tanaka, E. D., Simoes, Z. L., Hartfelder, K., Evans, J. D., Barribeau, S.
546 M., Johnson, R. M., Massey, J. H., Southey, B. R., Hasselmann, M., Hamacher, D.,
547 Biewer, M., Kent, C. F., Zayed, A., Blatti, C., 3rd, Sinha, S., Johnston, J. S., Hanrahan,
548 S. J., Kocher, S. D., Wang, J., Robinson, G. E., & Zhang, G. (2015). Social evolution.
549 Genomic signatures of evolutionary transitions from solitary to group living. *Science*,
550 348(6239), 1139-1143. <https://doi.org/10.1126/science.aaa4788>

551 Kaun, K. R., Chakabarty-Chatterjee, M., & Sokolowski, M. B. (2008). Natural variation in
552 plasticity of glucose homeostasis and food intake. *J Exp Biol*, 211(Pt 19), 3160-3166.
553 <https://doi.org/211/19/3160> [pii]
554 10.1242/jeb.010124

555 Kaun, K. R., Hendel, T., Gerber, B., & Sokolowski, M. B. (2007). Natural variation in
556 *Drosophila* larval reward learning and memory due to a cGMP-dependent protein
557 kinase. *Learn Mem*, 14(5), 342-349. <https://doi.org/14/5/342> [pii]
558 10.1101/lm.505807

559 Kaun, K. R., & Sokolowski, M. B. (2009). cGMP-dependent protein kinase: linking foraging
560 to energy homeostasis [Research Support, N.I.H., Extramural
561 Review]. *Genome*, 52(1), 1-7. <https://doi.org/10.1139/G08-090>

562 Kent, C. F., Daskalchuk, T., Cook, L., Sokolowski, M. B., & Greenspan, R. J. (2009). The
563 *Drosophila foraging* gene mediates adult plasticity and gene-environment interactions
564 in behaviour, metabolites and gene expression in response to food deprivation. *PLoS*
565 *Genet.*, 5(8), e1000609. <https://doi.org/10.1371/journal.pgen.1000609>

566 Kodaira, Y., Ohtsuki, H., Yokoyama, J., & Kawata, M. (2009). Size-dependent foraging gene
567 expression and behavioral caste differentiation in *Bombus ignitus*. *BMC Res Notes*, 2,
568 184. <https://doi.org/10.1186/1756-0500-2-184>

569 Kohlmeier, P., Alleman, A. R., Libbrecht, R., Foitzik, S., & Feldmeyer, B. (2019). Gene
570 expression is more strongly associated with behavioural specialization than with age or
571 fertility in ant workers. *Molecular Ecology*, 28(3), 658-670.

572 Kohn, N. R., Reaume, C. J., Moreno, C., Burns, J. G., Sokolowski, M. B., & Mery, F. (2013).
573 Social environment influences performance in a cognitive task in natural variants of the
574 *foraging* gene. *PLoS ONE*, 8(12), e81272.

575 Kohno, H., Suenami, S., Takeuchi, H., Sasaki, T., & Kubo, T. (2016). Production of
576 Knockout Mutants by CRISPR/Cas9 in the European Honeybee, *Apis mellifera* L.
577 *Zoolog Sci*, 33(5), 505-512. <https://doi.org/10.2108/zs160043>

578 Kroner, C., Boekhoff, I., Lohmann, S. M., Genieser, H. G., & Breer, H. (1996). Regulation of
579 olfactory signalling via cGMP-dependant protein kinases. *Trends Biochem. Sci.*, 22(8),
580 307-312.

581 Kuntz, S., Poeck, B., & Strauss, R. (2017). Visual Working Memory Requires Permissive and
582 Instructive NO/cGMP Signaling at Presynapses in the Drosophila Central Brain. *Curr*
583 *Biol*, 27(5), 613-623. <https://doi.org/10.1016/j.cub.2016.12.056>

584 L'Etoile, N. D., Coburn, C. M., Eastham, J., Kistler, A., Gallegos, G., & Bargmann, C. I.
585 (2002). The cyclic GMP-dependent protein kinase EGL-4 regulates olfactory adaptation
586 in *C. elegans*. *Neuron*, 36(6), 1079-1089. <https://doi.org/S0896627302010668> [pii]

587 Lattorff, H. M., & Moritz, R. F. (2013). Genetic underpinnings of division of labor in the
588 honeybee (*Apis mellifera*). *Trends Genet*, 29(11), 641-648.
589 <https://doi.org/10.1016/j.tig.2013.08.002>

590 Lee, J. I., O'Halloran, D. M., Eastham-Anderson, J., Juang, B. T., Kaye, J. A., Scott Hamilton,
591 O., Lesch, B., Goga, A., & L'Etoile, N. D. (2010). Nuclear entry of a cGMP-dependent
592 kinase converts transient into long-lasting olfactory adaptation. *Proc Natl Acad Sci U S*
593 *A*, 107(13), 6016-6021. <https://doi.org/10.1073/pnas.1000866107>

594 Lehner, B. (2013). Genotype to phenotype: lessons from model organisms for human
595 genetics. *Nat Rev Genet*, 14(3), 168-178. <https://doi.org/10.1038/nrg3404>

596 Leitner, N., & Dornhaus, A. (2019). Dynamic task allocation: how and why do social insect
597 workers take on new tasks? *Animal Behaviour*, 158, 47-63.
598 <https://doi.org/https://doi.org/10.1016/j.anbehav.2019.09.021>

599 Levy, S., & Bargmann, C. I. (2020). An Adaptive-Threshold Mechanism for Odor Sensation
600 and Animal Navigation. *Neuron*, 105(3), 534-548 e513.
601 <https://doi.org/10.1016/j.neuron.2019.10.034>

- Libbrecht, R., & Keller, L. (2013). Genetic compatibility affects division of labor in the Argentine ant *Linepithema humile*. *Evolution: International Journal of Organic Evolution*, 67(2), 517-524.
- Libbrecht, R., Oxley, P. R., Kronauer, D. J., & Keller, L. (2013). Ant genomics sheds light on the molecular regulation of social organization. *Genome Biol*, 14(7), 212.
<https://doi.org/10.1186/gb-2013-14-7-212>
- Lin, M., Nairn, A. C., & Guggino, S. E. (1992). cGMP-dependant protein kinase regulation of a chloride channel in T84 cells. *Am. J. Physiol.*, 262(5 Pt 1), 1304-1312.
- Lohmann, S. M., Vaandrager, A. B., Smolenski, A., Walter, U., & De Jonge, H. R. (1997). Distinct and specific functions of cGMP-dependent protein kinases. *Trends Biochem Sci*, 22(8), 307-312. [https://doi.org/10.1016/s0968-0004\(97\)01086-4](https://doi.org/10.1016/s0968-0004(97)01086-4)
- Lucas, C., Hughson, B. N., & Sokolowski, M. B. (2010). Job switching in ants: role of a kinase. *Communicative and Integrative Biology*, 3(1), 6-8.
<https://doi.org/10.4161/cib.3.1.9723>
- Lucas, C., Kornfein, R., Chakaborty-Chatterjee, M., Schonfeld, J., Geva, N., Sokolowski, M. B., & Ayali, A. (2010). The locust *foraging* gene. *Archives of Insect Biochemistry and Physiology*, 74(1), 52-66. <https://doi.org/10.1002/arch.20363>
- Lucas, C., Nicolas, M., & Keller, L. (2015). Expression of *foraging* and *Gp-9* are associated with social organization in the fire ant *Solenopsis invicta*. *Insect Molecular Biology*, 24(1), 93-104.
- Lucas, C., & Sokolowski, M. B. (2009). Molecular basis for changes in behavioral state in ant social behaviors. *Proceedings of the National Academy of Sciences of the United States of America*, 106(15), 6351-6356. <https://doi.org/doi:10.1073/pnas.0809463106>
- Ma, W., Jiang, Y., Meng, J., Zhao, H., Song, H., & Shen, J. (2018). Expression Characterization and Localization of the *foraging* Gene in the Chinese Bee, *Apis cerana cerana* (Hymenoptera: Apidae). *J Insect Sci*, 18(2): 1-5.
<https://doi.org/10.1093/jisesa/iey034>
- Malé, P.-J. G., Turner, K. M., Doha, M., Anreiter, I., Allen, A. M., Sokolowski, M. B., & Frederickson, M. E. (2017). An ant–plant mutualism through the lens of cGMP-dependent kinase genes. *Proceedings of the Royal Society B: Biological Sciences*, 284(1862), 20170896.
- Manfredini, F., Lucas, C., Nicolas, M., Keller, L., Shoemaker, D., & Grozinger, C. M. (2014). Molecular and social regulation of worker division of labour in fire ants. *Molecular Ecology*, 23(3), 660-672.

- McAfee, A., Pettis, J. S., Tarpy, D. R., & Foster, L. J. (2019). Feminizer and doublesex knock-outs cause honey bees to switch sexes. *PLoS Biol*, 17(5), e3000256.
<https://doi.org/10.1371/journal.pbio.3000256>
- Menzel, R., Lebouille, G., & Eisenhardt, D. (2006). Small brains, bright minds. *Cell*, 124(2), 237-239. <https://doi.org/Doi> 10.1016/J.Cell.2006.01.011
- Merchant, A., Song, D., Yang, X., Li, X., & Zhou, X. J. (2019). Candidate foraging gene orthologs in a lower termite, *Reticulitermes flavipes*. *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution*, 334(3), 168-177.
- Mery, F., Belay, A. T., So, A. K., Sokolowski, M. B., & Kawecki, T. J. (2007). Natural polymorphism affecting learning and memory in *Drosophila*. *Proc Natl Acad Sci U S A*, 104(32), 13051-13055. <https://doi.org/0702923104> [pii] 10.1073/pnas.0702923104
- Moon, C., Jaber, P., Otto-Bruc, A., Baehr, W., Palczewski, K., & Ronnet, G. V. (1998). Calcium-sensitive particulate guanylyl cyclase as a Modulator of cAMP in olfactory Receptor Neurons. *The J. of Neurosci.*, 18(9), 3195-3205.
- NCBI. Retrieved 9th April 2021 from [https://www.ncbi.nlm.nih.gov/protein/?term=cGMP-dependent+protein+kinase%5BAll+Fields%5D+AND+\(ant+%5BAll+Fields%5D+or+bce%5BAll+Fields%5D+or+termite%5BAll+Fields%5D+or+wasp%5BAll+Fields%5D+not+Nasonia+not+Amantichitinum+not+fungi+not+chordates+not+arachnids+not+blattella\)](https://www.ncbi.nlm.nih.gov/protein/?term=cGMP-dependent+protein+kinase%5BAll+Fields%5D+AND+(ant+%5BAll+Fields%5D+or+bce%5BAll+Fields%5D+or+termite%5BAll+Fields%5D+or+wasp%5BAll+Fields%5D+not+Nasonia+not+Amantichitinum+not+fungi+not+chordates+not+arachnids+not+blattella))
- Nygaard, S., Zhang, G., Schiøtt, M., Li, C., Wurm, Y., Hu, H., Zhou, J., Ji, L., Qiu, F., & Rasmussen, M. (2011). The genome of the leaf-cutting ant *Acromyrmex echinatior* suggests key adaptations to advanced social life and fungus farming. *Genome Research*, 21(8), 1339-1348.
- Oettler, J., Nachtigal, A.-L., & Schrader, L. (2015). Expression of the foraging gene is associated with age polyethism, not task preference, in the ant *Cardiocondyla obscurior*. *PLoS ONE*, 10(12), e0144699.
- Osborne, K., Robichon, A., Burgess, E., Butland, S., Shaw, R., Coulthard, A., Pereira, H., Greenspan, R., & Sokolowski, M. (1997). Natural behavior polymorphism due to a cGMP-dependent protein kinase of *Drosophila*. *Science*, 277(5327), 834-836.
- Oster, G. F., & Wilson, E. O. (1978). *Caste and ecology in the social insects*. Princeton, New Jersey: Princeton University Press.
- Page, R. E., Jr., Gadau, J., & Beye, M. (2002). The emergence of hymenopteran genetics. *Genetics*, 160(2), 375-379. <https://www.ncbi.nlm.nih.gov/pubmed/11861545>

670 Passera, L. (1985). Soldier determination in ants of the genus *Pheidole*. In J. A. L. Watson, B.
671 M. Okot-Kotber, & C. Noirot (Eds.), *Caste differentiation in social insects* (pp. 331-
672 346). Oxford: Pergamon Press.

673 Patel, N., Haydak, M., & Gochnauer, T. (1960). Electrophoretic components of the proteins in
674 honeybee larval food. *Nature*, 186(4725), 633-634.

675 Percy, M., Aron, S., Doums, C., & Keller, L. (2004). Conditional use of sex and
676 parthenogenesis for worker and queen production in ants. *Science*, 306(5702), 1780-
677 1783.

678 Raizen, D. M., Zimmerman, J. E., Maycock, M. H., Ta, U. D., You, Y.-j., Sundaram, M. V.,
679 & Pack, A. I. (2008). Lethargus is a *Caenorhabditis elegans* sleep-like state. *Nature*,
680 451(7178), 569-572.

681 Robertson, R. M., & Sillar, K. T. (2009). The nitric oxide/cGMP pathway tunes the
682 thermosensitivity of swimming motor patterns in *Xenopus laevis* tadpoles. *Journal of*
683 *Neuroscience*, 29(44), 13945-13951.

684 Robinson, G. E., Grozinger, C. M., & Whitfield, C. W. (2005). Sociogenomics: social life in
685 molecular terms. *Nat Rev Genet*, 6(4), 257-270. <https://doi.org/10.1038/nrg1575>

686 Rogers, S. M., Matheson, T., Despland, E., Dodgson, T., Burrows, M., & Simpson, S. J.
687 (2003). Mechanosensory-induced behavioural gregarization in the desert locust
688 *Schistocerca gregaria*. *Journal of Experimental Biology*, 206(Pt 22), 3991-4002.
689 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cita](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14555739)
690 [tion&list_uids=14555739](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14555739)

691 Ross, K. G., & Keller, L. (1995). Joint influence of gene flow and selection on a
692 reproductively important genetic polymorphism in the fire ant *Solenopsis invicta*. *The*
693 *American Naturalist*, 146, 325-348.
694 http://my.unil.ch/serval/document/BIB_6181B2F1C88C.pdf or available through:
695 Universite de Lausanne website http://www.unil.ch/dee/page6998_en.html

696 Roussel, E., Carcaud, J., Sandoz, J.-C., & Giurfa, M. (2009). Reappraising Social Insect
697 Behavior through Aversive Responsiveness and Learning. *PLoS ONE*, 4(1), e4197.
698 <https://doi.org/10.1371/journal.pone.0004197>

699 Rueffler, C., Hermisson, J., & Wagner, G. P. (2012). Evolution of functional specialization
700 and division of labor. *Proc Natl Acad Sci U S A*, 109(6), E326-335.
701 <https://doi.org/10.1073/pnas.1110521109>

- Ruppell, O., Pankiw, T., & Page, R. E., Jr. (2004). Pleiotropy, epistasis and new QTL: the genetic architecture of honey bee foraging behavior. *J Hered*, 95(6), 481-491.
<https://doi.org/10.1093/jhered/esh072>
- Sadd, B. M., Barribeau, S. M., Bloch, G., de Graaf, D. C., Dearden, P., Elsik, C. G., Gadau, J., Grimmelikhuijzen, C. J., Hasselmann, M., Lozier, J. D., Robertson, H. M., Smagghe, G., Stolle, E., Van Vaerenbergh, M., Waterhouse, R. M., Bornberg-Bauer, E., Klasberg, S., Bennett, A. K., Camara, F., Guigo, R., Hoff, K., Mariotti, M., Munoz-Torres, M., Murphy, T., Santesmasses, D., Amdam, G. V., Beckers, M., Beye, M., Biewer, M., Bitondi, M. M., Blaxter, M. L., Bourke, A. F., Brown, M. J., Buechel, S. D., Cameron, R., Cappelle, K., Carolan, J. C., Christiaens, O., Ciborowski, K. L., Clarke, D. F., Colgan, T. J., Collins, D. H., Cridge, A. G., Dalmay, T., Dreier, S., du Plessis, L., Duncan, E., Erler, S., Evans, J., Falcon, T., Flores, K., Freitas, F. C., Fuchikawa, T., Gempe, T., Hartfelder, K., Hauser, F., Helbing, S., Humann, F. C., Irvine, F., Jermin, L. S., Johnson, C. E., Johnson, R. M., Jones, A. K., Kadowaki, T., Kidner, J. H., Koch, V., Kohler, A., Kraus, F. B., Lattorff, H. M., Leask, M., Lockett, G. A., Mallon, E. B., Antonio, D. S., Marxer, M., Meeus, I., Moritz, R. F., Nair, A., Napflin, K., Nissen, I., Niu, J., Nunes, F. M., Oakeshott, J. G., Osborne, A., Otte, M., Pinheiro, D. G., Rossie, N., Rueppell, O., Santos, C. G., Schmid-Hempel, R., Schmitt, B. D., Schulte, C., Simoes, Z. L., Soares, M. P., Swevers, L., Winnebeck, E. C., Wolschin, F., Yu, N., Zdobnov, E. M., Aqrabi, P. K., Blankenburg, K. P., Coyle, M., Francisco, L., Hernandez, A. G., Holder, M., Hudson, M. E., Jackson, L., Jayaseelan, J., Joshi, V., Kovar, C., Lee, S. L., Mata, R., Mathew, T., Newsham, I. F., Ngo, R., Okwuonu, G., Pham, C., Pu, L. L., Saada, N., Santibanez, J., Simmons, D., Thornton, R., Venkat, A., Walden, K. K., Wu, Y. Q., Debyser, G., Devreese, B., Asher, C., Blommaert, J., Chipman, A. D., Chittka, L., Fouks, B., Liu, J., O'Neill, M. P., Sumner, S., Puiu, D., Qu, J., Salzberg, S. L., Scherer, S. E., Muzny, D. M., Richards, S., Robinson, G. E., Gibbs, R. A., Schmid-Hempel, P., & Worley, K. C. (2015). The genomes of two key bumblebee species with primitive eusocial organization. *Genome Biol*, 16, 76.
<https://doi.org/10.1186/s13059-015-0623-3>
- Scheiner, R., Sokolowski, M. B., & Erber, J. (2004). Activity of cGMP-dependent protein kinase (PKG) affects sucrose responsiveness and habituation in *Drosophila melanogaster*. *Learn Mem*, 11(3), 303-311. <https://doi.org/10.1101/lm.71604>

735 Schulte, C., Theilenberg, E., Muller-Borg, M., Gempe, T., & Beye, M. (2014). Highly
 736 efficient integration and expression of piggyBac-derived cassettes in the honeybee (*Apis*
 737 *mellifera*). *Proc Natl Acad Sci U S A*, *111*(24), 9003-9008.
 738 <https://doi.org/10.1073/pnas.1402341111>

739 Schwander, T., & Keller, L. (2008). Genetic compatibility affects queen and worker caste
 740 determination. *Science*, *322*(5901), 552-552.

741 Schwander, T., Lo, N., Beekman, M., Oldroyd, B. P., & Keller, L. (2010). Nature versus
 742 nurture in social insect caste differentiation. *Trends in Ecology & Evolution*, *25*(5), 275-
 743 282.

744 Shapira, M., Thompson, C. K., Soreq, H., & Robinson, G. E. (2001). Changes in neuronal
 745 acetylcholinesterase gene expression and division of labor in honey bee colonies. *J Mol*
 746 *Neurosci*, *17*(1), 1-12.
 747 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cita](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11665858)
 748 [tion&list_uids=11665858](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11665858)

749 Shpigler, H., Tamarkin, M., Gruber, Y., Poleg, M., Siegel, A. J., & Bloch, G. (2013). Social
 750 influences on body size and developmental time in the bumblebee *Bombus terrestris*.
 751 *Behavioral Ecology and Sociobiology*, *67*(10), 1601-1612.
 752 <https://doi.org/10.1007/s00265-013-1571-0>

753 Simola, D. F., Graham, R. J., Brady, C. M., Enzmann, B. L., Desplan, C., Ray, A., Zwiebel,
 754 L. J., Bonasio, R., Reinberg, D., Liebig, J., & Berger, S. L. (2016). Epigenetic
 755 (re)programming of caste-specific behavior in the ant *Camponotus floridanus*. *Science*,
 756 *351*(6268), aac6633. <https://doi.org/10.1126/science.aac6633>

757 Sinha, S., Ling, X., Whitfield, C. W., Zhai, C., & Robinson, G. E. (2006). Genome scan for
 758 cis-regulatory DNA motifs associated with social behavior in honey bees. *Proceedings*
 759 *of the National Academy of Sciences*, *103*(44), 16352-16357.
 760 <https://doi.org/10.1073/pnas.0607448103>

761 Smith, C. R., Toth, A. L., Suarez, A. V., & Robinson, G. E. (2008). Genetic and genomic
 762 analyses of the division of labour in insect societies. *Nature Reviews Genetics*, *9*(10),
 763 735-748. <https://doi.org/10.1038/Nrg2429>

764 Sokolowski, M. B. (1980). Foraging strategies of *Drosophila melanogaster*: a chromosomal
 765 analysis. *Behavior Genetics*, *10*(3), 291-302.

766 Stewart, A. M., Braubach, O., Spitsbergen, J., Gerlai, R., & Kalueff, A. V. (2014). Zebrafish
 767 models for translational neuroscience research: from tank to bedside. *Trends in*

768 *Neurosciences*, 37(5), 264-278.

769 <https://doi.org/http://dx.doi.org/10.1016/j.tins.2014.02.011>

770 Struk, A. A., Mugon, J., Huston, A., Scholer, A. A., Stadler, G., Higgins, E. T., Sokolowski,
771 M. B., & Danckert, J. (2019). Self-regulation and the *foraging* gene (PRKG1) in
772 humans. *Proceedings of the National Academy of Sciences of the United States of*
773 *America*, 116(10), 4434-4439.

774 Tarès, S., Arthaud, L., Amichot, M., & Robichon, A. (2013). Environment exploration and
775 colonization behavior of the pea aphid associated with the expression of the *foraging*
776 gene. *PLoS ONE*, 8(5), e65104.

777 Thamm, M., & Scheiner, R. (2014). PKG in honey bees: spatial expression, Amfor gene
778 expression, sucrose responsiveness, and division of labor. *J Comp Neurol*, 522(8), 1786-
779 1799. <https://doi.org/10.1002/cne.23500>

780 Tinbergen, N. (1963). On aims and methods of Ethology. *Zeitschrift für Tierpsychologie*,
781 20(4), 410-433. <https://doi.org/10.1111/j.1439-0310.1963.tb01161.x>

782 Tobback, J., Mommaerts, V., Vandersmissen, H. P., Smagghe, G., & Huybrechts, R. (2011).
783 Age- and task-dependent foraging gene expression in the bumblebee *Bombus terrestris*.
784 *Archives of Insect Biochemistry and Physiology*, 76(1), 30-42.
785 <https://doi.org/10.1002/arch.20401>

786 Tobback, J., Verlinden, H., Vuerinckx, K., Vleugels, R., Vanden Broeck, J., & Huybrechts, R.
787 (2013). Developmental-and food-dependent foraging transcript levels in the desert
788 locust. *Insect Science*, 20(6), 679-688.

789 Toth, A. L., & Robinson, G. E. (2007). Evo-devo and the evolution of social behavior. *Trends*
790 *Genet*, 23(7), 334-341. [https://doi.org/S0168-9525\(07\)00155-2](https://doi.org/S0168-9525(07)00155-2) [pii]
791 10.1016/j.tig.2007.05.001

792 Toth, A. L., Varala, K., Henshaw, M. T., Rodriguez-Zas, S. L., Hudson, M. E., & Robinson,
793 G. E. (2010). Brain transcriptomic analysis in paper wasps identifies genes associated
794 with behaviour across social insect lineages. *Proceedings of the Royal Society B:*
795 *Biological Sciences*, 277(1691), 2139-2148.

796 Toth, A. L., Varala, K., Henshaw, M. T., Rodriguez-Zas, S. L., Hudson, M. E., & Robinson,
797 G. E. (2010). Brain transcriptomic analysis in paper wasps identifies genes associated
798 with behaviour across social insect lineages. *Proc Biol Sci*, 277(1691), 2139-2148.
799 <https://doi.org/rspb.2010.0090> [pii]
800 10.1098/rspb.2010.0090

Toth, A. L., Varala, K., Newman, T. C., Miguez, F. E., Hutchison, S. K., Willoughby, D. A., Simons, J. F., Egholm, M., Hunt, J. H., Hudson, M. E., & Robinson, G. E. (2007). Wasp gene expression supports an evolutionary link between maternal behavior and eusociality. *Science*, 318(5849), 441-444. <https://doi.org/10.1126/science.1146647>

Trible, W., Olivos-Cisneros, L., McKenzie, S. K., Saragosti, J., Chang, N. C., Matthews, B. J., Oxley, P. R., & Kronauer, D. J. C. (2017). orco Mutagenesis Causes Loss of Antennal Lobe Glomeruli and Impaired Social Behavior in Ants. *Cell*, 170(4), 727-735 e710. <https://doi.org/10.1016/j.cell.2017.07.001>

Tschinkel, W. R. (2006). *The fire ants*. Cambridge: Harvard University Press.

Tsvetkov, N., Madani, B., Krimus, L., MacDonald, S. E., & Zayed, A. (2019). A new protocol for measuring spatial learning and memory in the honey bee *Apis mellifera*: effects of behavioural state and cGMP. *Insectes Sociaux*, 66(1), 65-71. <https://doi.org/10.1007/s00040-018-0641-8>

Vanengelsdorp, D., & Meixner, M. D. (2010). A historical review of managed honey bee populations in Europe and the United States and the factors that may affect them. *J Invertebr Pathol*, 103 Suppl 1, S80-95. <https://doi.org/10.1016/j.jip.2009.06.011>

Walton, A., Sheehan, M. J., & Toth, A. L. (2020). Going wild for functional genomics: RNA interference as a tool to study gene-behavior associations in diverse species and ecological contexts. *Hormones and Behavior*, 124, 104774.

Wang, S., & Sokolowski, M. B. (2017). Aggressive behaviours, food deprivation and the *foraging* gene. *Royal Society Open Science*, 4(4), 170042.

Wang, X., & Robinson, P. J. (1997). Cyclic GMP-dependent protein kinase and cellular signaling in the nervous system. *Journal of Neurochemistry*, 68(2), 443-456.

Wang, Y., Jorda, M., Jones, P. L., Maleszka, R., Ling, X., Robertson, H. M., Mizzen, C. A., Peinado, M. A., & Robinson, G. E. (2006). Functional CpG methylation system in a social insect. *Science*, 314(5799), 645-647. <https://doi.org/314/5799/645> [pii] 10.1126/science.1135213

Wang, Z., Pan, Y., Li, W., Jiang, H., Chatzimanolis, L., Chang, J., Gong, Z., & Liu, L. (2008). Visual pattern memory requires foraging function in the central complex of *Drosophila*. *Learn Mem*, 15(3), 133-142. <https://doi.org/10.1101/lm.873008>

Weinstock, G. M., Robinson, G. E., Gibbs, R. A., Worley, K. C., Evans, J. D., Maleszka, R., Robertson, H. M., Weaver, D. B., Beye, M., Bork, P., Elsik, C. G., Hartfelder, K., Hunt, G. J., Zdobnov, E. M., Amdam, G. V., Bitondi, M. M. G., Collins, A. M., Cristino, A. S., Lattorff, H. M. G., Lobo, C. H., Moritz, R. F. A., Nunes, F. M. F., Page, R. E.,

835 Simoes, Z. L. P., Wheeler, D., Carninci, P., Fukuda, S., Hayashizaki, Y., Kai, C.,
836 Kawai, J., Sakazume, N., Sasaki, D., Tagami, M., Albert, S., Baggerman, G., Beggs, K.
837 T., Bloch, G., Cazzamali, G., Cohen, M., Drapeau, M. D., Eisenhardt, D., Emore, C.,
838 Ewing, M. A., Fahrbach, S. E., Foret, S., Grimmelikhuijzen, C. J. P., Hauser, F.,
839 Hummon, A. B., Huybrechts, J., Jones, A. K., Kadowaki, T., Kaplan, N., Kucharski, R.,
840 Leboulle, G., Linial, M., Littleton, J. T., Mercer, A. R., Richmond, T. A., Rodriguez-
841 Zas, S. L., Rubin, E. B., Sattelle, D. B., Schlipalius, D., Schoofs, L., Shemesh, Y.,
842 Sweedler, J. V., Velarde, R., Verleyen, P., Vierstraete, E., Williamson, M. R., Ament, S.
843 A., Brown, S. J., Corona, M., Dearden, P. K., Dunn, W. A., Elekonich, M. M., Fujiyuki,
844 T., Gattermeier, I., Gempe, T., Hasselmann, M., Kadowaki, T., Kage, E., Kamikouchi,
845 A., Kubo, T., Kucharski, R., Kunieda, T., Lorenzen, M. D., Milshina, N. V., Morioka,
846 M., Ohashi, K., Overbeek, R., Ross, C. A., Schioett, M., Shippy, T., Takeuchi, H., Toth,
847 A. L., Willis, J. H., Wilson, M. J., Gordon, K. H. J., Letunic, I., Hackett, K., Peterson,
848 J., Felsenfeld, A., Guyer, M., Solignac, M., Agarwala, R., Cornuet, J. M., Monnerot, M.,
849 Mougél, F., Reese, J. T., Vautrin, D., Gillespie, J. J., Cannone, J. J., Gutell, R. R.,
850 Johnston, J. S., Eisen, M. B., Iyer, V. N., Iyer, V., Kosarev, P., Mackey, A. J., Solovyev,
851 V., Souvorov, A., Aronstein, K. A., Bilikova, K., Chen, Y. P., Clark, A. G., Decanini, L.
852 I., Gelbart, W. M., Hetru, C., Hultmark, D., Imler, J. L., Jiang, H. B., Kanost, M.,
853 Kimura, K., Lazzaro, B. P., Lopez, D. L., Simuth, J., Thompson, G. J., Zou, Z., De
854 Jong, P., Sodergren, E., Csuros, M., Milosavljevic, A., Osoegawa, K., Richards, S., Shu,
855 C. L., Duret, L., Elhaik, E., Graur, D., Anzola, J. M., Campbell, K. S., Childs, K. L.,
856 Collinge, D., Crosby, M. A., Dickens, C. M., Grametes, L. S., Grozinger, C. M., Jones,
857 P. L., Jorda, M., Ling, X., Matthews, B. B., Miller, J., Mizzen, C., Peinado, M. A., Reid,
858 J. G., Russo, S. M., Schroeder, A. J., St Pierre, S. E., Wang, Y., Zhou, P. L., Jiang, H.
859 Y., Kitts, P., Ruef, B., Venkatraman, A., Zhang, L., Aquino-Perez, G., Whitfield, C. W.,
860 Behura, S. K., Berlocher, S. H., Sheppard, W. S., Smith, D. R., Suarez, A. V., Tsutsui,
861 N. D., Wei, X. H., Wheeler, D., Havlak, P., Li, B. S., Liu, Y., Sodergren, E., Jolivet, A.,
862 Lee, S., Nazareth, L. V., Pu, L. L., Thorn, R., Stolc, V., Newman, T., Samanta, M.,
863 Tongprasit, W. A., Claudianos, C., Berenbaum, M. R., Biswas, S., de Graaf, D. C.,
864 Feyereisen, R., Johnson, R. M., Oakeshott, J. G., Ranson, H., Schuler, M. A., Muzny,
865 D., Chacko, J., Davis, C., Dinh, H., Gill, R., Hernandez, J., Hines, S., Hume, J.,
866 Jackson, L., Kovar, C., Lewis, L., Miner, G., Morgan, M., Nguyen, N., Okwuonu, G.,
867 Paul, H., Santibanez, J., Savery, G., Svatek, A., Villasana, D., Wright, R., & Consort, H.

- 868 G. S. (2006). Insights into social insects from the genome of the honeybee *Apis*
869 *mellifera*. *Nature*, 443(7114), 931-949. <https://doi.org/Doi> 10.1038/Nature05260
- 870 Wheeler, D. E. (1991). The developmental basis of worker caste polymorphism in ants. *The*
871 *American Naturalist*, 138(5), 1218-1238.
- 872 Whitfield, C. W., Ben-Shahar, Y., Brillet, C., Leoncini, I., Crauser, D., Leconte, Y.,
873 Rodriguez-Zas, S., & Robinson, G. E. (2006). Genomic dissection of behavioral
874 maturation in the honey bee. *Proc Natl Acad Sci U S A*, 103(44), 16068-16075.
875 <https://doi.org/10.1073/pnas.0606909103>
- 876 Whitfield, C. W., Cziko, A. M., & Robinson, G. E. (2003). Gene expression profiles in the
877 brain predict behavior in individual honey bees. *Science*, 302(5643), 296-299.
878 [http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cita](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14551438)
879 [tion&list_uids=14551438](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14551438)
- 880 Wilson, E. O. (1987). Causes of ecological success: the case of the ants. *The Journal of*
881 *Animal Ecology*, 1-9.
- 882 Ziegelberger, G., van den Berg, M. J., Kaissling, K. E., Klumppt, S., & Schultz, J. E. (1990).
883 Cyclic GMP levels and guanylate cyclase activity in pheromone-sensitive antennae of
884 slimoths *Antherae polyphemus* and *Bombyx mori*. *J. Neurosci.*, 10, 217-225.
- 885
- 886

Figure 1. cGMP-dependent protein kinase is a key node in the gene network that regulates behavioral states associated with division of labour. (A) Division of labour in the honey bee *Apis mellifera* is regulated solely by age-dependent processes. Young bees start their lives as nurses and typically start foraging when they are three weeks old. (B) Division of labour in the ant *Pheidole pallidula*. Tasks are regulated by developmental caste determination of “majors” and “minors”, as well by age-dependent regulation of behaviours such as nursing and foraging. Division of labour is also associated with task-specific sensory thresholds (A-B); and changes in neuronal anatomy and physiology (C). (D) Current models stipulate that *for* is a conserved regulatory node in the gene networks that regulate behavioral states across diverse social insect species. Therefore, changes in PKG activity can lead to shifts in the activity of peripheral and central neuronal populations that drive task-specific behavioral repertoires.

A

Age-dependent
Division of labour

Nursing state:

1. Negative phototaxis
2. Low response threshold to brood pheromone
3. High response threshold to sugar

Foraging state:

1. Positive phototaxis
2. High response threshold to brood pheromone
3. Low response threshold to sugar
4. Attraction to floral odours

B

Developmental
caste-dependent
Division of labour

Soldier state:

1. Guarding
2. Low response threshold to intruders
3. High response threshold to food
4. Process large food items

Forager state:

1. Food search
2. High response threshold to intruders
3. Low response threshold to food
4. Recruitment to food sources/intruders

C

Brain anatomy
and physiology

D

PKG-associated
gene network