



A machine Learning pipeline to track dynamics of a population of nanoparticles during in situ Environmental Transmission Electron Microscopy in gases

Khuram Faraz, Thomas Grenier, Christophe Ducottet, Thierry Epicier

► To cite this version:

Khuram Faraz, Thomas Grenier, Christophe Ducottet, Thierry Epicier. A machine Learning pipeline to track dynamics of a population of nanoparticles during in situ Environmental Transmission Electron Microscopy in gases. M&M2021, Microscopy Society of America (MSA), Aug 2021, Pittsburgh (virtual), United States. hal-03271239

HAL Id: hal-03271239

<https://hal.science/hal-03271239>

Submitted on 25 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A machine Learning pipeline to track dynamics of a population of nanoparticles during in situ Environmental Transmission Electron Microscopy in gases

Khuram Faraz^{1,2}, Thomas Grenier³, Christophe Ducottet², Thierry Epicier^{1,4}

¹Univ Lyon, INSA-Lyon, Université Claude Bernard Lyon 1, CNRS, MATEIS, UMR 5510, 69621 Villeurbanne Cedex, France

²Université Lyon, UJM-Saint-Etienne, CNRS, Institut Optique Graduate School, Laboratoire Hubert Curien, UMR 5516, 42023 Saint-Etienne, France

³Univ Lyon, INSA-Lyon, Université Claude Bernard Lyon 1, CNRS, CREATIS, UMR 5220, INSERM U1206, 69621 Villeurbanne Cedex, France.

⁴Univ Lyon, Université Claude Bernard Lyon 1, CNRS, IRCELYON, UMR 5526, 69626 Villeurbanne, France

Tracking the dynamics of supported nanoparticles (NPs) during the conditioning and under reactive conditions of heterogeneous nanocatalysts is an essential key for the optimization of their activity and durability [1]. Whereas Transmission Electron Microscopy (TEM) has always been a privileged technique owing to its spatial resolution, the availability of Environmental techniques, either in a dedicated ETEM or in close cells increases the potential of TEM experiments since they provide a way to follow directly such an evolution under gas and at high temperature. Features that are of significant importance are typically the mobility of NPs (surface diffusion, anchorage effects on their support), their size evolution (e.g. growth by coalescence or Ostwald Ripening, disappearance by dissolution or sublimation).

Probably one of the main concerns is, as always, the poor representativity of TEM results in terms of statistical meaning. This issue is of stringent importance when video sequences are acquired during in situ experiments, since the production of large sets of data renders their analysis and interpretation very tedious and time consuming. This is however a reasonable strategy to obtain more statistics.

Whereas detailed treatments can confidently be performed manually when analyzing a few NPs from relatively narrow regions of interest (see for example [2-3]), new automation approaches are required to achieve a reasonable statistical relevance. This becomes to be more and more possible owing to the constant development of easy-to-use machine learning routines as already performed in this field by a few groups (e.g. [4-5]).

This contribution will present a complete pipeline dedicated to the tracking of a NP population evolving under in situ gas and temperature conditions, with the aim of enabling a thorough analysis of their evolution according to the previously mentioned interactions and features regarding mobility and size evolution.

It consists in several steps which (see figure 1) will be detailed and illustrated in the case of Scanning TEM imaging of a Pd-delta-Alumina catalytic system followed during in situ calcination under oxygen in a dedicated FEI-Titan ETEM microscope [6]:

- Proper registration of successive images (frames) from continuous sequences
- Robust detection of NPs using the well-known Convolutional Neuronal Network (CNN) U-net [7] assisted by a verification of the 'Treacy-Rice' analysis of scattered intensities [8]
- Dedicated training and fine tuning of the CNN using large quantities of realistic annotated simulated images
- Identification of trajectories using an energy criteria-based approach (referred to as Multiple Object Tracking method [9]) derived from the continuous energy minimization tracking developed by [10]
- Automatic analysis of 'fusion' events (i.e. NPs coalescence) [11].




Figure 1: general synoptic of the NP tracking analysis.

References

- [1] Y. Zhou et al., Nano Today, 20 (2018) 101-120.
- [2] T.W. Hansen et al., Accounts of Chemical Research, 46 8 (2013) 1720-1730.
- [3] T.E. Martin et al., ChemCatChem, 7 (2015) 3705-3711.
- [4] J.P. Horwath et al., npj Computational Materials 6 (2020) #108.
- [5] L. Yao et al., ACS Cent. Sci. 6 (2020) 1421-1430.
- [6] T. Epicier et al., Catalysis Today, 334, 15 (2019), 68-78.
- [7] O. Ronneberger et al., Lect. Notes Comput. Sci. 9351 (2015) 234-241.
- [8] M.M.J. Treacy, S.B. Rice, J. Microsc. 156 (1989) 211-234.
- [9] Y. Xu et al., IET Computer Vision 13 4 (2019) 355-368.
- [10] A. Milan et al., IEEE Trans Pattern Analysis & Machine Intelligence, 38 10 (2016) 2054-2068.
- [11] The authors thank the support of the French National Research Agency (ANR) through the cooperative project '3DCLEAN' n° 15-CE09-0009-01 and of the EUR SLEIGHT (<https://manutech-sleight.com/>). CLYM (www.clym.fr) is acknowledged for the access to the ETEM and IFPEN (Solaize, F) for providing samples.