Highlights

Early Famennian MS values differ on N-Gondwana and on S-Laurussia margins

Famennian MS values correlate with conodont biofacies

MS is a powerful tool for geodynamics

Graphical Abstract

**North-Gondwana - Laurussia dynamic paleogeography challenged by
magnetic susceptibility through the Famennian**

**Girard, Catherine^{1*}; Feist, Raimund¹; Mossoni, Angelo²; Cornée, Jean-Jacques³;
Camps, Pierre³; Charruault, Anne-Lise¹; Corradini, Carlo⁴**

¹*Institut des Sciences de l'Evolution de Montpellier (ISEM), Univ Montpellier, CNRS, EPHE,
IRD, Montpellier, France.*

²*2 Kingsdown court, 56 Bolingbroke grove, London SW11 6HS, UK*

³*Géosciences Montpellier, Université de Montpellier, CNRS, and Université des Antilles,
Montpellier and Pointe à Pitre, (FWI), France*

⁴*Università degli Studi di Trieste, Dipartimento di Matematica e Geoscienze, Trieste, Italy*

*corresponding author (catherine.girard@umontpellier.fr)

Abstract

Large-scale magnetic susceptibility (MS) variations in ancient sediments are usually interpreted as related to sea-level and climate changes affecting the erosional regime and the amount of detrital input. To constrain such environmental changes during the Famennian, we compared MS records in representative sections of pelagic limestones of (1) the Avalonian margin of Laurussia [Sessacker (SES) and Beringhauser Tunnel (BHT), Rhenish Slate Mountains, Germany] and (2) in Gondwana related terranes [Erfoud (ERF), Tafilalt, Morocco; Col des Tribes (CT), Montagne Noire, France; Buschteich (BU), Thuringia, Germany; Corona Mizziu (CM I, II), Sardinia and Pizzul West (PZW) in the Carnic Alps, Italy]. Very low MS values throughout the Famennian characterize sections from Avalonia. In contrast, MS values are high and oscillating in Gondwana during the early Famennian; they drop significantly all together in the middle Famennian *Palmatolepis marginifera marginifera* Zone to match the pattern of low MS values of the Avalonian margin thereafter. To evaluate the importance of partial diagenetic overprint, MS data were compared to hysteresis parameters and geochemical proxies in sections BU, CM I, II and CT. The degree of thermal diagenesis is estimated taking into account the Color Alteration Indices (CAI) of conodonts. Striking dissimilarities in MS records between Laurussia and Gondwana are in favor for an existing remnant oceanic barrier between these continents during early Famennian times, when the amount of detrital supplies was obviously different. Uniformity is reached after the drop of MS values in N-Gondwana, when oceanic barriers could have vanished during the early middle Famennian T-R fluctuations of the sea-level. All sites remained under a rather similar regime of detrital supply through middle to uppermost Famennian. Comparison of MS trends with trends of conodont biofacies in five sections (CT, BU, CM I, II and SES) allows emphasizing the concordance of their relationship to sea level fluctuations.

43

44 Keywords: magnetic susceptibility, conodont biofacies, Famennian, N-Gondwana, Laurussia,
45 paleogeography

46

1. Introduction

Low-field magnetic susceptibility (MS) is currently considered a proxy to assess the amount of detrital inputs in relation to sea level changes, integrating mineralogical and sedimentological information on their nature and origin (e.g. Ellwood et al. 2006, 2007, 2008, Hladil 2002, Da Silva et al. 2009, 2010). As such it may constitute a powerful and independent proxy to control global trends of sea level changes on condition that the primary origin of low-field magnetic susceptibility is demonstrated. This can be done by comparing MS values with hysteresis parameters and/or elemental concentrations (Devleeschouwer et al. 2010); Riquier et al. 2010). Color Alteration Indices (CAI) of conodonts are taken into account to estimate the degree of thermal diagenesis of the rock samples (Epstein et al. 1977). Indeed, the color of the conodonts is related to the temperature that affected the rocks. With an average MS value for lithified marine deposits of $5.5 \times 10^{-8} \text{ m}^3/\text{kg}$ (Ellwood et al. 2011a) most of Paleozoic limestones range between 0.1 and $10 \times 10^{-8} \text{ m}^3/\text{kg}$ (Ellwood et al. 2006). This is the case in Devonian off-shore carbonate successions where a fair number of MS investigations (e.g. Crick et al. 2002, Da Silva et al. 2009, Ellwood et al. 2011b, Riquier et al. 2010) were conducted since the first global sea-level curve was established by Johnson et al. (1985). Comprehensive spatio-temporal investigations combining variations of litho- and biofacies with isotope data indicate a long term Famennian regressive trend (Johnson et al. 1985) that is punctuated by short transgressive episodes/events such as at the base of the Famennian (Sandberg et al. 2002). However, within-section MS evolution remains sometimes difficult to correlate with precision between sedimentary sequences from different paleogeographic entities (e.g., Crick et al. 2002). Attention is drawn to the possibility that MS signal in carbonate systems may be affected by diagenesis (Da Silva et al 2012, Devleeschouwer et al.

2015), or casually indicate rather particular local conditions (Davies et al. 2013) in spite of global eustasy.

Indeed, both the amount and nature of detrital material may possibly be disturbed by aeolian or volcanic inputs, heterogeneity of climatic influence in the continental source areas, and by the importance of physical action during gravitational transport (e.g., current-induced re-working, debris-massflows). On the contrary, slow sedimentation rates or absence of deposition leading to hardgrounds and hiatuses may result in different records of MS signals and their amplitudes. MS studies have been mainly focused on episodes of major extinction events such as the terminal Frasnian Kellwasser events that are particularly marked by instabilities of the sea-level (e.g., Averbuch et al. 2005, Riquier et al. 2010). According to these authors, the increase of detrital input related to the marked sea-level fall in the earliest Famennian post-event period is expressed by a significant increase in the MS signals. Noticeably, the values of this excursion is particularly less marked in sections on the Avalonian margin of Laurussia than in those of the Gondwana margin (Averbuch et al. 2005, fig. 4) (Fig. 1).

In this contribution we carry out investigations on the entire Famennian from representative sites of both N-Gondwanan and Avalonian terranes to allow the evaluation of the extent to which the environmental changes of approaching Laurussia/Gondwana margins may have influenced the MS signals (Fig. 1a). To this end, we conducted an integrated study of MS, hysteresis parameters and element concentrations, in some sections (CM I, CM II, PZW and partially in BU), in order to test whether MS can be considered as a reliable detrital proxy. Following Riquier et al. (2010), some elements characteristic of the terrigenous input are not affected by diagenesis (e.g., Zirconium) and can be used as a proxy of the bulk magnetic mineralogical component (Total Fe). Thereafter, we will compare MS measurements with analyses of conodont biofacies in sections that provide sufficient amounts of conodont

elements. Indeed, conodonts can also characterize palaeo-environments because of the different ecological preferences of the various genera (Sandberg 1976, Seddon and Sweet 1971, Girard et al. 2020), indicative of distance regarding the shoreline. According to this view, the changes in the relative proportion of the different genera, namely biofacies, have been proven useful to track environmental changes through time (e.g., Corradini 2003, Girard et al. 2014, 2017, 2020). As a prerequisite, the continuity of MS records is controlled by fine-scaled conodont biostratigraphy of Spalletta et al. (2017), updated in the latest Famennian by Corradini et al. (2021).

Whether trends in combined MS and biofacies curves allow constituting characteristic features of Famennian deposits in general, and to what extent they are permanent or are restricted to a definite duration, are prime subjects of our investigations. They allow pointing possible causes that we aim to discuss.

2. Famennian sections

The investigated sections, CT, BU, PZW, CM, ERF, BHT, SES (and DGHS for comparison) are situated between north Gondwana and Laurussia at tropical to subtropical paleolatitudes (Fig. 1a). The position of some samples, conodont zones and main lithofacies are shown in columnar sections Figure 1b.

2.1. North Gondwana sections

Col des Tribes (Montagne Noire, France).

This section (72 m-thick) was described by Girard et al. (2014). It is composed of (from bottom to top): - bioclastic limestones (17 m- thick) organized into dm- thick beds; - massive, red mudstones to wackestones/floatstones (15 m- thick) organized into dm- thick beds

yielding abundant cheiloceratid goniatites (Griottes facies); - massive micritic limestones (20 m- thick) organized into dm- thick beds; - pseudo-nodular limestones (15 m- thick) composed of cm- thick beds which have experienced pressure-dissolution processes; - massive micritic limestones (5 m- thick). These sediments were deposited in mid ramp to basin settings. Abundance of conodonts allows establishing a fine-scaled biozonation from the *Pa. rhenana* (late Frasnian) to the *Bi. ultimus* Zone (*Pr. meischneri* Subzone, latest Famennian). In this section conodonts are relatively well preserved, with a low Conodont Alteration Index (CAI) ~ 2-2.5 (temperature range = 60 to 110°C).

Buschteich (Thuringia, Germany).

This section (35 m- thick) was studied by Girard et al. (2017). From bottom to top, it is composed of: - cm bedded to pseudonodular wackstones (3 m- thick) (Frasnian); - dm to m- thick beds of mudstones to wackstones (7 m- thick); - mud-wackstones and pseudonodular mudstones (7 m- thick); mm to cm- bedded mud-wackstones (10 m- thick); - mm to cm bedded to pseudonodular wackstones, and some clay (8 m- thick). These sediments were deposited in outer ramp to basin settings. The section spans an interval from the *Pa. rhenana* to *Bi. ultimus* (*Pr. meischneri* Subzone) conodont Zones, but the lowest part of the Famennian is missing (from *Pa. subperlobata* to *Pa. minuta minuta* Zones). The CAI of conodonts is high, estimated as 4 (temperature range = 190 to 300°C).

Pizzul West (Carnic Alps, Italia).

This section (24 m- thick) was described by Mossoni et al. (2014), and Corradini et al. (2017). From bottom to top, it comprises: - massive to nodular grey mudstones to packstones (16.5 m- thick); - red nodular wackstones to packstones (7.5 m- thick) (Griottes facies). The limestones contain few fossils. The Griottes facies comprises nodules up to 1 cm of diameter

coated with hematite precipitations probably due to syngenetic diagenesis. The sediments were deposited in mid to outer ramp settings. The section spans an interval from the *Pa. glabra prima* to *Pa. marg. marginifera* Zones. The CAI of conodonts is around 4 (temperature range = 190 to 300°C).

Corona Mizziu I and Corona Mizziu II (Sardinia, Italy).

The Corona Mizziu sections have been studied by Corradini (1998, 2003).

The Corona Mizziu I (CM I) section (30 m- thick) comprises beds of poorly fossiliferous mudstones, with wackestone to packstone beds in the middle part of the succession.

Sediments were deposited in a pelagic setting. The conodont data allow discriminating twelve conodonts zones from the *Pa. rhomboidea* Zone up to the *Bi. ultimus* Zone.

The Corona Mizziu II (CM II) section (18 m-thick) crops out a hundred meters apart from the CM I section and exposes grey massive monotonous mudstones with poor fossil remains. The tectonic imprint is marked by stylolite structures and calcite recrystallization. Sediments were deposited in a pelagic setting. The section spans a time from the *Pa. crepida* to the *Pa. rugosa trachytera* Zones. For these two sections (CM I and CM II), the CAI of conodonts is very high, with values near 4.5 – 5 (temperature range = 300 to 400°C).

Erfoud (Tafilalt, SE Morocco).

The Erfoud section spanning the entire Late Devonian has been described by Buggisch & Clausen (1972). The considered lower to middle Famennian part of the section (8 m- thick) was re-sampled. It comprises from bottom to top: - 3 m-thick dark grey mud to wackestones and black shales, with dm-thick beds with cephalopods and bivalve remains (Kellwasser Limestone facies); - 0.45 m- thick platy red-brownish and grey-olive microsparitic calcilutites with bioclasts and Fe-oxid coated oncoids ; - 1.10 m-thick grey-brown cephalopod calcilutites

with Fe-oxid coated intraclasts; - 1.95 m-thick thin-bedded nodular calcilutites with cephalopods and trilobites ; -1.50 m- thick alternating cm- thick bedded beige argillaceous nodular limestones and marls. In the lower part of the section the *Pa. crepida* Zone is extremely condensed at the top of the dysoxic black Kellwasser-type limestone. The latter characterises a dysoxic bottom sea depositional environment. Above the Kellwasser limestones, the depositional conditions reflect low sedimentation rates in an open outer shelf environment within the photic zone as testified by prevailing large-eyed trilobites (deep inner to mid ramp). The section encompasses the interval of the *Pa. subperlobata* to the *Pa. rugosa trachytera* Zones with a gap between the *Pa. termini* to *Pa. gracilis gracilis* Zones (Fig. 1b). Conodonts have a CAI of 3-4 (temperature range = 150 to 300°C).

2.2. South Laurussia sections

Beringhauser Tunnel (Rhenish Slate Mountains).

This section (15 m- thick) was studied by Schülke & Popp (2005). It consists of mainly well-bedded massive to subordinately nodular cephalopod limestones. At the top of the section, facies change from massive limestones to nodular limestones. Sediments were deposited in mid to outer ramp setting. The considered section spans a time from the *Pa. crepida* to the *Pa. rugosa trachytera* Zones. The CAI of conodonts is high, about 4.5 (Joachimski et al. 2009), (temperature range around 300°C).

Sessacker (Rhenish Slate Mountains)

This highly condensed section (2.5 m- thick) was studied by Ziegler (1962) and Schülke (1999), and complemented by new data. It consists of red mudstones to wackestones with cephalopods, and some sparitic intercalations occur. The sediments were deposited in outer

ramp setting. The section spans from the *Pa. delicatula platys* to the *Pa. rugosa trachytera* Zones. The conodonts are well-preserved with a CAI around 2.5 (temperature range = 60 to 110°C).

Dupont GHS drillhole (Illinois basin, Eastern North America).

Drillcores in the Chattanooga Shale of Illinois revealed fine-grained siliciclastic sediments (9 m- thick) dating Famennian (Over et al. 2019). Conodont assemblages indicate that the section spans the *Pa. glabra prima* to *Bi. ac. aculeatus* Zones. MS values are low, in the - 0.25/-0.75 interval.

3. Methods

3.1. Magnetic Susceptibility

The samples studied are from previous collections for CT (Girard et al. 2014), BU (Girard et al. 2017), CMI and II (Mossoni 2014), PZW (Mossoni 2014). 129 new samples from sections SES, BHT and ERF have been analysed. We measured the low field magnetic susceptibility (χ_{LF}), abbreviates as magnetic susceptibility (MS) for the seven representatively distributed Famennian sections.

At Col des Tribes (CT, Montagne Noire, France) the field magnetic susceptibility (MS) was directly measured in situ with the Bartington MS2E sensor connected to the MS3 meter. To ensure a representative result, five measures were performed and then averaged for the 74 successive stratigraphic levels (Supplementary Data, Table S1). Ex-situ measurements were performed in the laboratory on block samples from German (63 samples for BU, 26 for SES, and 15 BHT) and Moroccan (25 samples for ERF) sections with the same device and the same protocol involving a minimum of 2 measurements per level (Supplementary Data, Tables S2,

S3, S4 and S5). Samples from the Italian sections (Pizzul West and Corona Mizziu I, II were measured with the KLY-3S Kappabridge at the University of Liège (Supplementary Data, Tables S6, S7, S8).

All the MS data figured represent an average of the measurements (Fig. 2) expressed in 10^{-8} m³/kg for readability. Following the age model published by Girard et al. (2020), the base of conodont zones was dated by means of absolute age estimates provided by Becker et al. (2012), and updated by Becker et al. (2020), and all MS data were figured based on the estimated absolute ages (Fig. 2).

3.2. Hysteresis parameters

The Italian sections (CM I, CM II and PZW) were prepared for hysteresis loops (Mossoni 2014) with the objective to roughly estimate the nature of the ferromagnetic (*sensu lato*) Fe-oxides. Indeed, as red nodular facies occur in Pizzul West but not in Corona Mizziu sections, we have suspected significant differences in the nature, the oxidation state, and the concentration quantity of strongly oxidized Fe-oxides such as hematite in the former section, and non-oxidized Fe-oxides such as magnetite in the later section. The hysteresis loops were measured with the J-Coercivity “rotation” magnetometer at Dourbes IRM Geophysical Centre, respectively (Supplementary Data, Tables S8, S9, S10).

For the hysteresis measurements, the specimens were prepared from blocks taken every 10 cm all along the studied cross-sections. We calculated from the hysteresis loops the high-field magnetic susceptibility (χ_{HF}) and the ferromagnetic susceptibility (χ_{ferro}) following methods developed in Riquier et al. (2010), and Da Silva et al. (2012, 2015). These two parameters allow assessing the respective contribution of the para/dia-magnetic minerals (χ_{HF}) and the ferromagnetic minerals to the initial susceptibility. At the end of the hysteresis loop

acquisition, the magnetic viscosity coefficient was calculated from the remanence decay, which was monitored for 100s after the field was removed. Additionally, the major elements (Al, Si, K and Ti) were considered to estimate the proportion of minerals in terms of their magnetic amounts for 17 samples in PZW, 13 samples in CM I and 14 samples in CM II (Tables PZW, CM I and CMII). These analyses on major elements were performed with an X-Ray Fluorescence (Panalytical MagiX PW2540) device at the University of Cagliari. The powder disks were prepared using at least 20 grams of the sample mixed with polyvinyl alcohol, on a base of boric acid. The total amount of oxides and LOI (loss on ignition) has been considered acceptable for samples with an error of $\pm 2\%$. In order to have comparable results between the studied sections, all analyses were normalized to 100. In this way a small error percentage has been distributed between all the measured parameters (Supplementary Data, Tables S8, S9, S10).

3.3. Detrital vs authigenic Fe content

To further constrain the diagenetic versus detrital imprints on MS, we compared the low-field magnetic susceptibility (χ_{HF}) variations with selected geochemical proxies measured by mean of the methods developed by Riquier et al. (2010). X-Ray Fluorescence analyses were performed on the samples of Buschteich which presents no signs of strong oxydation and is the most complete section of this study. The analyses were performed with the Niton XL3t-900 Gold X-ray fluorescence portable analyzer (Thermo Scientific®, Waltham, MA, USA) at the University of Montpellier. All the rock samples were carefully cleaned prior to all treatment, and weathered surfaces were removed.

We selected elements that can be used as proxies for the terrigenous input. The Zirconium (Zr) was selected here as the more reliable detrital proxy because it is not affected by diagenesis; and the total Fe (Fe_{tot}) content will be used as a proxy of the bulk magnetic

mineralogical component. The total Fe was separated into an inherited detrital part and a secondary authigenic part. Following Riquier et al. (2010), the detrital and authigenic fractions of the total iron content were estimated considering that the Fe/Al ratio of the detrital part of the studied rocks is supposed to be the same as that of the average shale value ($\text{Fe}/\text{Al}_{\text{average carbonate}} ; 0.55$ Clarkson et al. 2014). The detrital fraction of iron (Fe_{det}) is calculated as $\text{Fe}_{\text{det}} = \text{Al}_{\text{sample}} * \text{Fe}/\text{Al}_{\text{average carbonate}}$. Consequently, the iron fraction in excess relative to the detrital fraction, namely, $\text{Fe}_{\text{exc}} = \text{Fe}_{\text{tot}} - \text{Fe}_{\text{det}}$, may be considered to be of authigenic/diagenetic origin.

3.4. Biofacies

Five sections were sampled in detail for conodont biofacies based on conodont genera abundances. Conodont biofacies are advocated to be mostly driven by water depth variations due to the habitat preferences of the different conodont genera (e.g., Sandberg 1976, Seddon and Sweet 1971). In this regard, the distribution of conodont genera is thought to be controlled by water depth along a proximal-distal gradient (Klapper and Barrick 1978, Girard et al. 2020), and as such may constitute a proxy of sea-level variations and indirectly of detrital input. Several biofacies can be defined, based on the percentage of different genera present in the samples from biofacies characterized by distal surface dwellers (*Palmatolepis*) to proximal surface dwellers (*Icriodus*). The percentages of conodont elements per genus, later on called biofacies, were estimated for samples of each section: 74 samples for CT (Girard et al. 2014), 34 samples for BU (Girard et al. 2017), 12 samples for SES (Schülke 1995; this study), 13 samples in CM I and 13 samples for CM II (Corradini 2003, Mossoni 2014). All data are available in the Supplementary Data, Table S11. As MS is also known to be related to detrital input, we compared the MS trends of these five sections with trends in percentage of conodont genera (biofacies). A Principal Component Analysis (PCA on the

variance - covariance matrix) allows summarizing biofacies of the five considered sections (CT, BU, CM I and CM II, and SES) on few synthetic axes. This analysis also provides a representation of the records of CT, BU, CM I, CM II and SES on the same synthetic axes, allowing a direct comparison of the biofacies variations in the five outcrops. The contribution of each genus percentage to the axes allowed their interpretation in relation to the biofacies. Correlations between the MS and PC values are calculated using Pearson's product-moment correlation to investigate if MS and PC values are correlated or not.

4. Results

4.1. Magnetic susceptibility data

The relationship between the absolute age of the base of the biozone boundaries and sediment thickness allows establishing an age model for each section. The ages estimated for each sample were thereafter used to plot the variations of magnetic susceptibility through time. When we compare the evolution of magnetic susceptibility (MS) of Famennian sequences in the different sections, various types of record are documented (Fig. 2):

- In the investigated sections of the North Gondwana shelf (Morocco, Montagne Noire), MS values vary markedly and congruently during early Famennian times: initially high and fluctuating MS values shift all together to low ones within the *Pa. marginifera marginifera* Zone (initial mid-Famennian), and remain constantly low thereafter, lower than the common Paleozoic magnetic susceptibility values (around $5.5 \times 10^{-8} \text{ m}^3/\text{kg}$).
- In the most distal sections of the Gondwana shelf (Sardinia, Carnic Alps, Thuringia), the same trend is observed but fluctuations are less pronounced during the lower Famennian with temporary increases in MS values before the *Pa. marginifera marginifera* Zone.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

319 - In south Laurussia (DGHS section), the MS values remain low throughout the record, almost
320 always less than the average of $5.5 \times 10^{-8} \text{ m}^3/\text{kg}$, being rather close to the common Paleozoic
321 magnetic susceptibility values.

322 - In the Avalonian outer shelf areas of Laurussia (SES and BHT), the MS values are very low
323 during the early Famennian and are similar to those of both Laurentian (DGHS, Fig. 1) and
324 north Gondwanan terranes during mid-through late Famennian times.

325 In summary, in the north Gondwana margin a two-fold stage subdivision of the magnetic
326 susceptibility curve is observed. During the lower Famennian, increasing high values are first
327 recorded then followed by decreasing values that culminate during the *Pa. marginifera*
328 *marginifera* Zone. During the middle to upper Famennian, MS values remain lower than the
329 common Paleozoic magnetic susceptibility values (vertical dotted line in Figure 2). The major
330 shift from high to low MS values during the *Pa. marginifera marginifera* Zone in Gondwana
331 is not recorded in the Laurussia section where MS values remain low through the entire
332 Famennian.

334 4.2. Magnetic hysteresis data

335 For the Italian sections PZW, CM I and CM II, the correlations of oxides representing
336 terrigenous input with the MS data are shown in the Supplementary Data (Fig. S1), and
337 detailed in Table 1.

338 For the Pizzul West section, a significant correlation exists between the low field magnetic
339 susceptibility (χ_{LF}) and ferromagnetic contribution (χ_{ferro}), but not with the high field magnetic
340 susceptibility (χ_{HF}). A correlation between low field magnetic susceptibility (χ_{LF}) and the
341 geochemical parameters is not observed ($\text{Al}_2\text{O}_3=0.005$, $\text{SiO}_2=0.12$, $\text{TiO}_2=0.12$, $\text{K}_2\text{O}=0.04$)
342 (Table 1). However, there is a highly significant correlation between the high field magnetic

susceptibility (χ_{HF}) and geochemistry ($Al_2O_3=0.84$, $SiO_2=0.81$, $TiO_2=0.85$, $K_2O=0.83$), that demonstrates that the paramagnetic fraction is linked to Al, Si, K and Ti bearing minerals. As most of the Hcr values are lower than 60mT, the MS variations are probably related to low coercivity minerals as magnetite.

For the CM I section a strong correlation exists between the low field magnetic susceptibility (χ_{LF}) and the high field magnetic susceptibility (χ_{HF}), and the ferromagnetic contribution (χ_{ferro}). This demonstrates that the paramagnetic fraction is linked to Al, Si, K and Ti bearing minerals. A very good correlation also exists between the low field magnetic susceptibility and the four measured oxides ($Al_2O_3=0.68$, $TiO_2=0.70$, $K_2O=0.69$) (Table 1).

For the Corona Mizziu section II, the low field magnetic susceptibility has a good correlation with χ_{ferro} ($r = 0.85$) and with χ_{HF} ($r = 0.82$) (Table1). This suggests that the low field magnetic susceptibility of the Corona Mizziu II section is controlled by both the ferromagnetic and paramagnetic contributions. All the MS values are extremely low (lower than $3.0 \times 10^{-8} m^3/kg$, indicating the influence of the diamagnetic fraction, which may have diluted the potential detrital signal.

The magnetic signal for Pizzul West section would be mainly controlled by ferromagnetic contribution of low coercivity mineral (likely magnetite), whereas, for the Corona Mizziu I and II sections, the magnetic signal seems to be influenced by dia and/or paramagnetic contributions and, to a lesser extent, by ferromagnetic contribution.

4.3. MS correlation to Fe content

Both measured Fe_{tot} and calculated Fe_{det} correlate with the MS evolution ($r = 0.46$, $p = 0.002$ for Fe_{det} and $r = 0.41$; $p = 0.0005$ for total Fe). The Fe_{exc} does not show a correlation with the MS variations ($r = 0.19$, $p = 0.157$). A correlation exists between MS and the Zr content ($r=0.28$, $p = 0.037$) as well as between the MS and the Fe/Zr ratio ($r=0.73$, $p<0.001$).

(Supplementary Data, Fig. S2 and Supplementary Data, Table S2). Although this theoretical calculation gives an estimate of the Fe-bearing fractions only, Fe_{det} and Fe_{exc} , it supports the coexistence of a detrital component, controlling the MS evolution in Buschteich, and an authigenic component, that could have induced the widespread remagnetization process observed in these carbonates. This means that for the Buschteich section we cannot exclude that a part of the signal results from diagenetic processes, but this part is not as significant as the detrital part.

In summary, the geochemical data show that, although in part affected by diagenesis, as shown by correlation between MS and Fe/Zr, MS can be used as an indicator of the detrital input evolution in the studied Famennian sections. It is consequently possible to use it as a proxy of sea level variations which can be compared to the conodont biofacies proxy.

4.4. Comparison with conodont biofacies

The principal component analysis performed on the proportions of different genera allows summarizing the biofacies variations (Fig. 3) on main axes that express the total variance. The record of the five sections CT, BU, CM I, CM II (North Gondwana) and SES (South Laurussia) can be directly compared on these axes.

The first axis (PC1, 75.8% of the total variance) of the principal component analysis on conodont percentages shows a progressive trend to positive values (from 0 to 1) through the early Famennian. This trend reverses in the early mid – Famennian to negative values (Fig. 3b). The contribution of the variables (here, the percentages of the different genera) to the axes (Fig. 3b) shows that positive values along the first axis correspond to a high proportion of *Palmatolepis* whereas negative values correspond to a high proportion of *Bispathodus* (Supplementary Data, Table S11). A correlation ($N = 115$; $r = 0.26$; $p < 0.005$) exists between the PC1 and the MS data. The third axis represents only $< 4\%$ of variance, and mostly

corresponds to variations in the proportion of *Icriodus* in the early Famennian : peaks from *Pa. triangularis* to *Pa. crepida* Zones in CT, peak in the *Pa. crepida* Zone in BU, peak from the *Pa. glabra prima* to the *Pa. rhomboidea* Zones for CM, peak from *Pa. delicatula platys* to *Pa. minuta minuta* in SES (Supplementary Data, Table S1, and Fig. 3c), but values on the PC3 axis are not correlated with the MS values.

5. Discussion

Since pelagic carbonates are dominantly microsparites of diagenetic origin (Franke & Walliser 1983), one can only speculate on the ultimate source of the MS signals as these may be variously impacted by post-depositional processes. Following intervening aspects are discussed:

5.1. Impact of diagenesis

The magnetic susceptibility of a rock depends on both the mineralogical composition of the rock and the proportion of each mineral. The three main magnetic behaviours are: diamagnetic minerals (such as carbonates and quartz) displaying extremely weak negative MS values; paramagnetic minerals (e.g., clay minerals, particularly chlorite, smectite, illite and glauconite, ferromagnesian silicates, iron and manganese carbonates, pyrite) displaying weak positive values, and ferromagnetic minerals (mainly magnetite, pyrrhotite, maghemite, and hematite) displaying strong positive values (Da Silva et al. 2010). Diagenetic and post-diagenetic processes lead to mineralogical and chemical transformation of iron oxides that in turn influences the MS values. Riquier et al (2010) showed that although submitted to different burial histories and diagenetic conditions, consistent long-term trends can be observed in the MS evolution from sections through the Frasnian-Famennian boundary. Their data demonstrated that the MS evolution of the carbonate sections they studied can be

interpreted in terms of paleoenvironmental changes along the former margins of Laurussia and Gondwana.

In Pizzul West and Corona Mizziu, the conodont Color Alteration Index (CAI) ranges between 4.5 – 5.5. The temperature (200-400°C) which affected the rocks is far beyond the temperature of the goethite-hematite transition. Thus, the high coercivity values measured are related to the presence of hematite, visible also in the microfacies (Mossoni 2014). However, where high coercivity minerals control the MS (χ_{LF}), the presence of hematite is here interpreted to be of detrital origin because both the high field magnetic susceptibility (χ_{HF}) and the viscous decay do not correlate.

At Buschteich, the Fe in excess does not modify the general trend of MS, at least not enough to overprint the detrital signal.

In the CT section, CAI values of 2 – 2.5 were measured by Wiederer et al (2002), that corresponds to burial temperature of less than 100°C. Low illite crystallinity indicates that the site underwent anchizonal metamorphism at maximum. We assume therefore that a part of the MS signal, associated to iron-bearing primary clay minerals, has been preserved in the N-Gondwana sections despite different diagenetic stories.

Trends in MS are similar on the N-Gondwana margin (CAI estimated to 4-5 in PZW, CMI and II, BU and around 2-2.5 for CT), high values during the early Famennian and low values during mid-late Famennian. Similar patterns also occur in the S-Laurussia margin (CAI near 4 for BHT and only 2-2.5 for SES), but values remain low during the Famennian. This indicates that, despite different burial histories, the primary MS signature was not overprinted by the diagenetic signal, and it reflects in part its detrital origin.

5.2. Relations between depositional setting and MS

Variations in the amount of detrital ferromagnetic minerals in off-shore cephalopod limestones may be of different origins, and, in consequence, it is often difficult to precisely determine the cause of changing MS values and their amplitudes (e.g. Devleeschouwer et al. 2015). In this regard, the depositional conditions of the investigated sections could play a role on the accumulation of ferromagnetic minerals.

In the Rhenish Slate Mountains (SES, BHT), the Upper Devonian cephalopod limestones were deposited upon deep submarine rises beyond direct influence of sea-level changes (Franke & Walliser 1983). Besides *in situ* precipitation, pelagic carbonate mud originated from shallow-water areas. Fine-grained sand and clay fractions, originated from distal turbidites deposition, have also probably reached the rises but they mostly have been winnowed away by bottom currents. Consequently, rather low MS values might reflect presence low amount of transported ferro-magnetic elements, as coarse detrital influxes from northern shelf margins were concentrated in sandstone turbidite bodies that by-passed and were dispatched in troughs between the rises.

In the Montagne Noire (CT), very high MS values are recorded during the *Pa. marginifera marginifera* Zone prior to a sudden shift to low values under more stable environmental conditions in deeper mid-Famennian outer ramp setting. This shift is coincident with an abrupt change in facies: high though strongly oscillating values are found in the famous red-colored Griottes facies (Tucker 1974) and low values in the overlying compact light grey carbonate mudstones. The Griottes facies is a cephalopod-rich, nodular limestone with abundant iron-clay coatings and some ferruginous hard-grounds. The MS values in this facies are influenced by different chemical and/or physical constraints, not only the detrital inputs of ferro-magnetic minerals. Indeed, ferruginous mineral phases accumulations may also be influenced by changing both bottom sea oxygenation and rates of Fe-oxide –reduction related to buried organic matter and to diagenetic re-precipitation of Fe-oxide (Franke & Paul 1980).

This is the case, for example, of condensed pelagic carbonates across the Givetian/Frasnian boundary, where Devleeschouwer et al. (2015) drew attention to the fact that the ferromagnetic fraction may contain high coercivity minerals such as hematite and goethite of secondary origin that could affect the MS signal. This is probably also the case in the CT section, but no hysteresis analyses are available to confirm the assumption that the observed positive excursions may be solely related to hematization during diagenesis. Post-depositional transformation of paramagnetic clay minerals may also alter the detrital signal of MS. However, this is not the case because the burial temperature was less than 100° (CAI 2-2.5) in CT (Wiederer et al. 2002). It is noteworthy however, that, at CT, the Griotte limestones were deposited on an unstable slope under current activity and underwent repeated intraformational submarine sliding (debris flows, slumping...) (Girard et al. 2014), leading to local accumulation of Fe-rich sediments. These events may correspond to anomalously high positive peaks in the MS curve at the base of the *Pa. marginifera marginifera* Zone (Fig. 2).

At the basins scale, the scenario of MS positive excursions coinciding with hematite-rich early Famennian Griotte limestones has to be discussed. The Griottes limestones are commonly found in southern Europe including the Pyrenees, Montagne Noire and Carnic Alps. Nevertheless, they do not exist in contemporaneous strata in Sardinia despite a similar MS signal. Moreover, this facies is not present in the Thuringian Buschteich section where the MS variation pattern is the same. Conversely, at the beginning of the mid-Famennian, high MS values characterize also compact reddish and brown calcilutites with iron-oxide coatings on the shallow northern Tafilalt platform subjected to emergences (Buggisch & Clausen 1972, Wendt & Aigner 1985). It is noteworthy that a different scenario prevails in the Rhenish slate-mountains where condensed carbonate deposits on deep-water submarine rises upon volcanic substrates at Sessacker are red-coloured despite rather low MS values. At BHT similar low

values, after a marked positive excursion in the earliest Famennian (Riquier et al. 2010), occur in mud-mound carbonates which are not red-colored (Schülke & Popp 2005). In summary, the MS variations during the Famennian is independant from the presence of red facies (Griottes or others), and are consequently interpreted as primary. Consequently, the effect of post-depositional processes probably did not significantly overprint those of sea level changes during the Famennian to explain the differences between the records of Gondwana and Laurussia, especially the high MS values during the early Famennian in North Gondwana.

5.3. Influence of sea-level changes

Magnetic susceptibility and conodont biofacies are potentially interrelated: MS could be a proxy to evaluate detrital input if the signal is of primary origin, and conodont biofacies are interpreted as related to water depth due to habitat preferences of the different conodont genera along a proximal-distal gradient in the surface waters. Conodont PC1 displays a strong relationship with magnetic susceptibility in N-Gondwana terranes, and also on Laurussia, at least for Sessacker, and as such suggests constituting a real indicator of sea-level variations during the Famennian: low MS signals indicate a transgressive trend and correspond to distal biofacies, and, inversely, high MS signals indicate a regressive trend and correspond to proximal biofacies.

The shift towards lower values for MS and the trend to negative values on the PC1 in N-Gondwana terranes coincides with a local T-R (transgressive - regressive) couplet called “Enkeberg event” described in Laurussia (Rhenish Slate Mountains; House 1985). In Avalonia (Laurussia), there is no major significant change in MS values but the Enkeberg event was identified in the faunal record (Becker 1993). This event has not been yet identified

in the North Gondwana terranes. Consequently, the differences in the MS records of N-Gondwana and S-Laurussia rely on other factors, such as paleoclimate and paleogeography.

5.4. *Paleoclimate and paleo-latitude*

During the Famennian, SES, BHT, BU, and PZW sections were located in the subtropical zone of the southern hemisphere (Fig. 1). A long-term discrete cooling occurred during the Famennian, but no major changes in temperatures of the sea-surface waters was recorded (Joachimski et al. 2009); this may exclude a major role of physico-chemistry parameters of sea water conditions on MS values. Following the draw-down of atmospheric CO₂ concentrations and burial of organic carbon during the end-Frasnian Kellwasser crises, oxygen isotope values in conodont apatite indicate a global mean temperature drop of low latitude surface-water in post-event early Famennian times (Balter et al. 2008; Joachimski & Buggisch 2002; Joachimski et al. 2009). Mean sea-surface temperatures estimated from conodont apatite $\delta^{18}\text{O}$ values in the Rhenish Slate Mountains (Beringhauser Tunnel section) seem to be slightly higher than in Gondwana related terranes (Thuringia, Carnic Alps, Southern France) (Joachimski et al. 2009). Concerning climate, the main difference between Gondwana and Laurussia could rely on the latitudinal position between sites on the subtropical Laurussia and those of North Gondwana that are farther south in a more temperate position. Indeed, emerged parts of North Gondwana would have accentuated seasonality of climate and enhanced continental weathering (Percival et al. 2019; Herman et al. 2013). Therefore, emerged areas within the North Gondwana domain might have provided larger amounts of detrital material than the emerged parts of Laurussia. The decrease of high MS values on the Gondwana shelf began during the *Pa. rhomboidea* highstand of sea level and was probably achieved during the following transgressive pulse of the Enkeberg Event during the early mid-Famennian (Sandberg et al. 2002).

5.5. *Dynamic paleogeography*

Though still debated, Late Devonian plate tectonic models postulate convergence and collisional events between Gondwana and Laurussia heralding both the onset and the development of the Variscan orogeny. Subduction of oceanic crusts between Gondwana-derived continents led successively to the closure from S to N of several oceanic basins (including, Saxo-Thuringian, Rheno-Hercynian oceans...) that previously separated Armorican terranes (e.g., Thuringia, Bohemia) from the southern Laurussian margin (Franke et al. 2017, Golonka 2020) (Fig. 1). This resulted in the formation of large carbonate platforms on distal marine shelves on the Gondwana margins (parts of NW Africa to Montagne Noire). Opposite, on the southern margin Laurussia (Thuringia microcontinent), condensed pelagic limestones were restricted to tectonic or volcanic submarine highs.

During the early Famennian, only the narrowing oceanic interspaces of the Rheno-hercynian basin, assumed width was 1000 km during the Frasnian, separated Laurussia from outer-shelves of Gondwana-derived terranes (Franke et al. 2017). The age of the closure of the Rheic Ocean remains debated. After Franke (2019) it closed during the Early Devonian times. Other models maintained the Rheic ocean open between Saxo-Thuringia and Avalonia until the Carboniferous (Eckelmann et al. 2014; Golonka 2020, fig. 4). Nevertheless, all these models consider that the northern Gondwana shelf was much more extensive during the Famennian than those of South Laurussia which were restricted to narrow submarine highs and troughs. In such a setting, the detrital supplies from emerged source areas to carbonate platforms were naturally more homogeneously dispatched in the N-Gondwana shelf than in the Laurussia margin. Opposite, the accumulation of the detrital material in the Laurussia margin, even if locally important, was in fact located in troughs (e.g., Rhenish area) and low MS values are recorded on the rises. Consequently, the N-Gondwana and South Laurussia

margins clearly underwent a different paleogeographic behavior, and thus a different setting for detrital accumulation.

The separation between the two margins by the narrowing Rheno-Hercynian Ocean is effective at least until the beginning of the mid-Famennian, even if some local pelagic deposits can be found till the early Carboniferous (Franke et al. 2017). From the *Pa. marginifera marginifera* Zone onwards to the latest Famennian the effectiveness of the separation apparently vanished when climate became cooler (Girard et al. 2020). MS values considerably dropped and became conform to median values throughout the entire region between the Avalonian margins of Laurussia and N-Gondwana related terranes. Consequently, more uniform conditions and smaller amounts of detrital supply prevailed throughout. Concomitantly, the conjunction of vicinity of the two margins and similar climatic conditions is also corroborated by the increasing identity of stenotopic level-bottom biotas such as phacopid trilobites (Feist 2019) in both areas. Consequently, during this time interval, we consider that mid-late Famennian deposits with low MS values seems to be controlled by first an evolving paleogeographic setting, and, at a lesser extent, eustasy, current activity and climate. This paleogeographic setting results from converging margins prior to closure, and questions the existence, during mid-late Famennian, of wide oceanic domains.

Conclusions

The MS study of five sections in the N-Gondwana margin shows high and oscillating values during the early Famennian, and low values during the mid-late Famennian. In the S-Laurussia margin, MS values remain low through the whole Famennian. Geochemical investigations, and hysteresis parameters in three N-Gondwana sections indicate that the diagenetic processes did not significantly overprint the primary signal of MS. This is

emphasized by the degree of burial, thus the temperature during diagenesis (CAI of conodonts) and the presence or absence of red facies, as the MS values remain consistent from one section to other.

For the first time, we demonstrate a high correlation between these MS and conodont biofacies, confirming their validity for sea level reconstructions. Taking into account all the above mentioned studies, MS can now be used for paleogeographic reconstructions in the presently dismembered areas of the Variscan Belt.

During the early Famennian, the difference in the MS trends and the high values in N-Gondwana compared to S-Laurussia can be explained by two main paleogeographic constraints: i) the Rheic Ocean was not closed; ii) the North-Gondwana margin was wide whilst the South Laurussia margin was fragmented into rises and troughs. During the mid-late Famennian, the uniformization of MS toward low values is interpreted as indicative of rather similar conditions in both margins, linked to plate convergence and vanishing influence of oceanic spaces.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This contribution profited substantially from critical remarks and constructive suggestions provided by W. Franke and an anonymous reviewer for which we are most grateful. We acknowledge the assistance of Thomas Tirard, Lucas Tortarolo, and Lisa Zaharias for

measurements of Buschteich and Col des Tribes during their third-year university degree.

Authors thank Eberhart Schindler who provided samples from Sessacker, and Bernard Orth

for his assistance during the field-work in Morocco. A.M. thanks Anne Christine Da Silva and

Dr Simo Spassov for their help with the issues of the magnetic susceptibility and for allowing

him to work in their laboratories. This research was funded by the ANR project ECODEV

(ANR-13-BSV7-005). This is contribution ISEM 2021-111.

References

Averbuch, O., Tribovillard, N., Devleeschouwer, X., Riquier, L., Mistiaen, B., Van Vliet-Lanoe, B., 2005. Mountain building-enhanced continental weathering and organic carbon burial as major causes for climatic cooling at the Frasnian-Famennian boundary (c. 376 Ma)? *Terra Nova* 17, 25-34.

Balter, V., Renaud, S., Girard, C., Joachimski, M.M., 2008. Record of climate-driven morphological changes in 376 Ma Devonian fossils. *Geology* 36, 907-910.

Becker, R.T., 1993. Anoxia, eustatic changes, and upper Devonian to lowermost Carboniferous global ammonoid diversity, In: House, M.R. (Ed.), *The Ammonoidea: Environment, Ecology, and Evolutionary Change*. Systematics Assoc. Spec. Vol., Oxford, pp. 115-163.

Becker, R.T., Gradstein, F.M., Hammer, O., 2012. The Devonian Period, in: Gradstein, F.M., Ogg, J.G., Schmitz, M.D., Ogg, G.M. (Eds.), *The Geologic Time Scale 2012*. Elsevier, Oxford, pp. 559-602.

Becker, R.T., Marshall, J.E.A., Da Silva, A.-C., 2020. The Devonian Period. In *Geological Time Scale 2020*. doi: <https://doi.org/10.1016/B978-0-12-824360-2.00022-X>

Buggisch, W., Clausen, C.-D., 1972. Conodonten- und Goniatiten-Faunen aus dem oberen Frasnium und unteren Famennium Marokkos (Tafilalt, AntiAtlas). *Neues Jahrb. Geol. Palaontol. Abh.* 141, 137-167.

Clarkson, M.O., Poulton, S.W., Guilbaud, R., Wood, R.A., 2014. Assessing the utility of Fe/Al and Fe-speciation to record water column redox conditions in carbonate-rich sediments. *Chem. Geol.* 382, 111-122.

Corradini, C., 1998. Famennian conodonts from two sections near Villasalto. *Giorn. Geol. Spec. Issue*, 122-135.

Corradini, C., 2003. Late Devonian (Famennian) conodonts from the Corona Mizziu Sections near Villasalto (Sardinia, Italy). *Palaeontogr. Ital.* 89, 65-116.

Corradini C., Mossoni A., Corrigan M.G., Spalletta C., 2021. The Devonian/Carboniferous Boundary in Sardinia (Italy). *Paleobiodivers. Paleoenv.* 101. doi: 10.1007/s12549-019-00411-5

Corradini, C., Mossoni, A., Pondrelli, M., Simonetto, L., 2017. Famennian conodonts in the Mt. Pizzul West (PZW) section. *Ber. Inst. Erdwiss. K.F.-Univ. Graz*, 23, 246-250.

- Crick, R.E., Ellwood, B.B., Feist, R., El Hassani, A., Schindler, E., Dreesen, R., Over, D.J., Girard, C., 2002. Magnetostratigraphy susceptibility of the Frasnian/Famennian boundary. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 181, 67-90.
- Da Silva A.C., Mabillic C., Boulvain F., 2009. Influence of sedimentary setting on the use of magnetic susceptibility: examples from the Devonian of Belgium. *Sedimentology* 56, 1292-1306.
- Da Silva A.C., Yans J., Boulvain F., 2010. Sedimentology and magnetic susceptibility during the “punctata” event of the Ardenne area (Belgium): identification of severe and rapid sea level fluctuations. In: Da Silva, A.C. and Boulvain, F. (eds): magnetic susceptibility, correlations and Palaeozoic environments. *Geol. Belg.* 13/4, 319-332.
- Da Silva, A.-C., Whalen, M.T., Hladil, J., Chadimova, L., Chen, D., Spassov, S., Boulvain, F., Devleeschouwer, X., 2015. Magnetic susceptibility application: a window onto ancient environments and climatic variations: foreword. *Geol. Soc. Spec. Publ.*, London 414, 1-13.
- Da Silva, A.C., Dekkers, M.J., Mabillic, C., Boulvain, F., 2012. Magnetic signal and its relationship with paleoenvironments and diagenesis - examples from the Devonian carbonates of Belgium. *Studia Geophys. Geod.* 56, 677-704.
- Davies, E.J., Ratcliffe, K.T., Montgomery, P., Pomar, L., Ellwood, B.B., Wray, D.S., 2013. Magnetic Susceptibility (χ) Stratigraphy and Chemostratigraphy Applied to an Isolated Carbonate Platform Reef Complex; Lluçmajor Platform, Mallorca, In: Verwer, K., Playton, T.E., Harris, P.M.M. (Eds.), *Deposits, Architecture, and Controls*
- Devleeschouwer, X., Petitclerc, E., Spassov, S., Prat, A., 2010. The Givetian-Frasnian boundary at Nismes parastratotype (Belgium): the magnetic susceptibility signal controlled by ferromagnetic minerals. *Geol. Belg.*, 14/4, 345-360.
- Devleeschouwer, X., Riquier, L., Bbek, O., Devleeschouwer, D., Petitclerc, E., Sterckx, S., Spassov, S., 2015. Magnetization carriers of grey to red deep-water limestones in the GSSP of the Givetian-Frasnian boundary (Puech de la Suque, France): signals influenced by moderate diagenetic overprinting, In: Da Silva, A.C., Whalen, M.T., Hladil, J., Chadimova, L., Chen, D., Spassov, S., Boulvain, F., Devleeschouwer, X. (Eds.), *Magnetic Susceptibility Applications: A Window onto Ancient Environments and Climatic Variations*. *Geol. Soc. Spec. Publ.*, London 414 (1), 157-180.
- Eckelmann, K., Nesbor, H.-D., Knigshof, P., Linnemann, U., Hofmann, M., Lange, J.-M., Sagawe, A., 2014. Plate interactions of Laurussia and Gondwana during the formation of Pangaea — Constraints from U–Pb LA–SF–ICP–MS detrital zircon ages of Devonian and Early Carboniferous siliciclastics of the Rhenohercynian zone, Central European Variscides. *Gondwana Res.* 25, 1484-1500.
- Ellwood B.B., Garcia-Alcalde J.L., El Hassani A., Hladil J., Soto, F.M., Truyols-Massoni, M., Weddige, K., Koptikova, L., 2006. Stratigraphy of the Middle Devonian Boundary: Formal Definition of the Susceptibility Magnetostratotype in Germany with comparisons to Sections in the Czech Republic, Morocco and Spain. *Tectonophysics* 418, 31-49.
- Ellwood B.B., Tomkin J., Richards B., Benoist S.L., Lambert L.L., 2007. MSEC Data Sets Record Glacially Driven Cyclicity: Examples from the Arrow Canyon Mississippian-Pennsylvanian GSSP and Associated Sections. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 255, 377-390. doi: 10.1016/j.palaeo.2007.08.006.

- Ellwood B.B., Tomkin J.H., Ratcliffe K.T., Wright M., Kafafy A.M., 2008. High-resolution magnetic susceptibility and geochemistry for the Cenomanian/Turonian boundary GSSP with correlation to time equivalent core. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 261, 105-126.
- Ellwood, B.B., Algeo, T., El Hassani, A., Tomkin, J.H., Rowe, H., 2011a. Defining the Timing and Duration of the Kačák Interval within the Eifelian/Givetian Boundary GSSP, Mech Irdane, Morocco, Using Geochemical and Magnetic Susceptibility Patterns. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 304, 74–84.
- Ellwood, B.B., Balsam, W.L., Roberts, H.H., 2006. Gulf of Mexico Sediment Sources and Sediment Transport Trends from Magnetic Susceptibility Measurements of Surface Samples. *Mar. Geol.* 230, 237–248.
- Ellwood, B.B., Tomkin, J.H., El Hassani, A., Bultynck, P., Brett, C.E., Schindler, E., Feist, R., Bartholomew, A.J., 2011b. A climate-driven model and development of a floating point time scale for the entire Middle Devonian Givetian Stage: A test using magnetostratigraphy susceptibility as a climate proxy. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 304, 85-95.
- Epstein, A., Epstein, J., Harris, L. 1977. Conodont Color Alteration - an Index to Organic Metamorphism. *Geol. Surv. Prof. Pap.* 995, 1-27. [doi:10.3133/pp995](https://doi.org/10.3133/pp995).
- Feist, R., 2019. Post-Kellwasser event recovery and diversification of phacopid trilobites in the early Famennian (Late Devonian). *Bull. Geosci.* 94, 1-22.
- Franke, W. Walliser, O.H. 1983. “Pelagic” carbonates in the Variscan Belt – their sedimentary and tectonic environments, In: Martin H. and Eder F. W., *Intercontinental Fold Belts*. Springer, Berlin, pp. 77-92.
- Franke, W., Cocks, L.R.M., Torsvik, T.H., 2017. The Palaeozoic Variscan oceans revisited. *Gondwana Res.* 48, 257-284.
- Franke, W., Cocks, L.R.M., Torsvik, T.H., 2019. Detrital zircons and the interpretation of palaeogeography, with the Variscan Orogeny as an example. *Geol. Mag.* 157 (4), 1-5. doi.org/10.1017/S0016756819000943
- Franke, W., Paul, J. 1980. Pelagic redbeds in the Devonian of Germany – deposition and diagenesis. *Sediment. Geol.* 25, 231-256.
- Girard, C., Cornée, J.-J., Charruault, A.-L., Corradini, C., Weyer, D., Bartsch, K., Joachimski, M.M., Feist, R., 2017. Conodont biostratigraphy and palaeoenvironmental trends during the Famennian (Late Devonian) in the Thuringian Buschteich section (Germany). *Newsl. Stratigr.* 50, 71-89.
- Girard, C., Cornée, J.-J., Corradini, C., Fravallo, A., Feist, R., 2014. Palaeoenvironmental changes at Col des Tribes (Montagne Noire, France), a reference section for the Famennian of north Gondwana-related areas. *Geol. Mag.* 151, 864-884.
- Girard, C., Cornée, J.-J., Joachimski, M.M., Charruault, A.-L., Dufour, A.-B., Renaud, S., 2020. Paleogeographic differences in temperature, water depth and conodont biofacies during the Late Devonian. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 549, 108852. <https://doi.org/10.1016/j.palaeo.2018.06.046>
- Golonka, J., 2020. Late Devonian paleogeography in the framework of global tectonics. *Global Planet. Change* 186. <https://doi.org/10.1016/j.gloplacha.2020.103129>

- Herman, F., Seward, D., Valla, P.G., Carter, A., Kohn, B., Willett, S.D., Ehlers, T.A., 2013. Worldwide acceleration of mountain erosion under a cooling climate. *Nature* 504.
- Hladil, J., 2002. Geophysical records of dispersed weathering products on the Frasnian carbonate platform and early Famennian ramps in Moravia, Czech Republic: proxies for eustasy and palaeoclimate. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 181, 213-250.
- House, M.R., 1985. Correlation of mid-Palaeozoic ammonoid evolutionary events with global sedimentary perturbations. *Nature* 313, 17-22.
- Joachimski, M., Buggisch, W., 2002. Conodont apatite $\delta^{18}\text{O}$ signatures indicate climatic cooling as a trigger of the Late Devonian mass extinction. *Geology* 30, 711-714.
- Joachimski, M.M., Breisig, S., Buggisch, W., Talent, J.A., Mawson, R., Gereke, M., Morrow, J.R., Day, J., Weddige, K., 2009. Devonian climate and reef evolution: Insights from oxygen isotopes in apatite. *Earth Planet. Sci. Lett.* 284, 599-609.
- Johnson, J.G., Klapper, G., Elrick, M., 1996. Devonian transgressive-regressive cycles and biostratigraphy, Northern Antelope Range, Nevada: establishment of reference horizons for global cycles. *Palaaios* 11, 3-14.
- Johnson, J.G., Klapper, G., Sandberg, C.A., 1985. Devonian tatic fluctuations in Euramerica. *Geol. Soc. Am. Bull.* 96, 567-587.
- Klapper, G., Barrick, J.E., 1978. Conodont ecology: pelagic versus benthic. *Lethaia* 11, 15-23.
- Mossoni, A., 2014. Selected Famennian (Late Devonian) events (Condroz, Annulata, Hangenberg) in Sardinia and in the Carnic Alps: conodont biostratigraphy, magnetic susceptibility and geochemistry, Università degli Studi di Cagliari. Cagliari, p. 171 (unpublished thesis).
- Over, D.J., Hauf, E., Wallace, J., Chiarello, J., Over, J.-S., Gilleaudeau, G.J., Song, Y., Algeo, T.J., 2019. Conodont biostratigraphy and magnetic susceptibility of Upper Devonian Chattanooga Shale, eastern United States: Evidence for episodic deposition and disconformities. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 524, 137-149. <https://doi.org/10.1016/j.palaeo.2019.03.017>.
- Percival, L.M.E., Selby, D., Bond, D.P.G., Rakociński, M., Racki, G., Marynowski, L., Adatte, T., Spangenberg, J.E., Föllmi, K.B., 2019. Pulses of enhanced continental weathering associated with multiple Late Devonian climate perturbations: Evidence from osmium-isotope compositions. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 524, 240-249.
- Riquier, L., Averbuch, O., Devleeschouwer, X., Tribovillard, N., 2010. Diagenetic versus detrital origin of the magnetic susceptibility variations in some carbonate Frasnian-Famennian boundary sections from Northern Africa and Western Europe: implications for paleoenvironmental reconstructions. *Int. J. Earth Sci.* 99, 57-73.
- Sandberg, C.A., 1976. Conodont biofacies of late Devonian *Polygnathus styriacus* Zone in western United States, In: Barnes, C.R. (Ed.), *Conodont Paleocology*. Geol. Assoc. Canada spec. Pap. Montreal, pp. 171-186.
- Sandberg, C.A., Morrow, J.R., Ziegler, W., 2002. Late Devonian sea-level changes, catastrophic events, and mass extinctions. *Geol. Soc. Am. Spec. Pap.* 356, 473-487.
- Schülke, I., 1995. Evolutive Prozesse bei *Palmatolepis* in der frühen Famenne-Stufe (Conodonta, Ober-Devon). *Göttinger Arbeiten zur Geologie und Paläontologie*, Göttingen.

- Schülke, I., 1999. Conodont multielement reconstructions from the the early Famennian (Late Devonian) of the Montagne Noire (southern France). *Geol. Palaeontol.* 31, 1-123.
- Schülke, I., Popp, A., 2005. Microfacies development, sea-level change, and conodont stratigraphy of Famennian mid- to deep platform deposits of the Beringhauser Tunnel section (Rheinisches Schiefergebirge, Germany). *Facies* 50, 647-664.
- Seddon, G., Sweet, W.C., 1971. An ecologic model for conodonts. *J. Paleontol.* 45, 869-880.
- Spalletta, C., Perri, M.C., Over, D.J., Corradini, C. 2017. Famennian (Upper Devonian) conodont zonation: revised global standard. *Bull. Geosci.* 92 (1), 31-57.
- Tucker, E., 1974. Sedimentology of Palaeozoic pelagic limestones: the Devonian Griotte (Southern France) and Cephalopodenkalk (Germany). *Spec. Pub., Intern. Assoc. Sedimentologists* 1, 71-92.
- Wendt, J., Aigner, T., 1985. Facies patterns and depositional environments of Palaeozoic cephalopod limestones. *Sediment. Geol.* 44, 263-300. <http://doi.org/10.1144/SP414.15>
- Wiederer, U., Königshof, P., Feist, R., Franke, W. and Doublier M. P. 2002. Low-grade metamorphism in the Montagne Noire (S-France): Conodont Alteration Index (CAI) in Palaeozoic carbonates and implications for the exhumation of a hot metamorphic core complex. *Schweiz. mineral. petrogr. Mitt.* 82, 393-407.
- Ziegler, W., 1962. Taxonomy und Phylogenie oberdevonischer Conodonten und ihre stratigraphische Bedeutung. *Abh. hess. Landesamt. Bodenforsch. Wiesbaden.*

Figure captions

Figure 1. a) Early Famennian (-370.3 Ma) paleogeographic map (after Franke et al., 2017 and Golonka, 2020) showing the locations of the studied sections.

Abbreviations: SES: Sessacker (Rhenish Slate Mountains, Germany), BHT: Beringhauser Tunnel (Rhenish Slate Mountains, Germany), PZW: Pizzul West (Carnic Alps, Italy), CT: Col des Tribes (Montagne Noire, France), CM: Corona Mizziu (Sardinia, Italy), BU: Buschteich (Thuringia, Germany), ERF: Erfoud (Tafilalt, SE Morocco), DGHS (DuPont GHS Core, Eastern United States, after Over et al. 2019). **b)** Biostratigraphic correlation of the the studied sections.

Figure 2. Magnetic susceptibility curves through the Famennian (Late Devonian) for sections of S-Laurussia and N-Gondwana. Ages based on Girard et al. (2020), and updated after Becker et al. (2020). The curves represent the average of the MS values obtained with 2 to 5 measurements with the standard error in grey. Vertical dotted line: average MS values for lithified marine deposits estimated to 5.5×10^{-8} m³/kg (Ellwood et al. 2011a). Data of DGHS (Eastern United States) are after Over et al. (2019) and data of uppermost Frasnian to lower Famennian in Coumiac (CUQ*) and Berinhausen Tunnel (BHT*) after Riquier et al. (2010). Conodont Zones after Spalletta et al. (2017).

Figure 3. a) Magnetic susceptibility curves through the Famennian (Late Devonian) at Col des Tribes (CT, France) in green, Buschteich (BU, Germany) in red and Corona Mizziu (CMI and CM II, Italia) in orange, Sessacker (SES) in black. **b)** Conodont biofacies through time depicted as variations along the first axis of the PCA on conodont genera percentages of CT (in green), BU (in red), CM (dotted line in orange), and (SES) in black. **c)** Simplified

transgressive-regressive cycles after Johnson et al. (1985, 1996). Conodont Zones after Spalletta et al. (2017). Ages based on Girard et al. (2020), modified after Becker et al. (2020).

Supplementary Figure 1. Low field magnetic susceptibility (χ_{LF}) and geochemical data (Al_2O_3 , SiO_2 , Fe_2O_3 , TiO_2 , K_2O) for the PZW, CM II and CM I sections.

Supplementary Figure 2. Scatterplots reporting the MS versus selected geochemical indexes for samples from the Buschteich section (Level BU03 not figured). **a)** The zirconium (Zr) concentration (ppm); **b)** the iron (Fe) concentration (%); **c)** Fe_{exc} (%), **d)** Fe_{det} (%). The number of samples, the correlation coefficient r , and the p are reported for each diagram. Open red circles: levels after the “Enkeberg event”, Red coloured circles: levels before the “Enkeberg event”.

Supplementary Table 1. Magnetic susceptibility (MS) for the Col des Tribes (CT) section. MS in m^3/kg . Depth in cm. Conodont Zones after Spalletta et al. (2017). Ages based on Girard et al. (2020), and modified after Becker et al. (2020).

Supplementary Table 2. Magnetic susceptibility (MS) for the Buschteich (BU) section. MS in m^3/kg . Depth in cm. Conodont Zones after Spalletta et al. (2017). Ages based on Girard et al. (2020), and modified after Becker et al. (2020). Concentrations in Fe, Al, Zr (in ppm), and associated errors.

Supplementary Table 3. Magnetic susceptibility (MS) for the Sessacker (SES) section. MS in m^3/kg . Depth in cm. Conodont Zones after Spalletta et al. (2017). Ages based on Girard et al. (2020), and modified after Becker et al. (2020).

Supplementary Table 4. Magnetic susceptibility (MS) for the Beringhauser Tunnel (BHT) section. MS in m^3/kg . Depth in cm. Conodont Zones after Spalletta et al. (2017). Ages based on Girard et al. (2020), and modified after Becker et al. (2020).

Supplementary Table 5. Magnetic susceptibility (MS) for the Erfoud (ERF) section. MS in m³/kg. Depth in cm. Conodont Zones after Spalletta et al. (2017). Ages based on Girard et al. (2020), and modified after Becker et al. (2020).

Supplementary Table 6. Magnetic susceptibility (MS) for the Pizzul West (PZW) section. MS in m³/kg. Depth in cm. Conodont Zones after Spalletta et al. (2017). Ages based on Girard et al. (2020), and modified after Becker et al. (2020).

Supplementary Table 7. Magnetic susceptibility (MS) for the Corona Mizziu II (CM II) section. MS in m³/kg. Depth in cm. Conodont Zones after Spalletta et al. (2017). Ages based on Girard et al. (2020), and modified after Becker et al. (2020).

Supplementary Table 8. Magnetic susceptibility (MS) and low field magnetic susceptibility (χ_{LF}), ferromagnetic contribution (χ_{Ferro}), hysteresis parameters (high field magnetic susceptibility (χ_{HF})), saturation magnetization (MS), coercitive force (Hc), coercivity of remanence (Hcr), highfield remanence and viscosity decay; geochemical data for Corona Mizziu I (CM I) section.

Supplementary Table 9. Low field magnetic susceptibility (χ_{LF}), ferromagnetic contribution (χ_{Ferro}), hysteresis parameters (high field magnetic susceptibility (χ_{HF})), saturation magnetization (MS), coercitive force (Hc), coercivity of remanence (Hcr), highfield remanence and viscosity decay, geochemical data for the Corona Mizziu II (CM II) section.

Supplementary Table 10. Low field magnetic susceptibility (χ_{LF}), ferromagnetic contribution (χ_{Ferro}), hysteresis parameters (high field magnetic susceptibility (χ_{HF})), saturation magnetization (MS), coercitive force (Hc), coercivity of remanence (Hcr), highfield remanence and viscosity decay and geochemical data for the Pizzul West (PZW) section.

Supplementary Table 11. Mean MS (m³/kg), and conodont genera percentages (pc). Names of the samples: CT (Col des Tribes), BU (Buschteich), SES (Sessacker), CM (Corona Mizziu). Ages are given by the conodont zones after Spalletta et al. (2017) and by the age model (in My). Proportions of the different genera: pcPa: *Palmatolepis*; pcPo: *Polygnathus*; pcAn: *Ancyrognathus* and *Ancyrodella*; pcIc: *Icriodus*; pcBi: *Bispathodus*, *Branmehla* and *Mehlina*; pcSc: *Scaphignathus* and *Alternognathus*, pcSi: *Siphonodella*.

Table 1. Correlations between MS (χ_{LF}), and ferromagnetic contribution (χ_{Ferro}), high field magnetic susceptibility (χ_{HF}), coercivity of remanence (Hcr), and geochemical data using linear regression. Number of samples (N), coefficient of correlation (r) and probability (p) are given. Significant correlations marked in bold ($p < 0.05$).

			CM I			CM II			PZW		
			N	r	p	N	r	p	N	r	p
Xlf	Xferro		32	0.69	0.000	14	0.85	0.000	20	0.87	0.000
	Xhf		32	0.82	0.000	14	0.82	0.000	21	0.03	0.454
	Hcr		32	0.68	0.000	14	0.74	0.002	21	0.03	0.422
Xlf	Al2O3		13	0.68	0.010	11	0.25	0.463	17	0.005	0.785
Xhf			13	0.48	0.095	11	0.27	0.915	17	0.84	0.000
Xlf	SiO2		13	0.51	0.072	11	0.47	0.146	17	0.12	0.653
Xhf			13	0.55	0.050	11	0.49	0.122	17	0.81	0.000
Xlf	Fe2O3		13	0.70	0.008	11	0.67	0.023	17	0.13	0.613
Xhf			13	0.50	0.081	11	0.39	0.239	17	0.92	0.000
Xlf	TiO2		13	0.70	0.008	11	0.30	0.361	17	0.12	0.655
Xhf			13	0.48	0.096	11	0.31	0.350	17	0.85	0.000
Xlf	K2O		13	0.70	0.008	11	0.08	0.805	17	0.04	0.890
Xhf			13	0.45	0.123	11	0.46	0.158	17	0.83	0.000

Girard et al., Fig. 1

Figure 2

[Click here to access/download;Figure;Fig2_R#1_apercu.jpg](#)

Figure3

[Click here to access/download;Figure;Fig3_R#1_corrige_apercu.tif](#)

[Click here to access/download](#)
e-component
[Suppl_Fig1_R#2_corrige_aperçu.jpg](#)

[Click here to access/download](#)
e-component
[Suppl_Fig2_#1_corrige.jpg](#)

TableS1

[Click here to access/download](#)
e-component
TableS1.docx

[Click here to access/download](#)
e-component
TableS2.docx

[Click here to access/download](#)
e-component
TableS3.docx

[Click here to access/download](#)
e-component
TableS4.docx

[Click here to access/download](#)
e-component
TableS5.docx

[Click here to access/download](#)
e-component
TableS7.docx

Click here to access/download
e-component
Suppl_Table8_CMI_geoch#1.xlsx

[Click here to access/download](#)

e-component

[Suppl_Table9_CMII_mean_geoch#1.xls](#)

[Click here to access/download](#)

e-component

[Suppl_Table10_PZW_meangeoch#1bis.xlsx](#)