Morphogenesis of the human preimplantation embryo: bringing mechanics to the clinics

Julie Firmin and Jean-Léon Maître

Institut Curie, PSL Research University, Sorbonne Université, CNRS UMR3215, INSERM U934, Paris, France.

Correspondence to jean-leon.maitre@curie.fr

Abstract

During preimplantation development, the human embryo forms the blastocyst, the structure enabling uterine implantation. The blastocyst consists of an epithelial envelope, the trophectoderm, encompassing a fluid-filled lumen, the blastocoel, and a cluster of pluripotent stem cells, the inner cell mass. This specific architecture is crucial for the implantation and further development of the human embryo. Furthermore, the morphology of the human embryo is a prime determinant for clinicians to assess the implantation potential of in vitro fertilized human embryos, which constitutes a key aspect of assisted reproduction technology. Therefore, it is crucial to understand how the human embryo builds the blastocyst. As any material, the human embryo changes shape under the action of forces. Here, we review recent advances in our understanding of the mechanical forces shaping the blastocyst. We discuss the cellular processes responsible for generating morphogenetic forces that were studied mostly in the mouse and review the literature on human embryos to see which of them may be conserved. Based on the specific morphological defects commonly observed in clinics during human preimplantation development, we discuss how mechanical forces and their underlying cellular processes may be affected. Together, we propose that bringing tissue mechanics to the clinics will advance our understanding of human preimplantation development, as well as our ability to help infertile couples to have babies.

Introduction:

In the first days after fertilization, the human embryo needs to acquire implantation competencies [1–4]. The structure enabling the human embryo to implant is called the blastocyst, which has a very characteristic architecture [5,6]. It consists of an epithelium, the trophectoderm (TE), enveloping a fluid-filled lumen, the blastocoel, and a cluster of pluripotent stem cells, the inner-cell mass (ICM). The TE invades the maternal uterus and implants the embryonic tissues deriving from the ICM [7]. Until implantation, the human embryo can develop without intervention from the mother. As a consequence, the self-organisation of the human embryo into the blastocyst can be studied *ex vivo* and offers unique opportunities to study human embryonic development.

To form the blastocyst, the human embryo initially undergoes cleavage divisions, *i.e.* there is no cell growth during the interphase (Fig 1). Essentially, cells halve their volume with each cleavage and cells rearrange themselves to shape the blastocyst without changing the total cellular volume of the embryo, as measured accurately in the mouse [8]. The series of cellular rearrangements sculpting the blastocyst can be broken down into three steps: compaction, internalisation and lumen formation [5]. Compaction is the process by which the loosely attached blastomeres enlarge their cell-cell contacts and reduce their surface exposed to the outside medium [9–12]. Internalisation occurs when a subset of cells become entirely surrounded by neighbouring blastomeres and isolated from the outside medium [13–17]. This differential positioning of cells is a prerequisite to the first lineage specification into ICM for inner cells and TE for those remaining at the embryo surface [18–21]. Finally, the blastocoel appears when TE cells pump fluid, which pushes the ICM into one quadrant of the blastocyst [22–24]. The lumen breaks the radial symmetry of the embryo, which is key to the formation of the axes of symmetry after implantation [23,25,26]. The lumen also serves as a new interface for the differentiation of the ICM into primitive endoderm (PrE) and epiblast (Epi) and of the TE into mural and polar TE (mTE and pTE respectively) [27–30]. Sandwiched between the pTE and PrE, the Epi will provide all cells of the human body while the PrE contributes to most of the yolk sac [31,32]. The pTE is thought to mediate uterine implantation in human whereas this role seems devoted to the mTE in mice [28,29,31,33,34].

Since the advent of *in vitro* culture, the morphology of the human embryo has been observed and described in numerous studies. In fact, together with cell number, the morphology of the human embryo is one of the prime determinant in clinicians assessment of the implantation potential of human embryos [35,36]. For example, poor compaction or slow lumen growth are associated with lower implantation rates [10,37–39]. Therefore, the current and future efforts in developing algorithms able to better predict the implantation potential of human embryos grown *in vitro* will undoubtedly rely on morphological criteria [40–42].

Despite the importance of human embryo morphology for its development, we know little of the mechanisms responsible for shaping the human embryo [1–3]. When it comes to shaping any materials, forces are necessarily in action [43,44]. In the past decades, major discoveries regarding the nature of forces that shape animals were made using model organisms such as the worm, fly, fish, frog and, more recently, mouse [11,45–48]. The same forces are most likely responsible for the shaping of the human embryo. Of particular relevance to the shaping of the human blastocyst are surface tension [49–51], adhesive coupling between cells [52–54] and, osmotic pressure [55,56]. In animal cells, surface tension is governed by the contractility of the acto-myosin cortex, a thin layer of cross-linked actin filaments underneath the plasma membrane that is put under tension by the action of non-

muscle myosin II motors [57,58]. The mechanical stresses exerted by the acto-myosin cortex is a prime determinant of cell shape as it drives cell rounding and cytokinetic cleavage in half during cytokinesis, or cell body retraction during migration or blebbing [59–63]. Adhesive coupling between cells resists their separation [23,64,65]. It is mediated by cadherin adhesion molecules and associated proteins, the catenins, which anchor the adhesion complex to the actin cytoskeleton. By doing so, the adhesion complex connects the acto-myosin cortex of contacting cells and transmits its tension throughout the tissue [66,67]. Osmotic pressure dictates the movement of water between sealed compartments such as the outside medium, the cytoplasm and fluid-filled lumens. Osmotic pressure relies on tight junctions, which seal intercellular spaces, and ion pumps, which actively change the osmolarity of sealed compartments [68,69]. Osmotic gradients will then trigger the passive movement of water between compartments through aquaporin water channels [70].

In this review, we discuss how learning about the mechanics of tissues can help understanding the shaping of the human blastocyst. We consider how defective morphologies could be explained by aberrant force patterns and point to specific cellular process underlying them. In particular, we examine defects in cleavage divisions, compaction, internalisation and lumen formation. Since defects in the morphology of human embryos is a prime determinant of their health, bringing mechanics to the clinics will be key to improve assisted reproduction technologies. For reviews covering the patterning of blastocyst lineages or general clinical aspects, we recommend alternative reviews [1–4].

Cleavages

Before human morphogenesis begins, cleavage divisions provide the only changes in the morphology of the embryo (Fig 1). The first division takes place after a day and the second half a day later [71,72]. Then, sister blastomeres divide every day in waves of progressively decreasing synchrony (Fig 1). The early cleavages are notoriously error prone [4,73]. They frequently lead to chromosome segregation errors [74] and are at the origin of aneuploidy that can lead to developmental arrest [73]. Alternatively, these errors can be corrected by excluding aneuploid cells from the embryonic tissue, which are later found in the placenta [75]. Aneuploid cells can also contribute to the epiblast with or without further consequences on embryonic development [76]. In addition to chromosome mis-segregation, cytokinesis can also be faulty. Cytokinesis should split cells in two equal volumes, forming embryos with even cell number and size. However, this is frequently not the case and has dire consequences for the embryo. Indeed, human embryos with unequally sized blastomeres at the 4- and 8-cell stages were found to have more frequent multinucleation and ploidy issues, which eventually reduces their implantation rate [77–79]. Therefore, unequally sized blastomeres can be used as an indicator of the genome integrity of the embryo and of its health in general.

Unequal blastomere sizes can result from distinct events (Fig 2). Cytokinesis itself can produce sister cells with uneven sizes. Skewing the cleavage furrow during cytokinesis, for example by contractility unbalance, will result in uneven cleavage [62]. Alternatively, blastomeres can undergo “reverse cleavages”, fusing sister cells back together after the cleavage furrow completed the separation of the cellular volumes of the sister cells [72,80]. In addition to reducing the total expected number of cell, this results in the formation of polyploid blastomeres with twice the size of their neighbouring cells, which can easily be detected. Incomplete cytokinesis could result from defects in abscission, the process sealing apart the cytoplasm of sister cells and leading to the formation of the midbody [81]. Abscission requires

the coordination of complex cellular machineries and often fails when chromosomes are mis-segregated. Often referred to as “direct cleavage”, blastomeres can cleave into 3 similarly sized daughter cells instead of 2. After the first and second cleavages, embryos can show 3 cells and 5 cells respectively [72,82]. This increases the number of expected cells and results in the formation of blastomeres that are $1/3^{\text{rd}}$ smaller than expected, which can be difficult to notice with insufficient resolution. Direct cleavages are associated with fewer embryos reaching blastocyst stage and reduced implantation rate [82]. Nevertheless, blastocysts produced after direct cleavages can show normal ploidy, suggesting some repairing mechanism, possibly involving the exclusion of defective cells [72,83]. Such abnormal divisions can result from multipolar spindle and supernumerary centrioles caused by defective sperms [84,85]. Finally, human blastomeres often produce cellular fragments during cytokinesis, which wastes cellular material [73,86]. Importantly, fragments sometimes entrap chromosomes, making their cell of origin aneuploid [73,87]. Unsurprisingly, fragmentation is associated with poor prognosis for implantation [88,89]. Fragmentation is observed frequently in human embryos but rarely in mice, which makes the process difficult to study. Why human blastomeres produce so many fragments during cytokinesis is unclear. Blocking the actin and microtubule cytoskeletons reduces fragmentation, suggesting that these generate the forces driving fragmentation [90]. Cellular fragments have been compared to blebs due to their spherical shape. However, blebs are short-lived (around a minute), while fragments are stable over tens of hours [60]. Blebs could persist for longer time if contractility would be hyper-activated [91]. However, there is not enough information available on fragment dynamics to understand how they form and persist. Alternatively, membrane threads connecting the zona pellucida to blastomeres were proposed to pull on blastomeres during cytokinesis [92].

Compaction

Starting as early as during the 8-cell stage, compaction is the first morphogenetic process associated with the formation of the blastocyst (Fig 1). During this process, blastomeres get closer together, forming a tighter structure (Fig 3). This developmentally regulated adhesion process requires the calcium-dependent cell-cell adhesion molecule CDH1 (formerly known as E-cadherin or uvomorulin) [90,93–95]. Removing extracellular calcium prevents CDH1 binding and causes embryos to de-compact [96–98]. The compaction process was long thought to be driven by increased adhesion of cells via modifications of the CDH1-dependent adhesion machinery [1,99]. However, mechanical measurements in the mouse embryo revealed that the forces driving compaction are located at the embryo surface rather than at cell-cell contacts [11]. Micropipette aspiration of mouse blastomeres throughout compaction uncovers raising tensions at the surface of the embryo as a result from the action of the actomyosin cortex (Toolbox 1). Cells literally pull themselves together using their intracellular muscles. The adhesion molecule CDH1 enables compaction by anchoring the actin cytoskeleton of contacting cells so that they can effectively pull onto each other [67]. CDH1 also provides local signals lowering contractility at cell-cell contacts and effectively relaxing them [100]. Whether such mechanism is also responsible for human compaction remains to be tested. Identifying the nature of the forces driving human embryo compaction would help understanding what causes this process to fail in some instances.

Clinical studies found that compaction can be defective in several different ways (Fig 3). Compaction can be globally reduced with all blastomeres simply failing to grow their cell-cell contacts, which is associated with fewer embryos reaching blastocyst stage and lower

implantation rates [38,101]. Several cellular processes could be at fault, such as cell adhesion or contractility. Micropipette aspiration on embryos failing to compact would narrow down the options. Low tensions at the embryo surface would point towards contractility problems, such as in mouse mutants for myosin paralogs [16,102]. High tensions at the embryo surface would instead suggest defective adhesion, such as in mouse mutants for cadherin adhesion molecules [11].

Alternatively, compaction can be delayed. While a minority of embryo start compaction as early as during the 8-cell stage, the majority of human embryos begin compaction during the 4th wave of cleavages, with at least 9 blastomeres and most finish compaction during the 16-cell stage [10]. Embryos compacting less than 80 h after fertilization show higher rate of implantation and live birth than embryos compacting after the 80 h mark [12,101,103]. What controls the timing of compaction remains unclear though. In the mouse, activation of PKC can induce a premature (and transient) compaction [104,105], which has been proposed to promote contractility at the embryo surface [106].

Finally, compaction can be unequal (also referred to as partial compaction), with some blastomeres forming a compact mass while other cells appear excluded from this group (Fig 3). Excluded cells may later on contribute to the TE or be excluded entirely from the blastocyst. The time of exclusion can occur before or after compaction is complete, with higher survival rates for embryos excluding cells during the compaction process [9]. Interestingly, excluded cells tend to be aneuploid while the rest of the embryo seems euploid [72]. This indicates that exclusion of cells during compaction could serve as a repair mechanism ensuring that defective cells do not participate to the blastocyst or to the embryonic tissues. Indeed, cells of the placenta show a high rate of aneuploidy and have long been thought to act as a sink eliminating defective cells from embryonic tissues [75]. However, the mechanism underlying unequal compaction and exclusion of defective cells remain unknown. Again, excluded cells could have lower tension at their surfaces or higher tensions at their cell-cell contacts, which, if measured, would point at the underlying cellular process at fault, contractility and/or adhesion (Toolbox 1).

Internalisation of the ICM precursors

The positioning of cells within the human embryo away from the cell-medium interface is a critical morphogenetic step for the formation of the TE and ICM lineages (Fig 1). Experimental manipulation of cell position reveals that outside cells from early human blastocyst can adopt ICM fate if transplanted onto the ICM, away from the cell-medium interface [18]. This apparent plasticity is explained by the fact that human lineages are definitely set in the late blastocyst stage [20,31,107]. Until then, the position of cells within the human embryo will guide their differentiation into either TE or ICM. How do blastomeres adopt different positions within the human embryo in the first place?

Regardless of the molecular and cellular mechanisms that may control cell internalisation, with sufficient cleavages, blastomeres could end up on the inside of the embryo for geometrical reasons only [108]. Indeed, the packing of spherical objects into a sphere would necessarily position cells on the inside when they become sufficiently small as compared to the size of the embryo. If human embryos would be relying on geometrical packing only, then reducing the embryo size would prevent the formation of an ICM. Dissociating human embryos at the 4-cell stage reveals that 4-cell stage blastomeres are able to form smaller blastocyst [21]. Quarter embryos compact, grow their lumen and contain inner cells expressing the ICM

marker NANOG. This implies that mechanisms other than geometrical packing drive cell internalisation in human embryos.

In the mouse, at least two distinct mechanisms can internalise cells: oriented cell divisions and contractility-mediated cell sorting. Oriented cell division consists in aligning the mitotic spindle perpendicularly to the embryo surface so that one daughter cell is pushed inward after cytokinesis [15,109,110]. Cell sorting occurs during the interphase and relies on differences in surface tensions between cells, which, if sufficiently high, allow the strongest cells to pull their way inside the embryo [16,17]. Both positioning mechanisms rely on the establishment of apicobasal polarity with the apical domain facing the cell-medium interface [13–16]. Division orientation is guided by the tethering of one of the spindle pole to the apical domain [15]. This also facilitates the asymmetric inheritance of the apical domain, including sub-apical components such as intermediate filaments when present [111]. The asymmetric division of apical components, such as the apical kinases PRKCz and PRKCi (also known as aPKC_i and aPKC_z) that downregulate actomyosin contractility, among sister cells is key to generate cell populations with different mechanical properties and drive cell sorting (Toolbox 1) [16,112]. Last but not least, apicobasal polarity governs the positional signals guiding the differentiation of TE and ICM lineages (Toolbox 2). Apical signals promote TE lineages by enforcing the nuclear localisation of the co-transcriptional activator YAP while cell-cell contacts favour its cytoplasmic degradation [112,113]. Therefore, the apical domain is a master regulator of lineage positioning and specification in the mouse embryo.

How much of this mechanism is conserved in human embryos is still unclear. Recently, the presence and function of the core components of the apical machinery controlling TE fate were confirmed in human and cow embryos [20]. Apical markers such as PRKCz, PARD6B or AMOT are observed at the apical domain of human embryos (Toolbox 2). The accumulation of PARD6B also seems to occur about 90 h post fertilization [114]. The human embryo then consists of at least 16 cells and compaction has completed [10,12,103]. This is later than in the mouse, which compacts and polarises during the 8-cell stage [11,20]. Similarly to the mouse [112], inhibition of PRKCz using a chemical inhibitor or TRIM-AWAY, which targets proteins to the proteasome using antibodies [115], impacts lineage specification in human embryos [20]. When apical signals are reduced, YAP nuclear localisation is compromised and the level of the TE-specific transcription factor GATA3 in outer cells is reduced [20]. Therefore, the TE specification module by apical signals seems conserved in human embryos. However, whether human embryos also rely on apicobasal polarity to position inner and outer cells remains to be determined.

At that point, the human embryo forms what is called the morula (Fig 1). The morula is compact, surface cells have an apical domain facing the outside medium, which initiate their TE differentiation, while inner cells do not have apical material and start adopting an ICM fate. Importantly, compaction is functionally independent from polarisation (and cell internalisation). This is best illustrated with mouse mutant embryos in which the processes can be impaired selectively. Lacking essential apical kinases, *Prkcz*; *Prkci* mutants fail to form normal apical domains, which prevents the correct specification of TE and ICM lineages but show a normal, or even enhanced, compaction [112]. On the other hand, *Myh9* or *Myh9*; *Myh10* mutants fail to compact due to the absence of sufficient actomyosin contractility but polarise correctly and form ICM and TE lineages in the correct proportions [102]. Therefore, despite their apparent synchrony, compaction and polarisation (and cell internalisation) can occur independently from

one another in mouse mutants. This also seems to be the case for human embryos, which appear to compact normally when apical polarisation is impaired using inhibitors or TRIM-AWAY [20]. Whether compaction is required for human embryo polarisation remains to be tested.

Lumen formation

Apicobasal polarisation is also key for the formation of the blastocoel, the first mammalian lumen, which is the last morphogenetic step shaping the blastocyst (Fig 1). Fluid accumulates within the morula and inflates the embryo to almost 10 times its original volume [39]. Fluid is pumped into the intercellular space through the polarised surface epithelium while inner cells cluster into a bud against the epithelium. This bud forms the embryonic pole of the blastocyst, where the human embryo will mediate the uterine implantation [28,29,33,34]. In the expanded blastocyst, surface cells are now differentiated into TE and inner cells into ICM, which further specifies into PrE and Epi as the blastocyst prepares for implantation.

The formation of the blastocoel relies on the tight sealing of the TE, polarised transport and the ordered detachment of the ICM into one pole. Recent studies in the mouse provided mechanistic details on how these architectural changes proceed. Initially forming a domain in the centre of the cell-medium interface of surface cells, the apical domain expands [15]. When the apical domain hits the apical edge of cell-cell junctions, tight junctions seal the TE [24]. The expansion of the apical domain relies on coordinated cytoskeletal actions of the microtubules signalling to exclude acto-myosin from the apical domain, as observed in PRKC mutants [16,24]. This sealing can be challenged during cell divisions, which requires cells to round up and pull on their neighbouring cells, putting junctions under mechanical stress [22,116]. Therefore, tight junctions must reinforce themselves to prevent the embryo from collapsing under the increasing pressure of the blastocoel and during the waves of divisions of the TE. Polarised transport occurs through the cells rather than via junctions, as suggested by mouse mutants that, after failing all successive divisions, form a single-celled embryo, which nevertheless initiates blastocoel formation [102]. Indeed, as long as osmolytes transporters, such as the Na/K pump, and aquaporins are polarised along the apicobasal axis, blastomeres pump fluid from their apical to their basolateral compartment. This basolateral compartment is enriched in adhesion molecules, such as CDH1, and mechanically opposes detachment and fluid accumulation [23]. However, the fluid pressure, between 5-10 times higher than that of cells, is large enough to fracture cell-cell contacts. This includes contacts between ICM cells, which are transiently broken [23]. Contacts then repair themselves thanks to the action of their actomyosin cytoskeleton [117]. Interestingly, patterning the contractility of cells is sufficient to direct the positioning of the lumen and ICM, effectively dictating the first axis of symmetry of the mammalian embryo, which guides its implantation.

How much of these mechanisms are conserved in human embryos is unclear. The presence of key components from the tight junctions and fluid pumping machineries are likely [19,118–121]. More generally, the differentiation of the TE into a functional epithelium is required for the integrity of the human blastocoel, which fails when key transcription factor POU5F1 is mutated [19]. If conserved in human embryos, these mechanisms could have important clinical implications for better assessing embryos. The rate of blastocyst expansion is thought to provide information on its implantation potential, with implanting embryos showing higher expansion rates than non-implanting embryos [37,39]. Interestingly, the expansion rate is also an indicator of the ploidy state of the embryo, with aneuploid embryos inflating slower

than euploid ones [122]. Aneuploidies seem to be more frequent in the TE directly lying over the ICM, the polar TE, which attaches to the uterus [123]. The reason for the slower expansion rate could be linked to leaky TE, for example as a result of poor sealing of tight junctions or could be due to weaker polarised transport. Non-invasive methods allowing measuring the sealing and transport of human embryos would be very helpful to assess these properties and better understand the causes underlying slower expansion. Fluorescence based assays can be used to detect leakage [22,68]. Measuring volume changes after osmotic shocks would allow assessing the transport properties [124]. Besides, analysing blastocyst growth rates with higher temporal resolution could give more information on sudden collapses [55,116]. Several studies investigated the relationship between collapses and implantation rates, with sometimes contradictory conclusions [125–127]. As collapses may result from cell divisions transiently disrupting the TE seal, they may report on a healthy dividing TE [22,116]. Alternatively, if collapses are too frequent, they may reflect a poorly sealed and therefore dysfunctional TE. Finally, reports of human blastocyst with multiple clusters of inner cells suggest that monozygotic twinning may originate from the blastocyst [128]. If conserved in human embryos, hydraulic fracturing of contact between inner cells during blastocoel initiation may be responsible for the formation of multiple ICM. However, the formation of multiple ICM was not observed in the mouse, even when the fracturing or repair mechanisms were genetically manipulated [23]. Further investigations in the adhesive properties of ICM and TE cells will be needed to identify how the ICM could split.

Perspectives

Based on recent studies in mouse and human embryos, we have detailed how key steps of the shaping of the human blastocyst could proceed and fail. Since morphology serves as the basis for selecting the most suitable human embryos during assisted reproduction technology procedures [35,36], we believe there is much to be learned from the mechanisms of blastocyst morphogenesis. Current and future studies are taking exciting directions with the development of artificial intelligence (AI) based image analysis and non-invasive mechanical measurements. AI already shows performances equivalent to or even better than clinicians in predicting implantation rates, based on retrospective studies [40–42]. Most importantly, AI will allow more systematic procedures between clinics, as embryo diagnostic will rely less on the subjectivity of clinician eyes. However, algorithms, especially those based on machine learning, do not necessarily permit knowing the precise criteria used for predicting the implantation potential of embryos. Therefore, we should not rely on AI to uncover mechanistic understanding underlying developmental defects. On the other hand, non-invasive mechanical measurements offer the possibility to better understand the mechanisms for a given morphogenetic defects. For example, micropipette aspiration has been used on mouse and human oocytes and provided improved prediction of developmental potential [129]. Micropipette aspiration was performed through the zona pellucida, a porous glycoprotein shell of high elasticity compared to the oocyte. This aspiration measurement is therefore likely to reflect how porous and elastic the zona pellucida is, rather than probing the mechanics of the oocyte. Nevertheless, similar measurements could reveal how compaction fails in human embryos or even specific blastomeres. Such microaspiration could for example be performed as blastomeres are being removed during preimplantation genetic diagnostics. Together, better understanding the shaping of human blastocyst will improve the success rates of ART

351 procedures, which become more frequent as the age of first conception keeps on increasing
352 [130].

Acknowledgements

We thank Diane Pelzer and Markus Schliffka for critical reading of the manuscript. J.F. is funded by the Fondation pour la Recherche Médicale. Research in the lab of J.-L.M. is supported by the Institut Curie, the Centre National de la Recherche Scientifique (CNRS), the Institut National de la Santé Et de la Recherche Médicale (INSERM), and is funded by grants from the ATIP-Avenir program, the Fondation Schlumberger pour l'Éducation et la Recherche, the European Research Council Starting Grant ERC-2017-StG 757557, the European Molecular Biology Organization Young Investigator program (EMBO YIP), the INSERM transversal program Human Development Cell Atlas (HuDeCA), Paris Sciences Lettres (PSL) "nouvelle équipe" and QLife (17-CONV-0005) grants and Labex DEEP (ANR-11-LABX-0044) which are part of the IDEX PSL (ANR-10-IDEX-0001-02).

Figure 1: Human blastocyst morphogenesis

Starting from the zygote (Day 0), the human embryo undergoes cleavage divisions, which become progressively less synchronous with successive waves of cleavages, finishing here with the 6th wave on Day 5. During the cleavage stages, there is no morphogenesis. Compaction begins on Day 3, around the time of the 4th wave and ends on Day 4. The morula is complete when the embryo is compacted and contains inner cells (red). At the time of 5th cleavage, the outer cells (blue) pump fluid to inflate a lumen and turn the embryo into the blastocyst. Defective morphologies result from defects in cleavages (see Fig 2), compaction (see Fig 3) and/or lumen formation. Note that developmental time is not represented linearly.

Figure 2: Cleavage defects during human preimplantation development

Normal cleavage division splits the volume of the blastomere in equal parts (top). Cleavage defects can affect the volume of daughter cells in a noticeable way (right). Cytokinesis itself can be asymmetric and produce daughter cells of unequal size. Tripolar mitotic spindles can produce three daughter cells instead of two and result in what is referred to as direct cleavage. This often result in the production of aneuploid cells (marked by the presence of extra chromosomes in red). Cytokinesis or abscission can fail resulting in the fusion of sister cells and the formation of a polyploid or bi-nucleated cell (reverse cleavage). During cytokinesis, cellular fragment frequently form. Fragments waste material and sometimes entrap chromosomes leading to aneuploidy.

Figure 3: Compaction defects in human embryos

During normal compaction (top), the human embryo increases the area of contact between cells and reduces the surface exposed to the outside medium to form the compacted morula. Compaction can fail completely or be weak (bottom left). Compaction can also occur in subset of blastomeres only, excluding some blastomeres (bottom right). Blastomeres excluded from unequally compacted morula can be aneuploid. Excluded blastomeres could contribute to the TE and therefore placental tissues or be excluded from the blastocyst entirely.

Toolbox 1: Surface tensions during compaction and cell internalisation

Comparing the human embryo to contacting soap bubbles provides a general theoretical framework explaining compaction and cell internalisation [11,16,108]. Blastomeres are given a surface tension at their cell-medium (γ_{cm}) and cell-cell interfaces (γ_{cc}). Following the Young-Dupré equation, the level of compaction is simply given by the ratio α of tension between cell-cell and cell-medium interfaces ($\gamma_{cc}/2\gamma_{cm}$). The surface tension of blastomeres can be calculated using the Young-Laplace equation, which relates the surface tension, pressure and curvature of liquid-like materials. Micropipette aspiration can then be used to determine the pressure inside blastomeres [11,131]. This approach revealed that, during compaction of the mouse embryo, γ_{cm} doubles while γ_{cc} decreases by a third. Using the absolute values of surface tensions further uncovers that $\frac{3}{4}$ of mouse compaction results from the changes in surface tension γ_{cm} and $\frac{1}{4}$ from the relaxation of cell-cell contacts [11].

Considering the ratio of surface tensions at cell-medium interfaces of individual blastomeres $\delta = \gamma_{cm1}/\gamma_{cm2}$, internalisation occurs when cell 1 grows its tension above a threshold value. This threshold is set at $\delta > 1 + 2\alpha$, and therefore depends the level of compaction [16]. Interestingly, differences in size have no effect on internalisation in theory. For the mouse embryo, the internalisation threshold is at 1.5, i.e. cells need to grow their tension 50% higher than that of their neighboring cells to pull their way inside the embryo.

Toolbox 2: De novo apicobasal polarisation and TE-ICM specification

During preimplantation development, blastomeres form de novo a domain of apical material (red) at the surface of the embryo, away from cell-cell contacts (basolateral interfaces in blue). The apical domain forms in the center of the contact free surface and signals to the actomyosin cortex (green) to deplete it locally. The apical domain progressively expands until it reaches the apical edge of the cell junctions. The apical domain contains molecules such as **CDC42**, **PARD6B** or **PRKC** that prevent the LATS kinase from phosphorylating the co-transcriptional activator YAP. YAP can shuttle to the nucleus to interact with the transcription factor TEAD4 and activate the expression of TE specific genes such as GATA3. Without an apical domain, inner cells show less YAP in their nucleus as the LATS kinase, with the help of NF2 and AMOT localised at cell-cell contacts, can phosphorylate YAP and target it to the proteasome. The presence of some of the signalling molecules involved in the subcellular localisation of YAP has been confirmed in human embryos (labelled in bold font), while other have only been studied in mouse embryos.

References

- [1] G. Coticchio, C. Lagalla, R. Sturmey, F. Pennetta, A. Borini, The enigmatic morula: mechanisms of development, cell fate determination, self-correction and implications for ART, *Human Reproduction Update*. 25 (2019) 422–438. <https://doi.org/10.1093/humupd/dmz008>.
- [2] M.N. Shahbazi, Mechanisms of human embryo development: from cell fate to tissue shape and back, *Development*. 147 (2020) dev190629. <https://doi.org/10.1242/dev.190629>.
- [3] S.E. Wamaitha, K.K. Niakan, Human Pre-gastrulation Development, *Current Topics in Developmental Biology*. 128 (2018) 295–338. <https://doi.org/10.1016/bs.ctdb.2017.11.004>.
- [4] M. Popovic, L. Dhaenens, A. Boel, B. Menten, B. Heindryckx, Chromosomal mosaicism in human blastocysts: the ultimate diagnostic dilemma, *Human Reproduction Update*. 26 (2020) 313–334. <https://doi.org/10.1093/humupd/dmz050>.
- [5] J.-L. Maître, Mechanics of blastocyst morphogenesis., *Biology of the Cell*. 109 (2017) 323–338. <https://doi.org/10.1111/boc.201700029>.
- [6] M.D. White, J. Zenker, S. Bissiere, N. Plachta, Instructions for Assembling the Early Mammalian Embryo, *Developmental Cell*. 45 (2018) 667–679. <https://doi.org/10.1016/j.devcel.2018.05.013>.
- [7] E. Posfai, I. Rovic, A. Jurisicova, The mammalian embryo's first agenda: making trophoctoderm, *The International Journal of Developmental Biology*. 63 (2019) 157–170. <https://doi.org/10.1387/ijdb.180404ep>.
- [8] R. Niwayama, P. Moghe, Y.-J. Liu, D. Fabréges, F. Buchholz, M. Piel, T. Hiiragi, A Tug-of-War between Cell Shape and Polarity Controls Division Orientation to Ensure Robust Patterning in the Mouse Blastocyst, *Developmental Cell*. (2019) 1–18. <https://doi.org/10.1016/j.devcel.2019.10.012>.
- [9] G. Coticchio, K. Ezoe, C. Lagalla, K. Shimazaki, K. Ohata, M. Ninomiya, N. Wakabayashi, T. Okimura, K. Uchiyama, K. Kato, A. Borini, Perturbations of morphogenesis at the compaction stage affect blastocyst implantation and live birth rates, *Human Reproduction*. 36 (2021) 918–928. <https://doi.org/10.1093/humrep/deab011>.
- [10] K. Iwata, K. Yumoto, M. Sugishima, C. Mizoguchi, Y. Kai, Y. Iba, Y. Mio, Analysis of compaction initiation in human embryos by using time-lapse cinematography, *J Assist Reprod Genet*. 31 (2014) 421–426. <https://doi.org/10.1007/s10815-014-0195-2>.
- [11] J.-L. Maître, R. Niwayama, H. Turlier, F. Nédélec, T. Hiiragi, Pulsatile cell-autonomous contractility drives compaction in the mouse embryo., *Nature Cell Biology*. 17 (2015) 849–855. <https://doi.org/10.1038/ncb3185>.
- [12] L. Rienzi, D. Cimadomo, A. Delgado, M.G. Minasi, G. Fabozzi, R. del Gallego, M. Stoppa, J. Bellver, A. Giancani, M. Esbert, A. Capalbo, J. Remohí, E. Greco, F.M. Ubaldi, M. Meseguer, Time of morulation and trophoctoderm quality are predictors of a live birth after euploid blastocyst transfer: a multicenter study, *Fertility and Sterility*. 112 (2019) 1080-1093.e1. <https://doi.org/10.1016/j.fertnstert.2019.07.1322>.
- [13] S. Anani, S. Bhat, N. Honma-Yamanaka, D. Krawchuk, Y. Yamanaka, Initiation of Hippo signaling is linked to polarity rather than to cell position in the pre-implantation mouse embryo., *Development (Cambridge, England)*. 141 (2014) 2813–2824. <https://doi.org/10.1242/dev.107276>.
- [14] N. Dard, T. Le, B. Maro, S. Louvet-Vallée, Inactivation of aPKC λ reveals a context dependent allocation of cell lineages in preimplantation mouse embryos., *PLoS ONE*. 4 (2009) e7117. <https://doi.org/10.1371/journal.pone.0007117>.

- 482 [15] E. Korotkevich, R. Niwayama, A. Courtois, S. Friesse, N. Berger, F. Buchholz, T.
483 Hiiragi, The Apical Domain Is Required and Sufficient for the First Lineage
484 Segregation in the Mouse Embryo., *Dev Cell.* 40 (2017) 235–247.e7.
485 <https://doi.org/10.1016/j.devcel.2017.01.006>.
- 486 [16] J.-L. Maître, H. Turlier, R. Illukkumbura, B. Eismann, R. Niwayama, F. Nédélec, T.
487 Hiiragi, Asymmetric division of contractile domains couples cell positioning and fate
488 specification., *Nature.* 536 (2016) 344–348. <https://doi.org/10.1038/nature18958>.
- 489 [17] C.R. Samarage, M.D. White, Y.D. Álvarez, J.C. Fierro-González, Y. Henon, E.C.
490 Jesudason, S. Bissiere, A. Fouras, N. Plachta, Cortical Tension Allocates the First Inner
491 Cells of the Mammalian Embryo, *Developmental Cell.* 34 (2015) 435–447.
492 <https://doi.org/10.1016/j.devcel.2015.07.004>.
- 493 [18] C. De Paepe, G. Cauffman, A. Verloes, J. Sterckx, P. Devroey, H. Tournaye, I.
494 Liebaers, H. Van de Velde, Human trophoctoderm cells are not yet committed, *Human*
495 *Reproduction.* 28 (2013) 740–749. <https://doi.org/10.1093/humrep/des432>.
- 496 [19] N.M.E. Fogarty, A. McCarthy, K.E. Snijders, B.E. Powell, N. Kubikova, P. Blakeley,
497 R. Lea, K. Elder, S.E. Wamaitha, D. Kim, V. Maciulyte, J. Kleinjung, J.-S. Kim, D.
498 Wells, L. Vallier, A. Bertero, J.M.A. Turner, K.K. Niakan, Genome editing reveals a
499 role for OCT4 in human embryogenesis, *Nature.* 550 (2017) 67–73.
500 <https://doi.org/10.1038/nature24033>.
- 501 [20] C. Gerri, A. McCarthy, G. Alanis-Lobato, A. Demtschenko, A. Bruneau, S. Loubersac,
502 N.M.E. Fogarty, D. Hampshire, K. Elder, P. Snell, L. Christie, L. David, H. Van de
503 Velde, A.A. Fouladi-Nashta, K.K. Niakan, Initiation of a conserved trophoctoderm
504 program in human, cow and mouse embryos, *Nature.* 587 (2020) 443–447.
505 <https://doi.org/10.1038/s41586-020-2759-x>.
- 506 [21] H. Van de Velde, G. Cauffman, H. Tournaye, P. Devroey, I. Liebaers, The four
507 blastomeres of a 4-cell stage human embryo are able to develop individually into
508 blastocysts with inner cell mass and trophoctoderm, *Human Reproduction (Oxford,*
509 *England).* 23 (2008) 1742–1747. <https://doi.org/10.1136/vr.114.10.240>.
- 510 [22] C.J. Chan, M. Costanzo, T. Ruiz-Herrero, G. Mönke, R.J. Petrie, M. Bergert, A. Diz-
511 Muñoz, L. Mahadevan, T. Hiiragi, Hydraulic control of mammalian embryo size and
512 cell fate., *Nature.* 571 (2019) 112–116. <https://doi.org/10.1038/s41586-019-1309-x>.
- 513 [23] J.G. Dumortier, M. Le Verge-Serandour, A.F. Tortorelli, A. Mielke, L. de Plater, H.
514 Turlier, J.-L. Maître, Hydraulic fracturing and active coarsening position the lumen of
515 the mouse blastocyst., *Science.* 365 (2019) 465–468.
516 <https://doi.org/10.1126/science.aaw7709>.
- 517 [24] J. Zenker, M.D. White, M. Gasnier, Y.D. Álvarez, H.Y.G. Lim, S. Bissiere, M. Biro, N.
518 Plachta, Expanding Actin Rings Zipper the Mouse Embryo for Blastocyst Formation,
519 *Cell.* (2018) 1–34. <https://doi.org/10.1016/j.cell.2018.02.035>.
- 520 [25] H. Honda, N. Motosugi, T. Nagai, M. Tanemura, T. Hiiragi, Computer simulation of
521 emerging asymmetry in the mouse blastocyst., *Development (Cambridge, England).*
522 135 (2008) 1407–1414. <https://doi.org/10.1242/dev.014555>.
- 523 [26] Y. Kurotaki, K. Hatta, K. Nakao, Y. Nabeshima, T. Fujimori, Blastocyst axis is
524 specified independently of early cell lineage but aligns with the ZP shape., *Science.* 316
525 (2007) 719–723. <https://doi.org/10.1126/science.1138591>.
- 526 [27] C. Chazaud, Y. Yamanaka, Lineage specification in the mouse preimplantation
527 embryo, *Development (Cambridge, England).* 143 (2016) 1063–1074.
528 <https://doi.org/10.1242/dev.128314>.
- 529 [28] A. Deglincerti, G.F. Croft, L.N. Pietila, M. Zernicka-Goetz, E.D. Siggia, A.H.
530 Brivanlou, Self-organization of the in vitro attached human embryo, *Nature.* 533 (2016)
531 251–254. <https://doi.org/10.1038/nature17948>.

- [29] M.N. Shahbazi, A. Jedrusik, S. Vuoristo, G. Recher, A. Hupalowska, V. Bolton, N.M.E. Fogarty, A. Campbell, L.G. Devito, D. Ilic, Y. Khalaf, K.K. Niakan, S. Fishel, M. Zernicka-Goetz, Self-organization of the human embryo in the absence of maternal tissues, *Nature Cell Biology*. 18 (2016) 700–708. <https://doi.org/10.1038/ncb3347>.
- [30] C.S. Simon, S. Rahman, D. Raina, C. Schröter, A.-K. Hadjantonakis, Live Visualization of ERK Activity in the Mouse Blastocyst Reveals Lineage-Specific Signaling Dynamics, *Developmental Cell*. 55 (2020) 341–353.e5. <https://doi.org/10.1016/j.devcel.2020.09.030>.
- [31] D. Meistermann, S. Loubersac, A. Reignier, J. Firmin, V.F. Campion, S. Kilens, Y. Lelièvre, J. Lammers, M. Feyeux, P. Hulin, S. Nedellec, B. Bretin, S. Covin, G. Castel, A. Bihouée, M. Soumillon, T. Mikkelsen, P. Barrière, J. Bourdon, T. Fréour, L. David, Spatio-temporal analysis of human preimplantation development reveals dynamics of epiblast and trophectoderm, *Cell Biology*, 2019. <https://doi.org/10.1101/604751>.
- [32] L. Xiang, Y. Yin, Y. Zheng, Y. Ma, Y. Li, Z. Zhao, J. Guo, Z. Ai, Y. Niu, K. Duan, J. He, S. Ren, D. Wu, Y. Bai, Z. Shang, X. Dai, W. Ji, T. Li, A developmental landscape of 3D-cultured human pre-gastrulation embryos, *Nature*. 577 (2020) 537–542. <https://doi.org/10.1038/s41586-019-1875-y>.
- [33] A. Aberkane, W. Essahib, C. Spits, C. De Paepe, K. Sermon, T. Adriaenssens, S. Mackens, H. Tournaye, J.J. Brosens, H. Van de Velde, Expression of adhesion and extracellular matrix genes in human blastocysts upon attachment in a 2D co-culture system, *MHR: Basic Science of Reproductive Medicine*. (2018). <https://doi.org/10.1093/molehr/gay024>.
- [34] S. Lindenberg, 1 Ultrastructure in human implantation: Transmission and scanning electron microscopy, *Baillière's Clinical Obstetrics and Gynaecology*. 5 (1991) 1–14. [https://doi.org/10.1016/S0950-3552\(05\)80067-7](https://doi.org/10.1016/S0950-3552(05)80067-7).
- [35] Alpha Scientists in Reproductive Medicine and ESHRE Special Interest Group of Embryology, B. Balaban, D. Brison, G. Calderon, J. Catt, J. Conaghan, L. Cowan, T. Ebner, D. Gardner, T. Hardarson, K. Lundin, M. Cristina Magli, D. Mortimer, S. Mortimer, S. Munne, D. Royere, L. Scott, J. Smits, A. Thornhill, J. van Blerkom, E. Van den Abbeel, The Istanbul consensus workshop on embryo assessment: proceedings of an expert meeting, *Human Reproduction*. 26 (2011) 1270–1283. <https://doi.org/10.1093/humrep/deg489>.
- [36] D.K. Gardner, M. Lane, J. Stevens, T. Schlenker, W.B. Schoolcraft, Blastocyst score affects implantation and pregnancy outcome: towards a single blastocyst transfer, *Fertility and Sterility*. 73 (2000) 1155–1158. [https://doi.org/10.1016/S0015-0282\(00\)00518-5](https://doi.org/10.1016/S0015-0282(00)00518-5).
- [37] Q.-Y. Du, E.-Y. Wang, Y. Huang, X.-Y. Guo, Y.-J. Xiong, Y.-P. Yu, G.-D. Yao, S.-L. Shi, Y.-P. Sun, Blastocoele expansion degree predicts live birth after single blastocyst transfer for fresh and vitrified/warmed single blastocyst transfer cycles, *Fertility and Sterility*. 105 (2016) 910–919.e1. <https://doi.org/10.1016/j.fertnstert.2015.12.014>.
- [38] D. Feil, R.C. Henshaw, M. Lane, Day 4 embryo selection is equal to Day 5 using a new embryo scoring system validated in single embryo transfers, *Human Reproduction*. 23 (2008) 1505–1510. <https://doi.org/10.1093/humrep/dem419>.
- [39] C. Lagalla, M. Barberi, G. Orlando, R. Sciajno, M.A. Bonu, A. Borini, A quantitative approach to blastocyst quality evaluation: morphometric analysis and related IVF outcomes, *J Assist Reprod Genet*. 32 (2015) 705–712. <https://doi.org/10.1007/s10815-015-0469-3>.
- [40] C.L. Bormann, M.K. Kanakasabapathy, P. Thirumalaraju, R. Gupta, R. Pooniwala, H. Kandula, E. Hariton, I. Souter, I. Dimitriadis, L.B. Ramirez, C.L. Curchoe, J. Swain, L.M. Boehnlein, H. Shafiee, Performance of a deep learning based neural network in

- the selection of human blastocysts for implantation, *ELife*. 9 (2020) e55301.
<https://doi.org/10.7554/eLife.55301>.
- [41] Y. Kan-Tor, N. Zabari, I. Erlich, A. Szeskin, T. Amitai, D. Richter, Y. Or, Z. Shoham, A. Hurwitz, I. Har-Vardi, M. Gavish, A. Ben-Meir, A. Buxboim, Automated Evaluation of Human Embryo Blastulation and Implantation Potential using Deep-Learning, *Advanced Intelligent Systems*. 2 (2020) 2000080.
<https://doi.org/10.1002/aisy.202000080>.
- [42] P. Khosravi, E. Kazemi, Q. Zhan, J.E. Malmsten, M. Toschi, P. Zisimopoulos, A. Sigaras, S. Lavery, L.A.D. Cooper, C. Hickman, M. Meseguer, Z. Rosenwaks, O. Elemento, N. Zaninovic, I. Hajirasouliha, Deep learning enables robust assessment and selection of human blastocysts after in vitro fertilization, *Npj Digit. Med*. 2 (2019) 21.
<https://doi.org/10.1038/s41746-019-0096-y>.
- [43] C. Collinet, T. Lecuit, Programmed and self-organized flow of information during morphogenesis, *Nat Rev Mol Cell Biol*. (2021). <https://doi.org/10.1038/s41580-020-00318-6>.
- [44] E. Hannezo, C.-P. Heisenberg, Mechanochemical Feedback Loops in Development and Disease, *Cell*. 178 (2019) 12–25. <https://doi.org/10.1016/j.cell.2019.05.052>.
- [45] E.H. Barriga, K. Franze, G. Charras, R. Mayor, Tissue stiffening coordinates morphogenesis by triggering collective cell migration in vivo, *Nature*. (2018) 1–19.
<https://doi.org/10.1038/nature25742>.
- [46] C. Bertet, L. Sulak, T. Lecuit, Myosin-dependent junction remodelling controls planar cell intercalation and axis elongation, *Nature*. 429 (2004) 667–671.
<https://doi.org/10.1038/nature02590> nature02590 [pii].
- [47] M. Krieg, Y. Arboleda-Estudillo, P. Puech, J. Kafer, F. Graner, D. Muller, C. Heisenberg, Tensile forces govern germ-layer organization in zebrafish, *Nature Cell Biology*. 10 (2008) 429–436. <https://doi.org/ncb1705> [pii]10.1038/ncb1705.
- [48] E. Munro, J. Nance, J. Priess, Cortical flows powered by asymmetrical contraction transport PAR proteins to establish and maintain anterior-posterior polarity in the early *C. elegans* embryo, *Dev Cell*. 7 (2004) 413–424.
<https://doi.org/10.1016/j.devcel.2004.08.001>.
- [49] A.G. Clark, O. Wartlick, G. Salbreux, E.K. Paluch, Stresses at the cell surface during animal cell morphogenesis., *Curr Biol*. 24 (2014) R484–94.
<https://doi.org/10.1016/j.cub.2014.03.059>.
- [50] M. Kelkar, P. Bohec, G. Charras, Mechanics of the cellular actin cortex: From signalling to shape change, *Current Opinion in Cell Biology*. 66 (2020) 69–78.
<https://doi.org/10.1016/j.ceb.2020.05.008>.
- [51] M. Murrell, M. Lenz, M.L. Gardel, Forcing cells into shape: the mechanics of actomyosin contractility, *Nat Rev Mol Cell Biol*. 16 (2015) 486–498.
<https://doi.org/10.1038/nrm4012>.
- [52] G. Charras, A.S. Yap, Tensile Forces and Mechanotransduction at Cell–Cell Junctions, *Current Biology*. 28 (2018) R445–R457. <https://doi.org/10.1016/j.cub.2018.02.003>.
- [53] P.-F. Lenne, J.-F. Rupprecht, V. Viasnoff, Cell Junction Mechanics beyond the Bounds of Adhesion and Tension, *Developmental Cell*. 56 (2021) 202–212.
<https://doi.org/10.1016/j.devcel.2020.12.018>.
- [54] J.-L. Maître, C.-P. Heisenberg, Three Functions of Cadherins in Cell Adhesion, *Current Biology*. 23 (2013) R626–R633. <https://doi.org/10.1016/j.cub.2013.06.019>.
- [55] C.J. Chan, T. Hiiragi, Integration of luminal pressure and signalling in tissue self-organization, *Development*. 147 (2020) dev181297.
<https://doi.org/10.1242/dev.181297>.

- [56] M.F. Schliffka, J.-L. Maître, Stay hydrated: basolateral fluids shaping tissues., *Curr Opin Genet Dev.* 57 (2019) 70–77. <https://doi.org/10.1016/j.gde.2019.06.015>.
- [57] L. Cao, A. Yonis, M. Vaghela, E.H. Barriga, P. Chugh, M.B. Smith, J. Maufront, G. Lavoie, A. Méant, E. Ferber, M. Bovellan, A. Alberts, A. Bertin, R. Mayor, E.K. Paluch, P.P. Roux, A. Jégou, G. Romet-Lemonne, G. Charras, SPIN90 associates with mDia1 and the Arp2/3 complex to regulate cortical actin organization, *Nat Cell Biol.* 22 (2020) 803–814. <https://doi.org/10.1038/s41556-020-0531-y>.
- [58] P. Chugh, A.G. Clark, M.B. Smith, D.A.D. Cassani, K. Dierkes, A. Ragab, P.P. Roux, G. Charras, G. Salbreux, E.K. Paluch, Actin cortex architecture regulates cell surface tension, *Nature Cell Biology.* 19 (2017) 689–697. <https://doi.org/10.1364/AO.41.000685>.
- [59] M. Bergert, S.D. Chandradoss, R.A. Desai, E. Paluch, Cell mechanics control rapid transitions between blebs and lamellipodia during migration., *Proceedings of the National Academy of Sciences of the United States of America.* 109 (2012) 14434–14439. <https://doi.org/10.1073/pnas.1207968109>.
- [60] G. Charras, C. Hu, M. Coughlin, T. Mitchison, Reassembly of contractile actin cortex in cell blebs, *The Journal of Cell Biology.* 175 (2006) 477–490.
- [61] O.M. Lancaster, M. Le Berre, A. Dimitracopoulos, D. Bonazzi, E. Zlotek-Zlotkiewicz, R. Picone, T. Duke, M. Piel, B. Baum, Mitotic rounding alters cell geometry to ensure efficient bipolar spindle formation., *Dev Cell.* 25 (2013) 270–283. <https://doi.org/10.1016/j.devcel.2013.03.014>.
- [62] J. Sedzinski, M. Biro, J.-Y. Tinevez, G. Salbreux, E. Paluch, Polar actomyosin contractility destabilizes the position of the cytokinetic furrow., *Nature.* 476 (2011) 462–466. <https://doi.org/10.1038/nature10286>.
- [63] N. Taneja, M.R. Bersi, S.M. Baillargeon, A.M. Fenix, J.A. Cooper, R. Ohi, V. Gama, W.D. Merryman, D.T. Burnette, Precise Tuning of Cortical Contractility Regulates Cell Shape during Cytokinesis, *Cell Reports.* 31 (2020) 107477. <https://doi.org/10.1016/j.celrep.2020.03.041>.
- [64] Y.-S. Chu, W.A. Thomas, O. Eder, F. Pincet, E. Perez, J.-P. Thiery, S. Dufour, Force measurements in E-cadherin-mediated cell doublets reveal rapid adhesion strengthened by actin cytoskeleton remodeling through Rac and Cdc42., *The Journal of Cell Biology.* 167 (2004) 1183–1194. <https://doi.org/10.1083/jcb.200403043>.
- [65] A.C. Martin, M. Gelbart, R. Fernandez-Gonzalez, M. Kaschube, E.F. Wieschaus, Integration of contractile forces during tissue invagination, *The Journal of Cell Biology.* 188 (2010) 735–749. <https://doi.org/10.1083/jcb.200910099>.
- [66] W. Engl, B. Arasi, L.L. Yap, J.P. Thiery, V. Viasnoff, Actin dynamics modulate mechanosensitive immobilization of E-cadherin at adherens junctions, *Nature Cell Biology.* 16 (2014) 587–594. <https://doi.org/10.1038/ncb2973>.
- [67] J.-L. Maitre, H. Berthoumieux, S.F.G. Krens, G. Salbreux, F. Julicher, E. Paluch, C.-P. Heisenberg, Adhesion Functions in Cell Sorting by Mechanically Coupling the Cortices of Adhering Cells, *Science.* 338 (2012) 253–256. <https://doi.org/10.1126/science.1225399>.
- [68] R.E. Stephenson, T. Higashi, I.S. Erofeev, T.R. Arnold, Rho Flares Repair Local Tight Junction Leaks, *Developmental Cell.* (2019) 1–21. <https://doi.org/10.1016/j.devcel.2019.01.016>.
- [69] C. Zihni, K. Matter, Tight junctions: from simple barriersto multifunctional molecular gates, *Nat Rev Mol Cell Biol.* 17 (2016) 564–580. <https://doi.org/10.1038/nrm.2016.80>.
- [70] S. Sigurbjörnsdóttir, R. Mathew, M. Leptin, Molecular mechanisms of de novo lumen formation, *Nat Rev Mol Cell Biol.* 15 (2014) 665–676. <https://doi.org/10.1038/nrm3871>.

- [71] H.N. Ciray, A. Campbell, I.E. Agerholm, J. Aguilar, S. Chamayou, M. Esbert, S. Sayed, Time-Lapse User Group, Proposed guidelines on the nomenclature and annotation of dynamic human embryo monitoring by a time-lapse user group., *Human Reproduction* (Oxford, England). 29 (2014) 2650–2660. <https://doi.org/10.1093/humrep/deu278>.
- [72] C. Lagalla, N. Tarozzi, R. Sciajno, D. Wells, M. Di Santo, M. Nadalini, V. Distratis, A. Borini, Embryos with morphokinetic abnormalities may develop into euploid blastocysts, *Reproductive BioMedicine Online*. 34 (2017) 137–146. <https://doi.org/10.1016/j.rbmo.2016.11.008>.
- [73] S.L. Chavez, K.E. Loewke, J. Han, F. Moussavi, P. Colls, S. Munne, B. Behr, R.A. Reijo Pera, Dynamic blastomere behaviour reflects human embryo ploidy by the four-cell stage, *Nat Commun*. 3 (2012) 1251. <https://doi.org/10.1038/ncomms2249>.
- [74] E. Ford, C.E. Currie, D.M. Taylor, M. Erent, A.L. Marston, G.M. Hartshorne, A.D. McAinsh, The First Mitotic Division of the Human Embryo is Highly Error-prone, *Cell Biology*, 2020. <https://doi.org/10.1101/2020.07.17.208744>.
- [75] T.H.H. Coorens, T.R.W. Oliver, R. Sanghvi, U. Sovio, E. Cook, R. Vento-Tormo, M. Haniffa, M.D. Young, R. Rahbari, N. Sebire, P.J. Campbell, D.S. Charnock-Jones, G.C.S. Smith, S. Behjati, Inherent mosaicism and extensive mutation of human placentas, *Nature*. (2021). <https://doi.org/10.1038/s41586-021-03345-1>.
- [76] M.N. Shahbazi, T. Wang, X. Tao, B.A.T. Weatherbee, L. Sun, Y. Zhan, L. Keller, G.D. Smith, A. Pellicer, R.T. Scott, E. Seli, M. Zernicka-Goetz, Developmental potential of aneuploid human embryos cultured beyond implantation, *Nat Commun*. 11 (2020) 3987. <https://doi.org/10.1038/s41467-020-17764-7>.
- [77] T. Hardarson, Human embryos with unevenly sized blastomeres have lower pregnancy and implantation rates: indications for aneuploidy and multinucleation, *Human Reproduction*. 16 (2001) 313–318. <https://doi.org/10.1093/humrep/16.2.313>.
- [78] E.V. Royen, Multinucleation in cleavage stage embryos, *Human Reproduction*. 18 (2003) 1062–1069. <https://doi.org/10.1093/humrep/deg201>.
- [79] L. Scott, A. Finn, T. O’Leary, S. McLellan, J. Hill, Morphologic parameters of early cleavage-stage embryos that correlate with fetal development and delivery: prospective and applied data for increased pregnancy rates, *Human Reproduction*. 22 (2007) 230–240. <https://doi.org/10.1093/humrep/del358>.
- [80] Y. Liu, V. Chapple, P. Roberts, P. Matson, Prevalence, consequence, and significance of reverse cleavage by human embryos viewed with the use of the Embryoscope time-lapse video system, *Fertility and Sterility*. 102 (2014) 1295-1300.e2. <https://doi.org/10.1016/j.fertnstert.2014.07.1235>.
- [81] B. Mierzwa, D.W. Gerlich, Cytokinetic Abscission: Molecular Mechanisms and Temporal Control, *Developmental Cell*. 31 (2014) 525–538. <https://doi.org/10.1016/j.devcel.2014.11.006>.
- [82] I. Rubio, R. Kuhlmann, I. Agerholm, J. Kirk, J. Herrero, M.-J. Escribá, J. Bellver, M. Meseguer, Limited implantation success of direct-cleaved human zygotes: a time-lapse study, *Fertility and Sterility*. 98 (2012) 1458–1463. <https://doi.org/10.1016/j.fertnstert.2012.07.1135>.
- [83] N. Desai, J.M. Goldberg, C. Austin, T. Falcone, Are cleavage anomalies, multinucleation, or specific cell cycle kinetics observed with time-lapse imaging predictive of embryo developmental capacity or ploidy?, *Fertility and Sterility*. 109 (2018) 665–674. <https://doi.org/10.1016/j.fertnstert.2017.12.025>.
- [84] I. Kola, A. Trounson, G. Dawson, P. Rogers, Trippronuclear Human Oocytes: Altered Cleavage Patterns and Subsequent Karyotypic Analysis of Embryos, *Biology of Reproduction*. 37 (1987) 395–401. <https://doi.org/10.1095/biolreprod37.2.395>.

- [85] Q. Zhan, Z. Ye, R. Clarke, Z. Rosenwaks, N. Zaninovic, Direct Unequal Cleavages: Embryo Developmental Competence, Genetic Constitution and Clinical Outcome, *PLoS ONE*. 11 (2016) e0166398. <https://doi.org/10.1371/journal.pone.0166398>.
- [86] I. Halvaei, M.A. Khalili, N. Esfandiari, S. Safari, A.R. Talebi, S. Miglietta, S.A. Nottola, Ultrastructure of cytoplasmic fragments in human cleavage stage embryos, *J Assist Reprod Genet*. 33 (2016) 1677–1684. <https://doi.org/10.1007/s10815-016-0806-1>.
- [87] B.L. Daughtry, J.L. Rosenkrantz, N.H. Lazar, S.S. Fei, N. Redmayne, K.A. Torkenczy, A. Adey, M. Yan, L. Gao, B. Park, K.A. Nevenon, L. Carbone, S.L. Chavez, Single-cell sequencing of primate preimplantation embryos reveals chromosome elimination via cellular fragmentation and blastomere exclusion, *Genome Res*. 29 (2019) 367–382. <https://doi.org/10.1101/gr.239830.118>.
- [88] P. Eftekhari-Yazdi, M.R. Valojerdi, S.K. Ashtiani, M.B. Eslaminejad, L. Karimian, Effect of fragment removal on blastocyst formation and quality of human embryos, *Reproductive BioMedicine Online*. 13 (2006) 823–832. [https://doi.org/10.1016/S1472-6483\(10\)61031-0](https://doi.org/10.1016/S1472-6483(10)61031-0).
- [89] M.D. Keltz, J.C. Skorupski, K. Bradley, D. Stein, Predictors of embryo fragmentation and outcome after fragment removal in in vitro fertilization, *Fertility and Sterility*. 86 (2006) 321–324. <https://doi.org/10.1016/j.fertnstert.2006.01.048>.
- [90] M. Alikani, Epithelial cadherin distribution in abnormal human pre-implantation embryos., *Human Reproduction (Oxford, England)*. 20 (2005) 3369–3375. <https://doi.org/10.1093/humrep/dei242>.
- [91] V. Ruprecht, S. Wieser, A. Callan-Jones, M. Smutny, H. Morita, K. Sako, V. Barone, M. Ritsch-Marte, M. Sixt, R. Voituriez, C.-P. Heisenberg, Cortical Contractility Triggers a Stochastic Switch to Fast Amoeboid Cell Motility, *Cell*. 160 (2015) 673–685. <https://doi.org/10.1016/j.cell.2015.01.008>.
- [92] R. Derrick, C. Hickman, O. Oliana, T. Wilkinson, D. Gwinnett, L.B. Whyte, A. Carby, S. Lavery, Perivitelline threads associated with fragments in human cleavage stage embryos observed through time-lapse microscopy, *Reproductive BioMedicine Online*. 35 (2017) 640–645. <https://doi.org/10.1016/j.rbmo.2017.08.026>.
- [93] F. Hyafil, D. Morello, C. Babinet, F. Jacob, A cell surface glycoprotein involved in the compaction of embryonal carcinoma cells and cleavage stage embryos., *Cell*. 21 (1980) 927–934.
- [94] Y. Shirayoshi, T.S. Okada, M. Takeichi, The calcium-dependent cell-cell adhesion system regulates inner cell mass formation and cell surface polarization in early mouse development., *Cell*. 35 (1983) 631–638.
- [95] R.O. Stephenson, Y. Yamanaka, J. Rossant, Disorganized epithelial polarity and excess trophectoderm cell fate in preimplantation embryos lacking E-cadherin, *Development (Cambridge, England)*. 137 (2010) 3383–3391. <https://doi.org/10.1242/dev.050195>.
- [96] J.C.M. Dumoulin, M. Bras, E. Coonen, J. Dreesen, J.P.M. Geraedts, J.L.H. Evers, Effect of Ca²⁺/Mg²⁺-free medium on the biopsy procedure for preimplantation genetic diagnosis and further development of human embryos, *Human Reproduction*. 13 (1998) 2880–2883. <https://doi.org/10.1093/humrep/13.10.2880>.
- [97] ESHRE PGT Consortium and SIG-Embryology Biopsy Working Group, G. Kokkali, G. Coticchio, F. Bronet, C. Celebi, D. Cimadomo, V. Goossens, J. Liss, S. Nunes, I. Sfontouris, N. Vermeulen, E. Zakharova, M. De Rycke, ESHRE PGT Consortium and SIG Embryology good practice recommendations for polar body and embryo biopsy for PGT[†], *Human Reproduction Open*. 2020 (2020) hoaa020. <https://doi.org/10.1093/hropen/hoaa020>.

- 780 [98] E.E. Zakharova, V.V. Zaletova, A.S. Krivokharchenko, Biopsy of Human Morula-
781 Stage Embryos: Outcome of 215 IVF/ICSI Cycles with PGS, PLoS ONE. 9 (2014)
782 e106433. <https://doi.org/10.1371/journal.pone.0106433>.
- 783 [99] M.S. Steinberg, M. Takeichi, Experimental specification of cell sorting, tissue
784 spreading, and specific spatial patterning by quantitative differences in cadherin
785 expression., *Proc Natl Acad Sci U S A*. 91 (1994) 206–209.
- 786 [100] H.H. Yu, M.R. Dohn, N.O. Markham, R.J. Coffey, A.B. Reynolds, p120-catenin
787 controls contractility along the vertical axis of epithelial lateral membranes, *J Cell Sci*.
788 129 (2016) 80–94. <https://doi.org/10.1242/jcs.177550>.
- 789 [101] C. Skiadas, K. Jackson, C. Racowsky, Early compaction on day 3 may be associated
790 with increased implantation potential, *Fertility and Sterility*. 86 (2006) 1386–1391.
791 <https://doi.org/10.1016/j.fertnstert.2006.03.051>.
- 792 [102] M.F. Schliffka, A.-F. Tortorelli, Ö. Özgüç, L. de Plater, O. Polzer, D. Pelzer, J.-L.
793 Maître, Multiscale analysis of single and double maternal-zygotic Myh9 and Myh10
794 mutants during mouse preimplantation development, *Developmental Biology*, 2020.
795 <https://doi.org/10.1101/2020.09.10.291997>.
- 796 [103] Y. Mizobe, Y. Ezono, M. Tokunaga, N. Oya, R. Iwakiri, N. Yoshida, Y. Sato, N.
797 Onoue, K. Miyoshi, Selection of human blastocysts with a high implantation potential
798 based on timely compaction., *Journal of Assisted Reproduction and Genetics*. 34
799 (2017) 991–997. <https://doi.org/10.1007/s10815-017-0962-y>.
- 800 [104] C.M. Pauken, D.G. Capco, Regulation of cell adhesion during embryonic compaction
801 of mammalian embryos: roles for PKC and beta-catenin., *Mol Reprod Dev*. 54 (1999)
802 135–144. [https://doi.org/10.1002/\(SICI\)1098-2795\(199910\)54:2<135::AID-](https://doi.org/10.1002/(SICI)1098-2795(199910)54:2<135::AID-MRD5>3.0.CO;2-A)
803 [MRD5>3.0.CO;2-A](https://doi.org/10.1002/(SICI)1098-2795(199910)54:2<135::AID-MRD5>3.0.CO;2-A).
- 804 [105] G.K. Winkel, J.E. Ferguson, M. Takeichi, R. Nuccitelli, Activation of protein kinase C
805 triggers premature compaction in the four-cell stage mouse embryo., *Developmental*
806 *Biology*. 138 (1990) 1–15.
- 807 [106] M. Zhu, C.Y. Leung, M.N. Shahbazi, M. Zernicka-Goetz, Actomyosin polarisation
808 through PLC-PKC triggers symmetry breaking of the mouse embryo, *Nature*
809 *Communications*. (2017) 1–16. <https://doi.org/10.1038/s41467-017-00977-8>.
- 810 [107] S. Petropoulos, D. Edsgård, B. Reinius, Q. Deng, S.P. Panula, S. Codeluppi, A.P.
811 Reyes, S. Linnarsson, R. Sandberg, F. Lanner, Single-Cell RNA-Seq Reveals Lineage
812 and X Chromosome Dynamics in Human Preimplantation Embryos, *Cell*. 165 (2016)
813 1012–1026. <https://doi.org/10.1016/j.cell.2016.03.023>.
- 814 [108] N.S. Goel, C.F. Doggenweiler, R.L. Thompson, Simulation of cellular compaction and
815 internalization in mammalian embryo development as driven by minimization of
816 surface energy., *Bulletin of Mathematical Biology*. 48 (1986) 167–187.
- 817 [109] N. Dard, S. Louvet-Vallée, B. Maro, Orientation of mitotic spindles during the 8- to 16-
818 cell stage transition in mouse embryos., *PLoS ONE*. 4 (2009) e8171.
819 <https://doi.org/10.1371/journal.pone.0008171>.
- 820 [110] T. Watanabe, J.S. Biggins, N.B. Tannan, S. Srinivas, Limited predictive value of
821 blastomere angle of division in trophectoderm and inner cell mass specification,
822 *Development (Cambridge, England)*. 141 (2014) 2279–2288.
823 <https://doi.org/10.1242/dev.103267>.
- 824 [111] H.Y.G. Lim, Y.D. Alvarez, M. Gasnier, Y. Wang, P. Tetlak, S. Bissiere, H. Wang, M.
825 Biro, N. Plachta, Keratins are asymmetrically inherited fate determinants in the
826 mammalian embryo, *Nature*. (2020). <https://doi.org/10.1038/s41586-020-2647-4>.
- 827 [112] Y. Hirate, S. Hirahara, K. Inoue, A. Suzuki, V.B. Alarcon, K. Akimoto, T. Hirai, T.
828 Hara, M. Adachi, K. Chida, S. Ohno, Y. Marikawa, K. Nakao, A. Shimono, H. Sasaki,
829 Polarity-dependent distribution of angiomotin localizes Hippo signaling in

- preimplantation embryos., *Curr Biol.* 23 (2013) 1181–1194.
<https://doi.org/10.1016/j.cub.2013.05.014>.
- [113] E. Wicklow, S. Blij, T. Frum, Y. Hirate, R.A. Lang, H. Sasaki, A. Ralston, HIPPO Pathway Members Restrict SOX2 to the Inner Cell Mass Where It Promotes ICM Fates in the Mouse Blastocyst., *PLoS Genetics.* 10 (2014) e1004618.
<https://doi.org/10.1371/journal.pgen.1004618>.
- [114] M. Zhu, M.N. Shahbazi, A. Martin, C. Zhang, M. Borsos, R.S. Mandelbaum, R.J. Paulson, M. Esbert, R.T. Scott, A. Campbell, S. Fishel, V. Gradinaru, H. Zhao, K. Wu, E. Seli, Mechanism of cell polarisation and first lineage segregation in the human embryo, *BioRxiv.* (2020).
- [115] D. Clift, W.A. McEwan, L.I. Labzin, V. Konieczny, B. Mogessie, L.C. James, M. Schuh, A Method for the Acute and Rapid Degradation of Endogenous Proteins, *Cell.* 171 (2017) 1692–1695.e18. <https://doi.org/10.1016/j.cell.2017.10.033>.
- [116] K. Leonavicius, C. Preece, B. Davies, S. Srinivas, Mechanics of mouse blastocyst hatching revealed by a hydrogel-based microdeformation assay, *Proceedings of the National Academy of Sciences of the United States of America.* 115 (2018) 10375–10380. <https://doi.org/10.1016/j.ijnonlinmec.2004.05.006>.
- [117] L. Casares, R. Vincent, D. Zalvidea, N. Campillo, D. Navajas, M. Arroyo, X. Trepac, Hydraulic fracture during epithelial stretching, *Nature Materials.* 14 (2015) 343–351. <https://doi.org/10.1038/nmat4206>.
- [118] M.R. Ghassemifar, Gene expression regulating epithelial intercellular junction biogenesis during human blastocyst development in vitro, *Molecular Human Reproduction.* 9 (2003) 245–252. <https://doi.org/10.1093/molehr/gag033>.
- [119] R. Gualtieri, L. Santella, B. Dale, Tight junctions and cavitation in the human pre-embryo., *Mol Reprod Dev.* 32 (1992) 81–87. <https://doi.org/10.1002/mrd.1080320113>.
- [120] F.D. Houghton, P.G. Humpherson, J.A. Hawkhead, C.J. Hall, H.J. Leese, Na², K², ATPase activity in the human and bovine preimplantation embryo, *Developmental Biology.* (2003) 7.
- [121] Y. Xiong, Y.-J. Tan, Y.-M. Xiong, Y.-T. Huang, X.-L. Hu, Y.-C. Lu, Y.-H. Ye, T.-T. Wang, D. Zhang, F. Jin, H.-F. Huang, J.-Z. Sheng, Expression of Aquaporins in Human Embryos and Potential Role of AQP3 and AQP7 in Preimplantation Mouse Embryo Development, *Cell Physiol Biochem.* 31 (2013) 649–658.
<https://doi.org/10.1159/000350084>.
- [122] T.T. Huang, D.H. Huang, H.J. Ahn, C. Arnett, C.T. Huang, Early blastocyst expansion in euploid and aneuploid human embryos: evidence for a non-invasive and quantitative marker for embryo selection, *Reproductive BioMedicine Online.* 39 (2019) 27–39.
<https://doi.org/10.1016/j.rbmo.2019.01.010>.
- [123] T.H. Taylor, T. Stankewicz, S.L. Katz, J.L. Patrick, L. Johnson, D.K. Griffin, Preliminary assessment of aneuploidy rates between the polar, mid and mural trophoctoderm, *Zygote.* 28 (2020) 93–96. <https://doi.org/10.1017/S0967199419000637>.
- [124] M.L. Verge-Serandour, H. Turler, Physics of blastocoel formation by hydro-osmotic lumen coarsening, *Biophysics*, 2020. <https://doi.org/10.1101/2020.12.01.406991>.
- [125] D. Bodri, T. Sugimoto, J. Yao Serna, S. Kawachiya, R. Kato, T. Matsumoto, Blastocyst collapse is not an independent predictor of reduced live birth: a time-lapse study., *Fertility and Sterility.* 105 (2016) 1476–1483.e3.
<https://doi.org/10.1016/j.fertnstert.2016.02.014>.
- [126] J. Marcos, S. Pérez-Albalá, A. Mifsud, M. Molla, J. Landeras, M. Meseguer, Collapse of blastocysts is strongly related to lower implantation success: a time-lapse study, *Hum. Reprod.* 30 (2015) 2501–2508. <https://doi.org/10.1093/humrep/dev216>.

- [127] R. Sciorio, R. Herrer Saura, K.J. Thong, M. Esbert Algam, S.J. Pickering, M. Meseguer, Blastocyst collapse as an embryo marker of low implantation potential: a time-lapse multicentre study, *Zygote*. 28 (2020) 139–147. <https://doi.org/10.1017/S0967199419000819>.
- [128] L. Noli, A. Capalbo, C. Ogilvie, Y. Khalaf, D. Ilic, Discordant Growth of Monozygotic Twins Starts at the Blastocyst Stage: A Case Study, *Stem Cell Reports*. 5 (2015) 946–953. <https://doi.org/10.1016/j.stemcr.2015.10.006>.
- [129] L.Z. Yanez, J. Han, B.B. Behr, R.A.R. Pera, D.B. Camarillo, Human oocyte developmental potential is predicted by mechanical properties within hours after fertilization., *Nature Communications*. 7 (2016) 10809. <https://doi.org/10.1038/ncomms10809>.
- [130] C. De Geyter, C. Calhaz-Jorge, M.S. Kupka, C. Wyns, E. Mocanu, T. Motrenko, G. Scaravelli, J. Smeenk, S. Vidakovic, V. Goossens, The European IVF-monitoring Consortium (EIM) for the European Society of Human Reproduction and Embryology (ESHRE), O. Gliozheni, E. Hambartsoumian, H. Strohmer, E. Petrovskaya, O. Tishkevich, K. Bogaerts, C. Wyns, D. Balic, S. Sibincic, I. Antonova, M. Pelekanos, K. Rezabek, J. Markova, J. Lemmen, D. Söritsa, M. Gissler, S. Pelkonen, F. Pessione, J. de Mouzon, A. Tandler—Schneider, S. Kalantaridou, J. Urbancsek, G. Kosztolanyi, H. Bjorgvinsson, E. Mocanu, J. Cloherty, G. Scaravelli, R. de Luca, V. Lokshin, S. Karibayeva, V. Magomedova, R. Bausyte, I. Masliukaite, Z. Petanovski, J. Calleja-Agius, V. Moshin, T.M. Simic, D. Vukicevic, J.M.J. Smeenk, L.B. Romundstad, A. Janicka, C. Calhaz—Jorge, A.R. Laranjeira, I. Rugescu, B. Doroftei, V. Korsak, N. Radunovic, N. Tabs, I. Virant-Klun, I.C. Saiz, F.P. Mondéjar, C. Bergh, E. Berger-Menz, M. Weder, H. Ryan, R. Baranowski, M. Gryshchenko, ART in Europe, 2015: results generated from European registries by ESHRE†, *Human Reproduction Open*. 2020 (2020) hoz038. <https://doi.org/10.1093/hropen/hoz038>.
- [131] K. Guevorkian, Micropipette aspiration: A unique tool for exploring cell and tissue mechanics in vivo., *Methods in Cell Biology*. 139 (2017) 187–201. <https://doi.org/10.1016/bs.mcb.2016.11.012>.