**Phenological and elevational shifts of plants, animals and fungi
under climate change in the European Alps**

Yann Vitasse^{1,*}, Sylvain Ursenbacher^{2,3}, Geoffrey Klein^{1,4}, Thierry
Bohnenstengel², Yannick Chittaro², Anne Delestrade^{5,6}, Christian Monnerat²,
Martine Rebetez^{1,4}, Christian Rixen⁷, Nicolas Strebel⁸, Benedikt R. Schmidt^{2,9},
Sonja Wipf^{7,10}, Thomas Wohlgemuth¹, Nigel Gilles Yoccoz¹¹ and Jonathan
Lenoir¹²

¹WSL Swiss Federal Institute for Forest, Snow and Landscape Research, Zürcherstrasse 111,
CH-8903 Birmensdorf, Switzerland

²info fauna CSCF & karch, Avenue de Bellevaux 51, CH-2000 Neuchâtel, Switzerland

³Department of Environmental Sciences, Section of Conservation Biology, University of
Basel, St. Johannis-Vorstadt 10, CH-4056 Basel, Switzerland

⁴Institute of Geography, University of Neuchâtel, Espace Tilo-Frey 1, CH-2000 Neuchâtel,
Switzerland

⁵Centre de Recherches sur les Ecosystèmes d'Altitude, 67 Lacets du Belvédère, 74400
Chamonix Mont-Blanc, France

⁶Laboratoire d'Ecologie Alpine (LECA), CNRS, Université Grenoble Alpes, Université Savoie
Mont Blanc, Savoie Technolac, 73376 Le Bourget du Lac Cedex, France

⁷WSL Institute for Snow and Avalanche Research SLF, Group Mountain Ecosystems,
Flüelastrasse 11, CH-7260 Davos Dorf, Switzerland

⁸Schweizerische Vogelwarte, Seerose 1, CH-6204 Sempach, Switzerland

⁹*Institut für Evolutionsbiologie und Umweltwissenschaften, Universität Zürich,*

Winterthurerstrasse 190, 8057 Zürich, Switzerland

¹⁰*Swiss National Park, Chastè Planta-Wildenberg, Runatsch 124, 7530 Zernez, Switzerland*

¹¹*Department of Arctic and Marine Biology, Faculty of Biosciences, Fisheries and Economics,*

UiT The Arctic University of Norway, PO Box 6050 Langnes, N-9037 Tromsø, Norway

¹²*UMR CNRS 7058 « Ecologie et Dynamique des Systèmes Anthropisés » (EDYSAN),*

Université de Picardie Jules Verne, 1 Rue des Louvels, 80037 Amiens Cedex 1, France

31

*Author for correspondence (E-mail: yann.vitasse@wsl.ch; Tel.: ++41 (0)44 739 2073).

33

34 ABSTRACT

35 Mountain areas are biodiversity hotspots and provide a multitude of ecosystem services of

36 irreplaceable socio-economic value. In the European Alps, air temperature has increased at a

37 rate of about 0.36 °C decade⁻¹ since 1970, leading to glacier retreat and significant snowpack

38 reduction. Due to these rapid environmental changes, this mountainous region is undergoing

39 marked changes in spring phenology and elevational distribution of animals, plants and fungi.

40 Long-term monitoring in the European Alps offers an excellent natural laboratory to

41 synthesize climate-related changes in spring phenology and elevational distribution for a large

42 array of taxonomic groups. This review assesses the climatic changes that have occurred

43 across the European Alps during recent decades, spring phenological changes and upslope

44 shifts of plants, animals and fungi from evidence in published papers and previously

45 unpublished data. Our review provides evidence that spring phenology has been shifting

46 earlier during the past four decades and distribution ranges show an upwards trend for most of

47 the taxonomic groups for which there are sufficient data. The first observed activity of reptiles

48 and terrestrial insects (e.g. butterflies) in spring has shifted significantly earlier, at an average

1
2
3 49 rate of -5.7 and -6.0 days decade $^{-1}$, respectively. By contrast, the first observed spring activity
4
5 50 of semi-aquatic insects (e.g. dragonflies and damselflies) and amphibians, as well as the
6
7 51 singing activity or laying dates of resident birds, show smaller non-significant trends ranging
8
9 52 from -1.0 to $+1.3$ days decade $^{-1}$. Leaf-out and flowering of woody and herbaceous plants
10
11 53 showed intermediate trends with mean values of -2.4 and -2.8 days decade $^{-1}$, respectively.
12
13 54 Regarding species distribution, plants, animals and fungi ($N = 2,133$ species) shifted the
14
15 55 elevation of maximum abundance (optimum elevation) upslope at a similar pace (on average
16
17 56 between $+18$ and $+25$ m decade $^{-1}$) but with substantial differences among taxa. For example,
18
19 57 the optimum elevation shifted upward by $+36.2$ m decade $^{-1}$ for terrestrial insects and $+32.7$ m
20
21 58 decade $^{-1}$ for woody plants, whereas it was estimated to range between -1.0 and $+11$ m decade $^{-1}$
22
23 59 for semi-aquatic insects, ferns, birds and wood-decaying fungi. The upper range limit
24
25 60 (leading edge) of most species also shifted upslope with a rate clearly higher for animals (from
26
27 61 $+47$ to $+91$ m decade $^{-1}$) than for plants (from $+17$ to $+40$ m decade $^{-1}$), except for semi-aquatic
28
29 62 insects (-4.7 m decade $^{-1}$). Although regional land-use changes could partly explain some
30
31 63 trends, the consistent upward shift found in almost all taxa all over the Alps is likely reflecting
32
33 64 the strong warming and the receding of snow cover that has taken place across the European
34
35 65 Alps over recent decades. However, with the possible exception of terrestrial insects, the
36
37 66 upward shift of organisms seems currently too slow to track the pace of isotherm shifts
38
39 67 induced by climate warming, estimated at about $+62$ to $+71$ m decade $^{-1}$ since 1970. In the light
40
41 68 of these results, species interactions are likely to change over multiple trophic levels through
42
43 69 phenological and spatial mismatches. This nascent research field deserves greater attention to
44
45 70 allow us to anticipate structural and functional changes better at the ecosystem level.
46
47 71
48
49
50
51
52
53
54
55
56 72 *Key words:* biodiversity redistribution, climate change velocity, disequilibrium dynamics,
57
58 73 global warming, migration, mountain ecosystems, phenological mismatches, upslope shift.
59
60

74	
75	CONTENTS
76	I. Introduction
77	II. Methods
78	III. Synthesis
79	(1) Climate change in the European Alps
80	(a) Changes in air temperature, precipitation and snow cover
81	(i) Temperature
82	(ii) Precipitation and snow
83	(b) Predicted climate change in the European Alps
84	(2) Phenological shifts
85	(3) Elevational range shifts
86	(a) Terrestrial plants and fungi
87	(b) Terrestrial animals
88	(c) Freshwater ecosystems
89	(4) Plant–animal interactions mediated by species-specific phenological and spatial shifts
90	(5) Research gaps and ecosystem-based monitoring
91	IV. Conclusions
92	V. Acknowledgements, author contributions and data availability
93	VI. References
94	VII. Supporting information
95	
96	I. INTRODUCTION
97	Biodiversity loss and redistribution as well as extreme climatic events are probably the two
98	most prominent environmental challenges with strong socio-economic consequences that

99 humans have to face in the 21st century. Biodiversity loss, mainly caused by human alteration
100 of the landscape, has been occurring at an unprecedented rate since humans expanded across
101 the globe (Butchart *et al.*, 2010). The distributions of many terrestrial organisms are currently
102 shifting in response to global warming and land-use changes in most parts of the world,
103 poleward and upslope to cooler latitudes and elevations, respectively, thus generating spatially
104 structured patterns in species redistribution (Chen *et al.*, 2011; Lenoir & Svenning, 2015;
105 Lenoir *et al.*, 2020). However, the rate of ongoing global warming is such that numerous plant
106 and animal species may not be able to adapt genetically, or to track their climatic niche
107 requirements closely, ultimately leading to local extinctions (reviewed in Jump & Penuelas,
108 2005; Aitken *et al.*, 2008; Corlett & Westcott, 2013; Cang, Wilson & Wiens, 2016; Jezkova &
109 Wiens, 2016), particularly in terrestrial ecosystems as opposed to marine ecosystems (Lenoir
110 *et al.*, 2020).

111 Ongoing global warming is changing the geographic distribution of organisms on Earth
112 but their physiology and seasonal activities are also dramatically altered by climate change. In
113 fact, phenological changes, such as flowering or leaf senescence for plants, as well as mating,
114 breeding, spawning or hibernating for animals, are one of the most visible and sensitive
115 responses to ongoing global warming in natural ecosystems (Parmesan & Yohe, 2003; Cleland
116 *et al.*, 2007). For instance, in Europe, leaf-out of temperate trees has advanced by 8–15 days since
117 the 1950s (Fu *et al.*, 2019) and the migration dates of birds have advanced by about a week since
118 1959 in Canada and northern Europe (Lehikoinen *et al.*, 2019). In a temperate climate, spring
119 phenology reflects how the physiology and behaviour of organisms interacts with the
120 progressive return of favourable conditions for growth, i.e. mainly the increase in air
121 temperature and day length. Because animals are dependent on the development of plants, any
122 change in plant phenology may elicit a cascade of consequences in plant–animal interactions
123 (Beard *et al.*, 2019; Visser & Gienapp, 2019; Kharouba & Wolkovich, 2020; Rehnus, Peláez

& Bollmann, 2020). How the synchrony between animals and plant phenology will change under global warming is a challenging and timely question, but the consequences of such synchrony are still poorly understood at the population, community and ecosystem levels due to a lack of long-term ecosystem-based monitoring time series (but see Kharouba *et al.*, 2018; Visser & Gienapp, 2019; Peláez *et al.*, 2020).

The European Alps form an arc approximately 800 km in length from east to west and around 200 km in width in western Europe with numerous peaks rising over 4,000 m. This mountain system has been a prominent obstacle for plants and animals during the glacial and post-glacial periods of the Pleistocene, thus being at the crossroads of several biogeographical units (Taberlet *et al.*, 1998; Hewitt, 2000). The European Alps are a hotspot of biodiversity offering myriad habitats and climates over very short distances (Körner, 2003; Zimmermann *et al.*, 2013). About 13,000 vascular plant species (> 500 endemics) and about 30,000 animal species inhabit the European Alps (Agrawala, 2007). Plants, animals and fungi living in this region include generalist species found across all of Europe but also numerous specialist species that have developed adaptations to harsh environments at high elevations, and are found only in restricted areas such as mountain summits (i.e. sky islands). Angiosperm plant species are found up to 4,507 m (*Saxifraga oppositifolia* L.) together with fungi and arthropods (Körner, 2011), while bryophytes are found up to 4,559 m (Vaccari, 1914). In the European Alps, as in all mountainous areas, the distance that an organism has to move in order to cross isotherms is quite short compared to flatter regions, because of the sharp temperature adiabatic gradient induced by elevational gradients (Rolland, 2003). This may, to a certain extent, help species to track isotherm shifts better in mountainous regions (Brito-Morales *et al.*, 2018). However, species living close to mountain summits will progressively be limited in habitat availability because moving upward generally means less land surface available, higher habitat fragmentation, and an inevitable increase in competition among co-existing species,

1
2
3 149 eventually leading to local extinction events (Wiens, 2016; Giezendanner *et al.*, 2019). The
4
5 150 shift of species ranges in response to climate change is also increasingly hampered by human
6
7
8 151 activities such as tourism, urbanization or agriculture, all of which alter species' habitat
9
10 152 suitability and distribution by fragmenting natural habitats (Elsen, Monahan & Merenlender,
11
12 153 2020). A warmer climate associated with changes in rain regime and snow cover will alter the
13
14 154 physiology and phenology of many species, which in turn will affect their fitness and biotic
15
16
17 155 interactions. Numerous studies have shown tight relationships between snow conditions and
18
19 156 population dynamics of alpine and arctic plants (e.g. Inouye, 2000, 2008; Körner, 2003; Wipf
20
21 157 & Rixen, 2010; Berteaux *et al.*, 2016) as well as population dynamics of animal species (e.g.
22
23 158 Korslund & Steen, 2006; Helle & Kojola, 2008; Robinson & Merrill, 2012). Hence, any
24
25
26 159 changes in air temperature, precipitation and snow cover patterns in high mountain areas are
27
28 160 expected to elicit substantial modifications in plant–animal interactions. In the European Alps,
29
30 161 the pace of climate warming is occurring at a faster rate than the average warming in the
31
32 162 northern hemisphere (Rebetez & Reinhard, 2008), generating faster velocities of isotherm
33
34 163 shifts. As a result, substantial shifts in spring phenology as well as numerous elevational range
35
36 164 shifts have already been observed for plants, animals and even fungi over recent decades (see
37
38 165 studies in Tables 1 and 2). Long-term monitoring in the European Alps offers an excellent
39
40 166 natural laboratory to synthesize climate-related changes in spring phenology and elevational
41
42 167 distribution for a large array of taxonomic groups.

46
47 168 Here we aim at synthesizing the climatic changes that have occurred across the European
48
49 169 Alps during recent decades and gathering the observed biotic responses in terms of both spring
50
51 170 phenological changes (the best documented season in terms of phenological observations) and
52
53 171 species range shifts, reported for flora, fauna and fungi. We provide a comprehensive
54
55 172 synthesis of these changes and discuss the consequences of such phenological and spatial
56
57 173 shifts on species interactions across different trophic levels. Moreover, we point out
58
59
60

174 knowledge gaps that require more investigation in order to be able to predict future structural
175 and functional changes better at the ecosystem level under global warming.

176

177 II. METHODS

178 We searched the scientific literature for publications reporting either species elevational range
179 shifts or spring phenological trends over recent decades. For phenological changes, we
180 focused on the spring season because this is better documented in terms of phenological
181 observations compared to the autumn season (Gallinat, Primack & Wagner, 2015), allowing
182 for robust comparisons among taxa. We focused on any important phenological event such as
183 leaf emergence or flowering of plants, peaks of detectability for birds or observations of the
184 adult phase for amphibians and insects. For elevational range shifts, we focused on studies
185 reporting either shifts of the core distribution of a given species (hereafter ‘optimum’) or shifts
186 of the upper elevational limit (hereafter ‘leading edge’). We restricted our review to studies
187 relying on regular monitoring data covering at least 10 years across parts of the European Alps
188 (i.e. Austria, France, Germany, Italy, Lichtenstein, Slovenia and Switzerland) within the
189 period 1980–2020. For diachronic studies comparing historical and modern surveys, we used
190 only studies that conducted their modern survey(s) within the period 1980–2020 and where the
191 median year of the historical survey(s) was after 1950 [often called the ‘great acceleration’
192 period (Steffen *et al.*, 2015)] to avoid including decades with no or significantly lower
193 warming trends. Indeed, including studies using historical surveys prior to 1950 (e.g.
194 Holzinger *et al.*, 2008; Rumpf *et al.*, 2018) would inevitably underestimate the rate at which
195 biotic responses are taking place in response to ongoing climate change for recent decades.

196 We focused on the European Alps because this region is both a hotspot of biodiversity
197 and benefits from a long research history, allowing a long-term view and comprehensive
198 synthesis on climate change-related processes. We extended the study area to lowland regions

199 in direct contact with the mountain range of the European Alps or to neighbouring mountain
200 ranges (the Jura), provided that part of the European Alps was considered in the studied region
201 and assumed to play a significant role in the results (e.g. Kuhn *et al.*, 2016).

202 Raw data for phenological shift (per decade or per degree of warming) and migration
203 rate across elevations per unit time (per year or per decade) were extracted at the species level
204 from the publications reviewed herein, either from direct access *via* the journal webpage
205 (appendix or data depository), or indirectly by requesting species-level data from the study
206 authors. Depending on the study, different methods were used to generate species-level data
207 on phenological shift or elevational range shift per decade. For instance, some studies directly
208 reported the observed rates in phenological shift or elevational range shift per decade while
209 others reported the magnitude of the observed changes in phenology or elevational distribution
210 over a given time period. In the latter case we extracted information on the duration of the
211 time period covered by that study and computed the rates per decade. Phenological studies
212 reporting the phenological change per degree of warming generally include a pre-season
213 temperature window in their calculations, which is best correlated with the studied
214 phenological event, i.e. a couple of weeks or months before the spring phenological event. For
215 more details, readers should refer to the publications listed in Table 1.

216 All publications used for our synthesis are summarised in Tables 1 and 2, and the full
217 data are provided as online supporting information in Tables S1 and S2, respectively. For
218 more robust comparisons, taxonomic groups containing observations of phenological or range
219 shifts for less than five species were discarded from our quantitative analyses, but are still
220 included in the tables for completeness.

221 In addition, we analysed two sources of previously unpublished data collected through a
222 citizen science program in Switzerland: the *info fauna* database for both phenology and

223 altitudinal migration of insects, reptiles and amphibians and the *PhenoForest* database for
224 phenological trends of several tree species.

225 From the Swiss central database of *info fauna*, we extracted all observations for
226 amphibians, reptiles, lepidopterans, orthopterans, and odonates available between 1991 and
227 2019. Although data collected before 1990 are available, we restricted our analyses to after
228 1990 because far fewer observers were active before 1990, which could influence trend
229 analyses. Species with taxonomic uncertainty, with a limited number of observations per year
230 (variable depending on taxonomic group, left to the experts' discretion), that are migrants or
231 show other features that could impact the frequency of observations were also removed after
232 careful examination by expert co-authors of this review. To avoid a distorting influence of
233 extreme values when assessing elevational range shifts from the raw data, we computed, for
234 every year, the median (50%) and 95th percentile of the elevation as surrogates for the
235 optimum and leading edge, respectively, because these were less likely to be distorted than the
236 mean and maximum elevation values. Similarly, for spring phenology, we used the 5th
237 percentile of the distribution in the observations for a given species for every year. Linear
238 regressions were then conducted for all species of these parameters against years, and we
239 report the slope values expressed per decade. In total, for *info fauna*, 273 animal species were
240 included in our analyses, comprising 1,558,862 observations (average: 5,710
241 observations/species; median: 2,807 observations/species; 5–95% quantile: 256–21,301) for
242 the altitude analysis and 1,560,265 observations (average: 5,715 observations/species; median:
243 2,823 observations/species; 5–95% quantile: 196–20,380) for the phenology analysis. The *info*
244 *fauna* database gathers observations mainly from citizen volunteers and does not use a
245 systematic sampling design, rather representing haphazard sampling, which may introduce
246 statistical issues (Callaghan *et al.*, 2019). However, because we used only data from taxa with
247 sufficient sample sizes and that were carefully checked by experts before they were entered

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

248 into our database, we believe this data set should provide robust trends of how these species
249 are currently migrating along elevational gradients. Additionally, the data from *info fauna* do
250 not show identical trends among taxa; if a particular trend was due to observer bias, we would
251 expect to find similar patterns across various taxa, which was not the case.

252 The *PhenoForest* database consists of records of visual observations of the leaf-out
253 timing of eight species during the period 1998–2020 from volunteers in different sites in
254 Switzerland (one observer per site, 16 permanent sites in total, 1–5 species per site). At each
255 site, 8–10 trees per species were identified and monitored every year by the same observer. To
256 estimate trends in the leaf-out dates over this study period, we applied a linear mixed-effect
257 model for each species with leaf-out dates as the response variable, year as the explanatory
258 variable and sites as a random intercept term, and extracted the estimated slope for each
259 species separately.

260 Following collection/calculation of all the species-level biotic responses in days per
261 decade for shifts in spring phenology or meters per decade for elevational range shifts as
262 described above, we compared the estimated rates among taxonomic groups. We ran linear
263 mixed-effects models with species’ phenological shift per year or species’ elevational range
264 shift per decade as response variables, study/database as a random intercept term and
265 taxonomic group (with as many levels as taxonomic groups for which there were sufficient
266 data, i.e. groups containing at least five species) as the main fixed effect. Based on the model
267 estimates of phenological shift per year or elevational range shift per decade for each
268 taxonomic group separately, we computed the associated 95% confidence intervals. *Post-hoc*
269 Tukey tests were then applied to test for differences among the different taxonomic groups. In
270 the results, we report the mean shift of the phenology or migration of a given taxonomic group
271 across studies, using the estimated marginal mean value $\pm$ CI from the mixed-effect model. We

also report the mean $\pm$ SE phenological and elevational shifts per study/database and taxonomic group in Tables 1 and 2.

In order to visualize whether the rate of species migration lags behind the pace of climate warming we computed the likely isotherm shifts that have occurred in the Alps since 1970. We used the annual warming trends shown in Fig. 1 for six stations across the Swiss Alps of $0.36\text{ }^{\circ}\text{C decade}^{-1}$ (similar values are reported by the national offices of Meteorology and Climatology, e.g. MeteoSwiss). To calculate the corresponding theoretical altitudinal distance required to maintain the same air temperature, we applied an altitudinal lapse rate of $0.51\text{--}0.58\text{ }^{\circ}\text{C }100\text{ m}^{-1}$, as typically evaluated for annual mean temperature across different regions of the Alps [France, Germany, Austria, Italy and Switzerland (Rolland, 2003; Kirchner *et al.*, 2013)]. The isotherm shift induced by climate warming was estimated in this way to range between $+62$ and $+71\text{ m decade}^{-1}$ during the period 1970–2019.

All calculations and statistics were conducted using R 3.5.1 (R Core Team, 2017).

III. SYNTHESIS

(1) Climate change in the European Alps

(a) Changes in air temperature, precipitation and snow cover

(i) Temperature

Since the beginning of the 20th century, mean annual air temperature has increased by more than 1.5°C across the European Alps (Böhm *et al.*, 2001; Begert, Schlegel & Kirchhofer, 2005; Rebetez & Reinhard, 2008), with this trend accelerating strongly during the second half of the 20th century (Auer *et al.*, 2007; Acquaotta, Fratianni & Garzena, 2015) with an average increase of about $0.36\text{ }^{\circ}\text{C decade}^{-1}$ between 1970 and 2019 (Klein *et al.*, 2016, Fig. 1). Air temperature trends in the European Alps exceed the average trend observed over the Northern Hemisphere by 1.6–2.5 times (Keiler, Knight & Harrison, 2010; Foster & Rahmstorf, 2011).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The increase in air temperatures during the 20th century was most pronounced around the altitude of the 0 °C isotherm (Serquet, Marty & Rebetez, 2013), in connection with the snow-albedo feedback of snow-covered surfaces and glaciers (Pepin & Lundquist, 2008; Scherrer *et al.*, 2012). However, these trends hide strong seasonal disparities. During the 20th century in Western Europe, air temperatures warmed more in winter compared to summer (Moberg *et al.*, 2006), but this ratio reversed by the end of the 20th and early 21st century (Klein *et al.*, 2018). Hence, several studies have shown that summer and spring have warmed more than autumn and winter since the 1970s (Rebetez & Reinhard, 2008; Klein *et al.*, 2018; Vitasse *et al.*, 2018a). Since the end of the 1980s, maximum air temperatures have been increasing more than minimum temperatures in connection with the decrease in European air pollution and particulate matter, especially in spring and at mid and high elevations (Rebetez & Reinhard, 2008; Vitasse *et al.*, 2018a).

(ii) *Precipitation and snow*

Annual precipitation has not changed across the European Alps over recent decades (Diolaiuti *et al.*, 2012). Regionally, only a few increasing seasonal trends have been observed since the beginning of the 20th century, mostly during autumn and winter (Schmidli & Frei, 2005; Scherrer *et al.*, 2016). Winter precipitation sum has increased slightly, whereas trends towards more intense but less frequent precipitation were observed in summer with no change in overall precipitation sum (Moberg *et al.*, 2006; Appenzeller *et al.*, 2011; Fischer *et al.*, 2019). Recently, European alpine countries have experienced several extreme drought events in summer, associated with heat waves, which have led to forest dieback, vegetation damage and insect outbreaks in the lowlands, particularly in 2003, 2012, 2015, 2016 and 2018 (Corona-Lozada, Morin & Choler, 2019; Schuldt *et al.*, 2020). Intense hot drought events had also occurred during the 20th century (e.g. in 1911, 1921, 1947, 1949, 1976) but air temperature

was, on average, lower than today, so that the evapotranspiration in summertime was lower (Schär *et al.*, 2004) and consequently the vegetation was much less impacted by these events. Because air temperature decreases steadily with elevation (and generally precipitation increases), the climatic water balance of sites >1,200 m still remains positive in the central part of the European Alps (Vitasse *et al.*, 2019). However, severe droughts are predicted to become more frequent even at high elevations where organisms might be less adapted to cope with water limitation (Rosbakh *et al.*, 2017).

Snow cover has an important impact in plant and animal phenology in mountain ecosystems. A significant reduction in the number of days with snow precipitation has been reported at all elevations during the 20th century in the Alps, particularly since the 1980s and in relation to warmer air temperatures (Laternser & Schneebeli, 2003; Pellicciotti, Bauder & Parola, 2010; Valt & Cianfarra, 2010; Serquet *et al.*, 2011; Diolaiuti *et al.*, 2012; Marty & Blanchet, 2012; Serquet *et al.*, 2013). A similar reduction has been observed across the northern hemisphere since the 1970s (Hernández-Henríquez, Déry & Derksen, 2015), especially in spring (Brown & Mote, 2009; Brown & Robinson, 2011), although this trend varies widely at a global scale (Bormann *et al.*, 2018; Pulliainen *et al.*, 2020). Analyses of snow water equivalent have shown a clear decrease in snow mass all over the European Alps during the period 1973–2012, at all elevations and more pronounced in spring compared to winter (Marty, Tilg & Jonas, 2017b). Mean snow depth has also been significantly reduced during the same period in the European Alps at all elevations and for all seasons (Durand *et al.*, 2009; Schöner, Auer & Böhm, 2009; Pellicciotti *et al.*, 2010; Marty & Blanchet, 2012; Marty & Meister, 2012). Detailed analyses in the Swiss Alps have confirmed a general decrease in all snow parameters at all elevations, with a faster reduction of snow cover at the time of snowmelt compared to the time of snow onset (Klein *et al.*, 2016). Spring (March to May) was the season with the highest air temperature increase over the last five decades, with

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

a rate of 0.47 °C decade⁻¹ since 1970 in the Swiss Alps (Klein *et al.*, 2018, Fig. 1). This strong warming has led to earlier snowmelt, for example −5.2 days decade⁻¹ across six locations in the Swiss Alps ranging from 1,298 to 2,540 m between 1970 and 2019 (Fig. 1). In autumn (September to November), the air temperature warming rate has been less than in spring, although still sufficient to delay time of snow onset at an average rate of +3.9 days decade⁻¹ since 1970 for these six stations, leading to a significant increase in the snow-free period at an average rate of +9.1 days decade⁻¹ in the Swiss Alps (Fig. 1). However, regional differences across the European Alps exist with on average stronger trends of snow depth reduction in the southern Alps over recent decades (Matiu *et al.*, 2021).

(b) Predicted climate change in the European Alps

Air temperature will increase further in the coming decades if present emission rates of greenhouse gases are not substantially reduced (IPCC, 2013). The difference between predictions based on lower [representative concentration pathway (RCP) 2.6] or higher (RCP 8.5) greenhouse gas emission rates (IPCC, 2013) increases with time, particularly after 2050. Déqué *et al.* (2007) predicted that air temperature in the European Alps would increase by 0.30 to 0.45 °C decade⁻¹ until 2100, with a higher expected increase in summer and autumn and an increased frequency of summer heatwaves (Keiler *et al.*, 2010). Precipitations in summer are expected to decrease over coming decades, although with a higher frequency of extreme rainfall events (Rajczak & Schär, 2017; Hodnebrog *et al.*, 2019). Recent analyses nevertheless showed that regional climate models (RCM, from the EURO-Coordinated Regional Downscaling Experiment) could greatly underestimate the warming rate during the growing season (Schwingshackl *et al.*, 2019; Boé *et al.*, 2020). This was due to models omitting the plant physiological CO₂ response (Schwingshackl *et al.*, 2019) and underestimating both the increase in shortwave radiation and changes in solar radiation

connected with cloud cover and aerosols (Boé *et al.*, 2020). For the same reasons, regional models also tended to underestimate the decrease in summer precipitation (Boé *et al.*, 2020) as well as the increase in heavy rainfall events (Borodina, Fischer & Knutti, 2017).

A strong reduction of snow depth and snow cover is expected to continue at all elevations across the European Alps during the 21st century (Jasper *et al.*, 2004; Rousselot *et al.*, 2012; Schmucki *et al.*, 2015; Marty *et al.*, 2017a), particularly during the spring season (Magnusson *et al.*, 2010; Steger *et al.*, 2013). Permanent snow, glaciers and permafrost are expected to decline strongly or disappear, with an overall melt of the snow cover in summer (Magnusson *et al.*, 2010). At the end of the 21st century, continuous snow cover in winter may only be present above 2,000 m, while mid-elevations (1,000–1,700 m) should experience random and/or discontinuous snow cover in about one in two winters (Schmucki *et al.*, 2017). These future expected changes in snow cover duration and thickness correspond to an average upward elevation shift of 800 m of the snow cover by the end of the century, while snow cover seasons may be reduced by two or three months, depending on the scenario (Bavay, Grünewald & Lehning, 2013; Marty *et al.*, 2017a; Beniston *et al.*, 2018).

Due to these rapid changes in air temperature and snowpack, the European Alps will undergo marked changes in the phenology and distribution of plants, animals and fungi, which are synthesized in the following sections.

(2) Phenological shifts

Our review reveals an advance in spring phenological events for most of the studied taxa, i.e. leaf-out and flowering for plants, singing activity, laying dates, arrival and hatching dates for birds, and the first yearly observation for insects, amphibians and reptiles although these were statistically significant only for reptiles and insects. There were also significant differences among these taxonomic groups (Fig. 2A). Reptiles and terrestrial flying insects such as

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Lepidoptera and Orthoptera showed the largest advance in their first yearly observation, with a significantly earlier -5.7 ± 3.8 days decade⁻¹ for reptiles (estimated marginal mean $\pm$ CI; linear mixed-effects model) and -6.0 ± 1.0 days decade⁻¹ for terrestrial insects, whereas a non-significant trend towards a slight delay was found for the peak of singing activity, laying dates or hatching dates of resident birds ($+1.3 \pm 4.5$ days decade⁻¹) and first yearly observation of amphibians ($+0.2 \pm 3.6$ days decade⁻¹; Fig. 2A). Insects with aquatic life stages (Odonata) showed non-significant trends for an earlier first yearly observation of the adult stage (-1.0 ± 2.2 days decade⁻¹, Fig. 2A). Plants also showed non-significant trends for an advance in leaf-out or flowering of -2.8 ± 6.3 and -2.4 ± 2.7 days decade⁻¹ for herbaceous and woody plants, respectively (Fig. 2A). A significant difference was found between semi-aquatic and terrestrial insects (Table 1; Fig. 2A). Whereas resident birds showed a tendency for a slight delay in their singing activity or laying dates ($+1.3 \pm 4.5$ days decade⁻¹), an advance was detected for long- and short-distance migrant birds (-1.7 ± 3.1 and -3.2 ± 3.9 days decade⁻¹, respectively; Fig. 2A), although these differences did not reach statistical significance. When grouped by habitat type, a non-significant trend for a delay in singing activity or laying dates was found for birds inhabiting forest or open land ($+0.5 \pm 6.9$ and $+0.6 \pm 5.9$, respectively), whereas a significantly different advance of about -7.5 ± 7.8 days decade⁻¹ was found for birds living in mixed habitats (Fig. S1).

Regarding phenological sensitivity to air temperature, fewer studies were available for some taxa, potentially making it harder to detect statistically significant differences among taxa (Table 1, Fig. 2B). However, we also found significant advances in spring phenology per degree of warming of -5.1 ± 3.9 days °C⁻¹ for woody plants and -4.9 ± 2.8 days °C⁻¹ for terrestrial insects, while non-significant trends for an advance were found in the other taxa.

For plants, phenological events are largely mediated by temperature and have the potential to change considerably with climate warming. In alpine environments, the few long-

term time series of phenological observations show the same consistent trends, for instance between the Rocky Mountains (USA), Greenland (Iler *et al.*, 2013) and the Tibetan plateau (Chen *et al.*, 2015). For the European Alps, we are not aware of any long-term studies of the phenology of alpine plants. However, indirect data for the beginning of vegetation growth detected by ultrasonic sensors (used to detect snow depth) are now available for about 20 years across 30 alpine sites in the Swiss Alps (Vitasse *et al.*, 2017). These data reveal that vegetation onset in alpine plants occurred earlier at around -6.3 days $^{\circ}\text{C}^{-1}$, with no strong photoperiodic constraint, in agreement with findings from the Tibetan plateau (Chen *et al.*, 2015). At lower elevations in the Swiss Alps, long-term phenological observations are available from a citizen network. These data show a significant advance of leaf-out or flowering dates during the period 1970–2012 by about -2.8 to -7.7 days $^{\circ}\text{C}^{-1}$ for various tree and shrub species and by about -3.7 to -7.4 days $^{\circ}\text{C}^{-1}$ for herbaceous species (Güsewell, 2014). It is known that spring phenological plasticity to temperature varies strongly among tree species, which could alter competition among species in response to global warming. For example, leaf-out of European beech (*Fagus sylvatica*) is one of the least sensitive to air temperature change, with an advance of less than -3 days $^{\circ}\text{C}^{-1}$, whereas ash (*Fraxinus excelsior*), sessile oak (*Quercus petraea*) and larch (*Larix decidua*) are among the most sensitive at more than -6 days $^{\circ}\text{C}^{-1}$ (Migliavacca *et al.*, 2008; Vitasse *et al.*, 2013; Asse *et al.*, 2018; Bison *et al.*, 2019).

For animals, phenological changes are less documented and overall less pronounced for large herbivores and birds that can migrate over long distances to track food availability and quality (i.e. tracking the ‘Green Wave’, e.g. Bischof *et al.*, 2012; Rehnus *et al.*, 2020). In the lowland areas of the Swiss Alps (< 600 m), the date of highest singing activity of various birds (mostly passerines), a proxy for the start of breeding, shifted earlier for the majority of species, and higher spring temperatures generally led to an earlier singing activity peak of around -1.1 days decade $^{-1}$ for resident and short-distance migrant birds with slightly higher trends for

trans-Saharan migrants (Strebel *et al.*, 2014; Table 1). Not surprisingly, insects seem to respond strongly to warming with the emergence date occurring earlier, for example by -21 days decade $^{-1}$ for bark beetles (*Ips typographus*) over the period 1996–2005 in the south-eastern Alps (Faccoli, 2009; Table 1). Similarly, butterflies seem to respond strongly to warming, with the purple emperor (*Apatura iris*) emerging earlier (-10.5 days decade $^{-1}$) during the period 1982–2002 in northern Switzerland (Dell, Sparks & Dennis, 2005; Table 1). Altermatt (2012) reported a temperature sensitivity of -4.9 ± 2.0 days $^{\circ}\text{C}^{-1}$ for 28 butterfly species studied in the Swiss lowlands over the period 1998–2010 (Table 1). Interestingly, our review shows that within insects, species from Orthoptera and Lepidoptera have advanced their spring phenology to a larger extent than species from Odonata (Fig. 2A). This might be explained by their habitat use, because species of Odonata generally spend their larval stages in ponds. The microclimate of ponds may buffer the effects of warming temperatures, especially in spring due to the provision of water from the melting snowpack higher up, although more data will be necessary to allow robust comparisons of the phenological sensitivity of aquatic species to air temperature.

(3) Elevational range shifts

Based on the current scientific literature focusing exclusively or partly on the European Alps, animals, fungi and plants have all shifted their optimum elevation upslope, reaching average rates of $+18.4 \pm 12.1$ ($N = 535$), $+18.8 \pm 26.8$ ($N = 118$) and $+24.8 \pm 13.6$ m decade $^{-1}$ ($N = 1,480$), respectively (mean $\pm$ CI; Fig. 3A). All taxonomic groups, except semi-aquatic insects, showed trends for a positive shift in optimum elevation (Fig. 3A), although this was statistically significant only for terrestrial insects ($+36.2 \pm 20.7$ m decade $^{-1}$), woody plants ($+32.7 \pm 17.9$ m decade $^{-1}$) and herbaceous plants (23.1 ± 14.4 m decade $^{-1}$). The average upward shift obtained from *info fauna* data regarding the optimum elevation of butterflies ($+39.9$ m

decade⁻¹, Table 2) agrees well with the trend reported in a recent study in Switzerland (+38 m between 2003 and 2010, or 48 m decade⁻¹) that systematically monitored butterfly communities (Roth, Plattner & Amrhein, 2014), suggesting that no major bias was introduced by the haphazard sampling inherent in *info fauna* data. By contrast, ferns, birds and wood-decaying fungi showed small and non-significant changes in optimum elevation, ranging between +4 and +11 m decade⁻¹ (Fig. 3A).

For the leading edge (upper elevational range limit), upward shifts were most pronounced for animals (on average +72.3±17.8 m decade⁻¹) than for plants (+28.2±21.8 m decade⁻¹; Fig. 3B), with the exception of semi-aquatic insect species that showed a non-significant trend for the leading edge to shift downward (−4.7±33.8 m decade⁻¹; Fig. 3B and Table 2). Remarkably, only terrestrial insects (+90.5±14.0 m decade⁻¹) and reptiles (63.7±57.2 m decade⁻¹) showed an upward shift of their leading edge within the range or higher than the pace of climate warming (Fig. 3B), estimated at ~62–71 m decade⁻¹ based on a warming rate of 0.36 °C decade⁻¹ (Fig. 1) and a temperature lapse rate of 0.51–0.58 m 100m⁻¹ (Rolland, 2003; Kirchner *et al.*, 2013).

(a) Terrestrial plants and fungi

For plants, our review suggests that there may be substantial differences among the studied taxonomic or functional groups, although these did not reach statistical significance, with insignificant shifts in the optimal elevation of +12.0±26.9 and 4.2±33.7 m decade⁻¹ for ferns and alpine plants and significant upward shifts of +23.1±14.4 and 32.7±17.9 m decade⁻¹ for herbaceous and woody species growing below the treeline (Fig. 3A). Notably, this general pattern among the taxonomic groups is the same for changes at the leading edge (Fig. 3B). Lenoir *et al.* (2008) showed for different mountain areas throughout France, including the western Alps, that the optimum elevation of 171 forest plants and trees moved upward in

1
2
3 497 elevation at a mean rate of +29 m decade⁻¹ when comparing vegetation surveys conducted
4
5 498 between 1905 and 1985 to surveys conducted between 1986 and 2005. However, when
6
7 499 splitting the data between short-lived herbaceous species (e.g. grasses and sedges) and long-
8
9 500 lived woody species (trees and shrubs), Lenoir *et al.* (2008) found that only herbaceous plants
10
11 501 from the forest understorey have significantly shifted their elevational optimum upslope (mean
12
13 502 rate of +38 m decade⁻¹). Similarly, by revisiting historical relevés and species indicator values,
14
15 503 Küchler *et al.* (2015) found an upward shift of forest plant species in the herbaceous layer of
16
17 504 Swiss forests at +10 m decade⁻¹ since the 1950s, while the elevational range of species from
18
19 505 the shrub and tree layers remained rather stable over the same time period. By contrast, our
20
21 506 quantitative review across the European Alps suggests that the optimum range of long-lived
22
23 507 woody plants is shifting upward at a significant rate, at least at the same rate as non-alpine
24
25 508 herbaceous plants (Fig. 3A). This discrepancy may stem from the fact that we did not
26
27 509 constrain our review only to forest systems, unlike the above-mentioned studies which
28
29 510 specifically focused on range shifts of forest plants (Lenoir *et al.*, 2008; Küchler *et al.*, 2015).
30
31 511 Thus, by including data from studies on forest systems with those reporting range shifts for
32
33 512 woody plant encroachment beyond the treeline or following land abandonment below the
34
35 513 treeline, we have a greater opportunity to detect elevational range shifts for woody plants. By
36
37 514 missing the upper part of the elevational gradient beyond the forest limit, studies restricted to
38
39 515 forest systems may have underestimated the true rate at which woody plant species are
40
41 516 responding to temperature increases or abandonment of land in the Alps. The dynamics of
42
43 517 recruitment of forest tree species at the upper elevational limits seems to be changing at a
44
45 518 relatively fast rate, with seedlings and saplings currently occurring well beyond the upper
46
47 519 elevational limit of adult trees (Vitasse *et al.*, 2012). Similarly, across several mountain areas
48
49 520 in western Europe, including the western Alps, Lenoir *et al.* (2009) reported differences in
50
51 521 elevational distribution between seedlings and adult life stages of trees, suggesting an ongoing
52
53
54
55
56
57
58
59
60

upward shift for woody plants but with quite high heterogeneity among species which could be due to species-specific variability in demographics and competitiveness (Scherrer *et al.*, 2020).

The average rate at which alpine plant species shifted upslope appears to be lower than for non-alpine plants or woody species below the treeline for both the optimum range and the upper elevational limits (Fig. 3), although note that these values did not differ statistically significantly. This slower rate at which alpine plants may be shifting upslope could be due to lower dispersal ability, biotic interactions or possibly to the methods employed to assess such trends. For alpine plants, resurveys are often compared with historical surveys conducted before 1950, i.e. during a period with lower warming rates than at present, but these studies were not included in our analysis, making this unlikely to explain the lower rate of upslope migration for alpine plants.

While climate warming plays an undeniable role in the recent redistribution of plants in mountain ecosystems (Steinbauer *et al.*, 2018), it might not be the only driver behind the upslope shifts along elevational gradients: precipitation (Rapacciuolo *et al.*, 2014), land use (Gehrig-Fasel, Guisan & Zimmermann, 2007; Guo, Lenoir & Bonebrake, 2018) or ongoing recolonization after the ‘Little Ice Age’ (Kammer, Schöb & Choler, 2007) might be additional drivers, either accelerating or slowing down the rate of elevational range shifts. For example, the treeline ecotone has been influenced by farming activities for several thousand years in the Alps (Tinner & Theurillat, 2003), and a large part of this recent upward shift has been attributed to substantial changes in land use, including: (i) land abandonment in the Swiss and Austrian Alps (Gehrig-Fasel *et al.*, 2007; Gellrich *et al.*, 2007; Tasser *et al.*, 2007), which seems to explain the large upward shift of *Abies alba* observed since the 1950s in the west central Alps (Chauchard *et al.*, 2010), and (ii) a reduction of cattle grazing, especially in the Italian Alps (Piotti *et al.*, 2007). By excluding open habitats in their analyses, Bodin *et al.*

(2013) showed that the upward shift they initially found for understorey forest plants became insignificant, a result that may be related to forest microclimate dynamics potentially buffering the effects of warming for understorey plants compared with plant communities from more open habitats (von Arx *et al.*, 2013; De Frenne *et al.*, 2019; Zellweger *et al.*, 2020). In some studies, a considerable part of the analysed herbaceous species may also shift downward (Lenoir *et al.*, 2010; Bodin *et al.*, 2013), potentially explained by more intensive management (disturbance) at lower elevations compared to forests at higher elevations (Lenoir *et al.*, 2010).

Interestingly, fungi may also be shifting their elevation. Although data in the Alps originated from a single study, that study assessed the shift in optimum elevation of 118 species (Diez *et al.*, 2020). They found values ranging between +10.7 and +23.6 m decade⁻¹ for wood-decaying and soil-dwelling fungi with an intermediate value for ectomycorrhizal fungi (+21.0 m decade⁻¹) (Table 2, Fig. 3A). A slower response of wood-decaying fungi would be consistent with their dependence on wood resources and particularly dead wood, with the availability of woody plants likely delayed compared to the rate of climate change.

(b) Terrestrial animals

Except for birds, elevational range shifts of animal species in the Alps have been less documented than for plants, with direct evidence available only for some groups. Our review reveals that terrestrial insects have moved significantly upslope in terms of both the optimum and the leading edge (Fig. 3). In an analysis of four orders of insects, a significant upslope movement is seen for each of the groups Lepidoptera, Coleoptera and to a lesser extent Orthoptera in both measures of elevational range (Fig. 4). For semi-aquatic insects from the group Odonata, neither range measure has changed significantly (Fig. 4). Insects, like most ectotherms, are very sensitive to changes in air temperature and consequently are likely to track climate warming closely. Accordingly, numerous upward range shifts have been

documented for terrestrial insects, especially for pests which receive much more attention. For example, upward shifts in the elevational range of outbreaks of the European spruce bark beetle (*I. typographus*) were related to the increased frequency of drier summers in the Italian Alps (Marini *et al.*, 2012). The pine processionary moth (*Thaumetopoea pityocampa*) shifted its distribution upward between 1975 and 2004 at average rates of +70 m decade⁻¹ and +29 m decade⁻¹ on the southern and northern slopes of the Italian Alps, respectively, with this shift related to the increase in minimum winter air temperature (Battisti *et al.*, 2005). As ectotherms, insects do not regulate their body temperature, and thus climatic variation in temperature will have a direct effect on their physiology. For example, warmer air temperatures in summer generally reduce European bark beetle development time (i.e. increased voltinism) (Jakoby, Lischke & Wermelinger, 2019), and therefore are likely to increase infestation pressure (Marini *et al.*, 2012; Jakoby *et al.*, 2019). The reproductive cycle of insects is expected to be faster under warmer air temperatures, potentially allowing an extra generation per growing season at all elevations by the end of the century for European spruce bark beetle (Jakoby *et al.*, 2019). Even though molluscs have a relatively low capacity to move over large distances, an upward shift of the land snail *Arianta arbustorum* of about 164 m has been reported in a comparison of its distribution in 1916–1917 with its distribution in 2011–2012 in the eastern Alps, and this upward shift was most pronounced (around 233 m) on south-facing slopes (Baur & Baur, 2013). Both habitat and microclimate are important factors for these organisms and the authors stressed that, on some mountain slopes, snails have already reached natural barriers (vertical rock walls with no soil), that prevent further upward shifts.

The mean elevation of reptiles has also changed considerably showing a non-significant shift in optimum elevation of +24.2±25.4 m decade⁻¹ (Fig. 3A) and a significant shift in their leading edge elevation of +63.7±22.3 m decade⁻¹; Fig. 3B). Nevertheless, more investigation

597 should be conducted for this group as this result originates from a single network of citizen
 598 observations in the Alps. An increase in air temperature will allow reptiles to colonise higher
 599 elevations, although the extent to which this is possible may be limited by oxygen availability
 600 (Jacobsen, 2020).

601 Interestingly, we found no significant differences in the upwards shift of optimum
 602 elevation between plants, fungi and animals (Fig. 3A), whereas animals appear to be
 603 responding fastest in terms of the leading edge, especially for terrestrial insects (Fig. 3B). The
 604 distribution of mammals and birds is mainly driven by habitat and food availability, and a
 605 change in temperature may not affect their distribution directly if features of their habitat
 606 remain stable or depend more on the micro- than the macroclimate (Jähnig *et al.*, 2020) and on
 607 plant phenology (Peláez *et al.*, 2020). For instance, two studies conducted on European roe
 608 deer (*Capreolus capreolus*) identified a strong dependence on the quality of food resources,
 609 with individuals performing long-distance seasonal movements upward and downward to
 610 locate food resources within their home range and thus tracking plant phenology (Gaudry *et al.*,
 611 2015; Peláez *et al.*, 2020). Another study revealed an upward shift of about $+37.5 \pm 10.3$ m
 612 decade⁻¹ of four mountain ungulate species in the Swiss Alps based on hunting records,
 613 attributing this upward shift to the lengthening of the snow-free period (Büntgen *et al.*, 2017,
 614 Table 2).

615 The effects of climate warming on animal distribution are likely to be complex, and will
 616 include indirect effects. A good example is the Alpine chough (*Pyrrhocorax graculus*) that
 617 nests at high elevations in cavities and fissures on inaccessible rock faces and forages in open
 618 alpine meadows (Delestrade & Stoyanov, 1995). Climate and land use are affecting the
 619 distribution of plants and forests but obviously not the distribution of the cliffs it uses for
 620 nesting. While upward shifts in distribution of birds have been detected in some areas in the
 621 Alps such as the central Swiss Alps (Maggini *et al.*, 2011; Knaus, 2018), several other studies

documented only marginal changes or no change in spite of significant air temperature increases during the study period (Archaux, 2004; Popy, Bordignon & Prodon, 2010). Consistently, our results revealed rather low and insignificant rates of optimum elevational range shift for birds in general, on average $+10.3 \pm 17.5$ m decade⁻¹, but Table 2 shows that there was much variation in the calculated values, with a trend for higher values in the Eastern and Southern Alps. This may be partly a result of human-induced land-use changes (Knaus, 2018) as in these regions a strong upward shift of the treeline and reforestation of formerly cut or unforested areas has taken place, moving up the habitat of specialized alpine birds such as the rock ptarmigan (*Lagopus muta*) (Pernollet, Korner-Nievergelt & Jenni, 2015). Given their ecological role as consumers, it is likely that responses of bird communities to climate change will lag behind that of some other taxonomic groups (Devictor *et al.*, 2012). It is possible that most specific elevational range shifts in mobile species like birds are related to site-specific factors associated with food availability, while for forest plants (Bertrand, Gegout & Bontemps, 2011) biotic interactions and population dynamics are likely to be more important (Scherrer *et al.*, 2020).

(c) *Freshwater ecosystems*

For freshwater ecosystems, only a few studies are available worldwide (Table 2), with several of these from the Alps. Comte & Grenouillet (2013) reported, across different mountain ranges in France including the western Alps, that the elevational range of stream fishes shifted upslope at a mean rate of about $+13.1 \pm 40.8$ m decade⁻¹ during the period 1986–2006 (Fig. 3A). For amphibians, dragonflies and damselflies, which depend directly on lakes, ponds, bogs, fens, streams and streamlets for their habitats, no significant upward shift was found in their optimum or leading edge elevation. The slower migration rates of semi-aquatic (e.g. dragonflies) and aquatic (e.g. fishes) animals compared to terrestrial animals or plants might

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

647 be due to their strong dependence on freshwater habitats which are heavily fragmented by
648 human activities (e.g. dams) and are also likely to exert strong microclimate buffering on
649 warming rate, especially within mountain ranges where streams are fed by snow and glacier
650 melt (McLaughlin *et al.*, 2017). In addition, most of these organisms are dependent on the
651 vegetation that structures their habitat. Typically, dragonflies inhabiting subalpine and alpine
652 zones live in bogs, fens and their associated small water bodies and runoff. The vegetation of
653 these habitats is very specific and species requirements in terms of vegetation structure are not
654 met in the pioneer water bodies of the upper level. The slower migration rate of these species
655 could therefore partially be explained by their requirement for colonization by vegetation to
656 occur first to offer new potential habitats for them. The formation of new bogs, fens or
657 swamps may take several decades or even centuries.

659 **(4) Plant–animal interactions mediated by species-specific phenological and spatial shifts**

660 On the one hand, because plants are the first link of the food web, changes in plant phenology
661 and composition will affect food quantity and quality for higher trophic levels, and may thus
662 affect complex trophic interactions (Plard *et al.*, 2014; Thackeray *et al.*, 2016; Kharouba *et al.*,
663 2018; Renner & Zohner, 2018; Kharouba & Wolkovich, 2020), animal performance (Sheridan
664 & Bickford, 2011) and species migration (Büntgen *et al.*, 2017; Peláez *et al.*, 2020; Rehnus *et*
665 *al.*, 2020). Conversely, numerous seed plants are pollinated and dispersed by animals, and are
666 affected by their interactions with herbivores, thus animal range shifts will be intimately
667 involved with plant range shifts. Responses of herbivores to climatic changes are generally
668 mediated through plant quality and quantity (Myserud *et al.*, 2001, 2008; Peláez *et al.*, 2020;
669 Rehnus *et al.*, 2020). In temperate regions, vegetation quantity gradually increases in spring to
670 reach a maximum in the middle of summer, while vegetation quality is generally highest
671 shortly after the beginning of vegetation growth in spring. In mountain ecosystems, mobile

organisms can track these changes by migrating upslope, following the ‘green wave’ induced by the progress of the season (Bischof *et al.*, 2012; Büntgen *et al.*, 2017; Herfindal *et al.*, 2019). Migratory bird species have also been observed to follow this peak in nutrient availability during spring migration (Van der Graaf *et al.*, 2006). Our review found that birds showed relatively small phenological and spatial changes in response to warming, perhaps because birds are vagile and thus can adapt easily to changes in food availability. Similarly, the parturition date of the European roe deer was shown to correlate better with spring plant phenology than with elevation in Switzerland (Peláez *et al.*, 2020). By contrast, species that are less mobile may become desynchronized in relation to their food resources as these move upwards in response to warming.

An example of in which such desynchronization has had a positive biotic effect for trees is in the frequency of outbreaks of the larch budmoth (*Zeiraphera griseana*), which historically occurred over 12 centuries every 8–9 years but disappeared after 1982 in the Engadine (eastern Swiss Alps). This was suggested to be due to a mismatch between the insect life cycle and larch budbreak (Esper *et al.*, 2007), which might have depended on climate warming (Johnson *et al.*, 2010). A recent outbreak did take place in 2017–2018, which was correlated with cold winter temperatures that reduced energy consumption during diapause in the moths, allowing them to emerge more in synchrony with the new foliage (Büntgen *et al.*, 2020). Warmer winters may continue to decrease the fitness of this insect in the future and upward elevational shifts of outbreak epicentres are expected.

On the contrary, many trees may experience more frequent severe defoliation due to pest outbreaks because warmer temperatures are expected to accelerate the life cycles of insects, potentially increasing the number of generations per season (Colombari *et al.*, 2012; Jakoby *et al.*, 2019). For example, the warmer and drier conditions that occurred in the late 1980s in the Italian prealps reduced the mortality of the web-spinning sawfly (*Cephalcia arvensis*) and

697 accelerated its development, causing severe defoliation of Norway spruce (*Picea abies*)
 698 needles on which the larvae feed (Marchisio, Cescatti & Battisti, 1994). Similarly, pine shoot
 699 beetles (*Tomicus piniperda*) and other bark beetles are likely to weaken Scots pine (*Pinus*
 700 *sylvestris*) and Norway spruce in the inner valleys of the European Alps, where they already
 701 suffer from drought, under warmer climates that will permit additional insect generations
 702 during the growth season (Rigling & Cherubini, 1999; Rebetez & Dobbertin, 2004; Jakoby *et*
 703 *al.*, 2019; Schneider, Comte & Rebetez, 2021). Such differences in phenological sensitivity to
 704 temperature are therefore expected to modify food-web interactions under climate change
 705 (Vitasse *et al.*, 2011; Roberts *et al.*, 2015; Asse *et al.*, 2018). Phenological differences between
 706 low- and high-elevation trees have reduced considerably over recent decades in the Swiss Alps
 707 due to a stronger phenological advance at higher elevations (Vitasse, Signarbieux & Fu,
 708 2018b), putting these plants at higher risk of frost exposure (Vitasse *et al.*, 2018a) and
 709 reducing the total period of availability of high-quality food for herbivores that follow the
 710 ‘green wave’ (Mysterud *et al.*, 2008).

711 In alpine environments, the life cycles of many plant and animal species are reliant on
 712 the date of snowmelt and the rapid greening that follows. For example, variation in the
 713 snowpack accounts for a substantial part of variation in survival (about 16%) and reproduction
 714 (29%) of alpine marmots (*Marmota marmota*) (Tafani *et al.*, 2013). Snow cover phenology is
 715 the main driver of foraging habitat selection for the white-winged snowfinch (*Montifringilla*
 716 *nivalis*) during the breeding season (Resano-Mayor *et al.*, 2019). Earlier snowmelt leads to
 717 earlier vegetation onset which in turn increases the reproductive success of the rock ptarmigan
 718 (*Lagopus muta*) (Novoa *et al.*, 2008), but decreases the litter size of alpine marmots (Tafani *et*
 719 *al.*, 2013). Similarly, earlier vegetation onset due to warming can increase survival of ibex
 720 (*Capra ibex*) through earlier access to food resources in spring (Pettorelli *et al.*, 2007), but a
 721 reduction of the snowpack in winter can also reduce the quality of the vegetation available in

722 spring (Robinson & Merrill, 2012).

723 The above examples of species interactions that are affected by phenological synchrony
724 represent direct effects on a few specific taxonomic groups; indirect and more complex species
725 interactions involving several trophic levels (network analyses) still await detailed
726 investigation. How the synchrony between winter climatic conditions, insect diapause, bird
727 activity and spring vegetation is affected will determine the ability of many species to adapt to
728 climate change. This nascent research field should receive more attention to allow us to
729 anticipate structural changes across multiple levels of ecosystems (Kharouba *et al.*, 2018;
730 Visser & Gienapp, 2019).

731

732 **(5) Research gaps and ecosystem-based monitoring**

733 Alpine ecosystems and the factors influencing species distributions have a long research
734 history in the European Alps (Braun-Blanquet, 1954; Offner & Le Brun, 1956), for example
735 with common garden experiments at different altitudes being performed in the Mont-Blanc
736 range in the late 19th century (Bonnier, 1888). We now have much data on elevational and
737 phenological shifts for several taxa, and a better understanding of the factors influencing
738 phenology and species distribution. This review provides evidence that recent climate change
739 may be affecting a wide range of taxonomic groups. However, our review also emphasizes
740 major gaps in our knowledge, especially regarding biotic responses to climate change, and in
741 particular, to a lack of long-term and ecosystem-based monitoring that could allow us to
742 disentangle the direct and indirect effects of climate and land-use changes on different trophic
743 levels (Yoccoz, Delestrade & Loison, 2010; Ims & Yoccoz, 2017).

744 Long-term ecosystem research (LTER) programs exist in the Alps, but mostly with a
745 focus on vegetation and soil, often in connection to ecosystem services and landscape patterns
746 (Tappeiner, Borsdorf & Bahn, 2013; Lavorel *et al.*, 2017; Rogora *et al.*, 2018). They do not

1
2
3 747 tend to integrate other ecosystem components such as herbivores, granivores and carnivores
4
5 748 from groups as diverse as insects, birds and mammals. There is, however, evidence that
6
7 749 changes in the abundance and distribution of species may be related to ecological interactions
8
9
10 750 [e.g. the mismatch between larch leaf-out and budmoth emergence (Büntgen *et al.*, 2009;
11
12 751 Johnson *et al.*, 2010)], or involve multiple trophic levels (Burgess *et al.*, 2018). The data
13
14 752 available for different taxonomic groups often come from different time periods or places. The
15
16
17 753 absence of a common design makes it difficult to build causal models integrating direct and
18
19 754 indirect effects [e.g. Grace & Irvine, 2020; but see Thuiller *et al.* (2018) for a species
20
21 755 distribution modelling approach to plant–herbivore interactions]. Our analysis identified a
22
23 756 large heterogeneity in species responses to climate change, making it difficult to understand
24
25
26 757 and predict changes at the ecosystem level (Mouquet *et al.*, 2015). Although citizen science
27
28 758 has allowed the collection of a large amount of data from haphazard observations, there is a
29
30 759 clear need for more systematic monitoring. Joint analysis of these two sources of data would
31
32
33 760 allow unbiased inferences (Dorazio, 2014). Developing integrated monitoring programs
34
35 761 focusing on different trophic levels, and combining these with underlying conceptual models
36
37 762 of the effects of climate and land-use changes on ecosystem structure and functioning should
38
39
40 763 be a priority.
41
42 764

43
44 765 **IV. CONCLUSIONS**

46
47 766 (1) In the European Alps, spring phenology has advanced over recent decades for all taxa,
48
49 767 except birds and amphibians, with these advances being statistically significant for
50
51 768 ectothermic animals such as terrestrial insects and reptiles.
52
53
54 769 (2) We found a trend for almost all studied taxa to shift their distribution range upslope, with
55
56 770 significant changes in optimum elevation found for terrestrial insects, herbaceous and woody
57
58
59
60

771 plants and in leading edge elevation for terrestrial insects, reptiles, birds and herbaceous
772 plants.

773 (3) We suggest that the trend for upward shift likely reflects the strong warming and the
774 receding of snow cover that has occurred across the European Alps over recent decades and
775 reflects the tracking by these taxa of their climatic niche, although regional land-use changes
776 may also provide an explanation in some cases.

777 (4) With the possible exception of some groups of terrestrial insects, such as butterflies, the
778 trend for an upward shift in plants, fungi and animals appears currently too slow to track
779 isotherm shifts induced by climate warming. Indeed, an upward shift of about 330 m would be
780 necessary to match the air temperature rise of close to 1.8°C that has occurred since 1970 in
781 the European Alps region (i.e. 0.36 °C per decade). This roughly corresponds to a mean shift
782 of elevational range of about +62–71 m decade⁻¹, assuming an annual adiabatic lapse rate of
783 0.51–0.58 °C 100 m⁻¹ as measured across the Alps (Rolland, 2003; Kirchner *et al.*, 2013). The
784 mean upward shift of most of the species groups observed over the last decades clearly lags
785 behind this value (between –2 and +34 m decade⁻¹ for the optimum elevation and between –5
786 and +64 m decade⁻¹ for the upper elevational limit), except for terrestrial insects (+36.2 and
787 +90.5 m decade⁻¹ for the optimum and upper elevational limits, respectively).

788 (5) Species interactions mediated by phenological and spatial shifts are likely to be altered by
789 climate warming but very few studies currently focus on biotic interactions across multiple
790 trophic levels. Integrating monitoring programs at ecosystem levels should be fostered to
791 allow better predictions of changes in plant and animal communities in response to climate
792 change.

793 (6) This review provides a valuable synthesis for future research focusing on the responses of
794 organisms to climate warming in the European Alps.

795

V. ACKNOWLEDGMENTS, AUTHOR CONTRIBUTIONS AND DATA

AVAILABILITY

We are grateful to all volunteers who provided observations for the following databases used in this study: *MeteoSwiss*, *PhenoForest*, *Vogelwarte*, and *info fauna*. We are grateful to J. Diez for providing raw data and to A. Cooper for her careful revision of the manuscript.

Author contributions: Y.V. and M.R. conceptualized and planned the work. Y.V., J.L. and S.U. collected the data. Y.V. and J.L. synthesized the data. S.U. analysed the data from the *info fauna* database and Y.V. from the *PhenoForest* database. M.R. wrote most of Section III.1 with contributions from G.K., who also created Fig. 1. Y.V. and J.L. wrote most of the text with substantial inputs from all co-authors.

Data availability: all raw data at species-level resolution used in the quantitative analyses are provided in the online supporting information.

VI. REFERENCES

ACQUAOTTA, F., FRATIANNI, S. & GARZENA, D. (2015). Temperature changes in the North-Western Italian Alps from 1961 to 2010. *Theoretical and Applied Climatology* **122**(3-4), 619–634.

AGRAWALA, S. (2007). Climate change in the European Alps: adapting winter tourism and natural hazards management. Organisation for Economic Cooperation and Development (OECD).

AITKEN, S. N., YEAMAN, S., HOLLIDAY, J. A., WANG, T. & CURTIS-MCLANE, S. (2008). Adaptation, migration or extirpation: climate change outcomes for tree populations. *Evolutionary Applications* **1**(1), 95–111.

ALTERMATT, F. (2012). Temperature-related shifts in butterfly phenology depend on the habitat. *Global Change Biology* **18**(8), 2429–2438.

APPENZELLER, C., BEY, I., CROCI MASPOLI, M., FUHRER, J., KNUTTI, R., KULL, C. & SCHÄR, C. (2011). Swiss climate change scenarios CH2011. ETH Zurich.

ARCHAUX, F. (2004). Breeding upwards when climate is becoming warmer: no bird response in the French Alps. *Ibis* **146**(1), 138–144.

ASSE, D., CHUINE, I., VITASSE, Y., YOCCOZ, N. G., DELPIERRE, N., BADEAU, V., DELESTRADE, A. & RANDIN, C. F. (2018). Warmer winters reduce the advance of tree spring phenology induced by warmer springs in the Alps. *Agricultural and Forest Meteorology* **252**, 220–230.

- 826 AUER, I., BÖHM, R., JURKOVIC, A., LIPA, W., ORLIK, A., POTZMANN, R., SCHÖNER, W., UNGERSBÖCK,
827 M., MATULLA, C. & BRIFFA, K. (2007). HISTALP—historical instrumental climatological
828 surface time series of the Greater Alpine Region. *International journal of climatology* **27**(1),
829 17–46.
- 830 BANI, L., LUPPI, M., ROCCHIA, E., DONDINA, O. & ORIOLI, V. (2019). Winners and losers: How the
831 elevational range of breeding birds on Alps has varied over the past four decades due to
832 climate and habitat changes. *Ecology and Evolution* **9**(3), 1289–1305.
- 833 BATTISTI, A., STASTNY, M., NETHERER, S., ROBINET, C., SCHOPF, A., ROQUES, A. & LARSSON, S.
834 (2005). Expansion of geographic range in the pine processionary moth caused by increased
835 winter temperatures. *Ecological Applications* **15**(6), 2084–2096.
- 836 BAUER, H.-G., HEINE, G., SCHMITZ, D., SEGELBACHER, G. & WERNER, S. (2019). Starke
837 Bestandsveränderungen der Brutvogelwelt des Bodenseegebietes—Ergebnisse aus vier
838 flächendeckenden Brutvogelkartierungen in drei Jahrzehnten. *Vogelwelt* **139**, 3–29.
- 839 BAUR, B. & BAUR, A. (2013). Snails keep the pace: shift in upper elevation limit on mountain slopes as
840 a response to climate warming. *Canadian Journal of Zoology* **91**(8), 596–599.
- 841 BAVAY, M., GRÜNEWALD, T. & LEHNING, M. (2013). Response of snow cover and runoff to climate
842 change in high Alpine catchments of Eastern Switzerland. *Advances in water resources* **55**, 4–
843 16.
- 844 BEARD, K. H., KELSEY, K. C., LEFFLER, A. J. & WELKER, J. M. (2019). The missing angle: ecosystem
845 consequences of phenological mismatch. *Trends in Ecology & Evolution* **34**(10), 885–888.
- 846 BEGERT, M., SCHLEGEL, T. & KIRCHHOFFER, W. (2005). Homogeneous temperature and precipitation
847 series of Switzerland from 1864 to 2000. *International Journal of Climatology* **25**(1), 65–80.
- 848 BENISTON, M., FARINOTTI, D., STOFFEL, M., ANDREASSEN, L. M., COPPOLA, E., ECKERT, N., FANTINI,
849 A., GIACONA, F., HAUCK, C. & HUSS, M. (2018). The European mountain cryosphere: a review
850 of its current state, trends, and future challenges. *The Cryosphere* **12**(2), 759–794.
- 851 BERTEAUX, D., GAUTHIER, G., DOMINE, F., IMS, R. A., LAMOUREUX, S. F., LÉVESQUE, E. & YOCCOZ,
852 N. (2016). Effects of changing permafrost and snow conditions on tundra wildlife: critical
853 places and times. *Arctic Science* **3**(2), 65–90.
- 854 BERTRAND, R., GEGOUT, J.-C. & BONTEMPS, J.-D. (2011). Niches of temperate tree species converge
855 towards nutrient-richer conditions over ontogeny. *Oikos* **120**(10), 1479–1488.
- 856 BISCHOF, R., LOE, L. E., MEISINGSET, E. L., ZIMMERMANN, B., VAN MOORTER, B. & MYSTERUD, A.
857 (2012). A migratory northern ungulate in the pursuit of spring: jumping or surfing the green
858 wave? *The American Naturalist* **180**(4), 407–424.
- 859 BISON, M., YOCCOZ, N. G., CARLSON, B. Z. & DELESTRADE, A. (2019). Comparison of budburst
860 phenology trends and precision among participants in a citizen science program. *International*
861 *Journal of Biometeorology* **63**(1), 61–72.

- 862 BODIN, J., BADEAU, V., BRUNO, E., CLUZEAU, C., MOISSELIN, J. M., WALTHER, G. R. & DUPOUEY, J.
 863 L. (2013). Shifts of forest species along an elevational gradient in Southeast France: climate
 864 change or stand maturation? *Journal of Vegetation Science* **24**(2), 269–283.
- 865 BOÉ, J., SOMOT, S., CORRE, L. & NABAT, P. (2020). Large discrepancies in summer climate change
 866 over Europe as projected by global and regional climate models: causes and consequences.
 867 *Climate Dynamics* **54**(5), 2981–3002.
- 868 BÖHM, R., AUER, I., BRUNETTI, M., MAUGERI, M., NANNI, T. & SCHÖNER, W. (2001). Regional
 869 temperature variability in the European Alps: 1760–1998 from homogenized instrumental time
 870 series. *International Journal of Climatology* **21**(14), 1779–1801.
- 871 BONNIER, M. G. (1888). Étude expérimentale de l'influence du climat alpin sur la végétation et les
 872 fonctions des plantes. *Bulletin de la Société botanique de France* **35**(8), 436–439.
- 873 BORMANN, K. J., BROWN, R. D., DERKSEN, C. & PAINTER, T. H. (2018). Estimating snow-cover trends
 874 from space. *Nature Climate Change* **8**(11), 924–928.
- 875 BORODINA, A., FISCHER, E. M. & KNUTTI, R. (2017). Models are likely to underestimate increase in
 876 heavy rainfall in the extratropical regions with high rainfall intensity. *Geophysical Research*
 877 *Letters* **44**(14), 7401–7409.
- 878 BRAUN-BLANQUET, J. (1954). La végétation alpine et nivale des Alpes françaises. Imprimerie
 879 Bayeusaine.
- 880 BRITO-MORALES, I., MOLINOS, J. G., SCHOEMAN, D. S., BURROWS, M. T., POLOCZANSKA, E. S.,
 881 BROWN, C. J., FERRIER, S., HARWOOD, T. D., KLEIN, C. J. & McDONALD-MADDEN, E. (2018).
 882 Climate Velocity Can Inform Conservation in a Warming World. *Trends in Ecology &*
 883 *Evolution*.
- 884 BROWN, R. D. & MOTE, P. W. (2009). The response of Northern Hemisphere snow cover to a changing
 885 climate. *Journal of Climate* **22**(8), 2124–2145.
- 886 BROWN, R. D. & ROBINSON, D. A. (2011). Northern Hemisphere spring snow cover variability and
 887 change over 1922–2010 including an assessment of uncertainty. *The Cryosphere* **5**(1), 219–
 888 229.
- 889 BÜNTGEN, U., FRANK, D., LIEBHOLD, A., JOHNSON, D., CARRER, M., URBINATI, C., GRABNER, M.,
 890 NICOLUSSI, K., LEVANIC, T. & ESPER, J. (2009). Three centuries of insect outbreaks across the
 891 European Alps. *New Phytologist* **182**(4), 929–941.
- 892 BÜNTGEN, U., GREUTER, L., BOLLMANN, K., JENNY, H., LIEBHOLD, A., GALVÁN, J. D., STENSETH, N.
 893 C., ANDREW, C. & MYSTERUD, A. (2017). Elevational range shifts in four mountain ungulate
 894 species from the Swiss Alps. *Ecosphere* **8**(4), e01761.
- 895 BÜNTGEN, U., LIEBHOLD, A., NIEVERGELT, D., WERMELINGER, B., ROQUES, A., REINIG, F., KRUSIC, P.
 896 J., PIERMATTEI, A., EGLI, S. & CHERUBINI, P. (2020). Return of the moth: rethinking the effect
 897 of climate on insect outbreaks. *Oecologia*, 1–10.

- BURGESS, M. D., SMITH, K. W., EVANS, K. L., LEECH, D., PEARCE-HIGGINS, J. W., BRANSTON, C. J.,
BRIGGS, K., CLARK, J. R., DU FEU, C. R. & LEWTHWAITE, K. (2018). Tritrophic phenological
match–mismatch in space and time. *Nature Ecology & Evolution* **2**(6), 970–975.
- BUTCHART, S. H., WALPOLE, M., COLLEN, B., VAN STRIEN, A., SCHARLEMANN, J. P., ALMOND, R. E.,
BAILLIE, J. E., BOMHARD, B., BROWN, C. & BRUNO, J. (2010). Global biodiversity: indicators
of recent declines. *Science*, 1187512.
- CALLAGHAN, C. T., ROWLEY, J. J. L., CORNWELL, W. K., POORE, A. G. B. & MAJOR, R. E. (2019).
Improving big citizen science data: Moving beyond haphazard sampling. *Plos Biology* **17**(6),
e3000357.
- CANG, F. A., WILSON, A. A. & WIENS, J. J. (2016). Climate change is projected to outpace rates of
niche change in grasses. *Biology Letters* **12**(9), 20160368.
- CHAUCHARD, S., BEILHE, F., DENIS, N. & CARCAILLET, C. (2010). An increase in the upper tree-limit
of silver fir (*Abies alba* Mill.) in the Alps since the mid-20th century: a land-use change
phenomenon. *Forest Ecology and Management* **259**(8), 1406–1415.
- CHEN, I.-C., HILL, J. K., OHLEMÜLLER, R., ROY, D. B. & THOMAS, C. D. (2011). Rapid range shifts of
species associated with high levels of climate warming. *Science* **333**(6045), 1024–1026.
- CHEN, X., AN, S., INOUE, D. & SCHWARTZ, M. (2015). Temperature and snowfall trigger alpine
vegetation green-up on the world's roof. *Global Change Biology* **21**(10), 3635–46.
- CLELAND, E. E., CHUINE, I., MENZEL, A., MOONEY, H. A. & SCHWARTZ, M. D. (2007). Shifting plant
phenology in response to global change. *Trends in Ecology & Evolution* **22**(7), 357–365.
- COLOMBARI, F., BATTISTI, A., SCHROEDER, L. M. & FACCOLI, M. (2012). Life-history traits promoting
outbreaks of the pine bark beetle *Ips acuminatus* (Coleoptera: Curculionidae, Scolytinae) in the
south-eastern Alps. *European Journal of Forest Research* **131**(3), 553–561.
- COMTE, L. & GRENOUILLET, G. (2013). Do stream fish track climate change? Assessing distribution
shifts in recent decades. *Ecography* **36**(11), 1236–1246.
- CORLETT, R. T. & WESTCOTT, D. A. (2013). Will plant movements keep up with climate change?
Trends in ecology & evolution **28**(8), 482–488.
- CORONA-LOZADA, M., MORIN, S. & CHOLER, P. (2019). Drought offsets the positive effect of summer
heat waves on the canopy greenness of mountain grasslands. *Agricultural and Forest
Meteorology* **276**, 107617.
- DAINESE, M., AIKIO, S., HULME, P. E., BERTOLLI, A., PROSSER, F. & MARINI, L. (2017). Human
disturbance and upward expansion of plants in a warming climate. *Nature Climate Change*
7(8), 577–580.
- DE FRENNE, P., ZELLWEGER, F., RODRÍGUEZ-SÁNCHEZ, F., SCHEFFERS, B. R., HYLANDER, K., LUOTO,
M., VELLEND, M., VERHEYEN, K. & LENOIR, J. (2019). Global buffering of temperatures under
forest canopies. *Nature Ecology & Evolution*.

1
2
3 934 DELESTRADE, A. & STOYANOV, G. (1995). Breeding biology and survival of the Alpine Chough
4 935 *Pyrrhocorax graculus*. *Bird Study* **42**(3), 222–231.
5
6 936 DELL, D., SPARKS, T. H. & DENNIS, R. L. (2005). Climate change and the effect of increasing spring
7 937 temperatures on emergence dates of the butterfly *Apatura iris* (Lepidoptera: Nymphalidae).
8 938 *European Journal of Entomology* **102**(2), 161–167.
9
10 939 DÉQUÉ, M., ROWELL, D., LÜTHI, D., GIORGI, F., CHRISTENSEN, J., ROCKEL, B., JACOB, D.,
11 940 KJELLSTRÖM, E., DE CASTRO, M. & VAN DEN HURK, B. (2007). An intercomparison of
12 941 regional climate simulations for Europe: assessing uncertainties in model projections. *Climatic*
13 942 *Change* **81**(1), 53–70.
14
15 943 DEVICTOR, V., VAN SWAAY, C., BRERETON, T., BROTONS, L., CHAMBERLAIN, D., HELIÖLÄ, J.,
16 944 HERRANDO, S., JULLIARD, R., KUUSSAARI, M. & LINDSTRÖM, Å. (2012). Differences in the
17 945 climatic debts of birds and butterflies at a continental scale. *Nature Climate Change* **2**(2), 121–
18 946 124.
19
20 947 DIEZ, J., KAUSERUD, H., ANDREW, C., HEEGAARD, E., KRISAI-GREILHUBER, I., SENN-IRLET, B.,
21 948 HØILAND, K., EGLI, S. & BÜNTGEN, U. (2020). Altitudinal upwards shifts in fungal fruiting in
22 949 the Alps. *Proceedings of the Royal Society B: Biological Sciences* **287**(1919), 20192348.
23
24 950 DIOLAIUTI, G., BOCCHIOLA, D., VAGLIASINDI, M., D’AGATA, C. & SMIRAGLIA, C. (2012). The 1975–
25 951 2005 glacier changes in Aosta Valley (Italy) and the relations with climate evolution. *Progress*
26 952 *in Physical Geography* **36**(6), 764–785.
27
28 953 DORAZIO, R. M. (2014). Accounting for imperfect detection and survey bias in statistical analysis of
29 954 presence - only data. *Global Ecology and Biogeography* **23**(12), 1472–1484.
30
31 955 DURAND, Y., GIRAUD, G., LATERNER, M., ETCHEVERS, P., MÉRINDOL, L. & LESAFFRE, B. (2009).
32 956 Reanalysis of 47 years of climate in the French Alps (1958–2005): climatology and trends for
33 957 snow cover. *Journal of applied meteorology and climatology* **48**(12), 2487–2512.
34
35 958 ELSEN, P. R., MONAHAN, W. B. & MERENLENDER, A. M. (2020). Topography and human pressure in
36 959 mountain ranges alter expected species responses to climate change. *Nature communications*
37 960 **11**(1), 1–10.
38
39 961 ESPER, J., BÜNTGEN, U., FRANK, D. C., NIEVERGELT, D. & LIEBHOLD, A. (2007). 1200 years of regular
40 962 outbreaks in alpine insects. *Proceedings of the Royal Society of London B: Biological Sciences*
41 963 **274**(1610), 671–679.
42
43 964 FACCOLI, M. (2009). Effect of Weather on *Ips typographus* (Coleoptera Curculionidae) Phenology,
44 965 Voltinism, and Associated Spruce Mortality in the Southeastern Alps. *Environmental*
45 966 *Entomology* **38**(2), 307–316.
46
47 967 FISCHER, A., KOTLARSKI, S., STRASSMANN, K., SCHÄR, C., CROCI-MASPOLI, M., KNUTTI, R. &
48 968 SCHWIERZ, C. (2019). The new Swiss climate change scenarios CH2018. In *Geophysical*
49 969 *Research Abstracts*, vol. 21.
50
51
52
53
54
55
56
57
58
59
60

- FOSTER, G. & RAHMSTORF, S. (2011). Global temperature evolution 1979–2010. *Environmental Research Letters* **6**(4), 044022.
- FU, Y. H., GENG, X., HAO, F., VITASSE, Y., ZOHNER, C. M., ZHANG, X., ZHOU, X., YIN, G., PEÑUELAS, J. & PIAO, S. (2019). Shortened temperature - relevant period of spring leaf - out in temperate - zone trees. *Global Change Biology* **25**(12), 4282–4290.
- GALLINAT, A. S., PRIMACK, R. B. & WAGNER, D. L. (2015). Autumn, the neglected season in climate change research. *Trends in Ecology & Evolution* **30**(3), 169–176.
- GAUDRY, W., SAÏD, S., GAILLARD, J.-M., CHEVRIER, T., LOISON, A., MAILLARD, D. & BONENFANT, C. (2015). Partial migration or just habitat selection? Seasonal movements of roe deer in an Alpine population. *Journal of Mammalogy* **96**(3), 502–510.
- GEHRIG-FASEL, J., GUIBAN, A. & ZIMMERMANN, N. E. (2007). Tree line shifts in the Swiss Alps: Climate change or land abandonment? *Journal of Vegetation Science* **18**(4), 571–582.
- GELLRICH, M., BAUR, P., KOCH, B. & ZIMMERMANN, N. E. (2007). Agricultural land abandonment and natural forest re-growth in the Swiss mountains: a spatially explicit economic analysis. *Agriculture, Ecosystems & Environment* **118**(1), 93–108.
- GIEZENDANNER, J., BERTUZZO, E., PASETTO, D., GUIBAN, A. & RINALDO, A. (2019). A minimalist model of extinction and range dynamics of virtual mountain species driven by warming temperatures. *PLoS ONE* **14**(3), e0213775.
- GRACE, J. B. & IRVINE, K. M. (2020). Scientist's guide to developing explanatory statistical models using causal analysis principles. *Ecology* **101**(4), e02962.
- GUO, F., LENOIR, J. & BONEBRAKE, T. C. (2018). Land-use change interacts with climate to determine elevational species redistribution. *Nature communications* **9**(1), 1–7.
- GÜSEWELL, S. (2014). Phenological responses to changing temperatures: representativeness and precision of results from the Swiss Phenological Network. Master thesis, ETH Zurich.
- HELLE, T. & KOJOLA, I. (2008). Demographics in an alpine reindeer herd: effects of density and winter weather. *Ecography* **31**(2), 221–230.
- HERFINDAL, I., ANDERWALD, P., FILLI, F., ANDRI, S. C. & REMPFLE, T. (2019). Climate, competition and weather conditions drive vertical displacement and habitat use of an alpine ungulate in a highly topographic landscape. *Landscape Ecology* **34**(11), 2523–2539.
- HERNÁNDEZ-HENRÍQUEZ, M. A., DÉRY, S. J. & DERKSEN, C. (2015). Polar amplification and elevation-dependence in trends of Northern Hemisphere snow cover extent, 1971–2014. *Environmental Research Letters* **10**(4), 044010.
- HEWITT, G. (2000). The genetic legacy of the Quaternary ice ages. *Nature* **405**(6789), 907–913.
- HODNEBROG, Ø., MARELLE, L., ALTERSKJÆR, K., WOOD, R., LUDWIG, R., FISCHER, E., RICHARDSON, T., FORSTER, P., SILLMANN, J. & MYHRE, G. (2019). Intensification of summer precipitation with shorter time-scales in Europe. *Environmental Research Letters* **14**(12), 124050.

- HOLZINGER, B., HÜLBER, K., CAMENISCH, M. & GRABHERR, G. (2008). Changes in plant species richness over the last century in the eastern Swiss Alps: elevational gradient, bedrock effects and migration rates. *Plant Ecology* **195**(2), 179–196.
- HÖLZINGER, J. (1997). Die Vögel Baden-Württembergs: Muscicapidae (Fliegenschnäpper) und Thraupidae (Ammertangaren). Singvögel. Passeriformes-sperlingsvögel. Verlag Eugen Ulmer.
- ILER, A. M., HØYE, T. T., INOUE, D. W. & SCHMIDT, N. M. (2013). Nonlinear flowering responses to climate: are species approaching their limits of phenological change? *Philosophical Transactions of the Royal Society of London B: Biological Sciences* **368**(1624), 20120489.
- IMS, R. A. & YOCCOZ, N. G. (2017). Ecosystem-based monitoring in the age of rapid climate change and new technologies. *Current opinion in environmental sustainability* **29**, 170–176.
- INOUE, D. W. (2000). The ecological and evolutionary significance of frost in the context of climate change. *Ecology Letters* **3**(5), 457–463.
- INOUE, D. W. (2008). Effects of climate change on phenology, frost damage, and floral abundance of montane wildflowers. *Ecology* **89**(2), 353–362.
- IPCC (2013). Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- JACOBSEN, D. (2020). The dilemma of altitudinal shifts: caught between high temperature and low oxygen. *Frontiers in Ecology and the Environment* **18**(4), 211–218.
- JÄHNIG, S., SANDER, M. M., CAPRIO, E., ROSSELLI, D., ROLANDO, A. & CHAMBERLAIN, D. (2020). Microclimate affects the distribution of grassland birds, but not forest birds, in an Alpine environment.
- JAKOBY, O., LISCHKE, H. & WERMELINGER, B. (2019). Climate change alters elevational phenology patterns of the European spruce bark beetle (*Ips typographus*). *Global Change Biology* **25**(12), 4048–4063.
- JASPER, K., CALANCA, P., GYALISTRAS, D. & FUHRER, J. (2004). Differential impacts of climate change on the hydrology of two alpine river basins. *Climate Research* **26**(2), 113–129.
- JEZKOVA, T. & WIENS, J. J. (2016). Rates of change in climatic niches in plant and animal populations are much slower than projected climate change. *Proc. R. Soc. B* **283**(1843), 20162104.
- JOHNSON, D. M., BÜNTGEN, U., FRANK, D. C., KAUSRUD, K., HAYNES, K. J., LIEBHOLD, A. M., ESPER, J. & STENSETH, N. C. (2010). Climatic warming disrupts recurrent Alpine insect outbreaks. *Proceedings of the National Academy of Sciences* **107**(47), 20576–20581.
- JUMP, A. S. & PENUELAS, J. (2005). Running to stand still: adaptation and the response of plants to rapid climate change. *Ecology Letters* **8**(9), 1010–1020.
- KAMMER, P. M., SCHÖB, C. & CHOLER, P. (2007). Increasing species richness on mountain summits: Upward migration due to anthropogenic climate change or re-colonisation? *Journal of Vegetation Science* **18**(2), 301–306.

- 1043 KEILER, M., KNIGHT, J. & HARRISON, S. (2010). Climate change and geomorphological hazards in the
 1044 eastern European Alps. *Philosophical Transactions of the Royal Society A: Mathematical,*
 1045 *Physical and Engineering Sciences* **368**(1919), 2461–2479.
- 1046 KHAROUBA, H. M., EHRLÉN, J., GELMAN, A., BOLMGREN, K., ALLEN, J. M., TRAVERS, S. E. &
 1047 WOLKOVICH, E. M. (2018a). Global shifts in the phenological synchrony of species
 1048 interactions over recent decades. *Proceedings of the National Academy of Sciences* **115**(20),
 1049 5211–5216.
- 1050 KHAROUBA, H. M. & WOLKOVICH, E. M. (2020). Disconnects between ecological theory and data in
 1051 phenological mismatch research. *Nature Climate Change* **10**(5), 406–415.
- 1052 KIRCHNER, M., FAUS - KESSLER, T., JAKOBI, G., LEUCHNER, M., RIES, L., SCHEEL, H. E. & SUPPAN, P.
 1053 (2013). Altitudinal temperature lapse rates in an Alpine valley: trends and the influence of
 1054 season and weather patterns. *International Journal of Climatology* **33**(3), 539–555.
- 1055 KLEIN, G., REBETEZ, M., RIXEN, C. & VITASSE, Y. (2018). Unchanged risk of frost exposure for
 1056 subalpine and alpine plants after snowmelt in Switzerland despite climate warming.
 1057 *International Journal of Biometeorology* **62**(9), 1755–1762.
- 1058 KLEIN, G., VITASSE, Y., RIXEN, C., MARTY, C. & REBETEZ, M. (2016). Shorter snow cover duration
 1059 since 1970 in the Swiss Alps due to earlier snowmelt more than to later snow onset. *Climatic*
 1060 *Change* **139**(3-4), 637–649.
- 1061 KNAUS, P. (2018). Schweizer Brutvogelatlas 2013-2016: Verbreitung und Bestandesentwicklung der
 1062 Vögel in der Schweiz und im Fürstentum Liechtenstein, Swiss Ornithological Institute,
 1063 Sempach.
- 1064 KÖRNER, C. (2003). Alpine plant life (2nd ed.). Springer, Berlin.
- 1065 KÖRNER, C. (2011). Coldest places on earth with angiosperm plant life. *Alpine Botany* **121**(1), 11–22.
- 1066 KORSLUND, L. & STEEN, H. (2006). Small rodent winter survival: snow conditions limit access to food
 1067 resources. *Journal of Animal Ecology* **75**(1), 156–166.
- 1068 KÜCHLER, M., KÜCHLER, H., BEDOLLA, A. & WOHLGEMUTH, T. (2015). Response of Swiss forests to
 1069 management and climate change in the last 60 years. *Annals of Forest Science* **72**(3), 311–320.
- 1070 KUHNS, E., LENOIR, J., PIEDALLU, C. & GÉGOUT, J.-C. (2016). Early signs of range disjunction of
 1071 submountainous plant species: an unexplored consequence of future and contemporary climate
 1072 changes. *Global Change Biology* **22**(6), 2094–2105.
- 1073 LATERNER, M. & SCHNEEBELI, M. (2003). Long - term snow climate trends of the Swiss Alps (1931 -
 1074 99). *International Journal of Climatology* **23**(7), 733–750.
- 1075 LAVOREL, S., GRIGULIS, K., LEITINGER, G., KOHLER, M., SCHIRPKE, U. & TAPPEINER, U. (2017).
 1076 Historical trajectories in land use pattern and grassland ecosystem services in two European
 1077 alpine landscapes. *Regional Environmental Change* **17**(8), 2251–2264.

1
2
3 1078 LEHIKONEN, A., LINDÉN, A., KARLSSON, M., ANDERSSON, A., CREWE, T. L., DUNN, E. H., GREGORY,
4 1079 G., KARLSSON, L., KRISTIANSEN, V., MACKENZIE, S., NEWMAN, S., RØER, J. E., SHARPE, C.,
5 1080 SOKOLOV, L. V., STEINHOLTZ, Å., STERVANDER, M., TIRRI, I.-S. & TJØRNØV, R. S. (2019).
6 1081 Phenology of the avian spring migratory passage in Europe and North America: Asymmetric
7 1082 advancement in time and increase in duration. *Ecological Indicators* **101**, 985–991.
8 1083 LENOIR, J., BERTRAND, R., COMTE, L., BOURGEAUD, L., HATTAB, T., MURIENNE, J. & GRENOUILLET,
9 1084 G. (2020). Species better track climate warming in the oceans than on land. *Nature Ecology &*
10 1085 *Evolution*, 1–16.
11 1086 LENOIR, J., GEGOUT, J. C., GUIBAN, A., VITTOZ, P., WOHLGEMUTH, T., ZIMMERMANN, N. E.,
12 1087 DULLINGER, S., PAULI, H., WILLNER, W. & SVENNING, J. C. (2010). Going against the flow:
13 1088 potential mechanisms for unexpected downslope range shifts in a warming climate. *Ecography*
14 1089 **33**(2), 295–303.
15 1090 LENOIR, J., GEGOUT, J. C., MARQUET, P. A., DE RUFFRAY, P. & BRISSE, H. (2008). A significant
16 1091 upward shift in plant species optimum elevation during the 20th century. *Science* **320**(5884),
17 1092 1768–1771.
18 1093 LENOIR, J., GEGOUT, J. C., PIERRAT, J. C., BONTEMPS, J. D. & DHOTE, J. F. (2009). Differences
19 1094 between tree species seedling and adult altitudinal distribution in mountain forests during the
20 1095 recent warm period (1986–2006). *Ecography* **32**(5), 765–777.
21 1096 LENOIR, J. & SVENNING, J. C. (2015). Climate - related range shifts-a global multidimensional
22 1097 synthesis and new research directions. *Ecography* **38**(1), 15–28.
23 1098 MAGGINI, R., LEHMANN, A., KERY, M., SCHMID, H., BENISTON, M., JENNI, L. & ZBINDEN, N. (2011).
24 1099 Are Swiss birds tracking climate change? Detecting elevational shifts using response curve
25 1100 shapes. *Ecological Modelling* **222**(1), 21–32.
26 1101 MAGNUSSON, J., JONAS, T., LÓPEZ-MORENO, I. & LEHNING, M. (2010). Snow cover response to
27 1102 climate change in a high alpine and half-glacierized basin in Switzerland. *Hydrology research*
28 1103 **41**(3–4), 230–240.
29 1104 MARCHISIO, C., CESCATTI, A. & BATTISTI, A. (1994). Climate, soils and *Cephalcia arvensis* outbreaks
30 1105 on *Picea abies* in the Italian Alps. *Forest Ecology and Management* **68**(2–3), 375–384.
31 1106 MARINI, L., AYRES, M., BATTISTI, A. & FACCOLI, M. (2012). Climate affects severity and altitudinal
32 1107 distribution of outbreaks in an eruptive bark beetle. *Climatic Change* **115**(2), 327–341.
33 1108 MARTY, C. & BLANCHET, J. (2012). Long-term changes in annual maximum snow depth and snowfall
34 1109 in Switzerland based on extreme value statistics. *Climatic Change* **111**(3–4), 705–721.
35 1110 MARTY, C. & MEISTER, R. (2012). Long-term snow and weather observations at Weissfluhjoch and its
36 1111 relation to other high-altitude observatories in the Alps. *Theoretical and Applied Climatology*
37 1112 **110**(4), 573–583.

- MARTY, C., SCHLÖGL, S., BAVAY, M. & LEHNING, M. (2017a). How much can we save? Impact of different emission scenarios on future snow cover in the Alps. *The Cryosphere* **11**(1), 517–529.
- MARTY, C., TILG, A.-M. & JONAS, T. (2017b). Recent evidence of large-scale receding snow water equivalents in the European Alps. *Journal of Hydrometeorology* **18**(4), 1021–1031.
- MATIU, M., CRESPI, A., BERTOLDI, G., CARMAGNOLA, C. M., MARTY, C., MORIN, S., SCHÖNER, W., CAT BERRO, D., CHIOGNA, G., DE GREGORIO, L., KOTLARSKI, S., MAJONE, B., RESCH, G., TERZAGO, S., VALT, M., *ET AL.* (2021). Observed snow depth trends in the European Alps: 1971 to 2019. *The Cryosphere* **15**(3), 1343–1382.
- MCLAUGHLIN, B. C., ACKERLY, D. D., KLOS, P. Z., NATALI, J., DAWSON, T. E. & THOMPSON, S. E. (2017). Hydrologic refugia, plants, and climate change. *Global Change Biology* **23**(8), 2941–2961.
- MENÉNDEZ, R., GONZÁLEZ-MEGÍAS, A., JAY-ROBERT, P. & MARQUÉZ-FERRANDO, R. (2014). Climate change and elevational range shifts: evidence from dung beetles in two European mountain ranges. *Global Ecology and Biogeography* **23**(6), 646–657.
- MIGLIAVACCA, M., CREMONESE, E., COLOMBO, R., Busetto, L., GALVAGNO, M., GANIS, L., MERONI, M., PARI, E., ROSSINI, M., SINISCALCO, C. & DI CELLA, U. M. (2008). European larch phenology in the Alps: can we grasp the role of ecological factors by combining field observations and inverse modelling? *International Journal of Biometeorology* **52**(7), 587–605.
- MOBERG, A., JONES, P. D., LISTER, D., WALTHER, A., BRUNET, M., JACOBET, J., ALEXANDER, L. V., DELLA - MARTA, P. M., LUTERBACHER, J. & YIOU, P. (2006). Indices for daily temperature and precipitation extremes in Europe analyzed for the period 1901–2000. *Journal of Geophysical Research: Atmospheres* **111**(D22).
- MOUQUET, N., LAGADEUC, Y., DEVICTOR, V., DOYEN, L., DUPUTIÉ, A., EVEILLARD, D., FAURE, D., GARNIER, E., GIMENEZ, O. & HUNEMAN, P. (2015). Predictive ecology in a changing world. *Journal of Applied Ecology* **52**(5), 1293–1310.
- MYSTERUD, A., STENSETH, N. C., YOCOZ, N. G., LANGVATN, R. & STEINHEIM, G. (2001). Nonlinear effects of large-scale climatic variability on wild and domestic herbivores. *Nature* **410**(6832), 1096–1099.
- MYSTERUD, A., YOCOZ, N. G., LANGVATN, R., PETTORELLI, N. & STENSETH, N. C. (2008). Hierarchical path analysis of deer responses to direct and indirect effects of climate in northern forest. *Philosophical Transactions of the Royal Society B: Biological Sciences* **363**(1501), 2357–2366.
- NOVOA, C., ASTRUC, G., DESMET, J.-F. & BESNARD, A. (2016). No short-term effects of climate change on the breeding of Rock Ptarmigan in the French Alps and Pyrenees. *Journal of Ornithology* **157**(3), 797–810.

1
2
3 1148 NOVOA, C., BESNARD, A., BRENOT, J. F. & ELLISON, L. N. (2008). Effect of weather on the
4 1149 reproductive rate of Rock Ptarmigan *Lagopus muta* in the eastern Pyrenees. *Ibis* **150**(2), 270–
5 1150 278.
6
7
8 1151 OFFNER, J. & LE BRUN, P. (1956). Un siècle de floristique à travers les Alpes françaises. *Bulletin de la*
9 1152 *Société botanique de France* **103**(5-6), 298–375.
10
11 1153 PARMESAN, C. & YOHE, G. (2003). A globally coherent fingerprint of climate change impacts across
12 1154 natural systems. *Nature* **421**(6918), 37–42.
13
14 1155 PAROLO, G. & ROSSI, G. (2008). Upward migration of vascular plants following a climate warming
15 1156 trend in the Alps. *Basic and Applied Ecology* **9**(2), 100–107.
16
17 1157 PELÁEZ, M., GAILLARD, J. M., BOLLMANN, K., HEURICH, M. & REHNUS, M. (2020). Large scale
18 1158 variation in birth timing and synchrony of a large herbivore along the latitudinal and altitudinal
19 1159 gradients. *Journal of Animal Ecology*.
20
21
22 1160 PELLICCIOTTI, F., BAUDER, A. & PAROLA, M. (2010). Effect of glaciers on streamflow trends in the
23 1161 Swiss Alps. *Water Resources Research* **46**(10).
24
25 1162 PEPIN, N. & LUNDQUIST, J. (2008). Temperature trends at high elevations: patterns across the globe.
26 1163 *Geophysical Research Letters* **35**(14).
27
28 1164 PERNOLLET, C. A., KORNER-NIEVERGELT, F. & JENNI, L. (2015). Regional changes in the elevational
29 1165 distribution of the Alpine Rock Ptarmigan *Lagopus muta helvetica* in Switzerland. *Ibis* **157**(4),
30 1166 823–836.
31
32
33 1167 PETTORELLI, N., PELLETIER, F., HARDENBERG, A. V., FESTA-BIANCHET, M. & CÔTÉ, S. D. (2007).
34 1168 Early onset of vegetation growth vs. rapid green - up: Impacts on juvenile mountain ungulates.
35 1169 *Ecology* **88**(2), 381–390.
36
37
38 1170 PIOTTI, A., PIOVANI, P., SCALFÌ, M., LEONARDI, S. & MENOZZI, P. (2007). Global change and upward
39 1171 shift of treeline in the Alps: genetic consequences of pastures colonization. *Forest@ - Journal*
40 1172 *of Silviculture and Forest Ecology* **4**(1), 69–78.
41
42
43 1173 PLARD, F., GAILLARD, J.-M., COULSON, T., HEWISON, A. M., DELORME, D., WARNANT, C. &
44 1174 BONENFANT, C. (2014). Mismatch between birth date and vegetation phenology slows the
45 1175 demography of roe deer. *Plos Biology* **12**(4), e1001828.
46
47
48 1176 POPY, S., BORDIGNON, L. & PRODON, R. (2010). A weak upward elevational shift in the distributions
49 1177 of breeding birds in the Italian Alps. *Journal of Biogeography* **37**(1), 57–67.
50
51 1178 PULLIAINEN, J., LUOJUS, K., DERKSEN, C., MUDRYK, L., LEMMETYINEN, J., SALMINEN, M., IKONEN, J.,
52 1179 TAKALA, M., COHEN, J. & SMOLANDER, T. (2020). Patterns and trends of Northern
53 1180 Hemisphere snow mass from 1980 to 2018. *Nature* **581**(7808), 294–298.
54
55
56 1181 R CORE TEAM. (2017). R: A language and environment for statistical computing. R Foundation for
57 1182 Statistical Computing, Vienna, Austria. <http://www.R-project.org/>.
58
59
60

- 1183 RAJCZAK, J. & SCHÄR, C. (2017). Projections of future precipitation extremes over Europe: a
 1184 multimodel assessment of climate simulations. *Journal of Geophysical Research: Atmospheres*
 1185 **122**(20), 10,773–10,800.
- 1186 RAPACCIUOLO, G., MAHER, S. P., SCHNEIDER, A. C., HAMMOND, T. T., JABIS, M. D., WALSH, R. E.,
 1187 IKNAYAN, K. J., WALDEN, G. K., OLDFATHER, M. F., ACKERLY, D. D. & BEISSINGER, S. R.
 1188 (2014). Beyond a warming fingerprint: individualistic biogeographic responses to
 1189 heterogeneous climate change in California. *Global Change Biology* **20**(9), 2841–2855.
- 1190 REBETEZ, M. & DOBBERTIN, M. (2004). Climate change may already threaten Scots pine stands in the
 1191 Swiss Alps. *Theoretical and Applied Climatology* **79**(1), 1–9.
- 1192 REBETEZ, M. & REINHARD, M. (2008). Monthly air temperature trends in Switzerland 1901–2000 and
 1193 1975–2004. *Theoretical and Applied Climatology* **91**(1–4), 27–34.
- 1194 REHNUS, M., PELÁEZ, M. & BOLLMANN, K. (2020). Advancing plant phenology causes an increasing
 1195 trophic mismatch in an income breeder across a wide elevational range. *Ecosphere* **11**(6),
 1196 e03144.
- 1197 RENNER, S. S. & ZOHNER, C. M. (2018). Climate change and phenological mismatch in trophic
 1198 interactions among plants, insects, and vertebrates. *Annual Review of Ecology, Evolution, and*
 1199 *Systematics* **49**, 165–182.
- 1200 RESANO-MAYOR, J., KORNER-NIEVERGELT, F., VIGNALI, S., HORRENBERGER, N., BARRAS, A. G.,
 1201 BRAUNISCH, V., PERNOLLET, C. A. & ARLETTAZ, R. (2019). Snow cover phenology is the
 1202 main driver of foraging habitat selection for a high-alpine passerine during breeding:
 1203 implications for species persistence in the face of climate change. *Biodiversity and*
 1204 *Conservation* **28**(10), 2669–2685.
- 1205 RIGLING, A. & CHERUBINI, P. (1999). Wieso sterben die Waldföhren im «Telwald» bei Visp? Eine
 1206 Zusammenfassung bisheriger Studien und eine dendroökologische Untersuchung| What is the
 1207 Cause of the High Mortality Rates of the Scots Pines in the «Telwald» near Visp
 1208 (Switzerland)? A Summary of Previous Studies and a Dendroecological Study. *Schweizerische*
 1209 *Zeitschrift für Forstwesen* **150**(4), 113–131.
- 1210 ROBERTS, A. M., TANSEY, C., SMITHERS, R. J. & PHILLIMORE, A. B. (2015). Predicting a change in the
 1211 order of spring phenology in temperate forests. *Global Change Biology*.
- 1212 ROBINSON, B. G. & MERRILL, E. H. (2012). The influence of snow on the functional response of
 1213 grazing ungulates. *Oikos* **121**(1), 28–34.
- 1214 ROGORA, M., FRATE, L., CARRANZA, M., FREPPAZ, M., STANISCI, A., BERTANI, I., BOTTARIN, R.,
 1215 BRAMBILLA, A., CANULLO, R. & CARBOGNANI, M. (2018). Assessment of climate change
 1216 effects on mountain ecosystems through a cross-site analysis in the Alps and Apennines.
 1217 *Science of the Total Environment* **624**, 1429–1442.
- 1218 ROLLAND, C. (2003). Spatial and seasonal variations of air temperature lapse rates in Alpine regions.
 1219 *Journal of Climate* **16**(7), 1032–1046.

1
2
3 1220 ROSBAKH, S., LEINGÄRTNER, A., HOISS, B., KRAUSS, J., STEFFAN-DEWENTER, I. & POSCHLOD, P.
4
5 1221 (2017). Contrasting Effects of Extreme Drought and Snowmelt Patterns on Mountain Plants
6
7 1222 along an Elevation Gradient. *Frontiers in Plant Science* **8**(1478).
8 1223 ROTH, T., PLATTNER, M. & AMRHEIN, V. (2014). Plants, Birds and Butterflies: Short-Term Responses
9
10 1224 of Species Communities to Climate Warming Vary by Taxon and with Altitude. *PLoS ONE*
11 1225 **9**(1), e82490.
12
13 1226 ROUSSELOT, M., DURAND, Y., GIRAUD, G., MÉRINDOL, L., DOMBROWSKI-ETCHEVERS, I., DÉQUÉ, M.
14 1227 & CASTEBRUNET, H. (2012). Statistical adaptation of ALADIN RCM outputs over the French
15
16 1228 Alps-application to future climate and snow cover. *The Cryosphere* **6**, 785-805.
17 1229 RUBOLINI, D., AMBROSINI, R., CAFFI, M., BRICHETTI, P., ARMIRAGLIO, S. & SAINO, N. (2007). Long-
18
19 1230 term trends in first arrival and first egg laying dates of some migrant and resident bird species
20
21 1231 in northern Italy. *International Journal of Biometeorology* **51**(6), 553–563.
22 1232 RUMPF, S. B., HÜLBER, K., KLONNER, G., MOSER, D., SCHÜTZ, M., WESSELY, J., WILLNER, W.,
23
24 1233 ZIMMERMANN, N. E. & DULLINGER, S. (2018). Range dynamics of mountain plants decrease
25
26 1234 with elevation. *Proceedings of the National Academy of Sciences* **115**(8), 1848–1853.
27 1235 SCHÄR, C., VIDALE, P. L., LUTHI, D., FREI, C., HABERLI, C., LINIGER, M. A. & APPENZELLER, C.
28
29 1236 (2004). The role of increasing temperature variability in European summer heatwaves. *Nature*
30 1237 **427**(6972), 332–336.
31
32 1238 SCHERRER, D., VITASSE, Y., GUISAN, A., WOHLGEMUTH, T. & LISCHKE, H. (2020). Competition and
33
34 1239 demography rather than dispersal limitation slow down upward shifts of trees' upper elevation
35
36 1240 limits in the Alps. *Journal of Ecology* **108**(6), 2416–2430.
37 1241 SCHERRER, S., CEPPI, P., CROCI-MASPOLI, M. & APPENZELLER, C. (2012). Snow-albedo feedback and
38 1242 Swiss spring temperature trends. *Theoretical and Applied Climatology* **110**(4), 509–516.
39 1243 SCHERRER, S. C., BEGERT, M., CROCI - MASPOLI, M. & APPENZELLER, C. (2016). Long series of
40
41 1244 Swiss seasonal precipitation: regionalization, trends and influence of large - scale flow.
42
43 1245 *International Journal of Climatology* **36**(11), 3673-3689.
44
45 1246 SCHMIDL, J. & FREI, C. (2005). Trends of heavy precipitation and wet and dry spells in Switzerland
46
47 1247 during the 20th century. *International Journal of Climatology: A Journal of the Royal*
48 1248 *Meteorological Society* **25**(6), 753–771.
49 1249 SCHMUCKI, E., MARTY, C., FIERZ, C. & LEHNING, M. (2015). Simulations of 21st century snow
50
51 1250 response to climate change in Switzerland from a set of RCMs. *International journal of*
52
53 1251 *climatology* **35**(11), 3262-3273.
54 1252 SCHMUCKI, E., MARTY, C., FIERZ, C., WEINGARTNER, R. & LEHNING, M. (2017). Impact of climate
55
56 1253 change in Switzerland on socioeconomic snow indices. *Theoretical and Applied Climatology*
57
58 1254 **127**(3-4), 875-889.
59
60

- 1255 SCHNEIDER, L., COMTE, V. & REBETEZ, M. (2021). Increasingly favourable winter temperature
1256 conditions for major crop and forest insect pest species in Switzerland. *Agricultural and Forest*
1257 *Meteorology* **298**, 108315.
- 1258 SCHÖNER, W., AUER, I. & BÖHM, R. (2009). Long term trend of snow depth at Sonnblick (Austrian
1259 Alps) and its relation to climate change. *Hydrological Processes: An International Journal*
1260 **23**(7), 1052–1063.
- 1261 SCHULDT, B., BURAS, A., AREND, M., VITASSE, Y., BEIERKUHNLEIN, C., DAMM, A., GHARUN, M.,
1262 GRAMS, T. E., HAUCK, M. & HAJEK, P. (2020). A first assessment of the impact of the extreme
1263 2018 summer drought on Central European forests. *Basic and Applied Ecology* **45**, 86–103.
- 1264 SCHWINGSHACKL, C., DAVIN, E. L., HIRSCHI, M., SØRLAND, S. L., WARTENBURGER, R. &
1265 SENEVIRATNE, S. I. (2019). Regional climate model projections underestimate future warming
1266 due to missing plant physiological CO₂ response. *Environmental Research Letters* **14**(11),
1267 114019.
- 1268 SERQUET, G., MARTY, C., DULEX, J. P. & REBETEZ, M. (2011). Seasonal trends and temperature
1269 dependence of the snowfall/precipitation - day ratio in Switzerland. *Geophysical Research*
1270 *Letters* **38**(7).
- 1271 SERQUET, G., MARTY, C. & REBETEZ, M. (2013). Monthly trends and the corresponding altitudinal
1272 shift in the snowfall/precipitation day ratio. *Theoretical and Applied Climatology* **114**(3–4),
1273 437–444.
- 1274 SHERIDAN, J. A. & BICKFORD, D. (2011). Shrinking body size as an ecological response to climate
1275 change. *Nature Climate Change* **1**(8), 401.
- 1276 STEFFEN, W., BROADGATE, W., DEUTSCH, L., GAFFNEY, O. & LUDWIG, C. (2015). The trajectory of
1277 the Anthropocene: the great acceleration. *The Anthropocene Review* **2**(1), 81–98.
- 1278 STEGER, C., KOTLARSKI, S., JONAS, T. & SCHÄR, C. (2013). Alpine snow cover in a changing climate:
1279 a regional climate model perspective. *Climate dynamics* **41**(3–4), 735–754.
- 1280 STEINBAUER, M. J., GRYTNES, J.-A., JURASINSKI, G., KULONEN, A., LENOIR, J., PAULI, H., RIXEN, C.,
1281 WINKLER, M., BARDY-DURCHHALTER, M. & BARNI, E. (2018). Accelerated increase in plant
1282 species richness on mountain summits is linked to warming. *Nature* **556**(7700), 231–234.
- 1283 STREBEL, N., KÉRY, M., SCHAUB, M. & SCHMID, H. (2014). Studying phenology by flexible modelling
1284 of seasonal detectability peaks. *Methods in Ecology and Evolution* **5**(5), 483–490.
- 1285 TABERLET, P., FUMAGALLI, L., WUST-SAUCY, A. G. & COSSON, J. F. (1998). Comparative
1286 phylogeography and postglacial colonization routes in Europe. *Molecular Ecology* **7**(4), 453–
1287 464.
- 1288 TAFANI, M., COHAS, A., BONENFANT, C., GAILLARD, J.-M. & ALLAINÉ, D. (2013). Decreasing litter
1289 size of marmots over time: a life history response to climate change? *Ecology* **94**(3), 580–586.

1
2
3 1290 TAPPEINER, U., BORSODORF, A. & BAHN, M. (2013). Long-term socio-ecological research in mountain
4 1291 regions: perspectives from the Tyrolean Alps. In *Long Term Socio-Ecological Research*. (pp.
5 1292 505–525. Springer.
6
7
8 1293 TASSER, E., WALDE, J., TAPPEINER, U., TEUTSCH, A. & NOGGLER, W. (2007). Land-use changes and
9 1294 natural reforestation in the Eastern Central Alps. *Agriculture, Ecosystems & Environment*
10 1295 **118**(1), 115–129.
11
12 1296 THACKERAY, S. J., HENRYS, P. A., HEMMING, D., BELL, J. R., BOTHAM, M. S., BURTHE, S., HELAOUET,
13 1297 P., JOHNS, D. G., JONES, I. D., LEECH, D. I., MACKAY, E. B., MASSIMINO, D., ATKINSON, S.,
14 1298 BACON, P. J., BRERETON, T. M., *ET AL.* (2016). Phenological sensitivity to climate across taxa
15 1299 and trophic levels. *Nature* **535**, 241.
16
17
18 1300 THUILLER, W., GUÉGUEN, M., BISON, M., DUPARC, A., GAREL, M., LOISON, A., RENAUD, J. &
19 1301 POGGIATO, G. (2018). Combining point - process and landscape vegetation models to predict
20 1302 large herbivore distributions in space and time—a case study of *Rupicapra rupicapra*.
21 1303 *Diversity and Distributions* **24**(3), 352–362.
22
23 1304 TINNER, W. & THEURILLAT, J.-P. (2003). Uppermost limit, extent, and fluctuations of the timberline
24 1305 and treeline ecocline in the Swiss Central Alps during the past 11,500 years. *Arctic, Antarctic,*
25 1306 *and Alpine Research* **35**(2), 158–169.
26
27 1307 VACCARI, L. (1914). La sopraelevazione dei limiti superiori dei muschi in valle d’aosta. *Société de la*
28 1308 *Flore Valdotaïne* **9**, 62–84.
29
30 1309 VALT, M. & CIANFARRA, P. (2010). Recent snow cover variability in the Italian Alps. *Cold Regions*
31 1310 *Science and Technology* **64**(2), 146–157.
32
33 1311 VAN DER GRAAF, A., STAHL, J., KLIMKOWSKA, A., BAKKER, J. P. & DRENT, R. H. (2006). Surfing on a
34 1312 green wave-how plant growth drives spring migration in the Barnacle Goose *Branta leucopsis*.
35 1313 *ARDEA-WAGENINGEN*- **94**(3), 567.
36
37 1314 VISSER, M. E. & GIENAPP, P. (2019). Evolutionary and demographic consequences of phenological
38 1315 mismatches. *Nature Ecology & Evolution* **3**(6), 879–885.
39
40 1316 VITASSE, Y., BOTTERO, A., CAILLERET, M., BIGLER, C., FONTI, P., GESSLER, A., LEVESQUE, M.,
41 1317 ROHNER, B., WEBER, P., RIGLING, A. & WOHLGEMUTH, T. (2019). Contrasting resistance and
42 1318 resilience to extreme drought and late spring frost in five major European tree species. *Glob*
43 1319 *Chang Biol* **25**(11), 3781–3792.
44
45 1320 VITASSE, Y., FRANCOIS, C., DELPIERRE, N., DUFRENE, E., KREMER, A., CHUINE, I. & DELZON, S.
46 1321 (2011). Assessing the effects of climate change on the phenology of European temperate trees.
47 1322 *Agricultural and Forest Meteorology* **151**(7), 969–980.
48
49 1323 VITASSE, Y., HOCH, G., RANDIN, C. F., LENZ, A., KOLLAS, C. & KÖRNER, C. (2012). Tree recruitment
50 1324 of European tree species at their current upper elevational limits in the Swiss Alps. *Journal of*
51 1325 *Biogeography* **39**(8), 1439–1449.
52
53
54
55
56
57
58
59
60

- VITASSE, Y., HOCH, G., RANDIN, C. F., LENZ, A., KOLLAS, C., SCHEEPENS, J. F. & KÖRNER, C. (2013). Elevational adaptation and plasticity in seedling phenology of temperate deciduous tree species. *Oecologia* **171**(3), 663–678.
- VITASSE, Y., REBETEZ, M., FILIPPA, G., CREMONESE, E., KLEIN, G. & RIXEN, C. (2017). ‘Hearing’ alpine plants growing after snowmelt: ultrasonic snow sensors provide long-term series of alpine plant phenology. *International Journal of Biometeorology* **61**(2), 349–361.
- VITASSE, Y., SCHNEIDER, L., RIXEN, C., CHRISTEN, D. & REBETEZ, M. (2018a). Increase in the risk of exposure of forest and fruit trees to spring frosts at higher elevations in Switzerland over the last four decades. *Agricultural and Forest Meteorology* **248**, 60–69.
- VITASSE, Y., SIGNARBIEUX, C. & FU, Y. H. (2018b). Global warming leads to more uniform spring phenology across elevations. *Proceedings of the National Academy of Sciences U S A* **115**(5), 1004–1008.
- VON ARX, G., GRAF PANNATIER, E., THIMONIER, A. & REBETEZ, M. (2013). Microclimate in forests with varying leaf area index and soil moisture: potential implications for seedling establishment in a changing climate. *Journal of Ecology* **101**(5), 1201–1213.
- WIENS, J. J. (2016). Climate-related local extinctions are already widespread among plant and animal species. *Plos Biology* **14**(12), e2001104.
- WIPF, S. & RIXEN, C. (2010). A review of snow manipulation experiments in Arctic and alpine tundra ecosystems. *Polar Research* **29**(1), 95–109.
- YOCOZ, N. G., DELESTRADE, A. & LOISON, A. (2010). Impact of climatic change on alpine ecosystems: inference and prediction. *Journal of Alpine Research | Revue de géographie alpine* (98-4).
- ZELLWEGER, F., DE FRENNE, P., LENOIR, J., VANGANSBEKE, P., VERHEYEN, K., BERNHARDT-RÖMERMANN, M., BAETEN, L., HÉDL, R., BERKI, I., BRUNET, J., VAN CALSTER, H., CHUDOMELOVÁ, M., DECOCQ, G., DIRNBÖCK, T., DURAK, T., *ET AL.* (2020). Forest microclimate dynamics drive plant responses to warming. *Science* **368**(6492), 772.
- ZIMMERMANN, N. E., GEBETSROITHER, E., ZÜGER, J., SCHMATZ, D. & PSOMAS, A. (2013). Future climate of the European Alps. *Management Strategies to Adapt Alpine Space Forests to Climate Change Risks; Zimmermann, NE, Ed*, 27–36.

VII. SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1359 **Fig. S1.** Mean phenological shift of the singing activity or laying dates per decade for birds
1360 classified according to habitat.
1361 **Table S1.** Spring phenological change per decade or/and per degree for each studied species.
1362 **Table S2.** Elevational range shifts of the optimum distribution (i.e. elevation of maximum
1363 abundance) or/and the leading edge (i.e. the upper range limit) for each studied species.

For Review Only

Figure legends

Fig. 1. Long-term time series of snow parameters (snowmelt timing and length of the snow free period) and mean air temperature (spring temperature from March to May and annual temperature) at six sites spread across the Swiss Alps during the period 1970–2019. Data updated from Klein *et al.* (2016) representing yearly average anomalies over the period 1970–2019 across six sites in Switzerland located between 1,298 and 2,540 m (Arosa, Davos, Grächen, Grimsel Hospiz, Scuol and Weissfluhjoch). Estimated trends from the Theil-Sen estimator method across the six sites are reported with the associated standard errors (grey shading) for each snow and temperature parameter. Significance of the slopes was tested using Mann-Kendall tests, see Klein *et al.* (2016) for more details.

Fig. 2. Spring phenological change per decade (A) or per degree (B) for different plant and animal groups studied in the European Alps region. Values correspond to the estimated marginal mean of the mixed-effect model with study as a random factor; error bars are 95% confidence intervals. The period of time varied among studies but was always more than 10 years within the period 1980–2020. Only groups with more than five species are shown (all studies are reported in Table 1). Abbreviations: ldm, long-distance migrants; sdm, short-distance migrants; res., resident; semi-aq., semi-aquatic. Different letters among groups indicate significant differences (*post-hoc* Tukey tests at $\alpha = 0.05$).

Fig. 3. Empirical evidence for elevational range shifts of (A) the optimum distribution (i.e. elevation of maximum abundance) and (B) the leading edge (i.e. the upper range limit) reported for groups of taxa in the European Alps region. Values correspond to the estimate of the mixed-effect model with study as a random factor; error bars are 95% confidence intervals.

1
2
3 1389 The period of time investigated varied among studies but was always more than 10 years
4
5 1390 within the period 1980–2020, or for diachronic studies comparing historical and modern
6
7
8 1391 surveys, the modern survey was conducted within the period 1980–2020, and the median year
9
10 1392 of the historical survey(s) was after 1950. Only groups with more than five species are shown
11
12 1393 (all studies are reported in Table 2). Abbreviations: ecm, ectomycorrhizal; sapro., saprophytic;
13
14 1394 semi-aq., semi-aquatic. Different letters among groups indicate significant differences (*post-*
15
16
17 1395 *hoc* Tukey tests at $\alpha = 0.05$).
18
19 1396
20
21
22 1397 **Fig. 4.** Mean elevational shift per decade for four different orders of insects in terms of (A) the
23
24 1398 optimum elevation (i.e. elevation of maximum abundance) and (B) the leading-edge elevation
25
26 1399 (i.e. the upper range limit). Values correspond to the estimate of the mixed-effect model with
27
28
29 1400 study as a random factor; error bars are 95% confidence intervals. The period of time
30
31 1401 investigated varied among studies but was always higher than 10 years within the period
32
33 1402 1980–2020. Different letters among groups indicate significant differences (*post-hoc* Tukey
34
35 1403 tests at $\alpha = 0.05$).
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Documented spring phenological changes in animals and plants in the European Alps over the recent decades. Mean $\pm$ standard errors are reported where data are available for more than one species. For information on the *info fauna* and *PhenoForest* databases, see Section II. The data set used for this table is provided as supporting online information in Table S1. Abbreviations: ldm, long-distance migrants; sdm, short-distance migrants; res., resident.

Taxa	Study period	Area	Number of species	Phenological variable	Trends (days decade ⁻¹)	Temperature sensitivity (days °C ⁻¹)	References
PLANTS							
Woody plants	1970–2012	Alps and lowlands in Switzerland	14	Leaf-out or flowering	−3.4 $\pm$ 0.3	−5.1 $\pm$ 0.4	Güsewell (2014)
Herbaceous	1970–2012	Alps and lowlands in Switzerland	5	Flowering	−2.8 $\pm$ 0.4	−5.6 $\pm$ 0.6	Güsewell (2014)
Trees	1998–2020	Alps and lowlands in Switzerland	8	Leaf-out	−2.2 $\pm$ 0.7	NA	<i>PhenoForest</i>
Woody plants	2005–2016	French Alps	6	Leaf-out	−0.5 $\pm$ 2.3	NA	Bison <i>et al.</i> (2019)
ANIMALS							
Birds							
Birds (res.)	1995–2019	Lowland areas in Switzerland	8	Singing activity	+2.5 $\pm$ 3.6	−2.1 $\pm$ 0.8	Strebel <i>et al.</i> (2014)
Birds (sdm)	1995–2019	Lowland areas in Switzerland	12	Singing activity	−3.5 $\pm$ 3.8	−1.0 $\pm$ 0.5	Strebel <i>et al.</i> (2014)
Birds (ldm)	1995–2019	Lowland areas in Switzerland	14	Singing activity	−1.7 $\pm$ 1.5	−1.0 $\pm$ 0.5	Strebel <i>et al.</i> (2014)
<i>Passer italiae</i> (res.)	1982–2006	Northern Italy	1	Laying dates	−1.9	−1.7	Rubolini <i>et al.</i> (2007)
<i>Sturnus vulgaris</i> (sdm)	1982–2006	Northern Italy	1	Laying dates	−0.1	−0.2	Rubolini <i>et al.</i> (2007)
Birds (ldm)	1982–2006	Northern Italy	6	Arrival or laying dates	−1.8 $\pm$ 0.8	−1.0 $\pm$ 0.6	Rubolini <i>et al.</i> (2007)
<i>Ficedula hypoleuca</i> (ldm)	1960–1995	South Germany	1	Arrival dates	−0.7	0.06	Hölzinger (1997)
<i>Lagopus muta</i> (res.)	2000–2009	Northern French Alps	1	Median hatching date	−4.9	−4.2	Novoa <i>et al.</i> (2016)
Insects							
<i>Ips typographus</i>	1996–2005	Southeastern Alps	1	First observation	−21	NA	Faccoli (2009)
Lepidoptera, butterflies	1998–2010	Swiss lowlands	28	First observation	NA	−4.9 $\pm$ 2.0	Altermatt (2012)
<i>Apatura iris</i>	1982–2002	Forest near Basel, Switzerland	1	First observation	−10.5	−6.4	Dell <i>et al.</i> (2005)
Lepidoptera, butterflies	1991–2019	Switzerland	138	First observation	−6.6 $\pm$ 0.5	NA	<i>info fauna</i>
Odonata	1991–2019	Switzerland	40	First observation	−1.0 $\pm$ 0.5	NA	<i>info fauna</i>
Orthoptera	1991–2019	Switzerland	64	First observation	−4.3 $\pm$ 0.8	NA	<i>info fauna</i>
Amphibians							
Anura	1991–2019	Switzerland	10	First observation	−1.0 $\pm$ 2.2	NA	<i>info fauna</i>
Urodela	1991–2019	Switzerland	5	First observation	+2.5 $\pm$ 3.0	NA	<i>info fauna</i>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

1408

Reptiles

Squamata

1991–2019

Switzerland

14

First observation

−5.7±1.6

NA

info fauna

For Review Only

Table 2. Elevational range shifts reported for diverse taxa in the European Alps region. Values are means $\pm$ SE. In the ‘Limit’ column, ‘optimum’ is the elevation of maximum abundance and ‘leading edge’ is the upper elevational limit of a given species. These two metrics are calculated in different ways depending on the studies (see article references and Section II). For Dainese *et al.* (2017) only native species were included. For information on the *info fauna* database see Section II. The data set used for this table is provided as supporting online information in Table S2. *N*, number of species. Abbreviations: ecm, ectomycorrhizal; sapro., saprophytic.

Taxa	Study period	Area	<i>N</i>	Mean shift (m decade ⁻¹)	Limit	References
PLANTS						
Alpine species	1956–2004 ¹	Central Italian Alps	93	+18.1 $\pm$ 2.5	Leading edge	Parolo & Rossi (2008)
Alpine species	1989–2009	Northeast Italy	80	+26.1 $\pm$ 10.0	Optimum	Dainese <i>et al.</i> (2017)
Forest plant species	1977–2005 ¹	Western French Alps, and other French mountains	97	+38.1 $\pm$ 7.8	Leading edge	Kuhn <i>et al.</i> (2016)
Forest plant species	1977–2005 ¹	Western French Alps, and other French mountains	97	+25.7 $\pm$ 4.6	Optimum	Kuhn <i>et al.</i> (2016)
Forest plant species	1985–1999	Southeast France and Corsica	175	+13.8 $\pm$ 6.9	Optimum	Bodin <i>et al.</i> (2013)
Non-alpine plant species	1989–2009	Northeast Italy	1,128	+32.8 $\pm$ 3.2	Optimum	Dainese <i>et al.</i> (2017)
BIRDS						
Various species	1982–2017	Italian Alps	29	+31.5 $\pm$ 8.4	Optimum	Bani <i>et al.</i> (2019)
Various species	1982–2017	Italian Alps	29	+55.6 $\pm$ 9.5	Leading edge	Bani <i>et al.</i> (2019)
Various species	1993–2016	Switzerland	71	+12.0 $\pm$ 3.5	Optimum	Knaus (2018)
Various species	1993–2004 ¹	Italian Alps (Piedmont)	54	+7.2 $\pm$ 9.4	Optimum	Popy <i>et al.</i> (2010)
Forest species	1978–2002	French northern Alps	24	−7.8 $\pm$ 6.5	Optimum	Archaux (2004)
Forest species	1973–2001	French southern Alps	17	−6.3 $\pm$ 6.3	Optimum	Archaux (2004)
<i>Lagopus muta</i>	1998–2011	Northern French Alps	1	−24.7	Optimum	Novoa <i>et al.</i> (2016)
<i>Lagopus muta helvetica</i>	1984–2012	Swiss Alps	1	+37.8	Optimum	Pernollet <i>et al.</i> (2015)
MAMMALS						
Ungulate species	1991–2013	Swiss Alps	4	+37.5 $\pm$ 10.3	Optimum	Büntgen <i>et al.</i> (2017)
INSECTS						
<i>Lepidoptera</i>						

<i>Thaumetopoea pityocampa</i>	1975–2004	Italian Alps	1	+49.8	Leading edge	Battisti <i>et al.</i> (2005)
Various butterfly species	1991–2019	Switzerland	138	+39.9±6.5	Optimum	info fauna
Various butterfly species	1991–2019	Switzerland	138	+112.4±6.3	Leading edge	info fauna
Coleoptera						
Dung beetle species	1992–2007	South western Alps	30	+40.5±24.2	Optimum	Menéndez <i>et al.</i> (2014)
Dung beetle species	1992–2007	South western Alps	30	+80.6±39.8	Leading edge	Menéndez <i>et al.</i> (2014)
Odonata						
Various species	1991–2019	Switzerland	40	–2.6±5.9	Optimum	info fauna
Various species	1991–2019	Switzerland	40	–4.7±9.1	Leading edge	info fauna
Orthoptera						
Various species	1991–2019	Switzerland	64	+20.6±5.4	Optimum	info fauna
Various species	1991–2019	Switzerland	64	+46.9±6.9	Leading edge	info fauna
AMPHIBIANS						
Anura						
Various species	1991–2019	Switzerland	10	+13.9±4.8	Optimum	info fauna
Various species	1991–2019	Switzerland	10	+42.8±17.1	Leading edge	info fauna
Urodela						
Various species	1991–2019	Switzerland	7	+19.7±18.7	Optimum	info fauna
Various species	1991–2019	Switzerland	7	+53.5±16.7	Leading edge	info fauna
REPTILES						
Various species	1991–2019	Switzerland	14	+22.6±10.8	Optimum	info fauna
Various species	1991–2019	Switzerland	14	+63.7±12.7	Leading edge	info fauna
FISHES						
Stream fishes	1986–2006 ¹	Streams in France	32	+13.1±3.8	Optimum	Comte & Grenouillet (2013)
FUNGI						
ecm species	1960–2010	European Alps	56	+21.0±3.0	Optimum	Diez <i>et al.</i> (2020)
Litter/soil sapro. species	1960–2010	European Alps	29	+23.6±4.0	Optimum	Diez <i>et al.</i> (2020)
Wood sapro. species	1960–2010	European Alps	33	+10.7±2.6	Optimum	Diez <i>et al.</i> (2020)

¹Median years of the historical and modern survey.

Long-term time series of snow parameters (snowmelt timing and length of the snow free period) and mean air temperature (spring temperature from March to May and annual temperature) at six sites spread across the Swiss Alps during the period 1970–2019.

Data updated from Klein et al. (2016) representing yearly average anomalies over the period 1970–2019 across six sites in Switzerland located between 1,298 and 2,540 m (Arosa, Davos, Grächen, Grimsel Hospiz, Scuol and Weissfluhjoch). Estimated trends from the Theil-Sen estimator method across the six sites are reported with the associated standard errors (grey shading) for each snow and temperature parameter. Significance of the slopes was tested using Mann-Kendall tests, see Klein et al. (2016) for more details.

136x92mm (300 x 300 DPI)

Spring phenological change per decade (A) or per degree (B) for different plant and animal groups studied in the European Alps region. Values correspond to the estimated marginal mean of the mixed-effect model with study as a random factor; error bars are 95% confidence intervals. The period of time varied among studies but was always more than 10 years within the period 1980–2020. Only groups with more than five species are shown (all studies are reported in Table 1). Abbreviations: ldm, long-distance migrants; sdm, short-distance migrants; res., resident; semi-aq., semi-aquatic. Different letters among groups indicate significant differences (post-hoc Tukey tests at $\alpha = 0.05$).

527x944mm (72 x 72 DPI)

Empirical evidence for elevational range shifts of (A) the optimum distribution (i.e elevation of maximum abundance) and (B) the leading edge (i.e. the upper range limit) reported for groups of taxa in the European Alps region. Values correspond to the estimate of the mixed-effect model with study as a random factor; error bars are 95% confidence intervals. The period of time investigated varied among studies but was always more than 10 years within the period 1980–2020, or for diachronic studies comparing historical and modern surveys, the modern survey was conducted within the period 1980–2020, and the median year of the historical survey(s) was after 1950. Only groups with more than five species are shown (all studies are reported in Table 2). Abbreviations: ecm, ectomycorrhizal; sapro., saprophytic; semi-aq., semi-aquatic. Different letters among groups indicate significant differences (post-hoc Tukey tests at $\alpha = 0.05$).

583x1166mm (72 x 72 DPI)

Mean elevational shift per decade for four different orders of insects in terms of (A) the optimum elevation (i.e. elevation of maximum abundance) and (B) the leading-edge elevation (i.e. the upper range limit). Values correspond to the estimate of the mixed-effect model with study as a random factor; error bars are 95% confidence intervals. The period of time investigated varied among studies but was always higher than 10 years within the period 1980–2020. Different letters among groups indicate significant differences (post-hoc Tukey tests at $\alpha = 0.05$).

555x999mm (72 x 72 DPI)

Fig. S1. Mean phenological shift of the singing activity or laying dates per decade for birds classified according to their habitat. Values correspond to the estimated marginal mean of the mixed-effect model with study as a random factor; error bars are 95% confidence intervals. The period of time investigated varied among studies but was always more than 10 years within the period 1980–2020. Different letters among groups indicate significant differences (*post-hoc* Tukey tests at $\alpha = 0.05$).