Anti-Influenza Drug Discovery and Development: Targeting the Virus and Its Host by All Possible Means

Olivier Terrier, Anny Slama-Schwok

► To cite this version:

Olivier Terrier, Anny Slama-Schwok. Anti-Influenza Drug Discovery and Development: Targeting the Virus and Its Host by All Possible Means. Antiviral Drug Discovery and Development, pp.195-218, 2021, 10.1007/978-981-16-0267-2_8 . hal-03286916

HAL Id: hal-03286916

<https://hal.science/hal-03286916>

Submitted on 1 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Chapter X:

Anti-influenza drug discovery and development: *targeting the virus and its host by all possible means*

Olivier Terrier¹, Anny Slama-Schwok²

Abstract

Influenza infections remain a major and recurrent source of public health concern. Together with vaccines, antiviral drugs play a key role in the prevention and treatment of influenza virus infection and disease. Today, the number of antiviral molecules approved for the treatment of influenza is relatively limited and their use is threatened by the emergence of viral strains with resistance mutations. There is therefore a real need to expand the prophylactic and therapeutic arsenal. This chapter summarizes the state of the art in drug discovery and development for the treatment of influenza virus infections, with a focus on both virus-targeting and host-cell-targeting strategies. Novel antiviral strategies targeting other viral proteins or targeting the host cell, some of which are based on drug repurposing, may be used in combination to strengthen our therapeutic arsenal against this major pathogen.

Keywords: antiviral, drug repurposing, replication, entry, immune modulator

Abbreviations:

CoV: Coronavirus

COX: cyclo-oxygenase

HA: hemagglutinin,

IAV: Influenza A virus,

IFN: Interferon

M2: Matrix 2

NA: neuraminidase,

NOX: NADPH oxydase

NP: nucleoprotein,

p09: H1N1 2009-pandemic strain

PA: polymerase acidic subunit,

PB1: polymerase basic subunit 1,

PB2: polymerase basic subunit 2,

PPI: protein–protein interaction

RdRP: RNA-dependent ribonucleoprotein complex

RIG-I: retinoic acid inducible gene–I

TNF- α : Tumor necrosis factor- α

vRNP: viral ribonucleoproteins

-
1. CIRI, Centre International de Recherche en Infectiologie, (Team VirPath), Univ Lyon, Inserm, U1111, Université Claude Bernard Lyon 1, CNRS, UMR5308, ENS de Lyon, F-69007, Lyon, France. Email : olivier.terrier@univ-lyon1.fr
 2. Sorbonne Université, Centre de Recherche Saint-Antoine, INSERM U938, Biologie et Thérapeutique du Cancer, Paris, France. Email : Anny.Slama-Schwok@inserm.fr

1. Introduction

Influenza infections remain a major and recurrent source of public health concern. Influenza viruses are the causative agents of annual flu epidemics, marked by up to 1 billion infections and 300,000-650,000 deaths worldwide, with a huge economic burden in terms of hospitalization costs and work/school absenteeism (WHO, 2018; [56]). In addition, Influenza A viruses (IAV) have been the cause of several pandemics in recent human history, from the Spanish Flu H1N1 in 1918 to the more recent H1N1 2009 pandemic [71].

Together with vaccines, antiviral drugs play a vital part in the prevention and treatment of influenza virus infection and disease. During a normal influenza season, antiviral drugs are mainly used to treat critically ill patients, such as those hospitalized in intensive care. In a pandemic context, pending the availability of a vaccine, antiviral drugs are essential both to treat patients who have been infected and to prevent infection in those exposed, including healthcare workers. Today, the number of antiviral molecules approved for the treatment of influenza, based on the targeting of viral proteins, is relatively reduced and threatened by the emergence of strains with resistance mutations. There is therefore a real need to expand the prophylactic and reinforce the current therapeutic arsenal. This chapter summarizes the state of the art in drug discovery and development for the treatment of influenza virus infections, with a focus on both virus-targeting and host-cell-targeting strategies (**Figure 1**). Novel antiviral strategies targeting other viral proteins or targeting the host cell, some of which are based on drug repurposing, may be used in combination to strengthen our therapeutic arsenal against this major pathogen.

2- From existing classic antiviral drugs to new pre-clinical candidates

2.1 M2 ion channel blockers (*amantadine/rimantadine*)

Influenza A M2 is a multifunctional viral homo-tetramer protein [57]. Its transmembrane (TM) domain forms a proton channel. This channel is required for acidification of the viral endosome formed after fusion and endocytosis of the virus within the host cell. This process allows viral ribonucleoproteins (vRNPs) to dissociate from the matrix 1 (M1) protein. The proton conductance mechanism relies on the conserved H37XXXW41 sequence which is responsible for selectively gating H⁺ ions [89, 132], [52,

121]. Channel blockers interfere with the proton conductance mechanism by binding to the transmembrane pore [137] (**Figure 2**). When proton conductance through M2 is blocked by the adamantane drug, this dissociation is prevented and the virus is no longer able to replicate. In recent years, adamantane drug-resistant mutants have become prevalent in circulating viruses. The two most prevalent drug-resistant mutants are S31N, L26F and V27A, all of which are located in the transmembrane region of M2 [138]. **Figure 2A** shows the strong interaction of amantadine with V27 in the upper part of the pore. Upon drug-resistance V27A mutation, this interaction is lost. Recently-developed spiro-amantadyl amine effectively binds to A27 of the pore (**Figure 2B**) [136]. Recently, new amantadine derivatives effective against double mutants M2-S31N/L26I and M2-S31N/V27A viral strains have been developed by Musharrafieh et al [91]. The antiviral efficacy of such compounds is summarized In **Table 1**. M2 resistance mutations in H1N1/H3N2 circulating strains prompted the WHO to remove both amantadine and rimantadine from the list of recommended anti-influenza agents for clinical use in 2009 [34].

2.2 Neuraminidase (NA) and hemagglutinin (HA) inhibitors

2.2.1 NA inhibitors

NA inhibitors competitively inhibit and prevent cleavage of the terminal sialic acid residues from glycoproteins and carbohydrates displayed on the surface of mammalian cells and influenza virus particles. Binding of virions to uncleaved sialic acid then impairs virion release and dissemination. Among these NA inhibitors, peramivir, zanamivir, oseltamivir carboxylate are the most frequently prescribed drugs and considered standard-of-care for influenza management (**Table 1 and Figure 3**). Resistance to oseltamivir can develop rapidly in both experimental settings and the clinic, and typically originates from substitutions at signature resistance sites in the viral NA protein such as H274Y and I223R (predominant in H1N1 and H5N1 viruses), and E119V, R292K, or N294S (predominant in H3N2 viruses). These three NA inhibitors are currently licensed worldwide for the treatment of influenza A and B infections, oseltamivir being the most widely used. There is still a lot of debate about the effectiveness and real impact of inhibitors on the prevention and treatment of influenza. New oseltamivir

derivatives, targeting either multiple sites or different NA cavities (as the “430” or the “150” cavity) have been recently developed. Some of these derivatives are very potent against multiple IAV and IBV strains, including oseltamivir-resistant ones (**Table 1**).

2.2.2 Hemagglutinin inhibitors

The surface glycoprotein HA enables viral entry into host cells by binding to cell-surface, sialic-acid-containing glycans and mediating fusion between the viral and host membranes in endosomal compartments. HA is composed of head (HA1) and stem (HA2/HA1) domains. As the regions on HA involved in binding and fusion are highly conserved, they are attractive sites for the design of new antivirals (Table 2). The broad-spectrum antiviral drug arbidol shows efficacy against influenza viruses by targeting the hemagglutinin (HA) stem region [63]. This molecule is currently licensed in Russia and China for the treatment of influenza and other infections [8]. A challenging strategy aiming at mimicking antibodies binding sites was successfully developed by Wilson et al, targeting the conserved stem region and more recently at the interface of the trimeric head region [4, 141, 155] (**Figure 4A**). The binding sites of the binding sites for CBS1117 and JNJ4796 were both found in the stem region close to the fusion peptide, highlighting the possibility of further structure-based designed compounds [2]. De novo design of high-affinity trimeric proteins called “HA mini-binders” that bind influenza A hemagglutinin trimer at a conserved region binding site (**Figure 4B**) [129]. These molecules were developed as alternative to antibodies. These and other compounds are summarized in **Table 2**.

2.3 Polymerase – nucleoprotein- RNA inhibitors

2.3.1 Polymerase/endonuclease inhibitor (favipiravir, Baloxavir marboxil)

Influenza virus transcribe and replicate their genome in the nucleus of infected cells by the means of a hetero-trimeric polymerase, PA, PB1 and PB2. The polymerase complex function requires the nucleoprotein NP, a protein associated with- and protecting the segmented genomic RNA. Therefore, all four proteins are essential for replication. Whereas replication requires the generation of complementary positive polarity RNA intermediates (cRNA) that are then copied into progeny negative

polarity segments (vRNPs), viral message is directly synthesized from vRNPs. Since the influenza virus RdRP lacks enzymatic activity to form 5' mRNA cap structures, endonuclease activity of the PA subunit is necessary for the generation of viral mRNAs through transfer of 5'-capped RNA primers derived from host mRNAs in a cap-snatching mechanism. The endonuclease active site of PA-N terminal comprises a histidine and a cluster of three strictly conserved acidic residues (Glu80, Asp108, Glu119), which coordinate (together with Ile120) one or two manganese or magnesium ions [32] (**Figure 5A**). PB2 is involved in binding of the capped primers, whereas the PB1 subunit harbors enzymatic activity for phosphodiester bond formation. Several classes of inhibitors are in the clinics (**Figure 6**): baloxavir (PA), favipiravir (PB1) and pimodivir (PB2, **Figure 5B**).

2.3.2 Pre-clinical compounds targeting the polymerase PA-PB1 and PA subunits; escape mutations and resistance.

Pre-clinical candidates, some of them being listed in Tables 3 and 4, are in development, benefiting from the recent insight provided by the structures of PA-PB1, PB1-PB2 and whole polymerase complex with or without RNA by X-ray crystallography [26, 48, 95, 102, 110, 130, 146] and cryo-electron microscopy [23, 42, 44, 111, 144]. The error-prone nature of influenza viral replication can rapidly generate point mutants for the selection of resistance that have seriously compromised the efficacy of influenza therapeutics. Escape mutations were identified under the pressure selection of PA inhibitors: the signature hotspot for escape from baloxavir marboxil is PA residue 38, for which several substitutions (PA I38T/M/F) have been described [96]. Similarly, escape mutations from L-742.001[127] and RO-7 [70] treatments were also characterized although in laboratory resistance-assays, escape mutants were not detected after multiple passages for L-742.001. While very tight affinities have been achieved by designing metal binding inhibitors to block the active site of the endonuclease activity in PA N-terminal (**Table 2**), the appearance of escape mutants often rapidly decrease their efficacy. Several recent reviews focus on the development of PA and polymerase inhibitors [61, 62, 87, 170].

Different strategies have been undertaken to attempt overcoming induced resistance. Interfering with its proper assembly of the RdRP polymerase to inhibit function is pursued using protein–protein interaction (PPI) inhibitors. The advantage of such an approach is the relatively large interacting surface between the two proteins as compared to the binding site of an active-site ligand. Indeed, inducing simultaneous mutation of at least one residue on both proteins while maintaining their interaction is less likely to develop resistance and suggests that PPI inhibitors could be less prone to drug resistance than inhibitors of enzyme active sites. The recent identification of a single- domain antibody (nanobody) allowing to disrupt dimerization of FluA polymerase is among these lines [42]. PPI inhibitors have been developed based on the structural insight given by PA–PB1 crystal structures in 2012 [81]. The inhibition of the polymerase PA–PB1 subunit interface has become an active field of research with the goal of remaining active against resistant strains to amantadine and to oseltamivir (**Table 3**). Recently, compound 12 was identified by structure-based screening of compounds targeting the PA-PB1 structure. No resistant virus was selected in vitro under drug selection pressure of compound 12a [164]. Moreover, derivatives of cyclothiophene and R151785 were found active against multiple strains of Influenza A and B [31, 94, 165].

Based on the ability of PA-PB1 to bind viral RNA, it is likely that novel types of inhibitors could be developed by structure-based design [131]. Additionally, inhibitors targeting PA C-terminal [78] and its interactions with vRNA or with PolII could be effective targets, based on the accumulating wealth of structural data [42, 102, 110, 144] and deeper insight in the multi-protein assembly required for during replication / transcription.

2.3.4 Broad-spectrum inhibitors

Favipavir is a drug with broad-spectrum antiviral activity in cell culture, inhibiting RNA viruses of the arenavirus, bunyavirus, flavivirus, alphavirus, norovirus, picornavirus, paramyxovirus, and rhabdovirus families, in addition to influenza viruses [160]. This drug is incorporated into newly synthesized RNA by the viral polymerase in place of purines but not pyrimidines, resulting in increased frequencies of C-to-U and G-to-A transition mutations. Although the barrier for resistance is relatively high, this drug seems to present toxicity issues. N 4-hydroxycytidine (NHC) is also a broad-spectrum antiviral

candidate, which showed oral efficacy against RSV and both highly-pathogenic avian and seasonal influenza viruses as well as SARS-CoV-2 virus [140].

2.3.5 Pre-clinical compounds targeting the polymerase PB2 subunit

Crystal structure of the PB2 cap-binding domain have been exploited to develop different 7-methylguanine derivatives [100]. Pimodivir (VX-787) is an inhibitor targeting the polymerase PB2 subunit at the m⁷ GTP-binding site, forming extensive stacking interactions with several aromatic residues His (**Figures 5B and 6**). It inhibits influenza virus replication and reduced viral load in animal infection models of H3N2 and H1N1 viruses, although potency was highest against H1N1 strains [9, 20]. Phase-2 clinical studies indicated that this drug is well-tolerated, reduced viral load, and resulted in slightly faster resolve of clinical signs. Further derivatives of pimodivir have been designed [84]. Targeting the PB1-PB2 interface by PPI inhibitors has been challenging: although PP7 exhibited antiviral activities against influenza virus subtypes A pandemic H1N1, H7N9 and H9N2, resistances have been unexpectedly detected in laboratory assays [162].

2.3.6 Pre-clinical compounds targeting the nucleoprotein or the nucleoprotein-RNA interactions

The nucleoprotein associated with viral RNA and the polymerase complex is essential for transcription and replication [22, 145, 146]. The assembly of NP-RNA oligomers into RNP has been determined by cryo-electron microscopy studies [3, 22, 23, 146]. In the X-rays structures of the NP [156], the protein adopts a trimeric structure. NP self-association to achieve trimer formation is mediated by a flexible tail-loop that protrudes into a pocket of the adjacent subunit, via the formation of a critical interaction between R416 of one subunit and E339 of the adjacent subunit. The R416A mutant lacking this interaction adopts a monomeric structure [16]. The native protein can also be purified in a monomeric form at low salt and concentration conditions [15, 16, 133]. The ability to modify the oligomeric state of NP is the structural basis of most NP inhibitors presently developed. Nucleozin was the first NP inhibitor developed as a molecule impeding nuclear accumulation. Nucleozin enhanced higher order structures [64][46]. **Figure 5C** shows the interactions of one of the nucleozin ligands found in the X-

ray structure (PDB ID 5B7B) stabilizing the interface between two NP subunits [98]. Escape mutants to nucleozin have been identified in laboratory assays. The opposite approach to impede nucleoprotein self-association has also been pursued by disrupting the important salt bridge R416-E339 mediating NP oligomerization [123]. Recently, new compounds with high affinity for NP were designed stabilizing monomeric NP [150]. Impeding NP binding to viral RNA has been achieved by naproxen drug repurposing, naproxen being a known inhibitor of cyclo-oxygenase (COX)[73]. As NP oligomerization is enhanced by the presence of RNA, naproxen binding to NP reduced NP oligomers and favored monomeric NP. Docking and single mutations studies identified Tyr148, the only aromatic residue within the RNA binding groove and residues of the C-terminal part of NP R355, R361, Phe489 being involved in the interaction of naproxen with NP. Laboratory assays showed no resistance after 8 cell passages infected with Influenza A. Naproxen exhibited antiviral effects in mice models of Influenza A infection [33, 73] as well as Influenza B virus [168]. Further structure-based design yielded new naproxen derivatives with improved antiviral effects and selectivity for NP without COX inhibition (**Figures 5D and 6**) [33, 134] (**Table 4**). Some of these derivatives were found inhibiting NP-PA interactions [33, 143]. Naproxen derivatives also present antiviral properties against oseltamivir-resistant strains [33]. Additional compounds with some similarity of their hydroxyquinoline scaffold to the methoxy naphthalene scaffold of naproxen called NUD were designed and were also found to be resistant in escape mutation laboratory assays [79].

2.4 Drugs targeting the non-structural protein-1 (NS1)

NS1 has a plethora of strategies to inhibit the host immune response due to its ability to establish multiple protein–protein and protein–RNA interactions. NS1 hampers different pathways both in the cytoplasm and in the nucleus of infected cells. NS1 antagonizes interferon-mediated antiviral host response by binding to double-stranded (ds) viral RNA, thus protecting it from cellular factors, by blocking retinoic acid inducible gene-I (RIG-I) and NF- κ B activation. One pathway by which NS1 increases virulence is through the activation of phosphoinositide 3-kinase (PI3K) by binding to its p85 β subunit [17]. NS1 has two structural domains—RNA-binding domain (RBD) and the effector domain

(ED)—connected by a short linker (LR), and a disordered C-terminal tail. New drugs binding to NS1 effector domain have been designed with low micromolar antiviral efficacy [68] (**Table 4**).

3. Host-targeting & drug repurposing approaches for the treatment of influenza

Considerable progress has been made in understanding the interactions between influenza viruses and the host cell in recent years. In this context, and in light of the emerging problem of resistance to available classical antivirals, many studies have focused on targeting host factors to limit virus replication, but also to modulate host immune response. The targeting of host-factors and/or signaling pathways makes sense in the context of virally-induced hypercytokinemia (also known as “cytokine storm”), which is directly correlated with tissue injury and an unfavorable prognosis of severe influenza [76]. Indeed, approaches to control or attenuate this disproportionate immune response are of particular interest and are the subject of numerous pre-clinical and clinical studies. As with all viruses, influenza viruses depend on cellular machinery for their replication and propagation. Many cellular factors essential for the replication of influenza viruses have been uncovered through genome-wide RNA interference approaches [65, 69, 86, 128] but also more broadly through different integrated cell biology approaches using interactome and transcriptome data, for example [109, 148]. In order to list the different host-targeting strategies developed, a distinction can be made between molecules with a mode of action associated with a relatively well-defined stage of the viral cycle, and molecules associated with the modulation of signaling pathways. It is these two main classes that will be described in the following sections.

3.1 Drugs targeting host cell component at different stages of influenza replication cycle

The replication cycle of influenza viruses can be divided into several distinct phases, 1) entry 2) nuclear import of viral genome (viral ribonucleoprotein; vRNPs) 2) genome replication and protein synthesis, 3) Nucleo-cytoplasmic export of vRNPs, and 4) plasma membrane transport and budding of neo-virions (**Figure 1**). A number of molecules targeting host factors in these different steps, at different preclinical/clinical development stages, are known today.

Viral entry is a target of great interest, as it is likely to allow prophylactic approaches, by blocking the infection in its early stages. One of the most advanced strategy consists to target the viral receptor. DAS181 (**Table 5**) (Fludase, Ansun BioPharma) is a sialidase fusion protein that cleaves both the Neu5Ac $\alpha(2,3)$ - and Neu5Ac $\alpha(2,6)$ -Gal linkages of sialic acid on host cells. DAS181 is administered as an inhalable dry powder to deliver sialidase to the pulmonary epithelium for cleavage of sialic acids, which renders the cells inaccessible to infection by virus [80]. DAS181 was demonstrated to have broad-spectrum activity, given the conserved nature of influenza and parainfluenza viruses binding to respiratory epithelium. Preclinical in vitro and in vivo studies demonstrated that DAS181 has activity against a number of seasonal influenza strains including those containing the H274Y mutation (conferring resistance to oseltamivir), highly pathogenic avian influenza strains (H5N1), and pandemic 2009 influenza A (H1N1). This compound was assessed in different Phase I and Phase II clinical trials (NCT 00527865, NCT 01651494, NCT01037205) with results indicating a significant reduction of viral load in treated influenza patients [88] but with identification of respiratory adverse events and rapid clearance of the drug being consistent with the induction of antibodies against DAS-181 - this could be a limitation in the duration and dosages of such treatment [163]. Other approaches targeting viral entry have also been described (**Table 5**), e.g. targeting the endosome acidification step by inhibition of V-ATPase (ex: bafilomycin A1, concanamycin), or inhibition of the internalization (ex: Dynasore) or cleavage steps of haemagglutinin (ex: camostat). Most of these strategies were primarily evaluated at the preclinical stage and have not been further evaluated as their efficacy was either limited or accompanied by cytotoxicity. One exception is the protease inhibitor aprotinin, which was approved as anti-influenza drug in Russia [169].

The step of **nuclear import of vRNPs** is a crucial one, for which there are today very few molecules with antiviral potential described in literature. Interestingly, it has been shown in vitro that ivermectin (**Table 5**), a well-known anti-parasite drug, was able to inhibit viral replication via inhibition of importins (IMP α/β), and therefore the nuclear import of vRNPs [47].

Targeting the **replication** stage of the virus is one of the earliest host-targeting strategies, with pioneer works on the antiviral efficacy of ribavirin in the 1970s [38]. However, this nucleoside analogue or its

prodrug, less toxic, do not appear to be options being considered for the treatment of influenza virus infections. of influenza viruses, despite interesting preliminary in vitro and in vivo results [125]. (**Table 5**). Other, more recent strategies propose to target **mRNA splicing**. Influenza viruses are known to hijack cellular splicing machinery to their benefit, making them extremely dependent on it [36, 37]. Several studies show that the inhibition of Cdc2-like kinase 1 (CLK1), involved in the alternative splicing of M2 gene of influenza, appears to be an interesting antiviral option, with several molecules available (TG003, Clypearin, Corilagin, **Table 5**). Of all its molecules, Clypearin has relatively low EC50s and very low toxicity, making it an attractive potential antiviral candidate. [65, 171].

While strategies to prevent the nuclear import of vRNPs are relatively uncommon, paradoxically there are many more therapeutic approaches to block the **nuclear-cytoplasmic transport of vRNPs**. Indeed, in contrast to the inhibition of importins, the inhibition of exportin 1 (XPO1) by Verdinexor (XPO1 antagonist KPT-335) allows to significantly reduce viral production in vitro and in vivo [101]. Another compound, DP2392-E10, inhibits nuclear export of both viral NP and nuclear export protein (NEP). More specifically, in vitro pull-down assays revealed that DP2392-E10 directly binds cellular CRM1, which mediates nuclear export of NP and NEP – highlighting CRM1 as a target of interest [19]. With the same objective, other strategies consist to target the Raf/MEK/ERK signaling pathway, known to be involved in the export of vRNPs [120]. Several MEK inhibitor molecules have been studied for their ability to inhibit the replication of influenza viruses, such as CI-1040 or U0126 [24, 108]. Interestingly, Schröder and colleagues have demonstrated that Trametinib (GSK-1120212), a licensed MEK inhibitor used for the treatment of malignant melanoma, efficiently blocks influenza viral replication of different subtypes in vitro and in vivo [119] (**Table 5**).

Apical transport and budding, the last part of the last major step of the replication cycle is also the object of several antiviral strategies, notably by blocking the transport of viral proteins to the plasma membrane (ex: Clonidine; [82]), or the cholesterol pathway, which would reduce virion egress. (U18666A; [92]). One of the most advanced strategies is Nitazoxanide, initially licensed for the treatment of parasitic infections, for which anti- influenza properties were first documented by Rossignol et al. [113]. Interestingly, the proposed mode of action of nitazoxanide toward influenza is clearly distinct to that for which it was designed in its initial indication, acting at the post-translational

level by selectively blocking the maturation of the viral glycoprotein HA, with a consecutive impact on its intracellular trafficking and insertion into the host plasma membrane [112]. This drug presents potent antiviral activity against a large panel of circulating strains [139]. The effectiveness of nitazoxanide in treating patients with non-complicated influenza was successful in a Phase IIb/III trial [50] and is currently being assessed in a Phase III clinical trial (NCT01610245).

3.2 Drugs targeting host cell signaling pathways and host-response that are crucial for influenza replication cycle.

Our increased knowledge of signaling pathways that are crucial in the response to infection and/or those hijacked by the virus has allowed many research teams to explore complementary antiviral strategies that can be described here (**Table 6**). The targeting of the raf/MEK/ERK channel, mentioned above, could of course also have been listed here. At the crossroads of the regulatory pathways of the immune response and the stress response, the **NF-kB pathway** was one of the first to be studied (**Table 6**). In the context of cell biology approaches, it was initially shown that the anti-inflammatory drug acetylsalicylic acid (ASA) had interesting antiviral effect against influenza viruses in vitro and in vivo, via inhibition of the NF-kB activating IkkB kinase [83, 153, 159]. Several drugs targeting the NF-kB pathway have been evaluated since then, such as pyrrolidine dithiocarbamate or SC7574; with encouraging in vivo results [39, 49, 149]. BAY81-8781/LASAG (D, L-Lysine acetylsalicylate-glycine) (**Table 6**), a modified version of ASA demonstrates antiviral activity against several human and avian influenza viruses in vitro. In a mouse infection model, inhalation of LASAG resulted in reduced lung viral titers and protection of mice from lethal infection [35]. More recently, a Phase II proof-of-concept study comparing LASAG versus placebo in patients with severe influenza demonstrated that aerosolized LASAG improved the time to symptom alleviation compared to placebo, despite the absence of a statistically significant reduction of viral load in LASAG-treated group [116].

Based on clinical observations, hydroxyl methylglutaryl-coenzyme A (**HMG-CoA**) reductase inhibitors such as statins (**Table 6**), approved for their use as cholesterol metabolism regulators, have demonstrated pleiotropic anti-inflammatory and immunomodulatory properties, which could be of benefit to improve survival of patients with severe influenza [43, 85]. However, most in vivo studies reported so far failed

to clearly demonstrate such a beneficial effect in the specific context of influenza infection [6, 72, 115].

On the other hand, a few observational studies highlighted an association between statin treatment with a reduction of mortality in patients hospitalized with laboratory-confirmed seasonal influenza [40, 142].

A randomized placebo-controlled Phase II clinical trial (NCT02056340) aimed at evaluating the potential effect of atorvastatin to reduce the severity of illness in influenza-infected patients is currently undergoing. The combination of naproxen with clarithromycin and oseltamivir twice daily reduced the both 30- and 90-day mortality and length of hospital stay of patients hospitalized for A(H3N2) influenza [54]. Other approaches, at the preclinical validation stages, propose to target the **TNF-alpha** (Etanercept); or **NOX2** (Apocynin/Ebselen) or lipoygenase/COX pathway (Celecoxib/Mesalazine) pathways [12, 28, 74, 97, 124, 157, 167]. A phase III clinical trial is currently investigating the benefit of celecoxib for the treatment of severe influenza (NCT02108366). These molecules could be of interest to better control the inflammatory response, which is a very important aspect of the pathology.

Modulation of immune and inflammatory responses is a therapeutic avenue that has been much explored, but which may present risks given the ambivalent aspect of these pathways in relation to viral replication and the evolution of the pathology. Indeed, such treatment should stimulate induction of antiviral genes to control IAV spread, without driving immunopathology. In this context, **IFN-lambda** (**Table 6**) appears as a potent anti-influenza therapeutic, without the inflammatory side effects of IFN-alpha treatment [29]. Intranasal administration of IFN-λ2/3 was shown to significantly suppress infection of various influenza strains, including WS/33 (H1N1), PR (H1N1), and H5N1 in the mouse lung, and was accompanied by greater up-regulation of ISGs [67]. More recently, using a transcriptome-based screening approaches, we identified and validated diltiazem, a calcium channel blocker used as an anti-hypertensive drug, as a very promising host-targeted inhibitor of influenza infection. Interestingly, the study of the mode of action revealed that diltiazem was a strong induced or type III IFN [107]. An ongoing French multicenter randomized clinical trial is investigating the effect of diltiazem- oseltamivir bi-therapy compared with standard oseltamivir monotherapy for the treatment of severe influenza infections in intensive care units (FLUNEXT trial NCT03212716).

4. Perspectives and concluding remarks

Among all the molecules listed in this chapter, some are already available on the market for other therapeutic indications and fall within the scope of drug repurposing. This is the case for naproxen, diltiazem, LASAG or Nitazoxanide, for example. The basis of drug repurposing relies on bypassing the long, risky and expensive preclinical and early clinical evaluation stage conventionally used for de novo drug development and exploiting available extensive human clinical, pharmacokinetics and safety data as the starting point for the development [106]. All these aspects make the repositioning of drugs a very interesting approach, in particular to enable a rapid response to the need for new antiviral strategies in the context of the emergence of a virus with pandemic potential.

Another very interesting perspective is the interest in combining different antiviral approaches with each other, including classical approaches targeting the virus with those targeting the host cell. The concept of combining therapies has already been used successfully, notably in the design of antiretroviral treatments [13]. Combination therapy can have several objectives, such as reducing the risk of the emergence of resistance by simultaneously targeting several viral proteins and/or key host factors, but also increasing the effectiveness of the treatments by obtaining additive or synergistic effects.

While there is relatively little convincing evidence to support the use of conventional virus-targeting antivirals in combination [41, 103], there are interestingly a growing number of examples of combinations of combination host-targeted approaches with oseltamivir. For example, we have shown that the combination of diltiazem and oseltamivir provides a much greater reduction in viral titers in a reconstructed human epithelium model compared to single treatments [107]. More recently Schloer and colleagues have shown that a combination treatment of an antifungal molecule, itraconazole, with oseltamivir, achieves much greater antiviral activity compared to monotherapy, making it possible to consider reducing the concentrations of drugs used, and thus possibly reducing the problems of adverse effects and emergence of resistance mutations [117]. These results open up interesting prospects for the development of future therapeutic strategies, particularly for the treatment of severe forms of influenza. The potential arsenal for fighting influenza virus infections is potentially very extensive, in particular thanks to the combination of new molecules targeting the virus, resulting from docking and structure-based design strategies, with approaches targeting cellular factors and signaling pathways. In this

context, the quality and relevance of the preclinical models, as well as the quality of the tools for evaluating combinations of molecules, are important critical elements.

Beyond influenza viruses, many of the antiviral molecules described in this chapter have the potential for broader-spectrum use. Indeed, some virus-targeted strategies can target viral determinants with very strong similarities between different viruses. This is particularly the case with Naproxen for which we have previously demonstrated antiviral activity against both influenza viruses and SARS-CoV-2 [73, 135]. This property is explained by the fact that the nucleoproteins N of enveloped, positive-sense, single-stranded viruses Coronavirus (CoV) share with negative-sense single-stranded viruses such as Influenza A virus the ability to bind to- and protect genomic viral RNA without sequence specificity and to form self-associated oligomers. Despite their differences, viruses induce and divert many common cellular pathways. As a result, host-targeted approaches can identify molecules with a broad spectrum of antiviral activity. An example is diltiazem, for which we have shown antiviral activity against influenza viruses [107], but which has been shown to be effective against other respiratory viruses, such as SARS-CoV-2 [104, 105], due to its mode of action involving the type III interferon response. Efforts to identify anti-influenza molecules therefore open up very interesting prospects for the broader development of antivirals. In many ways, antiviral research on influenza viruses is pioneering in this area and provides a starting point for the study of other emerging viruses.

5. References

Influenza (seasonal). World Health Organization. Available online at :[http://www.who.int/fact-sheets/details/influenza-\(seasonal\)](http://www.who.int/fact-sheets/details/influenza-(seasonal)) (accessed July 18, 2020).

1. Ai W, Zhang J, Zalloum WA et al. (2020) Discovery of novel "Dual-site" binding oseltamivir derivatives as potent influenza virus neuraminidase inhibitors. *Eur J Med Chem* 191:112147
2. Antanasijevic A, Durst MA, Cheng H et al. (2020) Structure of avian influenza hemagglutinin in complex with a small molecule entry inhibitor. *Life Sci Alliance* 3
3. Arranz R, Coloma R, Chichon FJ et al. (2012) The structure of native influenza virion ribonucleoproteins. *Science* 338:1634-1637
4. Bangaru S, Lang S, Schotsaert M et al. (2019) A Site of Vulnerability on the Influenza Virus Hemagglutinin Head Domain Trimer Interface. *Cell* 177:1136-1152 e1118
5. Basu A, Komazin-Meredith G, McCarthy C et al. (2017) Molecular Mechanism Underlying the Action of Influenza A Virus Fusion Inhibitor MBX2546. *ACS Infect Dis* 3:330-335
6. Belser JA, Szretter KJ, Katz JM et al. (2013) Simvastatin and oseltamivir combination therapy does not improve the effectiveness of oseltamivir alone following highly pathogenic avian H5N1 influenza virus infection in mice. *Virology* 439:42-46
7. Bimbo LM, Denisova OV, Makila E et al. (2013) Inhibition of influenza A virus infection in vitro by saliphenylhalamide-loaded porous silicon nanoparticles. *ACS Nano* 7:6884-6893
8. Blaising J, Polyak SJ, Pecheur EI (2014) Arbidol as a broad-spectrum antiviral: an update. *Antiviral Res* 107:84-94
9. Byrn RA, Jones SM, Bennett HB et al. (2015) Preclinical activity of VX-787, a first-in-class, orally bioavailable inhibitor of the influenza virus polymerase PB2 subunit. *Antimicrob Agents Chemother* 59:1569-1582
10. Cady SD, Schmidt-Rohr K, Wang J et al. (2010) Structure of the amantadine binding site of influenza M2 proton channels in lipid bilayers. *Nature* 463:689-692
11. Carcelli M, Rogolino D, Gatti A et al. (2016) N-acylhydrazones inhibitors of influenza virus PA endonuclease with versatile metal binding modes. *Sci Rep* 6:31500
12. Carey MA, Bradbury JA, Reboloso YD et al. (2010) Pharmacologic inhibition of COX-1 and COX-2 in influenza A viral infection in mice. *PloS One* 5:e11610
13. Chaudhuri S, Symons JA, Deval J (2018) Innovation and trends in the development and approval of antiviral medicines: 1987-2017 and beyond. *Antiviral Res.* 155:76-88
14. Chen HW, Cheng JX, Liu MT et al. (2013) Inhibitory and combinatorial effect of diphyllin, a v-ATPase blocker, on influenza viruses. *Antiviral Res* 99:371-382
15. Chenavas S, Crepin T, Delmas B et al. (2013) Influenza virus nucleoprotein: structure, RNA binding, oligomerization and antiviral drug target. *Future Microbiol* 8:1537-1545
16. Chenavas S, Estrozi LF, Slama-Schwok A et al. (2013) Monomeric nucleoprotein of influenza A virus. *PLoS Pathog* 9:e1003275
17. Cho JH, Zhao B, Shi J et al. (2020) Molecular recognition of a host protein by NS1 of pandemic and seasonal influenza A viruses. *Proc Natl Acad Sci U S A* 117:6550-6558
18. Choi MS, Heo J, Yi CM et al. (2016) A novel p38 mitogen activated protein kinase (MAPK) specific inhibitor suppresses respiratory syncytial virus and influenza A virus

- replication by inhibiting virus-induced p38 MAPK activation. *Biochem Biophys Res Commun* 477:311-316
19. Chutiwitoonchai N, Mano T, Kakisaka M et al. (2017) Inhibition of CRM1-mediated nuclear export of influenza A nucleoprotein and nuclear export protein as a novel target for antiviral drug development. *Virology* 507:32-39
20. Clark MP, Ledeboer MW, Davies I et al. (2014) Discovery of a novel, first-in-class, orally bioavailable azaindole inhibitor (VX-787) of influenza PB2. *J Med Chem* 57:6668-6678
21. Collins PJ, Haire LF, Lin YP et al. (2008) Crystal structures of oseltamivir-resistant influenza virus neuraminidase mutants. *Nature* 453:1258-1261
22. Coloma R, Arranz R, De La Rosa-Trevin JM et al. (2020) Structural insights into influenza A virus ribonucleoproteins reveal a processive helical track as transcription mechanism. *Nat Microbiol* 5:727-734
23. Coloma R, Valpuesta JM, Arranz R et al. (2009) The structure of a biologically active influenza virus ribonucleoprotein complex. *PLoS Pathog* 5:e1000491
24. Courtin N, Fotso AF, Fautrad P et al. (2017) Antiviral activity of formyl peptide receptor 2 antagonists against influenza viruses. *Antiviral Res.* 143:252-261
25. Credille CV, Morrison CN, Stokes RW et al. (2019) SAR Exploration of Tight-Binding Inhibitors of Influenza Virus PA Endonuclease. *J Med Chem* 62:9438-9449
26. Crepin T, Dias A, Palencia A et al. (2010) Mutational and metal binding analysis of the endonuclease domain of the influenza virus polymerase PA subunit. *J Virol* 84:9096-9104
27. D'agostino I, Giacchello I, Nannetti G et al. (2018) Synthesis and biological evaluation of a library of hybrid derivatives as inhibitors of influenza virus PA-PB1 interaction. *Eur J Med Chem* 157:743-758
28. Davidson S (2018) Treating Influenza Infection, From Now and Into the Future. *Frontiers in Immunology* 9:1946
29. Davidson S, McCabe TM, Crotta S et al. (2016) IFN λ is a potent anti-influenza therapeutic without the inflammatory side effects of IFN α treatment. *EMBO Mol Med* 8:1099-1112
30. De Vries E, Tscherne DM, Wienholts MJ et al. (2011) Dissection of the influenza A virus endocytic routes reveals macropinocytosis as an alternative entry pathway. *PLoS Pathog* 7:e1001329
31. Desantis J, Nannetti G, Massari S et al. (2017) Exploring the cycloheptathiophene-3-carboxamide scaffold to disrupt the interactions of the influenza polymerase subunits and obtain potent anti-influenza activity. *Eur J Med Chem* 138:128-139
32. Dias A, Bouvier D, Crepin T et al. (2009) The cap-snatching endonuclease of influenza virus polymerase resides in the PA subunit. *Nature* 458:914-918
33. Dilly S, Fotso Fotso A, Lejal N et al. (2018) From Naproxen Repurposing to Naproxen Analogues and Their Antiviral Activity against Influenza A Virus. *J Med Chem* 61:7202-7217
34. Dong G, Peng C, Luo J et al. (2015) Adamantane-resistant influenza A viruses in the world (1902-2013): frequency and distribution of M2 gene mutations. *PLoS One* 10:e0119115
35. Droebner K, Haasbach E, Dudek SE et al. (2017) Pharmacodynamics, Pharmacokinetics, and Antiviral Activity of BAY 81-8781, a Novel NF-kappaB Inhibiting Anti-influenza Drug. *Front Microbiol* 8:2130
36. Dubois J, Terrier O, Rosa-Calatrava M (2014) Influenza viruses and mRNA splicing: doing more with less. *mBio* 5:e00070-00014

- 474 37. Dubois J, Traversier A, Julien T et al. (2019) The Nonstructural NS1 Protein of
475 Influenza Viruses Modulates TP53 Splicing through Host Factor CPSF4. *J. Virol.* 93
- 476 38. Durr FE, Lindh HF, Forbes M (1975) Efficacy of 1-beta-D-ribofuranosyl-1,2,4-triazole-
477 3-carboxamide against influenza virus infections in mice. *Antimicrob. Agents*
478 *Chemother.* 7:582-586
- 479 39. Ehrhardt C, Rückle A, Hrincius ER et al. (2013) The NF- κ B inhibitor SC75741
480 efficiently blocks influenza virus propagation and confers a high barrier for
481 development of viral resistance. *Cell. Microbiol.* 15:1198-1211
- 482 40. Enserink M (2005) Infectious disease. Old drugs losing effectiveness against flu; could
483 statins fill gap? *Science (New York, N.Y.)* 309:1976-1977
- 484 41. Escuret V, Cornu C, Boutitie F et al. (2012) Oseltamivir-zanamivir bitherapy compared
485 to oseltamivir monotherapy in the treatment of pandemic 2009 influenza A(H1N1) virus
486 infections. *Antiviral Res.* 96:130-137
- 487 42. Fan H, Walker AP, Carrique L et al. (2019) Structures of influenza A virus RNA
488 polymerase offer insight into viral genome replication. *Nature* 573:287-290
- 489 43. Fedson DS (2013) Treating influenza with statins and other immunomodulatory agents.
490 *Antiviral Res.* 99:417-435
- 491 44. Fodor E, Te Velthuis AJW (2019) Structure and Function of the Influenza Virus
492 Transcription and Replication Machinery. *Cold Spring Harb Perspect Med*
- 493 45. Gaisina IN, Peet NP, Cheng H et al. (2020) Optimization of 4-Aminopiperidines as
494 Inhibitors of Influenza A Viral Entry That Are Synergistic with Oseltamivir. *J Med*
495 *Chem* 63:3120-3130
- 496 46. Gerritz SW, Cianci C, Kim S et al. (2011) Inhibition of influenza virus replication via
497 small molecules that induce the formation of higher-order nucleoprotein oligomers.
498 *Proc Natl Acad Sci U S A* 108:15366-15371
- 499 47. Götz V, Magar L, Dornfeld D et al. (2016) Influenza A viruses escape from MxA
500 restriction at the expense of efficient nuclear vRNP import. *Scientific Reports* 6:23138
- 501 48. Guilligay D, Tarendeau F, Resa-Infante P et al. (2008) The structural basis for cap
502 binding by influenza virus polymerase subunit PB2. *Nat Struct Mol Biol* 15:500-506
- 503 49. Haasbach E, Reiling SJ, Ehrhardt C et al. (2013) The NF-kappaB inhibitor SC75741
504 protects mice against highly pathogenic avian influenza A virus. *Antiviral Res.* 99:336-
505 344
- 506 50. Haffizulla J, Hartman A, Hoppers M et al. (2014) Effect of nitazoxanide in adults and
507 adolescents with acute uncomplicated influenza: a double-blind, randomised, placebo-
508 controlled, phase 2b/3 trial. *The Lancet. Infectious Diseases* 14:609-618
- 509 51. Harada S, Yokomizo K, Monde K et al. (2007) A broad antiviral neutral glycolipid,
510 fattiviracin FV-8, is a membrane fluidity modulator. *Cell Microbiol* 9:196-203
- 511 52. Hu F, Luo W, Hong M (2010) Mechanisms of proton conduction and gating in influenza
512 M2 proton channels from solid-state NMR. *Science* 330:505-508
- 513 53. Hu Y, Zhang J, Musharrafieh RG et al. (2017) Discovery of dapivirine, a nonnucleoside
514 HIV-1 reverse transcriptase inhibitor, as a broad-spectrum antiviral against both
515 influenza A and B viruses. *Antiviral Res* 145:103-113
- 516 54. Hung IFN, To KKW, Chan JFW et al. (2017) Efficacy of Clarithromycin-Naproxen-
517 Oseltamivir Combination in the Treatment of Patients Hospitalized for Influenza
518 A(H3N2) Infection: An Open-label Randomized, Controlled, Phase IIb/III Trial. *Chest*
519 151:1069-1080
- 520 55. Hussein AFA, Cheng H, Tundup S et al. (2020) Identification of entry inhibitors with
521 4-aminopiperidine scaffold targeting group 1 influenza A virus. *Antiviral Res*
522 177:104782

- 523 56. Iuliano AD, Roguski KM, Chang HH et al. (2018) Estimates of global seasonal
524 influenza-associated respiratory mortality: a modelling study. *Lancet* (London,
525 England) 391:1285-1300
- 526 57. Jalily PH, Duncan MC, Fedida D et al. (2020) Put a cork in it: Plugging the M2 viral
527 ion channel to sink influenza. *Antiviral Res* 178:104780
- 528 58. Jia R, Zhang J, Ai W et al. (2019) Design, synthesis and biological evaluation of "Multi-
529 Site"-binding influenza virus neuraminidase inhibitors. *Eur J Med Chem* 178:64-80
- 530 59. Jones JC, Kumar G, Barman S et al. (2018) Identification of the I38T PA Substitution
531 as a Resistance Marker for Next-Generation Influenza Virus Endonuclease Inhibitors.
532 *mBio* 9
- 533 60. Ju H, Xiu S, Ding X et al. (2020) Discovery of novel 1,2,3-triazole oseltamivir
534 derivatives as potent influenza neuraminidase inhibitors targeting the 430-cavity. *Eur J*
535 *Med Chem* 187:111940
- 536 61. Ju H, Zhan P, Liu X (2019) Designing influenza polymerase acidic endonuclease
537 inhibitors via 'privileged scaffold' re-evolution/refining strategy. *Future Med Chem*
- 538 62. Ju H, Zhang J, Huang B et al. (2017) Inhibitors of Influenza Virus Polymerase Acidic
539 (PA) Endonuclease: Contemporary Developments and Perspectives. *J Med Chem*
540 60:3533-3551
- 541 63. Kadam RU, Wilson IA (2017) Structural basis of influenza virus fusion inhibition by
542 the antiviral drug Arbidol. *Proc Natl Acad Sci U S A* 114:206-214
- 543 64. Kao RY, Yang D, Lau LS et al. (2010) Identification of influenza A nucleoprotein as
544 an antiviral target. *Nat Biotechnol* 28:600-605
- 545 65. Karlas A, Machuy N, Shin Y et al. (2010) Genome-wide RNAi screen identifies human
546 host factors crucial for influenza virus replication. *Nature* 463:818-822
- 547 66. Kim JI, Lee S, Lee GY et al. (2019) Novel Small Molecule Targeting the Hemagglutinin
548 Stalk of Influenza Viruses. *J Virol* 93
- 549 67. Kim S, Kim M-J, Kim C-H et al. (2017) The Superiority of IFN- λ as a Therapeutic
550 Candidate to Control Acute Influenza Viral Lung Infection. *Am. J. Respir. Cell Mol.*
551 *Biol.* 56:202-212
- 552 68. Kleinpeter AB, Jureka AS, Falahat SM et al. (2018) Structural analyses reveal the
553 mechanism of inhibition of influenza virus NS1 by two antiviral compounds. *J Biol*
554 *Chem* 293:14659-14668
- 555 69. König R, Stertz S, Zhou Y et al. (2010) Human host factors required for influenza virus
556 replication. *Nature* 463:813-817
- 557 70. Kowalinski E, Zubieta C, Wolkerstorfer A et al. (2012) Structural analysis of specific
558 metal chelating inhibitor binding to the endonuclease domain of influenza pH1N1
559 (2009) polymerase. *PLoS Pathog* 8:e1002831
- 560 71. Krammer F, Smith GJD, Fouchier RaM et al. (2018) Influenza. *Nat Rev Dis Primers*
561 4:3
- 562 72. Kumaki Y, Morrey JD, Barnard DL (2012) Effect of statin treatments on highly
563 pathogenic avian influenza H5N1, seasonal and H1N1pdm09 virus infections in
564 BALB/c mice. *Future Virol* 7:801-818
- 565 73. Lejal N, Tarus B, Bouguyon E et al. (2013) Structure-based discovery of the novel
566 antiviral properties of naproxen against the nucleoprotein of influenza A virus.
567 *Antimicrob Agents Chemother* 57:2231-2242
- 568 74. Lejal N, Truchet S, Bechor E et al. (2018) Turning off NADPH oxidase-2 by impeding
569 p67(phox) activation in infected mouse macrophages reduced viral entry and
570 inflammation. *Biochim Biophys Acta Gen Subj* 1862:1263-1275
- 571 75. Li P, Du R, Wang Y et al. (2020) Identification of Chebulinic Acid and Chebulagic
572 Acid as Novel Influenza Viral Neuraminidase Inhibitors. *Front Microbiol* 11:182

- 573 76. Liu Q, Zhou YH, Yang ZQ (2016) The cytokine storm of severe influenza and
574 development of immunomodulatory therapy. *Cell Mol Immunol* 13:3-10
- 575 77. Liu T, Liu M, Chen F et al. (2018) A Small-Molecule Compound Has Anti-influenza A
576 Virus Activity by Acting as a "PB2 Inhibitor". *Mol Pharm* 15:4110-4120
- 577 78. Lo CY, Li OT, Tang WP et al. (2018) Identification of influenza polymerase inhibitors
578 targeting C-terminal domain of PA through surface plasmon resonance screening. *Sci*
579 *Rep* 8:2280
- 580 79. Makau JN, Watanabe K, Ishikawa T et al. (2017) Identification of small molecule
581 inhibitors for influenza a virus using in silico and in vitro approaches. *PLoS One*
582 12:e0173582
- 583 80. Malakhov MP, Aschenbrenner LM, Smee DF et al. (2006) Sialidase fusion protein as a
584 novel broad-spectrum inhibitor of influenza virus infection. *Antimicrob. Agents*
585 *Chemother.* 50:1470-1479
- 586 81. Massari S, Goracci L, Desantis J et al. (2016) Polymerase Acidic Protein-Basic Protein
587 1 (PA-PB1) Protein-Protein Interaction as a Target for Next-Generation Anti-influenza
588 Therapeutics. *J Med Chem* 59:7699-7718
- 589 82. Matsui K, Ozawa M, Kiso M et al. (2018) Stimulation of alpha2-adrenergic receptors
590 impairs influenza virus infection. *Scientific Reports* 8:4631
- 591 83. Mazur I, Wurzer WJ, Ehrhardt C et al. (2007) Acetylsalicylic acid (ASA) blocks
592 influenza virus propagation via its NF-kappaB-inhibiting activity. *Cell. Microbiol.*
593 9:1683-1694
- 594 84. McGowan DC, Balemans W, Embrechts W et al. (2019) Design, Synthesis, and
595 Biological Evaluation of Novel Indoles Targeting the Influenza PB2 Cap Binding
596 Region. *J Med Chem* 62:9680-9690
- 597 85. Mehrbod P, Omar AR, Hair-Bejo M et al. (2014) Mechanisms of action and efficacy of
598 statins against influenza. *Biomed Res Int* 2014:872370
- 599 86. Meliopoulou VA, Andersen LE, Birrer KF et al. (2012) Host gene targets for novel
600 influenza therapies elucidated by high-throughput RNA interference screens. *FASEB J.*
601 26:1372-1386
- 602 87. Monod A, Swale C, Tarus B et al. (2015) Learning from structure-based drug design
603 and new antivirals targeting the ribonucleoprotein complex for the treatment of
604 influenza. *Expert Opin Drug Discov* 10:345-371
- 605 88. Moss RB, Hansen C, Sanders RL et al. (2012) A phase II study of DAS181, a novel
606 host directed antiviral for the treatment of influenza infection. *J. Infect. Dis.* 206:1844-
607 1851
- 608 89. Mould JA, Li HC, Dudlak CS et al. (2000) Mechanism for proton conduction of the
609 M(2) ion channel of influenza A virus. *J Biol Chem* 275:8592-8599
- 610 90. Muller KH, Kainov DE, El Bakkouri K et al. (2011) The proton translocation domain
611 of cellular vacuolar ATPase provides a target for the treatment of influenza A virus
612 infections. *Br J Pharmacol* 164:344-357
- 613 91. Musharrafieh R, Ma C, Wang J (2020) Discovery of M2 channel blockers targeting the
614 drug-resistant double mutants M2-S31N/L26I and M2-S31N/V27A from the influenza
615 A viruses. *Eur J Pharm Sci* 141:105124
- 616 92. Musiol A, Gran S, Ehrhardt C et al. (2013) Annexin A6-balanced late endosomal
617 cholesterol controls influenza A replication and propagation. *mBio* 4:e00608-00613
- 618 93. Nacken W, Ehrhardt C, Ludwig S (2012) Small molecule inhibitors of the c-Jun N-
619 terminal kinase (JNK) possess antiviral activity against highly pathogenic avian and
620 human pandemic influenza A viruses. *Biol Chem* 393:525-534
- 621 94. Nannetti G, Massari S, Mercorelli B et al. (2019) Potent and broad-spectrum
622 cycloheptathiophene-3-carboxamide compounds that target the PA-PB1 interaction of

- influenza virus RNA polymerase and possess a high barrier to drug resistance. *Antiviral Res* 165:55-64
95. Obayashi E, Yoshida H, Kawai F et al. (2008) The structural basis for an essential subunit interaction in influenza virus RNA polymerase. *Nature* 454:1127-1131
96. Omoto S, Speranzini V, Hashimoto T et al. (2018) Characterization of influenza virus variants induced by treatment with the endonuclease inhibitor baloxavir marboxil. *Sci Rep* 8:9633
97. Oostwoud LC, Gunasinghe P, Seow HJ et al. (2016) Apocynin and ebselen reduce influenza A virus-induced lung inflammation in cigarette smoke-exposed mice. *Scientific Reports* 6:20983
98. Pang B, Cheung NN, Zhang W et al. (2016) Structural Characterization of H1N1 Nucleoprotein-Nucleozin Binding Sites. *Sci Rep* 6:29684
99. Patnaik S, Basu D, Southall N et al. (2019) Identification, design and synthesis of novel pyrazolopyridine influenza virus nonstructural protein 1 antagonists. *Bioorg Med Chem Lett* 29:1113-1119
100. Pautus S, Sehr P, Lewis J et al. (2013) New 7-methylguanine derivatives targeting the influenza polymerase PB2 cap-binding domain. *J Med Chem* 56:8915-8930
101. Perwitasari O, Johnson S, Yan X et al. (2014) Verdinexor, a novel selective inhibitor of nuclear export, reduces influenza a virus replication in vitro and in vivo. *J. Virol.* 88:10228-10243
102. Pflug A, Guilligay D, Reich S et al. (2014) Structure of influenza A polymerase bound to the viral RNA promoter. *Nature* 516:355-360
103. Pizzorno A, Abed Y, Rhéaume C et al. (2014) Oseltamivir-zanamivir combination therapy is not superior to zanamivir monotherapy in mice infected with influenza A(H3N2) and A(H1N1)pdm09 viruses. *Antiviral Res.* 105:54-58
104. Pizzorno A, Padey B, Dubois J et al. (2020) In vitro evaluation of antiviral activity of single and combined repurposable drugs against SARS-CoV-2. *Antiviral Res.*:104878
105. Pizzorno A, Padey B, Julien T et al. (2020) Characterization and treatment of SARS-CoV-2 in nasal and bronchial human airway epithelia. *bioRxiv*:2020.2003.2031.017889
106. Pizzorno A, Padey B, Terrier O et al. (2019) Drug Repurposing Approaches for the Treatment of Influenza Viral Infection: Reviving Old Drugs to Fight Against a Long-Lived Enemy. *Frontiers in Immunology* 10:531
107. Pizzorno A, Terrier O, Nicolas De Lamballerie C et al. (2019) Repurposing of Drugs as Novel Influenza Inhibitors From Clinical Gene Expression Infection Signatures. *Frontiers in Immunology* 10:60
108. Pleschka S, Wolff T, Ehrhardt C et al. (2001) Influenza virus propagation is impaired by inhibition of the Raf/MEK/ERK signalling cascade. *Nat. Cell Biol.* 3:301-305
109. Powell JD, Waters KM (2017) Influenza-Omics and the Host Response: Recent Advances and Future Prospects. *Pathogens* 6
110. Reich S, Guilligay D, Pflug A et al. (2014) Structural insight into cap-snatching and RNA synthesis by influenza polymerase. *Nature* 516:361-366
111. Robb NC, Te Velhuis AJW, Fodor E et al. (2019) Real-time analysis of single influenza virus replication complexes reveals large promoter-dependent differences in initiation dynamics. *Nucleic Acids Res* 47:6466-6477
112. Rossignol J-F (2014) Nitazoxanide: a first-in-class broad-spectrum antiviral agent. *Antiviral Res.* 110:94-103
113. Rossignol JF, La Frazia S, Chiappa L et al. (2009) Thiazolidines, a new class of anti-influenza molecules targeting viral hemagglutinin at the post-translational level. *J. Biol. Chem.* 284:29798-29808

114. Russell RJ, Haire LF, Stevens DJ et al. (2006) The structure of H5N1 avian influenza neuraminidase suggests new opportunities for drug design. *Nature* 443:45-49
115. Salomon R, Hoffmann E, Webster RG (2007) Inhibition of the cytokine response does not protect against lethal H5N1 influenza infection. *Proc. Natl. Acad. Sci. U.S.A.* 104:12479-12481
116. Scheuch G, Canisius S, Nocker K et al. (2018) Targeting intracellular signaling as an antiviral strategy: aerosolized LASAG for the treatment of influenza in hospitalized patients. *Emerging Microbes & Infections* 7:21
117. Schloer S, Goretzko J, Pleschka S et al. (2020) Combinatory Treatment with Oseltamivir and Itraconazole Targeting Both Virus and Host Factors in Influenza A Virus Infection. *Viruses* 12
118. Schnell JR, Chou JJ (2008) Structure and mechanism of the M2 proton channel of influenza A virus. *Nature* 451:591-595
119. Schröder T, Dudek SE, Schreiber A et al. (2018) The clinically approved MEK inhibitor Trametinib efficiently blocks influenza A virus propagation and cytokine expression. *Antiviral Res.* 157:80-92
120. Schreiber A, Boff L, Anhlán D et al. (2020) Dissecting the mechanism of signaling-triggered nuclear export of newly synthesized influenza virus ribonucleoprotein complexes. *Proc. Natl. Acad. Sci. U.S.A.* 117:16557-16566
121. Sharma M, Yi M, Dong H et al. (2010) Insight into the mechanism of the influenza A proton channel from a structure in a lipid bilayer. *Science* 330:509-512
122. Sheahan TP, Sims AC, Zhou S et al. (2020) An orally bioavailable broad-spectrum antiviral inhibits SARS-CoV-2 in human airway epithelial cell cultures and multiple coronaviruses in mice. *Sci Transl Med* 12
123. Shen YF, Chen YH, Chu SY et al. (2011) E339...R416 salt bridge of nucleoprotein as a feasible target for influenza virus inhibitors. *Proc Natl Acad Sci U S A* 108:16515-16520
124. Shi X, Zhou W, Huang H et al. (2013) Inhibition of the inflammatory cytokine tumor necrosis factor- α with etanercept provides protection against lethal H1N1 influenza infection in mice. *Crit Care* 17:R301
125. Sidwell RW, Bailey KW, Wong MH et al. (2005) In vitro and in vivo influenza virus-inhibitory effects of viramidine. *Antiviral Res.* 68:10-17
126. Slaine PD, Kleer M, Smith NK et al. (2017) Stress Granule-Inducing Eukaryotic Translation Initiation Factor 4A Inhibitors Block Influenza A Virus Replication. *Viruses* 9
127. Song MS, Kumar G, Shadrack WR et al. (2016) Identification and characterization of influenza variants resistant to a viral endonuclease inhibitor. *Proc Natl Acad Sci U S A* 113:3669-3674
128. Stertz S, Shaw ML (2011) Uncovering the global host cell requirements for influenza virus replication via RNAi screening. *Microbes Infect.* 13:516-525
129. Strauch EM, Bernard SM, La D et al. (2017) Computational design of trimeric influenza-neutralizing proteins targeting the hemagglutinin receptor binding site. *Nat Biotechnol* 35:667-671
130. Sugiyama K, Obayashi E, Kawaguchi A et al. (2009) Structural insight into the essential PB1-PB2 subunit contact of the influenza virus RNA polymerase. *EMBO J* 28:1803-1811
131. Swale C, Monod A, Tengo L et al. (2016) Structural characterization of recombinant IAV polymerase reveals a stable complex between viral PA-PB1 heterodimer and host RanBP5. *Sci Rep* 6:24727

132. Tang Y, Zaitseva F, Lamb RA et al. (2002) The gate of the influenza virus M2 proton channel is formed by a single tryptophan residue. *J Biol Chem* 277:39880-39886
133. Tarus B, Bakowicz O, Chenavas S et al. (2012) Oligomerization paths of the nucleoprotein of influenza A virus. *Biochimie* 94:776-785
134. Tarus B, Bertrand H, Zedda G et al. (2015) Structure-based design of novel naproxen derivatives targeting monomeric nucleoprotein of Influenza A virus. *J Biomol Struct Dyn* 33:1899-1912
135. Terrier O, Dilly S, Pizzorno A et al. (2020) Broad-spectrum antiviral activity of naproxen: from Influenza A to SARS-CoV-2 Coronavirus. *bioRxiv:2020.2004.2030.069922*
136. Thomaston JL, Konstantinidi A, Liu L et al. (2020) X-ray Crystal Structures of the Influenza M2 Proton Channel Drug-Resistant V27A Mutant Bound to a Spiro-Adamantyl Amine Inhibitor Reveal the Mechanism of Adamantane Resistance. *Biochemistry* 59:627-634
137. Thomaston JL, Polizzi NF, Konstantinidi A et al. (2018) Inhibitors of the M2 Proton Channel Engage and Disrupt Transmembrane Networks of Hydrogen-Bonded Waters. *J Am Chem Soc* 140:15219-15226
138. Thomaston JL, Wu Y, Polizzi N et al. (2019) X-ray Crystal Structure of the Influenza A M2 Proton Channel S31N Mutant in Two Conformational States: An Open and Shut Case. *J Am Chem Soc* 141:11481-11488
139. Tilmanis D, Van Baalen C, Oh DY et al. (2017) The susceptibility of circulating human influenza viruses to tizoxanide, the active metabolite of nitazoxanide. *Antiviral Res.* 147:142-148
140. Toots M, Yoon JJ, Hart M et al. (2020) Quantitative efficacy paradigms of the influenza clinical drug candidate EIDD-2801 in the ferret model. *Transl Res* 218:16-28
141. Van Dongen MJP, Kadam RU, Juraszek J et al. (2019) A small-molecule fusion inhibitor of influenza virus is orally active in mice. *Science* 363
142. Vandermeer ML, Thomas AR, Kamimoto L et al. (2012) Association between use of statins and mortality among patients hospitalized with laboratory-confirmed influenza virus infections: a multistate study. *J. Infect. Dis.* 205:13-19
143. Vidic J, Noiray M, Bagchi A et al. (2016) Identification of a Novel Complex between the Nucleoprotein and PA(1-27) of Influenza A Virus Polymerase. *Biochemistry* 55:4259-4262
144. Walker AP, Fodor E (2019) Interplay between Influenza Virus and the Host RNA Polymerase II Transcriptional Machinery. *Trends Microbiol* 27:398-407
145. Wandzik JM, Kouba T, Cusack S (2020) Structure and Function of Influenza Polymerase. *Cold Spring Harb Perspect Med*
146. Wandzik JM, Kouba T, Karuppasamy M et al. (2020) A Structure-Based Model for the Complete Transcription Cycle of Influenza Polymerase. *Cell* 181:877-893 e821
147. Wang Y, Ding Y, Yang C et al. (2017) Inhibition of the infectivity and inflammatory response of influenza virus by Arbidol hydrochloride in vitro and in vivo (mice and ferret). *Biomed Pharmacother* 91:393-401
148. Watanabe T, Kawaoka Y (2015) Influenza virus-host interactomes as a basis for antiviral drug development. *Curr Opin Virol* 14:71-78
149. Wiesener N, Zimmer C, Jarasch-Althof N et al. (2011) Therapy of experimental influenza virus infection with pyrrolidine dithiocarbamate. *Med. Microbiol. Immunol.* 200:115-126
150. Woodring JL, Lu SH, Krasnova L et al. (2020) Disrupting the Conserved Salt Bridge in the Trimerization of Influenza A Nucleoprotein. *J Med Chem* 63:205-215

770 151. Wright ZVF, Wu NC, Kadam RU et al. (2017) Structure-based optimization and
771 synthesis of antiviral drug Arbidol analogues with significantly improved affinity to
772 influenza hemagglutinin. *Bioorg Med Chem Lett* 27:3744-3748

773 152. Wu G, Yu G, Yu Y et al. (2020) Chemoreactive-Inspired Discovery of Influenza A
774 Virus Dual Inhibitor to Block Hemagglutinin-Mediated Adsorption and Membrane
775 Fusion. *J Med Chem* 63:6924-6940

776 153. Wurzer WJ, Ehrhardt C, Pleschka S et al. (2004) NF-kappaB-dependent induction of
777 tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) and Fas/FasL is
778 crucial for efficient influenza virus propagation. *J. Biol. Chem.* 279:30931-30937

779 154. Yamaya M, Shimotai Y, Hatachi Y et al. (2015) The serine protease inhibitor camostat
780 inhibits influenza virus replication and cytokine production in primary cultures of
781 human tracheal epithelial cells. *Pulm Pharmacol Ther* 33:66-74

782 155. Yao Y, Kadam RU, Lee CD et al. (2020) An influenza A hemagglutinin small-molecule
783 fusion inhibitor identified by a new high-throughput fluorescence polarization screen.
784 *Proc Natl Acad Sci U S A*

785 156. Ye Q, Krug RM, Tao YJ (2006) The mechanism by which influenza A virus
786 nucleoprotein forms oligomers and binds RNA. *Nature* 444:1078-1082

787 157. Ye S, Lowther S, Stambas J (2015) Inhibition of reactive oxygen species production
788 ameliorates inflammation induced by influenza A viruses via upregulation of SOCS1
789 and SOCS3. *J. Virol.* 89:2672-2683

790 158. Yeganeh B, Ghavami S, Kroeker AL et al. (2015) Suppression of influenza A virus
791 replication in human lung epithelial cells by noncytotoxic concentrations bafilomycin
792 A1. *Am J Physiol Lung Cell Mol Physiol* 308:L270-286

793 159. Yin MJ, Yamamoto Y, Gaynor RB (1998) The anti-inflammatory agents aspirin and
794 salicylate inhibit the activity of I(kappa)B kinase-beta. *Nature* 396:77-80

795 160. Yoon JJ, Toots M, Lee S et al. (2018) Orally Efficacious Broad-Spectrum
796 Ribonucleoside Analog Inhibitor of Influenza and Respiratory Syncytial Viruses.
797 *Antimicrob Agents Chemother* 62

798 161. Yuan S, Chu H, Singh K et al. (2016) A novel small-molecule inhibitor of influenza A
799 virus acts by suppressing PA endonuclease activity of the viral polymerase. *Sci Rep*
800 6:22880

801 162. Yuan S, Chu H, Ye J et al. (2017) Identification of a novel small-molecule compound
802 targeting the influenza A virus polymerase PB1-PB2 interface. *Antiviral Res* 137:58-66

803 163. Zenilman JM, Fuchs EJ, Hendrix CW et al. (2015) Phase 1 clinical trials of DAS181,
804 an inhaled sialidase, in healthy adults. *Antiviral Res.* 123:114-119

805 164. Zhang J, Hu Y, Foley C et al. (2018) Exploring Ugi-Azide Four-Component Reaction
806 Products for Broad-Spectrum Influenza Antivirals with a High Genetic Barrier to Drug
807 Resistance. *Sci Rep* 8:4653

808 165. Zhang J, Hu Y, Wu N et al. (2020) Discovery of Influenza Polymerase PA-PB1
809 Interaction Inhibitors Using an In Vitro Split-Luciferase Complementation-Based
810 Assay. *ACS Chem Biol* 15:74-82

811 166. Zhang J, Poongavanam V, Kang D et al. (2018) Optimization of N-Substituted
812 Oseltamivir Derivatives as Potent Inhibitors of Group-1 and -2 Influenza A
813 Neuraminidases, Including a Drug-Resistant Variant. *J Med Chem* 61:6379-6397

814 167. Zheng B-J, Chan K-W, Lin Y-P et al. (2008) Delayed antiviral plus immunomodulator
815 treatment still reduces mortality in mice infected by high inoculum of influenza A/H5N1
816 virus. *Proc. Natl. Acad. Sci. U.S.A.* 105:8091-8096

817 168. Zheng W, Fan W, Zhang S et al. (2019) Naproxen Exhibits Broad Anti-influenza Virus
818 Activity in Mice by Impeding Viral Nucleoprotein Nuclear Export. *Cell Rep* 27:1875-
819 1885 e1875

- 820 169. Zhirnov OP, Klenk HD, Wright PF (2011) Aprotinin and similar protease inhibitors as
821 drugs against influenza. *Antiviral Res* 92:27-36
- 822 170. Zhou Z, Liu T, Zhang J et al. (2018) Influenza A virus polymerase: an attractive target
823 for next-generation anti-influenza therapeutics. *Drug Discov Today* 23:503-518
- 824 171. Zu M, Li C, Fang J-S et al. (2015) Drug Discovery of Host CLK1 Inhibitors for
825 Influenza Treatment. *Molecules* 20:19735-19747
- 826

Figure legends

Figure 1. Influenza viral particle and viral cycle; current state of anti-influenza drug discovery and development. (A) Influenza A virus (IAV) particle. The IAV genome is composed of eight ribonucleoprotein complexes (vRNPs). Each one consists of single-stranded negative-sense viral RNA (vRNA) encapsidated by viral nucleoprotein (NP) and a viral polymerase complex (PA, PB1, and PB2) positioned at the extremity of the vRNA segment. Three viral proteins are embedded within the viral membrane, hemagglutinin (HA), neuraminidase (NA), and ion channel protein (M2). Matrix protein 1 (M1) underlies the viral envelope and holds the vRNPs inside the virion. (B) The viral particle binds to sialic acid receptors and enters the cell via receptor-mediated endocytosis. Acidification of the endocytic vesicles leads to virus uncoating mediated by the M2 ion channel. vRNPs are then released into the cytoplasm and transported into the nucleus. There, the viral RNA-dependent RNA polymerase complex snatches the host mRNA caps to initiate the negative vRNA transcription. Transcribed vRNAs then need to undergo an mRNA maturation phase, including the pre-mRNA splicing, before export to the cytoplasm to be translated. vRNAs are also replicated in the nucleus to generate new vRNPs in association with neosynthesized viral proteins. Progeny vRNPs are transported toward the cytoplasmic membrane with viral components to be packaged into new infectious particles which are formed by cellular envelope budding. Classic virus-targeting strategies are highlighted in red, and virus-host-targeted strategies in blue. Figure created by BioRender.com

Figure 2: Looking down the M2 channel in the presence of inhibitors: Structure of M2 WT and VA27 mutant in complex with amantadine and spiroamantadine. View down the pore channel in A- WT-amantadine (V27 is colored in yellow, PDB ID 6BKK[137]) and B- V27A-spiroamantadine complexes (A27 is colored in yellow, PDB ID 6NV1[136])

Figure 3: Structures of the approved NA inhibitors

Figure 4: Structure of some of the pre-clinical candidates targeting HA: A: Structure of HA in complex with JNJ4796 shown in orange (PDB ID 6CF7)[141] B: Structure of trimeric HA in complex with mini-binder highlighted in yellow (PDB ID 6KUY)[129].

Figure 5: Structure of some of the pre-clinical candidates targeting the polymerase A: Active-site PA N-terminal inhibitor compound 22[25]; B: PB2 inhibitor Pimodivir [20] (the numbering are associated with this structure corresponding to the full-length PB2, C: nucleozin-NP oligomeric complex PDB ID 5B7B, monomers A and B are in cyan and yellow, respectively; D: Naproxen F1-NP monomeric complex from docking studies[33].

Figure 6: Structures of the approved polymerase inhibitors and some pre-clinical candidates

Tables

Table 1. Summary of the activity and structures of the main antiviral compounds bound to their target, the proton channel M2 of Influenza A or the neuraminidase NA of Influenza A and B.

Target	Compound	IC ₅₀	PDB ID	Stage (year approval)	references
M2	Amantadine	100 µM (H1N1 WT) > 500µM (S31N) 15.7 µM (WT channel ^a) [155]	6BKK	Approved (1976)	[137], [10]
	Rimantadine	0.1 µM (H1N1 WT) > 200µM (S31N)	2RLF	Approved (1994)	[118]
	Spiro-adamantyl amine	18.7 µM (WT channel ^a) 0.2 µM (V27A ^a)	6BMZ 6NV1 6OUG	Pre-clinical	[136, 137]
NA	Oseltamivir (Tamiflu)	0.8 nM (N5 NA)	2HT7	Approved (1999)	[114]
	Peramivir	3.4 nM	2HTU	Approved (2014)	[114]
	Zanamivir	0.6 nM (N5 NA)	3CKZ	Approved (1999)	[21]
	Chebulinic acid Chebulagic Acid	1.36 ± 0.36 µM (H1N1 PR8) (Oseltamivir resistant and H1N1 pdm09 viruses) CC ₅₀ > 100µM		Pre-clinical	[75]
	Oseltamivir derivatives	0.66 µM (H5N1)	Docking 150/430 cavity	Pre-clinical	[1, 58, 166]
	Triazol oseltamivir derivatives C1-modified oseltamivir derivatives	0.05-0.15 µM (H5N1, H5N2 and H5N6) 0.1 µM (H5N1, H5N6) 0.7 µM (Oseltamivir resistant virus)	Docking 430 cavity	Pre-clinical	[60]

a: Patch clamp assays [136]

Table 2: Recent antiviral candidates targeting HA, their activity and structures of their complexes with HA

Target	Compound / binding site	IC ₅₀ / CC ₅₀	PDB ID	Stage	references
HA	Arbidol / Stem region	4-12 μ M CC ₅₀ = 59 μ M	5T6S, 5T6N	Pre-clinical and clinical NCT03787459	[63, 147, 151]
	F0045(S) / Stem region	0.5-2 μ M (H1 HA)	6WCR	Pre-clinical	[155]
	JNJ4795 / Stem region	0.01-0.07 μ M (H1 HA)	6CF7	Pre-clinical	[141]
	IY7640 / Stem region	0.5-7 μ M (H1 HA) CC ₅₀ > 800 μ M	Docking studies	Pre-clinical	[66]
	CBS1117 / Stem region	3 μ M For H5 HA	6VMZ	Pre-clinical	[2, 45, 55]
	MB2746 / Stem region	0.3 μ M (H1 HA) CC ₅₀ > 100 μ M	Docking studies	Pre-clinical	[5]
	De novo design of “Mini-binder” proteins	0.15-0.19 nM (H3 and H1 HA)	6KUY		[129]
	Peninddone		HA1 and HA2	Pre-clinical	[152]

Table 3: Inhibitors of PA, PA-PB1 interactions and PB1

Target	Compound	IC ₅₀ / CC ₅₀	PDB ID	Stage	References
PA	Baloxavir marboxil	0.3-1 μ M (H1N1/H3N2)	6FS6 6FS9	Approved (2019) NCT02954354 NCT0294901	[96]
	L-742,001	3 μ M (WT H1N1) 24 μ M (WT H1N1 pdm09) 236 μ M (H1N1 pdm09 PA F105S)	5CGV 5D9J	Clinical trial NCT01526785	[127]
	RO7	16 nM (WT H1N1) 3 nM (H1N1 pdm09)	5VPX	Pre-Clinical	[59, 70]
	Ana-0	0.8 μ M	Docking	Pre-clinical	[161]
	Compound 22	110 pM	6E6W	Pre-clinical	[25]
	N-acylhydrazone derivatives	11 μ M	5EGA	Pre-clinical	[11]
	"312"	37 μ M (H1N1, H2N2 and H3N2)	PA –C-terminal	Pre-Clinical	[78]
PA-PB1	Compound 12a	0.9-2.7 μ M (FluA amantadine-& oseltamivir resistant, FluB)	Docking	Pre-clinical	[164]
	Amino-acids adducts of diphenyl-pyridine derivatives	39 $\pm$ 2 μ M (H1N1)	Docking	Pre-clinical	[27]
	Cycloheptathiophene-3-carboxamide	0.2 μ M-0.7 μ M H1N1 pdm09, H1N1 oseltamivir-resistant, H3N2, Influenza B	Docking	Pre-clinical	[31],[94]
	R151785	2.5, 5.0 μ M p09, H1N1 oseltamivir- & amantadine resistant Influenza B	Docking	Pre-clinical	[165]
PB1	Favipiravir	Broad-spectrum		Approved (2014)	[160]
	β -d-N4-Hydroxycytidine/ EIDD-2801	Broad-spectrum Influenza, SARS-CoV2		Clinical trial NCT04405739	[140], [122]

Table 4. Inhibitors of PB2 cap-binding, PB1-PB2, NP and NS1

Target	Compound / binding site	IC ₅₀ / CC ₅₀	PDB ID	Stage	References
PB2	Pimodividir (VX787)	2.6 nM	4P1U	Approved (2017)	[9, 20]
	5,7-difluoroindole derivative of pimodivir	11 nM	6S5V	Pre-clinical	[84]
	D 715-2441	3.6-4.4 μ M (H1N1, H3N2, H5N1, H7N9)	Docking	Pre-clinical	[77]
	Cap analogs	7.5 μ M H3N2	4CB5	Pre-clinical	[100]
PB1-PB2	PP7	1.4-9.5 μ M (Strain-specific)	Docking	Pre-clinical	[162]
NP	Nucleozin	0.07 μ M (H1N1) 0.16 μ M (H3N2) 0.33 μ M (H5N1Y287H)	5B7B	Pre-clinical	[64, 98]
	Compound 3	0.1 μ M (H1N1 and H5N1)	3RO5	Pre-clinical	[46]
	2-(4-chloro-3,5-difluorophenylamino)thiazole-4-carboxamide derivatives	0.11 μ M	Docking	Pre-clinical	[123], [150]
	Naproxen	Broad-spectrum FluA & Sars-CoV2 16 $\pm$ 5 μ M (H1N) 2.9 $\pm$ 0.3 μ M (H1N1) 1.8 μ M (H1N1 pdm09)	Docking	Pre-clinical	[33, 73, 134]
	Naproxen C0 (Naproxen derivative 2)	1.3 $\pm$ 0.2 μ M (H1N1) 0.7 μ M (H1N1 pdm09, H3N2, resistant to oseltamivir)			
	Naproxen F1 (Naproxen derivative 4)				
NS1	Hydroquinolinone derivatives (NUD)	1.8-7.0 μ M (H1N1)	Docking	Pre-clinical	[79]
	A22	$\approx$ 1 μ M (H1N1 PR8)	Docking	Pre-clinical	[68]
	ML303	0.7-17 μ M (H1N1 pdm09, H3N2)	HTS	Pre-clinical	[99]

Table 5. Drugs targeting host-cell component at different level of viral cycle stages.

Viral cycle stage	Drug name	Mode of action	Research phase	References
Viral entry	DAS181	Sialidase – removes sialic acid receptors	Phase I/II	Moss <i>et al.</i> 2012[88] Zenilman <i>et al.</i> 2015[163]
	Bafilomycin A1	V-ATPase inhibitors – inhibits endosomal acidification	preclinical	Yeganeh <i>et al.</i> 2015[158]
	Concanamycin			Müller <i>et al.</i> 2011[90]
	Diphyllin			Chen <i>et al.</i> 2013[14]
	Saliphenylthalamide			Bimbo <i>et al.</i> 2013[7]
	Aprotinin	Protease inhibitors – inhibit HA0 cleavage	Approved (2011)	Zhirnov <i>et al.</i> 2011[169]
	Camostat		Preclinical	Yamaya <i>et al.</i> 2015[154]
	Dynasore	Inhibition of internalization		de Vries <i>et al.</i> 2011[30]
	EIPA			Harada <i>et al.</i> 2007[51]
	Fattiviracin			
Nuclear import of vRNP	Ivermectin	Inhibits importin- α/β		Gotz <i>et al.</i> 2016[47]
Genomic replication & protein synthesis	TG003	CLK1 inhibitors -Regulation of splicing – decrease in M2 mRNA expression		Karlas <i>et al.</i> 2010[65]
	Clypearin			Zu <i>et al.</i> 2015[171]
	Corilagin			
	Sylvestrol	eIF4A inhibitors – inhibit viral protein synthesis		Slaine <i>et al.</i> 2017[126]
	Pateamine			
	Ribavirin	Approved (1986)	Durr <i>et al.</i> 1975[38]	
	Viramidine (ribavirin prodrug)	Phase III (HCV)	Sidwell <i>et al.</i> 2005[125]	
vRNP nuclear export	Cyclosporin A	Inhibits host RNA polymerase II	preclinical	Liu <i>et al.</i> 2012[77]
		Inhibits nuclear export of vRNPs		
	Verdinexor	Exportin 1 inhibitors		Perwitasari <i>et al.</i> [101] 2014
	DP2392-E10			Chutiwitoonchai <i>et al.</i> 2017[19]
	CI-1040			Haasbach <i>et al.</i> 2017[49]
	UO126	MEK inhibitor – nuclear retention of VRNP complex		Pleschka <i>et al.</i> 2001[108]
	PBP10/BOC2			Formyl peptide receptor 2 antagonists – Raf/MEK/ERK inhibition
	Trametinib	MEK1/2 inhibitor – inhibition of vRNP export	Approved (cancer)	Schräder <i>et al.</i> 2018[119]
	Dapivirine	Reverse transcriptase inhibitor - inhibition of vRNP export	Phase III (HIV)	Hu <i>et al.</i> 2017[53]
Apical transport and budding	Nitazoxanide	Anti-parasitic – Inhibition of HA maturation & transport	Phase III	Rossignol <i>et al.</i> 2009[113]
	Ruxolitinib	Virion formation & vRNA incorporation inhibition	Approved (myelofibrosis)	Watanabe <i>et al.</i> 2015 [148]
	U18666A	Hydrophobic polyamine - Reduces plasma membrane cholesterol level and decreases virion egress	preclinical	Musiol <i>et al.</i> 2013[92]
	Clonidine	alpha2-adrenergic receptors inhibitor - Inhibits transport of HA transport to plasma membrane		Matsui <i>et al.</i> 2018[82]

Table 6. Drugs targeting host-cell signaling pathway and host-responses that are crucial for influenza replication cycle.

Host signalling pathway/response	Drug name	Mode of action	Research phase	References
NF-kB pathway	Acetylsalicylic acid	Immune dysregulation Inhibition of caspase/vRNP export inhibition	Approved	Mazur <i>et al.</i> 2007[83]
	pyrrolidine dithiocarbamate		preclinical	Wiesener <i>et al.</i> 2011[149]
	SC75741		preclinical	Ehrhardt <i>et al.</i> 2013[39] Haasbach <i>et al.</i> 2013[49]
	LASAG		Phase II	Droebeiner <i>et al.</i> 2017[35] Scheuch <i>et al.</i> 2018[116]
C-Jun-N-terminal-kinase	SP600125	C-Jun N-Terminal kinase inhibitor – Immune dysregulation	preclinical	Nacken <i>et al.</i> 2012[93]
p38 MAPK	NJK14047	Immune dysregulation	preclinical	Choi <i>et al.</i> 2016[18]
HMG-CoA	Statins	Immunomodulation	Phase II	Fedson <i>et al.</i> 2013[43] Mehrbod <i>et al.</i> 2014[85] Fedson <i>et al.</i> 2018[43]
TNF-alpha	Etanercept	Anti-inflammatory drug - Prevents TNF-mediated lung injury and edema	preclinical	Shi <i>et al.</i> 2013[124]
Nox2	Apocynin	ROS scavenger, inhibits Nox2 activity	preclinical	Ye <i>et al.</i> 2015[157] Oostwoud <i>et al.</i> 2016
	Ebselen	ROS scavenger and glutathione peroxidase mimetic, inhibits Nox2	preclinical	
Lipoxygenase & COX pathways	Celecoxib	Immune dysregulation	Phase III	Davidson <i>et al.</i> 2018[28] Carey <i>et al.</i> 2010[12] Zheng <i>et al.</i> 2008[167]
	Mesalazine	Immune dysregulation	preclinical	
Type III IFN response	Type III IFN	Induction of type III IFN response	preclinical	Davidson <i>et al.</i> 2016[29] Kim <i>et al.</i> 2017[67]
	Diltiazem		Phase II	Pizzorno <i>et al.</i> 2019[106, 107]

A

B

Figure 2

Oseltamivir

Peramivir

Zanamivir

Laninamivir

Ardibol

A

B

Figure 4

Figure 5

Baloxavir

Favipiravir

Pimodivir

Nucleozin

Naproxen