Cyclization Dynamics and Competitive Processes of Photochromic Perfluorocyclopentene Dithienylethylene in Solution

Ismail Hamdi, Guy Buntinx, Amit Kumar Tiwari, Stéphanie Delbaere, Michinori Takeshita, Stéphane Aloïse

► To cite this version:

Ismail Hamdi, Guy Buntinx, Amit Kumar Tiwari, Stéphanie Delbaere, Michinori Takeshita, et al.. Cyclization Dynamics and Competitive Processes of Photochromic Perfluorocyclopentene Dithienylethylene in Solution. *ChemPhysChem*, 2020, 21 (19), pp.2223-2229. 10.1002/cphc.202000516 . hal-03292762

HAL Id: hal-03292762

<https://hal.science/hal-03292762>

Submitted on 20 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Cyclization Dynamics and competitive processes of photochromic perfluorocyclopentene dithienylethylene in solution.

Ismail Hamdi,^{*[a]} Guy Buntinx,^[a] Amit Kumar Tiwari,^[a,b] Stéphanie Delbaere,^[a] Michinori Takeshita,^[c] and Stéphane Aloïse^{*[a]}

[a] Dr. Ismail Hamdi, Dr. Guy Buntinx, Dr. Amit Kumar Tiwari, Dr. Stéphanie Delbaere, Dr. Stéphane Aloïse
Univ. Lille, CNRS, UMR 8516 - LASIR - Laboratoire de Spectrochimie Infrarouge et Raman, F-59000 Lille, France
Email: ismail.hamdi@u-paris.fr, guy.buntinx@univ-lille.fr, amit.tiwari.chem@gmail.com, stephanie.delbaere@univ-lille.fr, stephane.aloise@univ-lille.fr

[b] Dr. Amit Kumar Tiwari
School of Advanced Sciences, Department of Chemistry, Vellore Institute of Technology-Vellore, Tamil Nadu, India, 632014
Email: amit.tiwari.chem@gmail.com

[c] Dr. Michinori Takeshita
Department of Chemistry and Applied Chemistry, Faculty of Science and Engineering, Saga University, Honjo 1, Saga 840-8502, Japan.
Email: michi@cc.saga-u.ac.jp

Abstract: Time-resolved absorption spectroscopy measurements were applied to study the dynamics of photochromic 1,2-Bis(2,4-dimethylthiophene-3-yl)perfluoro-cyclopentene (DMTPF) in chloroform including antiparallel conformer ring-closure reaction and parallel conformer photophysics. All characteristic times are given, discussed and compared to a previous publication concerning the close molecule substituted with phenyl rings. (Hamdi *et al*, pccp, 2016). Apart from the expected photocyclization process, a ring condensed by-product formation is observed and hypothesis concerning the origin of this by-product are presented.

1. Introduction

Designing of the photochromic materials which convert reversibly its color upon light stimulation^{1–4} have gained the attraction of material scientist due to its various applications in holography,⁵ information storage/processing,⁶ biosensors,^{7,8} fluorescent probe,^{9,10} smart materials^{11,12} etc. Among the different molecular switches necessary to elaborate such applications,^{1–4} diarylethenes (DAEs) and more specifically dithienylethenes (DTEs)^{13,14} are often used due to their good photochromic properties especially in terms of fatigue resistance.¹⁵ DTEs in solution are having property to undergo efficient photocyclization ($\phi_{cyc} \sim 0.2-0.5$) between uncolored open form (OF) and colored closed form (CF) in presence of UV light, whereas photoreversion reactions are known to be less efficient ($\phi_{rev} \sim 0.01$) when expose towards the visible light.^{13,14} During the last decades, the quantum yield of carbon-carbon photoelectrocyclization limited to 0.5 was a subject of interest explained in term of coexistence of two OF isomers having parallel (P) and anti-parallel (AP) conformations in equal proportions but only the latter being photoactive.^{16,17} To overpass such limitation some researchers focused their efforts on exploring purposed-designed synthetic approach^{18–21} or specific excited-state reactivity.^{22,23} Indeed, the rationalization of complex excited state mechanisms are highly desired to still improve photochromic performances and so considerable investigations have been devoted worldwide to explore the excited state topologies of DTEs (or DAEs) via advanced optical

Scheme 1. UV/Vis Photochromic reactions of DMTPF (R= H) and DTE (R=C₆H₅) between the OF and CF. Thermal Interconversion between AP and P conformers is indicated.

spectroscopies^{24–27} and/or computational chemistry.^{28–30} In this perspective, our group has actively investigated the spectrokinetic features on the series of bridged DAEs,³¹ bridged DTEs^{29,32,33} or DTEs dedicated to supramolecular chemistry³⁴ using ultrafast spectroscopies combined with (TD)-DFT calculations. Our leitmotiv was the attempt to detangle the photochemistry of the antiparallel (AP) conformers from the photophysics of parallel (P) conformers by comparing bridged/constrained vs free analog molecules (the initial purpose of the former being the blockade of the molecule into a photoactive conformation). Due to its commercial availability, normal dithienylethene 1,2-bis(2,4-dimethyl-5-phenyl-3-thienyl) perfluorocyclopentene (denoted as

DTE, hereafter) has become a photoswitch of reference,^{26,35–38} reason why we reinvestigated its photochemistry together with a series of polyether bridged analogs. The main outcomes³² were pointed as follow: (i) for the P* conformer, an Intersystem crossing (ISC) occurred (40-150 ps) to produce a first triplet state (³P) after ultrafast relaxation toward S₁ relaxed state. Later on the ³P state relaxed within few tens of μs to the ground state; (ii) for the AP* conformer, the photocyclization reaction occur immediately after light absorption (100-200 fs) starting from the Franck-Condon (FC) region while the relaxed singlet state is populated in parallel way and evolves through an ISC reaction (1-40 ps) and generate a second triplet state, ³AP with a lifetime depending on the structural constraints; (iii) interconversion between the two triplet states has been experimentally identified, e.g. ³AP ↔ ³P only for some polyether bridged DTE (DTE-O₆ in ref ³³); (iv) an unidentified species (denoted as “X” hereafter) have been reported with variable statistical weight (0-50%) along with the two triplet states within μs regime.

In this paper, we decided to reinvestigate a smaller structural analogue of DTE without the two terminal phenyl groups, the 1,2bis(2,5-dimethyl-3-thienyl)-perfluorocyclopentene noticed in the following DMTPF. Such molecule has been studied in the past in crystalline phase^{35,36} and in solution^{37,38} with the main features to present a photocyclization more efficient in crystalline phase ($\phi_{cycl}=0.5$) as compared to hexane solution ($\phi_{cycl}=0.21$) while photoreversion yields are almost identical ($\phi_{cycl}=0.12$) in both phases. Knowing that AP population was restricted to 17% in the solid state (compared to almost 50% in solution), the authors concluded for an excited state energy migration during photoconversion, the P conformer acting as a sensitizer.

So, focalizing our attention on photocyclization process, we have applied transient absorption spectroscopy on the micro-femtosecond timescale in order to elucidate the excited states dynamics of DMTPF in solution with a systematic comparison with the previous study dealing with DTE molecule. We will pay attention on the possible existence of triplet or X species and discuss the previous results found in the literature under the light on our new results.

2. Results and Discussion

2.1 General photochemical features in solution.

The photochromism of DMTPF in chloroform is illustrated in Figure 1a through absorption spectra under continuous UV irradiation for less than 10 min. For uncoloured solution, OF has a peak at ~ 235 nm. Upon UV irradiation with 302 nm light, the solution becomes pink and the spectrum

	λ_{max}^{abs} (nm)		Population % (P:AP)	Quantum yields	
	OF	CF		Cyclization	Triplet
DMTPF	235	535	80:20	0.22	0.75
DTE	273	575	52:48	0.46	0.4

Table 1. Experimental photochromic parameters of DMTPF and DTE (ref ³²) in chloroform.

Figure 1. (a) Absorption spectral change of DMTPF in chloroform (50 μM) upon UV irradiation (302 nm) up to 10 minutes and (b) up to 1 hour (c) Absorbance evolution @ 532 nm with time for an UV irradiation starting at 60s after the beginning of the recording.

dramatically changes and has a peak at 351 nm and broad absorption band at 535 nm corresponding to the absorption of CF. After 350 s irradiation time with continuous UV, the absorbance of CF does not increase further and a plateau is reached (see Figure 1b) which indicates the photo-stationary state (PSS). However, with further UV irradiation, the solution turns to pale yellow and as seen in Figure 1c, a new band appears at ~300 nm with a concomitant decrease of the band at 532 nm responsible for the decrease of the curve in Figure 1b after PSS plateau. In contrast, as seen in Figure S.1, the PSS plateau for DTE is very stable (more than 50 hours) evidencing the lack of any photodegradation. In the case of DMTPF, by analogy with previous work dealing on neighbouring molecules,^{39,40} such drastic spectral modifications are probably due to the well-known condensed rings by-product formation resulting from UV excitation of the CF (see inset of Figure 1c). Such by-product species is well characterized by the band around 300 nm and the vanishing contribution in the visible region. This result indicates that the presence of the phenyl groups drastically increases the stability of the CF of DTE compared to DMTPF. Note that for the time-resolved studies here below, we have chosen experimental conditions in order to neglect as much as possible the by-product contamination.

Some important photochemical parameters were measured in chloroform and gathered in Table 1. The P:AP ratio for DMTPF has been determined based on ¹H-NMR measurements to be 80:20 at ambient temperature in d-chloroform (Figure S.2). For DTE, we have recover the same 52:48 ratio that was previously reported by Irie and co-workers.³⁷

The main indicative results concern the proportion of P conformers from 52% for DTE to 80% for DMTPF. It immediately rationalizes the quantum yields of the cyclization for DMTPF (0.22) being more than 2 times lower compared to DTE (0.46) paralleling the decrease of AP population from 48 to 20%. Another important findings concerns the triplet quantum yields measured by the method of energy transfer with beta-carotene we already applied to DTE³² indicated in the table 1. For DMTPF this value is as high as 0.75 while for DTE, it decreases down to 0.4. Actually, it is tempting to correlate the high triplet formation efficiency with the high proportion of parallel conformer. Indeed, a quantitative estimation for the intrinsic cyclization yield gives almost an unit value ($\approx 0.22/0.2$) indicating that almost all photoexcited AP is converted to CF with no population accumulated in the triplet state ³AP. First, such efficient mechanism is consistent with a ring closing process from the Franck-Condon region in proximity with conical intersection for which competitive routes are not taken. Then, the apparent

	τ_1	τ_2	τ_3	τ_4
DMTPF	0.09 (1)	0.30 (4)	1.2 (6)	80 (1)
DTE³²	0.10 (2)	0.20 (5)	1.0 (2)	55 (8)

Table 2. Characteristic times (in ps) deduced from the global fitting method applied to femtosecond transient absorption data for DMTPF and DTE.

Figure 2. Femto-picosecond time-resolved absorption spectra of DMTPF in chloroform ($2 \cdot 10^{-4}$ M) using 320 nm excitation wavelength. Transient data are separated into 4 different temporal windows for improved clarity. All spectra are corrected from the GVD. The assignments of the different bands are indicated (see text).

discrepancy between the high intrinsic cyclization yield and high triplet formation yield can be explained in term of ¹AP species

mainly produced through 1P species upon rapid interconversion. Computational estimation of the potential barrier between 1AP and 1P will be published elsewhere. By the way, we will discuss this point below with the aid of transient spectroscopy. Indeed, in the following we will describe the ultrafast dynamics of this molecule knowing that the photocyclization mechanism and the competitive process (P photophysics) are very similar to DTE molecule. For that reason, we will keep the same notations compare to our first article (we also present the results in a short manner) but we will present the transient species following the natural deexcitation timescale, femtosecond/picosecond regime and then nanosecond regime for which the two molecules display some different features.

2.2. Excited States dynamics in the pico-femtosecond regime.

The early photophysics just after laser excitation is now investigated. Figure 2 shows time-resolved transient absorption spectra obtained after the excitation of the DMTPF OF (for DTE, see Figure 6 of ref³²) in chloroform solution with a femtosecond 320 nm laser pulse, recorded in the 380-700 nm spectral range within four temporal windows (-1-0.25, 0.28-0.5, 0.52-1 and 2-500 ps). Global fitting of transient signal with four-exponential functions, convoluted with a Gaussian profile simulating the pump-probe correlation function are shown in supporting information (see Figure S.3). The characteristic times for DMTPF compared to DTE are represented in Table 3. As already said, because of similar results, the interpretations made for DMTPF data of will stick those made for DTE molecule in our previous work. The transient data results are summarized on Figure 4.

For short delay times (-1-0.25 ps), just after laser excitation, the spectra present essentially two intense bands peaking at 385 nm and 450 nm, respectively. By analogy with our previous work, these bands are attributed to the excited state absorption (ESA) of S_1 states on the Franck-Condon region (FC) for both AP and P conformers, i.e. $S_1^{FC}(AP)$ and $S_1^{FC}(P)$. For the delay times between 0.28-0.5 ps on the second panel, we observed a decrease of the band peaking at 385 nm while, in the same time, the 550-750 nm edge is maintained constant. Indeed, in this spectral range, the vanishing signal is compensated in the same extent by a rising component which matches perfectly the CF visible band. The related characteristic time can thus be ascribed to the photocyclization of DMTPF (OF $\rightarrow$ CF) which takes place with a time constant of 90 fs. For the third panel (0.52-1 ps), the ESA peaking at 450 nm decreases with a characteristic time of 300 fs. This evolution is attributed to the relaxation of the FC region for P conformer ($(S_1^{FC}(P) \rightarrow S_1^{rel}(P))$). Note that this spectral

Figure 3. (a) Transient absorption spectra of DMTPF in chloroform ($5 \cdot 10^{-4}$ M) for different time delays after nanosecond laser excitation at 290 nm and (b) Kinetic traces measured at 440 nm for solutions initially saturated with N_2 gas (blue line) or O_2 gas (red line). (c) Stern-Volmer plots concerning experimental decays k_{exp} of two triplet states and X species obtained from kinetics at 440 nm vs diene quencher concentration $[Q]$.

evolution was more difficult to detect for DTE, and can be related to the higher percent of P conformer population for DMTPF than for DTE (80% vs 52%). Finally, for pump-probe delay times higher than 2 ps, we observed a decrease of both transient signals with two different characteristic times 1.2 ps and 80 ps respectively attributed to the intersystem-crossing (ISC) of the AP conformer ($S_1^{rel}(AP) \rightarrow ^3AP$), and P conformer ($S_1^{rel}(P) \rightarrow ^3P$). Interestingly, the effect of introducing phenyl groups into the photochromic core does not affect the ISC of the AP species (1.2 ps vs 1.0 ps) contrary to the P species. This effect of structure can be nicely rationalized in terms of unpaired electrons well localized on the

	$\tau_1(^3\text{P})$	$\tau_2(^3\text{AP})$	$\tau_3(\text{X})$	k_1^Q	k_2^Q
DMTPF	1.80(6) (35%)	4 (1) (15%)	13 (1) (50%)	0.09 (2)	0.70 (5)
DTE	2.60(3) (80%)	25.5 (4) (20%)	...	0.10 (4)	2.23 (1)

Table 3. Lifetimes (in μs) deduced from multi-exponential fitting of transient kinetics obtained from the laser-flash photolysis experiment (the relative contributions are given into parenthesis) for DMTPF and DTE (ref³²). Quenching rate constants ($10^8 \text{ M}^{-1}\text{s}^{-1}$) of AP, P and CF species are also indicated (see text).

Figure 4. Photochemical processes (illustrations) including characteristic times derived according transient spectroscopy.

reactive carbons suggesting also a possible steric effect for the parallel species. Indeed, the faster ISC for DTE P conformer (55 ps instead of 80 ps) has to be related with torsional effect induced by the phenyl groups.

In summary, the ultrafast photocyclization from the FC region gives birth to the CF in competition with the ISC of both AP and P conformers. The deactivation pathways of both triplet states are now under examination.

2.3. Excited state dynamics in the nano-microsecond regime.

Two triplet states. The transient absorption spectra of the OF of DMTPF in chloroform excited at 290 nm and recorded in the 350-650 nm spectral range within a time windows of 0.15-50 μs are shown in Figure 3a. The spectra present essentially three bands: two intense bands peaking near 380-390 nm and 440 nm and a low intensity band at ~540 nm. In the 0.15-50 μs time domain, spectral evolution shows a subsequent decrease of the red bands letting appear the CF spectra (see PPS spectra in black dashed curved). Those transient data have to be compared with DTE case previously reported (see Figure 2 in ref³²).

As seen in Figure 3b, the differential absorption kinetic curve recorded at 440 nm (blue color curve) is well-fitted with a gaussian convoluted tri-exponential function (to take into account the irf) with decay time constants of 1.8 μs (35%), 4 μs (15%) and 13 μs (50%). Note that, these values are lower than those reported for DTE molecule (2.55 μs and 25.5 μs). After addition of oxygen in the solution, those three times are totally quenched (see red curve in the same figure) which suggests that transient species are either biradicals or triplet states. According to our previous work published for DTE,³² we attributed τ_1 and τ_2 to the ^3P and ^3AP respectively, while the third time is related with an unknown species X. To confirm such attribution, the effect of the addition of a quencher (2, 5-dimethyl-2, 4-hexadiene) on the kinetics at 440 nm has been tested with the resulting Stern-Volmer plot displayed in Figure 3c. As expected, τ_1 and τ_2 depend linearly on the quencher concentration which confirms the existence of both triplet states ^3P and ^3AP with a different quenching rate probably due to the difference in conformational arrangement of the two conformers. In contrast, the third species X is insensitive to the diene addition which is an evidence for its singlet nature with biradical structure (oxygen quenching). We will reconsider this point just hereafter. At this point, comparing DTE and DMTPF, it is worth to notice a clear structural effect concerning the lifetimes of triplet states. The substitutions of terminal hydrogens by phenyl rings induce a clear enhancement of the lifetimes for the AP (25 vs 4 μs) species compared to P species (2.6 vs 1.8 μs). Furthermore, in total consistency, the quenching rate constant are unaffected for P species while the value is reduced by a factor 3 for the AP species. Those two results bring a strong evidence for a delocalization of the unpaired electrons over the phenyl rings in the case of the AP species while such delocalization is lacking for the P species.

Origin of X species. We are now concerned by the identification of X species. Such species have been reported in more or less extent for the different DTEs we studied in the past³²⁻³⁴ but without noticeable photodegradation. In the case of DMTPF, we know that the sample has suffered some non-negligible photodegradation during transient experiment as proved by the presence of the by-product absorption band after the nanosecond measurements as shown in Figure S.4 and Figure S.5. Therefore, it seems reasonable to correlate the X species with a by-product precursor being in the singlet state with probable biradical structure. Following the admitted mechanism would be that the X species is due to the re-excitation of the CF and the X species would be a biradical species either in singlet or triplet states. By the way, among the two propositions emitted by Irie *et al.*, (the

direct precursors of the by-product in scheme 4 and scheme 5 in ref ⁴⁰),

Scheme 2. Structure with two unpaired electrons.

we validate the structure with the two unpaired electrons far away compatible with a singlet nature. Such species would have a 13 μ s lifetime.

At present time, this transient species responsible for the photodegradation is thought to originate from the CF excited state after UV irradiation. A second hypothesis is valuable as well involving the P conformer. Indeed, if we consider the relative contribution of triplet states species in the table 4, the results are instructive as well keeping in mind that the following discussion is qualitative. From NMR experiment, we know that the proportion of P conformer (see table 1) is much higher for DMTPF compared to DTE. Furthermore, it is worth to precise that molar extinction coefficient for the triplet states are expected to be in the same order of magnitude for both molecules (for DTE, spin density of 3 AP is not delocalized along lateral phenyls implying probable similar optical properties for DMTPF; see Figure S.24 of ref³²) So, upon inspection of table 3, the quite similar relative amplitudes for the decays (at around 15-20%) suggest a similar 3 AP triplet population. In contrast, among the remaining 3 P population (75-80%), one can notice that 50% of the P conformers are turned into X species. Based on those results, we can emit a new hypothesis concerning the nature of the X species: rather than originating from CF reexcitation, the by-product reported for diarylethene could come from the triplet of the P conformer of the OF. As an evidence, advanced CASSCF/CASPT2 calculations already performed for model DTEs²⁹ are now desired to explore the pathways starting from either CF* or from P* species evolving through the X singlet biradical structure to finally giving birth to the by-product molecule. Just as challenging, the interconversion between AP and P at both singlet and triplet manifold have to be included.

Concerning the exact role of the triplet species in solid state DMTPF photochromism, although it exists constraints due to crystal lattice, we do not expect any photochromism from the triplet route (3 AP $\rightarrow$ CF) because such a route is not observed in solution unlike the polyether bridged DTEs analogs. Indeed, the nanosecond transient signal acquired @ 550 nm CF maximum absorption (see Figure S.6) is a plateau for the long time, the

height of this plateau being proportional to the photo induced CF. Such signal is insensitive to oxygen addition bringing evidence for a pure photocyclization route in the singlet manifold.

3. Conclusion

In this article, we investigated new insights into the ultrafast dynamics of the ring-closure reaction for DMTPF in solution by means of steady state, nanosecond and femtosecond absorption spectroscopy. The general mechanism is summarized in Figure 4. For the P conformer, a vibrational relaxation takes place at $\sim$ 300 fs an expected ISC occurs (80 ps) to produce 3 P. The latter relaxes to the ground state with a time constant $\sim$ 1.8 μ s. Concerning the AP conformer, just after pump excitation a very rapid (90 fs) reaction channel for the ring closure was found. Then a triplet state is produced by the relaxed singlet state within $\sim$ 1.2 ps, the latter relaxes to the ground state within 4 μ s. In addition, we reported the by-product formation for which a singlet precursor has a 13 μ s lifetime. The possible origin of this species is either re-excitation of CF species by UV light or related with the triplet state of the P conformer. These two routes are now under investigations in our laboratories combining *ab initio* calculations, HPLC and RPE measurements.

4. Experimental Section

DMTPF was synthesized following already published protocol.³⁷ Chloroform solvents (spectroscopic grade, Sigma-Aldrich) were used as received. The stationary absorption spectra were measured with CARY 100bio absorption spectrometer. 1 H-NMR spectra were recorded on Bruker Avance-300 (300 MHz) in $CDCl_3$ solvent. The photocyclization yields have already been published for DTE³² and similar measurements have been performed for DMTPF. For time-resolved measurements, adjusted concentrations were chosen around 10^{-4} mol dm $^{-3}$ to get an OD around unity at the excitation wavelength. Our nanosecond transient absorption spectroscopy setup has been described elsewhere.⁴¹ Briefly, the third harmonic of a nanosecond Nd:YAG laser is used to pump a frequency doubled OPO laser tuned at 290 nm. 1mJ, 5 ns, 290 nm pulses were used as excitation light and a pulsed Xe lamp was utilized as the probe light. Transient absorption spectra were obtained from the transient absorption decays recorded at various wavelengths by sampling the absorbance changes for different given delay-times. Femtosecond time-resolved experiments were performed using a pump-probe spectrometer^{42,43} based on Ti:sapphire laser system (Coherent Mira-900-D oscillator and Libra-S regenerative

amplifier) delivering 800 nm, 1 mJ and 90 fs pulses with a repetition rate of 1 kHz. The pump pulses adjusted at 320 nm were generated by frequency quadrupling the output of a Quantronix Palitra OPA pumped at 800 nm and the energy at the sample was about 2 μJ (0.2 mJ cm^{-2}). Note that, we decided to excite the molecule far from the absorption peak which is localized near 270 nm to avoid short-time signal coming from the solvent. The probe pulses were obtained by focusing 1 μJ , 800 nm pulses into a 1 mm CaF_2 plate to generate a white light continuum. The pump-probe polarization configuration was set at the magic angle (54.7°). Transient absorbance was obtained by comparing signal and reference spectra with and without pump pulses for different delay times. The delay time between the pump and probe was varied up to 0.5 ns using an optical delay line. Sample solutions were circulated in a cell equipped with 200 μm thick CaF_2 entrance window and having 2 mm optical path length. The sample solution was circulated during measurements and was irradiated by CW Xe lamp visible light to prevent the accumulation of CF. In this configuration, for 320 nm excitation, the instrumental response function determined by fitting the Stimulated Raman transient spectra of acetonitrile was determined to be 180 fs (see Figure S.7.). All the transient spectra presented in this paper are GVD corrected according to the typical extrapolation method^{44,45} (the temporal chirp over the white-light 350-800 nm range was about 400 fs). The characteristic times deduced from kinetics are obtained by fitting the data with the result of a multi-exponential function convolved with a Gaussian pulse mimicking the pump-probe cross-correlation function (FWHM at 180 fs).

Acknowledgements

Chevreul Institute (FR 2638), Ministère de l'Enseignement Supérieur et de la Recherche, Région Nord-Pas de Calais and FEDER are acknowledged for supporting and funding this work.

This article is dedicated to the memory of Dr. Olivier Poizat, CNRS researcher at LASIR laboratory, who passed away much too soon. Olivier Poizat led the "photochemistry" team at LASIR for many years with great involvement, kindness and modesty. He was able to unite energies around numerous fundamental research projects dealing with photochemistry, photophysics and spectroscopy. He was at the origin of many research themes and of the development of time-resolved spectroscopy experiments in the laboratory. Throughout his career, he has participated in the training of many PhD students, post-doctorants and junior researchers. The authors of this paper are infinitely grateful to him for his unfailing support in the

development of researches on the elucidation of reaction mechanisms of photochromic molecules and for the many fruitful discussions they had with him on this subject and many others. Rest in peace Olivier.

Keywords: Photochromism, Photophysics, Photochemistry, Transient Absorption, Ultrafast dynamics.

- (1) Crano, J. C.; Guglielmetti, R. J. *Organic Photochromic and Thermochromic Compounds*; J.C Crano, R. J. G., Series Ed.; Plenum: New York, 1999; Vol. 1,2.
- (2) Feringa, B. L. *Molecular Switches*; Wiley-VCH: Darmstadt, 2001.
- (3) Bouas-Laurent, H.; Durr, H. *Organic Photochromism - (IUPAC Technical Report)*; Durr, H., Bouas-Laurent, H., Series Eds.; Photochromism: Molecules and Systems; Elsevier Science Bv: Amsterdam, 2003. <https://doi.org/10.1016/b978-044451322-9/50004-x>.
- (4) Tian, H.; Zhang, J. *Photochromic Materials: Preparation, Properties and Applications*; Photochromic Materials: Preparation, Properties and Applications; 2016. <https://doi.org/10.1002/9783527683734>.
- (5) Pu, S. Z.; Yang, T. S.; Yao, B. L.; Wang, Y. L.; Lei, M.; Xu, J. K. Photochromic Diarylethene for Polarization Holographic Optical Recording. *Mater. Lett.* **2007**, *61* (3), 855–859.
- (6) Andréasson, J.; Pischel, U. Storage and Processing of Information Using Molecules: The All-Photonic Approach with Simple and Multi-Photochromic Switches. *Isr. J. Chem.* **2013**, *53* (5), 236–246. <https://doi.org/10.1002/ijch.201300014>.
- (7) Avella-Oliver, M.; Morais, S.; Puchades, R.; Maquieira, A. Towards Photochromic and Thermochromic Biosensing. *TrAC - Trends Anal. Chem.* **2016**, *79*, 37–45. <https://doi.org/10.1016/j.trac.2015.11.021>.
- (8) Cheng, H.; Yoon, J.; Tian, H. Recent Advances in the Use of Photochromic Dyes for Photocontrol in Biomedicine. *Coord. Chem. Rev.* **2018**, *372*, 66–84. <https://doi.org/10.1016/j.ccr.2018.06.003>.
- (9) Metivier, R.; Amengual, R.; Leray, I.; Michelet, V.; Genet, J. P. Novel Fluorophores: Efficient Synthesis and Photophysical Properties. *Org. Lett.* **2004**, *6* (5), 739–742.
- (10) Metivier, R.; Leray, I.; Valeur, B. A Highly Sensitive and Selective Fluorescent Molecular Sensor for Pb(II) Based on a Calix 4 Arene Bearing Four Dansyl Groups. *Chem. Commun.* **2003**, No. 8, 996–997.
- (11) Goulet-Hanssens, A.; Eisenreich, F.; Hecht, S. Enlightening Materials with Photoswitches. *Adv. Mater.* **2020**. <https://doi.org/10.1002/adma.201905966>.
- (12) Russew, M. M.; Hecht, S. Photoswitches: From Molecules to Materials. *Adv. Mater.* **22** (31), 3348–3360. <https://doi.org/10.1002/adma.200904102>.
- (13) Irie, M. Diarylethenes for Memories and Switches. *Chem. Rev.* **2000**, *100* (5), 1685–1716.
- (14) Irie, M.; Fukaminato, T.; Matsuda, K.; Kobatake, S. Photochromism of Diarylethene Molecules and Crystals: Memories, Switches, and Actuators. *Chem. Rev.* **2014**, *114* (24), 12174–12277. <https://doi.org/10.1021/cr500249p>.
- (15) Herder, M.; Schmidt, B. M.; Grubert, L.; Paetzel, M.; Schwarz, J.; Hecht, S. Improving the Fatigue Resistance of Diarylethene Switches. *J. Am. Chem. Soc.* **2015**, *137* (7), 2738–2747. <https://doi.org/10.1021/ja513027s>.
- (16) Uchida, K.; Tsuchida, E.; Aoi, Y.; Nakamura, S.; Irie, M. Substitution Effect on the Coloration Quantum Yield of a Photochromic Bisbenzothienylethene. *Chem. Lett.* **1999**, No. 1, 63–64.
- (17) Morimitsu, K.; Kobatake, S.; Irie, M. Large Geometrical Structure Changes of Photochromic Diarylethenes upon Photoirradiation. *Tetrahedron Lett.* **2004**, *45* (6), 1155–1158. <https://doi.org/10.1016/j.tetlet.2003.12.005>.
- (18) Takeshita, M.; Nagai, M.; Yamato, T. A Photochromic Thiophenophan-1-Ene. *Chem. Commun.* **2003**, No. 13, 1496–1497. <https://doi.org/10.1039/b303916a>.
- (19) Takeshita, M.; Jin-nouchi, H.; Ishikawa, J. Syntheses and Photochromic Properties of Metacyclothiophenophan-1-Enes. *Chem. Lett.* **2009**, *38* (10), 982–983. <https://doi.org/10.1246/cl.2009.982>.
- (20) Takeshita, M.; Yamato, T. Enantioselective Photochromic Reaction of a 2.2 Metacyclophan-1-Ene. *Angew. Chem.-Int. Ed.* **2002**, *41* (12), 2156–2157. [https://doi.org/10.1002/1521-3773\(20020617\)41:12<2156::aid-anie2156>3.0.co;2-0](https://doi.org/10.1002/1521-3773(20020617)41:12<2156::aid-anie2156>3.0.co;2-0).
- (21) Fukumoto, S.; Nakashima, T.; Kawai, T. Intramolecular Hydrogen Bonding in a Triangular Dithiazolyl-Azaindole for Efficient Photoreactivity in Polar and Nonpolar Solvents. *Eur. J. Org. Chem.* **2011**, No. 26, 5047–5053. <https://doi.org/10.1002/ejoc.201100676>.
- (22) Ishibashi, Y.; Okuno, K.; Ota, C.; Umesato, T.; Katayama, T.; Murakami, M.; Kobatake, S.; Irie, M.; Miyasaka, H. Multiphoton-Gated Cycloreversion Reactions of Photochromic Diarylethene Derivatives with Low Reaction Yields upon One-Photon Visible Excitation. *Photochem. Photobiol. Sci.* **2010**, *9* (2), 172–180. <https://doi.org/10.1039/b9pp00116f>.
- (23) Ishibashi, Y.; Mukaida, M.; Falkenstrom, M.; Miyasaka, H.; Kobatake, S.; Irie, M. One- and Multi-Photon Cycloreversion Reaction Dynamics of Diarylethene Derivative with Asymmetrical Structure, as Revealed by Ultrafast Laser Spectroscopy. *Phys. Chem. Chem. Phys.* **2009**, *11* (15), 2640–2648. <https://doi.org/10.1039/b818591c>.
- (24) Kumpulainen, T.; Lang, B.; Rosspeintner, A.; Vauthey, E. Ultrafast Elementary Photochemical Processes of Organic Molecules in Liquid Solution. *Chem. Rev.* **2017**, *117* (16), 10826–10939. <https://doi.org/10.1021/acs.chemrev.6b00491>.
- (25) Sotome, H.; Kitagawa, D.; Nakahama, T.; Ito, S.; Kobatake, S.; Irie, M.; Miyasaka, H. Cyclization Reaction Dynamics of

- an Inverse Type Diarylethene Derivative as Revealed by Time-Resolved Absorption and Fluorescence Spectroscopies. *Phys. Chem. Chem. Phys.* **2019**, *21* (17), 8623–8632. <https://doi.org/10.1039/c8cp07393g>.
- (26) Jarota, A.; Pastorczak, E.; Abramczyk, H. A Deeper Look into the Photocycloreversion of a Yellow Diarylethene Photoswitch: Why Is It so Fast? *Phys. Chem. Chem. Phys.* **2020**, *22* (10), 5408–5412. <https://doi.org/10.1039/c9cp05452a>.
- (27) Seo, K.; Eom, I.; Shim, S.; Kim, C. H.; Joo, T. Ring Closure Reaction Pathway of a Diarylethene in Solution Using Femtosecond Time-Resolved Fluorescence Spectra. *Bull. Korean Chem. Soc.* **2019**, *40* (4), 352–358. <https://doi.org/10.1002/bkcs.11689>.
- (28) Nakamura, S.; Yokojima, S.; Uchida, K.; Tsujioka, T.; Goldberg, A.; Murakami, A.; Shinoda, K.; Mikami, M.; Kobayashi, T.; Kobatake, S.; Matsuda, K.; Irie, M. Theoretical Investigation on Photochromic Diarylethene: A Short Review. *J. Photochem. Photobiol. Chem.* **2008**, *200* (1), 10–18. <https://doi.org/10.1016/j.jphotochem.2008.05.005>.
- (29) Perrier, A.; Aloise, S.; Olivucci, M.; Jacquemin, D. Inverse versus Normal Dithienylethenes: Computational Investigation of the Photocyclization Reaction. *J. Phys. Chem. Lett.* **2013**, *4* (13), 2190–2196. <https://doi.org/10.1021/jz401009b>.
- (30) Boggio-Pasqua, M.; Ravaglia, M.; Bearpark, M. J.; Garavelli, M.; Robb, M. A. Can Diarylethene Photochromism Be Explained by a Reaction Path Alone? A CASSCF Study with Model MMVB Dynamics. *J. Phys. Chem. A* **2003**, *107* (50), 11139–11152. <https://doi.org/10.1021/jp036862e>.
- (31) Aloise, S.; Sliwa, M.; Pawlowska, Z.; Rehault, J.; Dubois, J.; Poizat, O.; Buntinx, G.; Perrier, A.; Maurel, F.; Yamaguchi, S.; Takeshita, M. Bridged Photochromic Diarylethenes Investigated by Ultrafast Absorption Spectroscopy: Evidence for Two Distinct Photocyclization Pathways. *J. Am. Chem. Soc.* **2010**, *132* (21), 7379–7390.
- (32) Hamdi, I.; Buntinx, G.; Perrier, A.; Devos, O.; Jaidane, N.; Delbaere, S.; Tiwari, A. K.; Dubois, J.; Takeshita, M.; Wada, Y.; Aloise, S. New Insights into the Photoswitching Mechanisms of Normal Dithienylethenes. *Phys. Chem. Chem. Phys.* **2016**. <https://doi.org/10.1039/C6CP03471C>.
- (33) Hamdi, I.; Buntinx, G.; Poizat, O.; Delbaere, S.; Perrier, A.; Yamashita, R.; Muraoka, K.; Takeshita, M.; Aloise, S. Unraveling Ultrafast Dynamics of the Photoswitchable Bridged Dithienylethenes under Structural Constraints. *Phys. Chem. Chem. Phys.* **2019**, *21* (12), 6407–6414. <https://doi.org/10.1039/C8CP07100D>.
- (34) Hamdi, I.; Buntinx, G.; Poizat, O.; Perrier, A.; Le Bras, L.; Delbaere, S.; Barrau, S.; Louati, M.; Takeshita, M.; Tokushige, K.; Takao, M.; Aloise, S. Excited-State Dynamics of Dithienylethenes Functionalized for Self-Supramolecular Assembly. *J. Phys. Chem. A* **2018**, *122* (14), 3572–3582. <https://doi.org/10.1021/acs.jpca.7b10767>.
- (35) Irie, M. Photochromic Diarylethenes for Photonic Devices. *Pure Appl. Chem.* **1996**, *68* (7), 1367–1371.
- (36) Kobatake, S.; Muto, H.; Irie, M. Excited State Energy Migration and Photochromic Reaction in 1,2-Bis(2,4-Dimethyl-3-Thienyl)Perfluorocyclopentene Single Crystal. *Chem. Lett.* **2005**, *35* (1), 102–103. <https://doi.org/10.1246/cl.2006.102>.
- (37) Irie, M.; Sakemura, K.; Okinaka, M.; Uchida, K. Photochromism of Dithienylethenes with Electron-Donating Substituents. *J. Org. Chem.* **1995**, *60* (25), 8305–8309.
- (38) Irie, M.; Eriguchi, T.; Takada, T.; Uchida, K. Photochromism of Diarylethenes Having Thiophene Oligomers as the Aryl Groups. *Tetrahedron* **1997**, *53* (36), 12263–12271.
- (39) Higashiguchi, K.; Matsuda, K.; Yamada, T.; Kawai, T.; Irie, M. Fatigue Mechanism of Photochromic 1,2-Bis(3-Thienyl)Perfluorocyclopentene. *Chem. Lett.* **2000**, *29* (12), 1358–1359. <https://doi.org/10.1246/cl.2000.1358>.
- (40) Higashiguchi, K.; Matsuda, K.; Kobatake, S.; Yamada, T.; Kawai, T.; Irie, M. Fatigue Mechanism of Photochromic 1,2-Bis(2,5-Dimethyl-3-Thienyl)Perfluorocyclopentene. *Bull. Chem. Soc. Jpn.* **2000**, *73* (10), 2389–2394. <https://doi.org/10.1246/bcsj.73.2389>.
- (41) Buntinx, G.; Poizat, O.; Leygue, N. TRANSIENT ABSORPTION AND RESONANCE RAMAN INVESTIGATION OF THE PHOTOCHEMICAL REACTIVITY OF 2,2'-BIPYRIMIDINE IN SOLUTION. *J. Phys. Chem.* **1995**, *99* (8), 2343–2352. <https://doi.org/10.1021/j100008a016>.
- (42) Buntinx, G.; Naskrecki, R.; Poizat, O. Subpicosecond Transient Absorption Analysis of the Photophysics of 2,2'-Bipyridine and 4,4'-Bipyridine in Solution. *J. Phys. Chem.* **1996**, *100* (50), 19380–19388. <https://doi.org/10.1021/jp960900v>.
- (43) Moine, B.; Rehault, J.; Aloise, S.; Micheau, J. C.; Moustrou, C.; Samat, A.; Poizat, O.; Buntinx, G. Transient Absorption Studies of the Photochromic Behavior of 3H-Naphtho[2,1-b]Pyrans Linked to Thiophene Oligomers via an Acetylenic Junction. *J. Phys. Chem. A* **2008**, *112* (21), 4719–4726. <https://doi.org/10.1024/jp711865j>.
- (44) Nakayama, T.; Amijima, Y.; Ibuki, K.; Hamanoue, K. Construction of a Subpicosecond Double-Beam Laser

Photolysis System Utilizing a Femtosecond Ti:Sapphire Amplifiers (a Regenerative Amplifier and Two Double Passed Linear Amplifiers), and Measurements of the Transient Absorption Spectra by a Pump-Probe Method. *Rev Sci Instrum* **1997**, 68 (12), 4364–4371.

- (45) Ziolk, M.; Lorenc, M.; Naskrecki, R. Determination of the Temporal Response Function in Femtosecond Pump-Probe Systems. *Appl. Phys. B-Lasers Opt.* **2001**, 72 (7), 843–847.