Break-and-Build Strategy for the Synthesis of 2-Benzoylbenzoxazoles from o-Aminophenols and Acetophenones

Le Anh Nguyen, Quoc Anh Ngo, Pascal Retailleau, Thanh Binh Nguyen

► To cite this version:

Le Anh Nguyen, Quoc Anh Ngo, Pascal Retailleau, Thanh Binh Nguyen. Break-and-Build Strategy for the Synthesis of 2-Benzoylbenzoxazoles from o-Aminophenols and Acetophenones. *Advanced Synthesis and Catalysis*, 2021, 363 (8), pp.2098-2103. 10.1002/adsc.201 . hal-03325810

HAL Id: hal-03325810

<https://hal.science/hal-03325810>

Submitted on 25 Aug 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Break-and-Build Strategy for the Synthesis of 2-Benzoylbenzoxazoles from *o*-Aminophenols and Acetophenones

Le Anh Nguyen,^{b,c} Quoc Anh Ngo,^{*b,c} Pascal Retailleau,^a and Thanh Binh Nguyen^{*a}

^a Institut de Chimie des Substances Naturelles, CNRS UPR 2301, Université Paris-Sud, Université Paris-Saclay, 1 avenue de la Terrasse, 91198 Gif-sur-Yvette, France

^b Institute of Chemistry, Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet, Cau Giay, Hanoi, Vietnam

^c Graduate University of Science and Technology, Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet, Cau Giay, Hanoi, Vietnam

E-mail: thanh-binh.nguyen@cnrs.fr; ngoqanh@ich.vast.vn

Received: ((will be filled in by the editorial staff))

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/adsc.201#####>. ((Please delete if not appropriate))

Abstract. Although compounds with a 2-benzoylbenzoxazole motif are biologically relevant, there are only a few methods for synthesizing them, most of which relied on multistep process or required substrates bearing activating groups. Herein, we report an efficient method for the synthesis of such compounds by direct reactions of *o*-aminophenols with acetophenones promoted by sulfur in DMSO. The reaction was found to proceed via a Willgerdt rearrangement-type benzoxazolation of acetophenones followed by a benzylic oxidation to reinstall the carbonyl function. This method has a broad substrate scope and good tolerance for sensitive functional groups.

Keywords: sulfur; 2-benzoylbenzoxazole; DMSO

Benzoxazole derivatives are prevalent in biologically and pharmacologically important molecules as well as in functional materials.^[1] A subgroup of this family is 2-benzoylbenzoxazoles, which are well-known for their significant biological activities such as antiproliferative activity (compound I),^[2] inhibition of fatty acid amide hydrolase FAAH^[3] (compound II) (Scheme 1). Consequently, several methods have been developed for the synthesis of 2-benzoylbenzoxazoles,^[4] most of them were based on functionalization of simpler benzoxazole cores IX^[5] or using complex starting materials which required multi-step syntheses.^[6]

The intuitive approach based on the construction of the benzoxazole ring by condensation of *o*-aminophenols **1** were described with phenylglyoxalic derivatives such as dithioester III^[7] or imidoyl cyanide IV^[8] via non redox condensation. Alternatively, when oxidation conditions were applied, condensation of **1** with derivatives of lower oxidation states such as α,α -dihaloacetophenones V^[9] or 2-bromophenylacetylenes VI^[10] were known. In this context, a direct use of

acetophenones **2**, which are inexpensive and readily available in a wide range of structure, in a selective oxidative condensation with *o*-aminophenols **1** would provide a convenient and cost-effective approach to a library of 2-benzoylbenzoxazoles for drug discovery study.

Representative bioactive 2-benzoylbenzoxazoles:

Known synthetic methods: complex starting materials

Our method: oxidative coupling of 1 with 2 using elemental sulfur and DMSO

advantages compared to known methods:

- + inexpensive, readily available, user friendly acetophenone substrates
- + inexpensive S and DMSO
- + simple reaction conditions

Scheme 1. 2-Benzoylbenzoxazoles: Bioactivity and Syntheses

We have recently reported a Willgerdt rearrangement-type benzoxazolation of acetophenones

2 with *o*-aminophenols **1** promoted by sulfur^[11] and *N*-methylpiperidine (Scheme 2).^[12] In this reaction leading to **4aa**, while the methyl group of acetophenone **2a** was oxidized and benzoxazolized with 2-aminophenol **1a**, the carbonyl group was reduced to a methylene group. Since this methylene group is surrounded by the phenyl group and the newly installed benzoxazole moiety, this new situation could open up opportunities for oxidation of **4aa** into a carbonyl in **3aa**. Such operation was previously performed on separated 2-benzylbenzoxazole derivatives using relatively complex reaction conditions involving strong oxidizing agents in the presence of transition metal catalysts.^[5m] We reason that a direct transformation from **1a** and **2a** to **3a** could be performed in one-pot in the presence of a suitable oxidant in the reaction medium. As DMSO is well known as an excellent solvent and a mild and selective oxidizing agent, especially in the presence of sulfur,^[13] we focused our attention to develop this strategy.

Scheme 2. Break-and Build Strategy for the Synthesis of **3**

For this purpose, we first performed the benzoxazolation of acetophenone **2a** with 2-aminophenol **1a** under the standard conditions (80 °C, 16h) using sulfur (3 equiv), *N*-methylpiperidine (NMP) (1 equiv) in the presence of DMSO (3 equiv) (Table 1, entry 1).

To our delight, this initial test resulted in a 80% conversion of 2-aminophenol **1a** into a nearly 1:1 mixture of **4aa**:**3aa**. Oxidation of the methylene moiety was significantly accelerated by heating at higher temperature (110 °C, entry 2). At this stage, we realized that DMSO could act as oxidant to regenerate sulfur from H₂S byproduct of the first step of Willgerodt-type benzoxazolation, the amount of sulfur could be lowered to stoichiometric amount, i.e. 1 equiv (entry 3).

Moreover, we noticed that the amount of NMP, which acted as a basic sulfur activator, could be further reduced to 0.5 equiv without lowering the reactivity (entries 4-5). Interestingly, other nitrogen bases such as pyridine, *N*-methylmorpholine (NMM), diisopropylethylamine (DIPEA), tri-*n*-propylamine or 1,4-diazabicyclo[2.2.2]octane (DABCO) (entries 6-10) could be used in place of NMP. It is noteworthy that the expected product **3aa** could be formed even in the absence of a base, despite in lower yield (entry 11). The reaction in the absence of sulfur resulted in recovery of unchanged starting materials, which confirmed clearly the crucial role of this element in promoting the cascade transformation (entry 12).

Finally, lowering the DMSO amount (entry 13) of replacing DMSO by another polar aprotic solvent such as DMF (entry 14) prevented partially or totally the formation of **3aa**, which confirmed the importance of DMSO in this oxidative condensation.

Table 1. Screening of the Reaction Conditions

Entry ^[a]	x	additive	y	yield (%) ^[b]
1 ^[c]	3	<i>N</i> -methylpiperidine	1	30
2	3	<i>N</i> -methylpiperidine	1	65
3	1	<i>N</i> -methylpiperidine	1	68
4	1	<i>N</i> -methylpiperidine	0.5	72
5	0.5	<i>N</i> -methylpiperidine	0.5	54
6	1	pyridine	0.5	70
7	1	<i>N</i> -methylmorpholine	0.5	77
8	1	DIPEA	0.5	71
9	1	tri- <i>n</i> -propylamine	0.5	70
10	1	DABCO	0.5	73
11	1	-	0	52
12	0	<i>N</i> -methylmorpholine	0.5	0
13 ^e	1	<i>N</i> -methylmorpholine	0.5	36
14 ^f	1	<i>N</i> -methylmorpholine	0.5	0 ^[d]

^[a] Reaction conditions: *o*-aminophenol **1a** (1 mmol), acetophenone **2a**, sulfur (x mmol, 32 mg.mmol⁻¹), additive (y mmol), DMSO (0.5 mL), 110 °C, 16 h unless otherwise noted. ^[b] isolated yield unless otherwise noted. ^[c] Reaction temperature 80 °C. ^[d] Determined by ¹H NMR of the crude mixture. ^[e] DMSO (0.3 mL) was used. ^[f] DMF (0.5 mL) was used in place of DMSO.

With this set of optimized reaction conditions, we next studied the scope and limitations of our strategy. In general, starting from 2-aminophenol **1a** and acetophenones **2** bearing a wide range of substituents of different electronic nature or positions, the reactions led to the expected 2-benzoylbenzoxazole products **3** in reasonable yields (Scheme 3). Acetophenones bearing electron donating groups such as methyl, methoxy, acetamido displayed good reactivity, provide the expected benzoxazoles **3ab-3aj** in moderate to good yields. We emphasized that the product **3ai** (compound **I** in Scheme 1) was previously demonstrated to be biologically active.^[2] Interestingly, the reaction conditions could be applied to acetophenones bearing unprotected hydroxy groups in good to excellent yields. It should be noted that **3al** could serve as a direct precursor to provide a wide range of bioactive derivative via acylation of its hydroxy group (compound **II** in Scheme 1). Acetophenones bearing halogen groups in different positions such as *p*-Cl, *m*-Cl, *o*-F and *m*-Br (**2m-2p**) displayed good reactivities and gave the products **3am-3ap** in good yields. Gratifyingly, acetophenones substrates having an electron withdrawing group such as CF₃, CN, CO₂Me

or NO₂, showed also to be competent substrates. The structure of nitro **3aw** was unambiguously confirmed by X-ray diffraction analysis.

Scheme 3. Scope of Methyl Aryl Ketones 2

Although CN and NO₂ groups were known to react with H₂S,^[14] which is a possible intermediate/byproduct of this transformation, via nucleophilic addition and reduction, respectively, the expected benzoxazole products **3as**, **3at**, **3av** and **3ax** were obtained in reasonable yields. In case of *p*-nitroacetophenone **2v**, the superiority of NMM compared to other basic additive such as NMP and pyridine or without base in assisting the

benzoxazolation process was clearly demonstrated as NMM was the only base that could afford the expected benzoxazole **3av**, despite in low yield.

2-Acetonaphthone as well as heterocyclic substrates derived from pyridine, thiophene and benzofuran gave moderate to good yields of the corresponding benzoxazole products **3ax-3aaa**. The twofold reaction could be performed when *p*-diacetylbenzene (0.6 equiv) was used.

The reaction conditions could also be applied successfully to substituted 2-aminophenols (Scheme 4). Benzoxazoles bearing a methyl group **3ba** and **3ca** were obtained in yields of 63% and 73%, respectively. Chloro substituents were also compatible with the reaction conditions: the oxidative coupling products **3da** and **3ea** were formed in moderate yields (63% and 57%, respectively). The presence of an electron withdrawing groups such as sulfone, sulfonamide or nitro on the phenyl ring of 2-aminophenol could reduce the nucleophilicity of the hydroxy group, they had little effect on the yields. In the case of 4-nitro-2-aminophenol **1g**, thanks to low solubility in methanol of product **3ga**, it was easily isolated from the reaction mixture by a simple trituration/filtration operation with methanol. The structure of sulfone **3fa** obtained in 61% yield was confirmed by X-ray diffraction analysis. *o*-Aminophenol substrate bearing an electron donating group MeO **1g** showed also good reactivity under our conditions, leading to benzoxazole **3ga** in good yield.

Scheme 4. Scope of *o*-Aminophenols 1

Our reaction conditions were however not applicable to aliphatic methyl ketones such as pinacolone, 3-methyl-2-butanone or 2-heptanone (Scheme 5, eq. 1). In these cases, the reactions mixtures were found to be complex, possibly because both step of benzoxazolation of methyl ketones and methylene oxidation occurred more difficultly in the absence of an aryl group on the methyl ketone substrates. Indeed, the presence of the products issued from the first step could be detected in the crude

mixtures, the subsequent oxidation of their methylene moiety failed. The reaction conditions applied to homologous of acetophenones such as propiophenone, butyrophenone or valerophenone led to complex mixtures since the intermediates of Willgerodt rearrangement of the longer chains underwent uncontrollable oxidation by DMSO (Scheme 5, eq. 2).

Scheme 5. Control Experiments

In order to support the proposed reaction mechanism, the primary Willgerodt-type product **4aa** was prepared separately and allowed to react with DMSO (0.5 equiv) (Scheme 5, eq. 3). The product **3aa** was formed in high yield under the standard conditions. On the other hand, performing the similar reaction in the absence of sulfur resulted in recovery of **4aa**. This confirmed unambiguously that the oxygen atom of 2-benzoylbenzoxazole **3aa** product comes from DMSO.

Additionally, we found that diphenylmethane could not be oxidized into benzophenone and remained unchanged under these oxidizing conditions (eq. 4). These results suggested the important role of both aryl and 2-benzoxazolyl substituents as well as the participation of sulfur to the oxidation of methylene moiety.

Scheme 6. Proposed Reaction Mechanism

On the basis of these results and our previous works,^[13] a plausible mechanism is proposed in Scheme 6. The reaction commenced with a Willgerodt-type reaction as described previously to provide benzoxazole **4aa**.

An easily tautomerization **4aa** $\rightleftharpoons$ **4aa'** facilitated by the presence of both benzoxazole and phenyl moieties would favor the formation of polysulfide salt **5**. Sulfur extrusion from **5** followed by a sulfur-promoted oxidation of thiol **6** to thione **8** would proceed via polysulfide **7** and its tautomer **7'**. Hydrolysis of thione **8** would lead to **3aa**.^[15] Since the oxygen atom of a carbonyl function of a ketone was known to be exchanged rapidly with water, it is not conclusive to use labelled water to confirm if the ketone oxygen atom **3aa** was issued from water or not via the hydrolysis step **8** $\rightarrow$ **3aa**.^[16] Water involving in this step is generated from the recycling of sulfur by oxidation of hydrogen (poly)sulfide with DMSO.^[17] At the present stage, we cannot exclude the pathway involving direct functionalization of the methyl group of acetophenone **1a** without the intermediacy of a Willgerodt-type reaction.

In conclusion, we have reported an efficient method for a direct access to 2-benzoylbenzoxazoles from 2-aminophenols and acetophenones via a cascade of sulfur-promoted Willgerodt type benzoxazolation and methylene oxidation in DMSO. The reactions conditions are compatible with many functional groups and heterocycles. The method is highlighted by the fact that both starting materials *o*-aminophenols and acetophenones are inexpensive and readily available in a wide range of structures. Moreover, since sulfur and DMSO were used as oxidant in the presence of substoichiometric amount of *N*-methylmorpholine as an additive, our method is inarguably the cheapest and most convenient method to provide a library of 2-benzoylbenzoxazoles.

Experimental Section

A mixture of 2-aminophenol **1** (1 mmol), acetophenone (1.2 mmol), elemental sulfur (32 mg, 1 mmol), *N*-methylmorpholine (56 mg, 0.5 mmol), and DMSO (0.5 mL) was heated under an argon atmosphere in a 7-mL test tube at 110 °C for 16 h. The product was purified by column chromatography on silica gel (heptane:EtOAc 1:0 to 5:1 or dichloromethane:heptane 2:1 to 1:0). For experimental details, full characterization, and copies of NMR spectra of all compounds, see Supporting Information.

CCDC 2045870 and 2042799 contain the supplementary crystallographic data for **3aw** and **3fa**, respectively in this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

Acknowledgements

We thank Vietnam Academy of Science and Technology for valuable financial support (KHCBHH.01/18-20). We thank Dr. A. Baron, C. Moriou, S. Beaupierre and S. Thoret for their valuable assistance.

References

- [1] For a selected review on benzoxazoles, see: a) C. S. Demmer, L. Bunch, *Eur. J. Med. Chem.* **2015**, *97*, 778.
- [2] J. Chen, C.-M. Li, J. Wang, S. Ahn, Z. Wang, Y. Lu, J. T. Dalton, D. D. Miller, W. Li, *Bioorg. Med. Chem.* **2011**, *19*, 4782.
- [3] M. Seierstad, J. G. Breitenbucher, *J. Med. Chem.* **2008**, *51*, 7327; b) A. Minkkilä, S. M. Saario, T. Nevalainen, *Cur. Top. Med. Chem.* **2010**, *10*, 828; c) M. J. Myllymäki, H. Kaesnaenen, A. O. Kataja, M. Lahtela-Kakkonen, S. M. Saario, A. Poso, A. M. P. Koskinen, *Eur. J. Med. Chem.* **2009**, *44*, 4179; d) M. J. Myllymäki, S. M. Saario, A. O. Kataja, J. A. Castillo-Melendez, T. Nevalainen, R. O. Juvonen, T. Järvinen, A. M. P. Koskinen, *J. Med. Chem.* **2007**, *50*, 4236; e) H. Kaesnaenen, M. J. Myllymäki, A. Minkkilä, A. O. Kataja, S. M. Saario, T. Nevalainen, A. M. P. Koskinen, A. Poso, *ChemMedChem*, **2010**, *5*, 213; f) K. Otrubova, C. Ezzili, D. L. Boger, *Bioorg. Med. Chem. Lett.* **2011**, *21*, 4674.
- [4] N. Aljaar, R. Gujjarappa, M. Al-Refai, M. Shtaiwi, C. C. J. Malakar, *Heterocyclic Chem.* **2019**, *56*, 2730.
- [5] For benzylation of 2-*H* benzoxazoles, see: a) N. K. Harn, C. J. Gramer, B. A. Anderson, *Tetrahedron Lett.* **1995**, *36*, 9453; b) X.-F. Wu, P. Anbarasan, H. Neumann, M. Beller, *Angew. Chem., Int. Ed.* **2010**, *49*, 7316; c) J. Chen, C.-M. Li, J. Wang, S. Ahn, Z. Wang, Y. Lu, J. T. Dalton, D. D. Miller, W. Li, *Bioorg. Med. Chem.* **2011**, *19*, 4782; d) P. Lassalas, F. Marsais, C. Hoarau, *Synlett*, **2013**, *24*, 2233; e) S. Sharma, I. A. Khan, A. K. Saxena, *Adv. Synth. Catal.* **2013**, *355*, 673; f) K. Yang, C. Zhang, P. Wang, Y. Zhang, H. Ge, *Chem. - Eur. J.* **2014**, *20*, 7241; g) K. Yang, X. Chen, Y. Wang, W. Li, A. A. Kadi, H. K. Fun, H. Sun, Y. Zhang, G. Li, H. Lu, *J. Org. Chem.* **2015**, *80*, 11065; h) S. Karthik, T. Gandhi, *Org. Lett.* **2017**, *19*, 5486; i) W. Jia, Y. Jian, B. Huang, C. Yang, W. Xia, *Synlett*, **2018**, *29*, 1881; j) Y. Ogiwara, Y. Iino, N. Sakai, *Chem. Eur. J.* **2019**, *25*, 6513. For transformation of 2-substituted benzoxazoles, see: (k) Q. Y. Toh, A. McNally, S. Vera, N. Erdmann, M. J. Gaunt, *J. Am. Chem. Soc.* **2013**, *135*, 3772; l) Z. Yin, Z. Zhang, J. F. Kadow, N. A. Meanwell, T. Wang, *J. Org. Chem.* **2004**, *69*, 1364; m) H. Sterckx, J. De Houwer, C. Mensch, I. Caretti, K. A. Tehrani, W. A. Herrebout, S. Van Doorslaer, B. U. W. Maes, *Chem. Sci.* **2016**, *7*, 346; n) H. Wang, J. Liu, J. P. Qu, Y. B. Kang, *J. Org. Chem.* **2020**, *85*, 3942; o) J. Kumar, A. Gupta, S. Bhadra, *Org. Biomol. Chem.* **2019**, *17*, 3314; p) W. A. Skinner, F. Gualtiere, G. Brody, A. H. Fieldsteel, *J. Med. Chem.* **1971**, *14*, 546.
- [6] For an example of redox condensation between *o*-nitrosophenols and phenacyl bromides, see: a) N. Aljaar, C. C. Malakar, J. Conrad, U. Beifuss, *J. Org. Chem.* **2015**, *80*, 10829. For an example of cyclization of densely functionalized *N*-substituted *o*-aminophenols, see: (b) S. S. K. Boominathan, W.-P. Hu, G. C. Senadi, J. K. Vandavasi, J.-J. Wang, *Chem. Commun.* **2014**, *50*, 6726.
- [7] S. M; Anil, K. R; Kiran, K. S; Rangappa, M. P; Sadashiva, J. B; Shruthi, K. P; Sukrutha, T. R. Swaroop, *Synthesis* **2019**, *51*, 4205.
- [8] L. Zhao, B. Liu, Q. Tan, C. Ding, B. Xu, *Org. Lett.* **2019**, *21*, 9223.
- [9] C. Wu, X. Xin, Z. Fu, L. Xie, K. Liu, Z. Wang, W. Li, Z. Yuan, W. He, *Green Chem.* **2017**, *19*, 1983; b) J. Jiang, H. Zou, Q. Dong, R. Wang, L. Lu, Y. Zhu, W. He, *J. Org. Chem.* **2016**, *81*, 51; c) X. Zhang, Y. Wu, Y. Zhang, H. Liu, Z. Xie, S. Fu, F. Liu, *Tetrahedron*. **2017**, *73*, 4513.
- [10] L. Cui, Y. He, X. Fan, *Chinese J. Chem.* **2012**, *30*, 992.
- [11] For recent reviews on the organic transformations involving elemental sulfur, see: a) T. B. Nguyen, *Adv. Synth. Catal.* **2020**, *362*, 3448; b) T. B. Nguyen, *Adv. Synth. Catal.* **2017**, *359*, 1106; c) T. B. Nguyen, *Asian J. Org. Chem.* **2017**, *6*, 477.
- [12] T. B. Nguyen, P. Retailleau, *Org. Lett.* **2017**, *19*, 3887.
- [13] For our recent results on the development of this pair of oxidants, see: a) T. B. Nguyen, D. H. Mac, P. Retailleau, *J. Org. Chem.* **2020**, *85*, 13508; b) L. A. Nguyen, T. D. Dang, Q. A. Ngo, T. B. Nguyen, *Eur. J. Org. Chem.* **2020**, 3818; d) T. T. T. Nguyen, V. A. Le, P. Retailleau, T. B. Nguyen, *Adv. Synth. Catal.* **2020**, *362*, 160; For selected references on chemistry of DMSO, see: e) X. F. Wu, K. Natte, *Adv. Synth. Catal.* **2016**, *358*, 336; f) E. Jones-Mensah, M. Karki, J. Magolan, *Synthesis* **2016**, *48*, 1421, g) D. Fu, J. Dong, H. Du, J. Xu, *J. Org. Chem.* **2020**, *85*, 2752.
- [14] M. L. Boys, V. L. Downs, *Synth. Commun.* **2006**, *36*, 295.
- [15] R. Okazaki, N. Tokitoh, *Acc. Chem. Res.* **2000**, *33*, 625.
- [16] M. Byrn, M. Calvin, *J. Am. Chem. Soc.* **1966**, *88*, 1916; (b) M. Cohn, H. C. Urey, *J. Am. Chem. Soc.* **1938**, *60*, 679.
- [17] Y. Mehmet, J. B. Hyne, *Phosphorus Sulfur Silicon Relat. Elem.* **1976**, *1*, 47.

Break-and-Build Strategy for the Synthesis of 2-Benzoylbenzoxazoles from *o*-Aminophenols and Acetophenones

Adv. Synth. Catal. **Year**, *Volume*, Page – Page

Le Anh Nguyen,^{b,c} Quoc Anh Ngo,^{*b,c} Pascal Retailleau,^a and Thanh Binh Nguyen^{*a}