**New insights in bacterial and eukaryotic diversity of microbial mats
inhabiting exploited and abandoned salterns at the Ré Island (France)**

Camille Mazière^{a,b}, Hélène Agogué^b, Cristiana Cravo-Laureau^a, Christine Cagnon^a, Isabelle Lanneluc^b, Sophie Sablé^b, Ingrid Fruitier-Arnaudin^b, Christine Dupuy^b, Robert Duran^{a*}

^a Université de Pau et des Pays de l'Adour, E2S UPPA, CNRS, IPREM, Pau, France

^b UMR 7266 LIENSs (Littoral Environnement et Sociétés), CNRS - La Rochelle Université – 2, rue Olympe de Gouges, 17000 LA ROCHELLE, France

* Corresponding author: robert.duran@univ-pau.fr, +33 5 59 40 74 68, Bâtiment IBEAS - Avenue de l'Université, BP 1155 64013 PAU CEDEX

ABSTRACT

In order to understand the effect of human practices on microbial mats organisation, the study aimed to investigate the biodiversity within microbial mats from exploited and abandoned salterns. Despite several attempts, archaeal 16S rRNA gene fragment sequences were not obtained, indicating that microbial mats were probably dominated by Bacteria with very low abundance of Archaea (< 1%). Thus, the study compared the bacterial and meiofaunal diversity of microbial mats from abandoned and exploited salterns. The higher salinity (101 ± 3.7 psu vs. 51.1 ± 0.7 psu; Welch t-test $p < 0.05$) of the exploited site maintained lower bacterial diversity in comparison to the abandoned site where the salinity gradient was no longer maintained. However, the microbial mats exhibited similar bacterial class composition while the eukaryotic diversity was significantly higher in the exploited saltern. The abandoned saltern was dominated by sulfate-reducing bacteria and Nematoda, while the exploited saltern was characterized by the presence of halophilic bacteria belonging to *Marinobacter*, *Salinivibrio* and *Rhodohalobacter* genera, and the larger abundance of Hypotrichia (ciliates). Such bacterial and eukaryotic diversity difference might be explained by human actions for salt recovery in exploited salterns such as scraping the surface of microbial mat and increasing salinity renewing the microbial mat each year. Such action decreases the bacterial diversity changing the food web structure that favour the presence of a larger diversity of eukaryotic organisms. Our study provides new insights on microbial mat communities inhabiting salterns, especially the consequences of abandoning saltern exploitation.

Keywords: Hypersaline environments, marine solar salterns, microbial diversity, meiofaunal diversity, human exploitation and perturbation

Funding: C.M. was supported by a PhD grant from E2S-UPPA program and the Région Nouvelle-Aquitaine. We thank the funding support from the ACI U. La Rochelle and the European programme ERANETMED AQUASALT (NMED-0003-01).

Conflicts of interest/Competing interests: No conflict of interest.

Availability of data and material: Sequencing data has been deposited in sequence database.

48 **Code availability:** Not applicable

1. Introduction

Microbial mats develop at the sediment/water interface in various habitats (Prieto-Barajas *et al.* 2018), including coastal areas (White *et al.* 2018, 2021), sand beaches (Bolhuis and Stal 2011), estuaries (Mir *et al.* 1991), lakes (Jørgensen and Cohen 1977), hot springs (Dobretsov *et al.* 2011), Andin glaciers (Fleming and Prufert - Bebout, 2010; Schmidt *et al.* 2017), polar region (Valdespino-Castillo *et al.* 2018), freshwater Lagunas (Falcón *et al.* 2020; Yanez-Montalvo *et al.* 2020), and salterns (Giani *et al.* 1989; Fourçans *et al.* 2004; Kolesnikov *et al.* 2017). They are complex systems of multi-layered microbial communities vertically stratified where microorganisms are organized according to physical-chemical gradients of light, oxygen and sulfur (Caumette *et al.* 1994), playing an important role in the major biogeochemical cycles (Jørgensen and Cohen 1977; Wieland *et al.* 2003; Sánchez-Baracaldo *et al.* 2021). In microbial mats, different functional microbial groups with various metabolic capacities coexist at a microscale (van Gemerden 1993). They experience daily fluctuations of environmental parameters resulting in variations of their vertical distribution (Fourçans *et al.* 2006, 2008; Martinez *et al.* 2019). Therefore, microbial mats constitute an ideal model to study microbial interactions and their response to environmental changes.

Microbial mats developing in marine solar salterns are among the most studied (Oren 2009) for examining microbial composition and their dynamics (Plominsky *et al.* 2018). Solar salterns, generally found in coastal regions, are man-made habitats for salt production through solar evaporation of seawater. Seawater evaporates through a series of shallow ponds, gradually increasing salinity constituting extreme hypersaline environments. The resulting brine, saturated with sodium chloride, reaches crystallizer ponds where salt deposits and is then collected by scraping the surface of the sediment. The salt production depends on physical-chemical processes and activities of the microbial mats developing in such extreme habitats, which could result in the production of several valuable chemicals by microorganisms with biotechnological potentials (Gómez-Villegas, Vigara and León 2018). Historically, several solar salterns have been exploited along the French Atlantic coast. Nowadays, many of these solar salterns have been abandoned because of loss of economic profitability. At the Ré Island (France), exploited and abandoned solar salterns coexist allowing the

comparison of microbial mats developing under different saline fluctuating conditions but with similar environmental parameters. The salt production is artisanal, handwork following traditional practices with salt harvest by hands in summer. In winter the salterns are flooded with seawater, the exploitation starting in spring by increasing progressively the salinity, especially in crystallizer ponds where large variations in salinity are observed until reaching concentrations where salt precipitates. In contrast, the salinity variation is limited in the abandoned salterns because the water entry is no regulated. It has been demonstrated that the salinity fluctuations during salt production disturb the natural habitats (Tran *et al.* 2019), influencing the development and the structure of the microbial mats that also depend on the seasonal variations (Boujelben *et al.* 2012; Cardoso *et al.* 2019). Because only few studies have investigated abandoned salterns (Cvitković *et al.* 2011; Lee *et al.* 2020), the information on how the ecosystem is affected by the modifications of human practices is still scarce. The comparison of the diversity within microbial mats subjected to large saline variations (exploited saltern) with that of microbial mats inhabiting ponds where the saline variations are limited (abandoned saltern) within the same hydrological system will allow to describe the modifications of microbial communities in response to a drastic change on the human practices, which are affecting the salinity fluctuations. Understanding how the human practices affect the microbial community of microbial mats in solar salterns is of paramount importance providing useful information for the management of microbial communities in the context of global changes.

In this study, we investigated the diversity within microbial mats from exploited and abandoned hypersaline ponds in salterns located at the Ré Island (France). We adopted a holistic view, investigating the microbial communities by 16S and 18S rRNA genes barcoding, completed by microscopic observations for the eukaryotic community, in order to describe the biodiversity in microbial mats and how it is affected by human-made drastic changes. Indeed, although bacterial communities inhabiting microbial mats have been widely studied (Fourçans *et al.* 2006, 2008; Bolhuis and Stal, 2011; Boujelben *et al.* 2012; Bolhuis *et al.*, 2014; Stal *et al.* 2019), only few studies have reported on their eukaryotic community composition (Cvitković *et al.* 2011; Edgcomb *et al.* 2014), resulting in a lack of knowledge in the whole diversity of microbial mats. Our results bring new

103 insights on microbial mats biodiversity modifications in response to a perturbation due to human
104 activity.

105 2. Material and methods

107 2.1 Sampling site

108 The microbial mats were collected in crystallizer ponds of salterns located in two different sites, but within
109 the same hydrological system, close to Ars-en-Ré (Ré Island, France; Fig. 1, the map was drawn with the R
110 library ggmap (Kahle and Wickham 2013)), allowing the investigation of abandoning saltern exploitation
111 considering similar physical-chemical parameters. The Abandoned site (46°12'44.928"N 1°30'20.123"W)
112 located in an abandoned saltern for 15 years, and the Exploited site (46°12'41.5"N 1°30'36.9"W) located in
113 an exploited saltern. The microbial mats were sampled in spring 2017. In each site, microbial mats were
114 sampled in independent biological triplicates (R1, R2 and R3) corresponding to three distinct ponds (Fig. 1).
115 Microbial mats were sampled using a PVC tube (15 cm diameter) as corer. From each core, the first 0.5 cm
116 was pushed out, from the bottom to the up, with a piston in order to collect the microbial mat of the expected
117 thickness. After homogenisation by mixing, subsamples for DNA microbial analysis were dispatched in 2 mL
118 cryotubes flash frozen in liquid nitrogen and then stored at -80°C. Meiofauna abundance and group
119 composition were obtained from another core. Sediment (60 mL) was preserved in absolute ethanol
120 (vol/vol).

121 The water salinity (Practical Salinity Unit, psu corresponding to g/L) and the water temperature (°C) were
122 measured in triplicate in each sampling spot with a multiparameter probe (pHenomenal® MU 6100H,
123 VWRTM, USA).

125 2.2 DNA extraction and Illumina sequencing

126 Microbial mat samples were ground in liquid nitrogen with a mortar and a pestle as previously described
127 (Fourçans et al., 2008). Then, DNA was extracted from 0.25 g subsamples using the DNeasy® PowerSoil kit
128 (Qiagen) according to the manufacturer's instructions, with a slight modification at the homogenization step,
129 by using a Precellys homogenizer (Bertin Instruments).

130 The V3-V4 region of the bacterial 16S rRNA genes (460 bp) was amplified using the primers PCR1F_460
131 (5'-ACGGRAGGCAGCAG-3') and PCR1R_460 (5'-TACCAGGGTATCTAATCCT-3') (Klindworth *et al.*
132 2013). The Primers 519F (5'-GCCGCCGCGGTAA-3') and 915R (5'-GTGCTCCCCCGCCAATTC-3') were
133 used to amplify the V3–V5 region (400 pb) of the archaeal 16S rRNA gene (Hugoni *et al.* 2015). Polymerase

chain reaction (PCR) mix consisted in 5 µL of MTP Taq DNA polymerase (5 U/µL) with 1 µL MTP Taq Buffer (10X) (Sigma-Aldrich), 1 µL of dNTP (10 mM), 1.25 µL of each primer (20 µM) and 10 ng of genomic DNA, in a final volume of 50 µL (adjusted with distilled water). All amplifications were performed on a Labcycler (SensoQuest) using the following PCR program: 2 min at 94°C, 30 cycles of 60 s at 94°C, 40 s at 65°C for Bacteria (40s at 57°C for Archaea), and 30 s at 72°C, and finally, 10 min at 72°C.

The V4 region (390 bp) of the 18S rRNA genes was amplified using the universal primer 515F (5'-GTGYCAGCMGCCGCGGTA-3') (Caporaso *et al.* 2011) and the eukaryotic primer 951R (5'-TTGGYRAATGCTTTCGC-3') (Lepère *et al.* 2016). PCR mix consisted in 27.5 µL of AmpliTaq Gold® 360 master mix (Applied Biosystems), 1.1 µL of each primer (20 µM) and 5.5 µL of genomic DNA, in a final volume of 55 µL (adjusted with distilled water). All amplifications were performed on a Veriti 96 Well Thermal Cycler (Applied Biosystem) using the following PCR program: 10 min at 95°C, 35 cycles of 30 s at 95°C, 30 s at 55°C and 45 s at 72°C, and finally, 10 min at 72°C.

All PCR were performed in triplicate for each sample and sequenced separately in order to obtain technical replicates in order to consider the PCR bias. Illumina sequencing was performed by the NED team (UMR1388 GenPhySE) and the GeT core facility (Toulouse, France), using Illumina MiSeq technology (paired-end 2 x 250 bp). The complete dataset was deposited in the NCBI Sequence Read Archive (SRA) database under SRA accession number PRJNA627371 for eukaryotic dataset and PRJNA627173 for bacterial dataset.

2.3 Sequence processing

Bioinformatic processing for DNA sequences were performed using QIIME 2 2019.4 (Bolyen *et al.* 2019). The same process was followed for 16S (Bacteria) and 18S (Eukaryote) rRNA gene sequences. Raw sequences were demultiplexed and then filtered, denoised, merged and grouped in non-chimeric sequences with DADA 2 (Callahan *et al.* 2016) followed by a singleton filtering. The taxonomic affiliation was performed against the Silva database v132 (Quast *et al.* 2012; Yilmaz *et al.* 2013) with 97% of similarities in order to compare the biodiversity obtained in previous studies, as it is used in most studies exploring biodiversity in microbial mats (Cardoso *et al.* 2019; Sierra *et al.* 2020; Vogt *et al.* 2018). Additionally, Silva database allows to analyse the three domains of life (Bacteria, Archaea, and Eukarya) while the other databases (e.g. Greengenes, RDP) will not (Balvočiūtė and Huson 2017). The non-affiliated sequences were

163 excluded. The rarefaction was done at 5,530 and 22,274 sequences per sample for bacteria and eukaryote
164 respectively (Supplementary materials, Fig. S1.), which correspond to the lowest number of sequences
165 considering the PCR replicates (data not shown).

166

167 **2.4 Meiofauna characterization**

168 Microbial mat was sieved through 50 μm before staining with rose Bengal and observation under a binocular
169 loupe (magnification x30, Zeiss). A sample splitter (Motoda box as Rzeznik-Orignac *et al.*, 2003) was used
170 to obtain an aliquot containing at least 100 individual nematodes for abundance estimation. The abundance
171 of other meiobenthic taxa (i.e. copepods and ostracodes) was too low to be evaluated in split samples and,
172 therefore, was quantified using whole samples. Abundances were expressed as individuals per cubic
173 centimetre (ind. cm^{-3}).

174

175 **2.5 Statistical analysis**

176 Statistical analyses were performed using Rstudio software (R version 3.6.3, (2020-02-29), R Core Team,
177 2020). The water temperature between the two sites was compared with a Student t-test because the
178 conditions of application (normality of data and variance independence) were verified and with a Welch two
179 sample t-test for salinity because the independence of variances was not verified. Then, the difference in the
180 relative microbial abundance was investigated between the abandoned and exploited sites. The biom file
181 produced by the bioinformatic analysis and the table of environmental data were merged into a single R
182 package using “phyloseq” (McMurdie and Holmes 2013). The statistical analyses were performed on the
183 average of the PCR triplicates (technical replicates) for each sample in order to consider the PCR bias. The
184 alpha diversity was calculated by two indices, the richness (based on presence/absence of Amplicon
185 Sequence Variants (ASVs)) and the Shannon index (according to the relative abundance of ASVs). The
186 abundances were compared at different taxonomic levels with a Student t-test if the conditions of application
187 (normality of data and variance independence) were verified, or with a Welch two sample t-test if the
188 normality was verified but not the independence of variances and with a Wilcoxon-Mann-Whitney test if any
189 condition was respected. The beta diversity between sites was then analysed with a principal coordinate
190 analysis (PCoA) of Bray-Curtis distance matrices. A dendrogram of each replicate of the sites was also done
191 with Bray-Curtis distance matrices. The contribution of each ASV explaining the differences between sites

192 was defined by SIMPER (SIMilarity PERcentages) based on Bray-Curtis distance measure. Linear
193 discriminant analysis effect size (LEfSe) (Segata *et al.* 2011) on the 50 more abundant ASVs was done on
194 Galaxy web application to determine bacterial genera or eukaryotic families biomarkers of each site. The
195 non-parametric Kruskal-Wallis sum-rank test ($\alpha = 0.05$) was performed to detect taxa with significant
196 differential abundance. The biological consistency was investigated by performing a pairwise Wilcoxon test
197 ($\alpha = 0.05$). A linear discriminant analysis (LDA) threshold score of 2.0 was applied.

198

199

200 **3. Results and discussion**

201

202 **3.1 Description of the environmental parameters and the microbial mats**

203 The water temperature and salinity were significantly higher in exploited site ($25 \pm 1^\circ\text{C}$; 101 ± 3.7 psu) than
204 in the abandoned site ($23 \pm 0.4^\circ\text{C}$; 51.1 ± 0.7 psu) (Student t-test for temperature, $p < 0.05$; Welch two sample
205 t-test for salinity, $p < 0.05$), while in both sites the water was at pH 8. Such results were expected as salinity
206 increases in ponds for salt production without being subjected to desiccation, while seawater entering into
207 the abandoned site maintains a lower salinity. The microbial mats showed different morphological aspects,
208 particularly in their thickness that may be explained by the scraping during salt recovery, resulting in a
209 thinner microbial mat in the exploited site. Also, it has been shown that salinity affect exopolysaccharide
210 (EPS) production, that in turn might affect the mat structure and properties (Decho and Gutierrez 2017).
211 Such observation suggested that the environmental conditions may affect the microbial mats as observed in
212 diverse coastal ecosystems including sediments (Pringault *et al.* 2008; Duran *et al.* 2015), estuarine mudflats
213 (Chronopoulou *et al.* 2013) and hypersaline microbial mats (Bordenave *et al.* 2004).

214

215 **3.2 Microbial and meiofaunal community diversity in abandoned and exploited salterns**

216 Despite several attempts, we were unable to obtain archaeal 16S rRNA gene fragment sequences. Although
217 DNA extraction and PCR amplification bias cannot be excluded, this result was in accordance with previous
218 studies showing that benthic mats in hypersaline ponds are dominated by bacteria and are even almost
219 exclusively bacterial (Bolhuis and Stal, 2011; Bolhuis *et al.*, 2014; Stal *et al.*, 2019), with very low
220 abundance of Archaea below 1%, especially within the upper part (0.5 cm) of the mat (Bolhuis and Stal,

221 2011) as analysed in this study. The flooding each winter, after salt recovery, induces the fluctuation of
222 environmental parameters resulting on the renewing of the microbial mats that may also explain the fact that
223 Archaea were not detected. Indeed, Archaea have been demonstrated to be sensitive to drastic modifications
224 of environmental parameters (Zhao *et al.*, 2020) probably preventing their development within microbial
225 mats. Because Archaea have been shown to play a critical role in ecosystem function (Wong *et al.* 2017),
226 further efforts with optimised molecular methods and tools will be beneficial to describe the archaeal
227 diversity in microbial mats. Particularly, it is worth to note that our attempt to target Archaea was based on
228 the utilisation of only a set of primers, several sets of primers have been developed allowing to extend the
229 capacity to reveal Archaea diversity (Bahram *et al.* 2019) that will help to detect Archaea in the microbial
230 mats. Thus, our study focuses on the bacterial and eukaryotic components of the microbial mats inhabiting
231 the abandoned and exploited saltern sites.

232 The composition of bacterial and eukaryotic communities was determined by high throughput sequencing in
233 order to compare microbial mats of the abandoned salterns to that of exploited salterns. The number of raw
234 sequences ranged between $30,243 \pm 4,898$ (abandoned site) and $19,200 \pm 2,626$ (exploited site) for Bacteria,
235 and between $50,119 \pm 720$ (abandoned site) and $39,684 \pm 1,408$ (exploited site) for Eukaryota
236 (Supplementary materials, Table S1). After filtering, the rarefaction provided a random subsample of 5,530
237 bacterial sequences and 22,274 eukaryotic sequences per sample (Supplementary materials, Table S1).

238 The bacterial richness observed for the microbial mats inhabiting the abandoned site was in the same range
239 (between 200-700 ASVs) to that observed for other coastal ecosystems (Dillon *et al.* 2013; Cardoso *et al.*
240 2019). The bacterial Shannon indexes of the abandoned and the exploited sites (5.75 and 4.6, respectively)
241 were higher than that reported for prokaryotic communities from solar salterns (Dillon *et al.* 2013) and
242 spring coastal microbial mats (Cardoso *et al.* 2019) (Fig. 2).

243 For Bacteria, the richness and the Shannon index were significantly higher (Student t-test, $p < 0.05$) in the
244 abandoned site (643 ± 60 ASVs and 5.73 ± 0.09) than in the exploited site (265 ± 75 ASVs and 4.77 ± 0.27 ;
245 Fig. 2A, 2B). Interestingly, for Eukaryota, the opposite was observed, the richness and the Shannon index
246 being significantly higher (Student t-test, $p < 0.05$) in the exploited site (187 ± 26 ASVs and 3.03 ± 0.29) than
247 in the abandoned site (121 ± 25 ASVs and 1.53 ± 0.33 ; Fig. 2C, 2D). Such observation was consistent with

previous reports indicating that salinity is a main driver for bacterial community diversity (Fei Xi *et al.* 2014; Jeanbille *et al.* 2016a; Ben Salem *et al.* 2019) and for the meiofaunal abundance and species composition (Coull 1999). It has been demonstrated that the richness and abundance of bacteria and eukaryotes decrease along a gradient of salinity because the conditions become extreme with increasing salinity (Nubel *et al.* 2000; Pedrós-Alió *et al.* 2000; Benlloch *et al.* 2002; Estrada *et al.* 2004). For eukaryotes, some populations have been reported dominant in high salt concentration (above seawater salinity, 35 psu) ecosystems (Elloumi *et al.* 2009). Interestingly, the eukaryotic diversity was higher in the more salted site (exploited) in our study.

The human actions for salt exploitation, which include regular scraping for salt recovery and microbial mat removal together with the fluctuations of environmental parameters (Wieland *et al.* 2003), are probably responsible for the decreased bacterial diversity observed in the exploited site in comparison to the abandoned site. Indeed, the abandoned site is not disturbed by human actions, which results to an ecosystem with weak fluctuations that is considered as an ecosystem with less selective pressure for bacterial communities (Graham *et al.* 2016). In contrast, the increasing of eukaryotic diversity with salinity has never been reported before for hypersaline environments (Pillay and Perissinotto 2009; Heidelberg *et al.* 2013). Other factors have been reported to influence the composition of the eukaryotic population like hydrodynamic (Kapusta *et al.* 2005), sediment stability (Kapusta *et al.* 2005), sediment particle size (Coull 1999), temperature (Coull 1999; Kapusta *et al.* 2005), depth (Baguley *et al.* 2006), physical disturbance (Austen and Widdicombe 2006), nutrient enrichment (Austen and Widdicombe 2006) and long-term anthropogenic impacts (Boldina, Beninger and Coz 2014). Nevertheless, we assume that physical disturbance and long-term anthropogenic actions were the most probable factors driving eukaryotic communities in our study since salt workers in Ré Island follow a traditional exploitation removing the accumulated mud each winter as described for the Aveiro salterns (Rodrigues *et al.* 2011). More information about the physical-chemical parameters such as pH, dissolved oxygen concentration and water renewable rates will be necessary to determine the factor influencing the eukaryotic composition.

273

274 **3.3 Bacterial and meiofaunal community composition of microbial mats inhabiting abandoned and** 275 **exploited salterns**

276 The first 0.5 cm of microbial mats from both sites were dominated by the same seven bacterial phyla with
 277 high abundance of Proteobacteria and Bacteroidetes (Fig. 3). This phyla composition was comparable to that
 278 of coastal microbial mats where Cyanobacteria, Proteobacteria and Bacteroidetes are dominant (Cardoso *et al.*
 279 *et al.* 2019) as well as in most hypersaline microbial mats (Pal *et al.* 2020). Bacteroidetes and Proteobacteria,
 280 described as dominant phyla in the water and sediment of other salterns (Boujelben *et al.* 2012; Lee *et al.*
 281 2020), are known to play important roles in carbon and nitrogen cycles (Bernhard, Marshall and Yiannos
 282 2012; Wong *et al.* 2015). Only Gammaproteobacteria showed significant differences between the studied
 283 sites: they were significantly more abundant in the exploited site than in the abandoned site (Student t-test,
 284 $p < 0.05$). Such Gammaproteobacteria abundance discrepancy between both sites might be explained by the
 285 fact that Gammaproteobacteria class has been reported to be composed of pioneering microorganisms during
 286 the biofilm formation (Lee *et al.* 2008), which is in accordance with the fact that the microbial mat is
 287 renewed every year in the exploited site. Generally, Cyanobacteria are found dominant in microbial mat
 288 composition (Fourçans *et al.* 2004, 2006, 2008; Bolhuis and Stal, 2011; Boujelben *et al.* 2012; Bolhuis *et al.*,
 289 2014; Stal *et al.* 2019). Surprisingly, they accounted for less than 1% of the total phyla abundance in our
 290 study although that underestimation of Cyanobacteria cannot be excluded due to technical bias such as DNA
 291 extraction and primers efficiencies as previously reported (Fourçans *et al.* 2004).
 292 Assuming that the physical-chemical parameters of the abandoned saltern, particularly the salinity, tend to be
 293 similar to those found at the near oceanic coast, we expect that the microbial mat composition will be close
 294 to that observed in microbial mats inhabiting such coastal areas. The microbial composition of the coastal
 295 sediment from Ré Island has not yet been described. However, Proteobacteria and Planctomycetes were
 296 found to be dominant in the first centimetre of sediments from Marennes-Oléron Bay, located about 50 km
 297 from Ré Island, while Bacteroidetes was the fourth dominant phylum (Lavergne 2014), which differs from
 298 the bacterial phyla composition of microbial mats from both abandoned and exploited sites. Such observation
 299 suggested that the abandonment period (15 years) was not long enough for erasing the effect of saltern
 300 exploitation to reach the bacterial composition observed in coastal sediments, probably because the salinity
 301 in both abandoned and exploited sites exceeds regularly the seawater salinity (> 35 psu). To the best of our
 302 knowledge, only two studies have reported the reorganization of microbial communities after the
 303 abandonment of salterns exploitation (Bernhard, Marshall and Yiannos 2012; Lee *et al.* 2020). These studies
 304 concluded that the physical-chemical parameters were similar to that observed at the coast only one year

305 after the natural collapse of embankments, but the bacterial composition remained different for more than 30
 306 years (Bernhard, Marshall and Yiannos 2012; Lee *et al.* 2020). It is important to mention that in our study the
 307 salterns were still intact maintaining the salinity gradient, which drives bacterial community structure.
 308 Further long time-series analyses comparing microbial communities in different locations at Ré Island are
 309 still required to determine whether the microbial community reach the structure observed in the coast.
 310 The eukaryotic community in the abandoned site was dominated by only three phyla (abundance above 1%)
 311 while the exploited site was characterized by a large eukaryotic diversity with six main phyla identified (Fig.
 312 4). Metazoa dominated the eukaryotic community in the abandoned site, being significantly more abundant
 313 (Student t-test, $p < 0.05$) than in the exploited site (Fig. 4). The presence of Metazoa in both sites was not
 314 surprising since they are often found in marine sediments, which are usually dominated by Nematoda
 315 followed by Platyhelminthes and Arthropoda (Fonseca *et al.* 2014; López-Escardó *et al.* 2018). All the phyla
 316 identified have been ever seen in other exploited salterns (Feazel *et al.* 2008; Stock *et al.* 2012) and in
 317 abandoned salterns (Cvitković *et al.* 2011) but, to our knowledge, no studies compared the eukaryotic
 318 composition in these two types of salterns.
 319 The more important eukaryotic diversity observed in the exploited site, in comparison to that of the
 320 abandoned site, might be explained by the human action of scratching the surface of the microbial mat,
 321 which probably prevents colonization by a single eukaryotic population.
 322 Because it is known that molecular analysis has inherent bias derived from DNA extraction efficiency and
 323 primers specificity (Tedersoo, Tooming-Klunderud and Anslan 2018), the eukaryotic diversity was
 324 completed by meiofaunal microscopic observations. The presence of Acari, Polychaeta, insect larvae and
 325 *Copepoda nauplii* was observed that were not detected by molecular analysis (Supplementary materials, Fig.
 326 S2). Consistently with the molecular analysis, Nematoda largely dominated (>99%) the observable
 327 meiofaunal community of the abandoned site (Supplementary materials, Fig. S2). But, in contrast to the
 328 molecular analysis, Nematoda were also observed (around 69%) together with adult Copepods (26%), Acari
 329 (4%), insect larvae (2%) and Polychaeta (2%) in the exploited site (Supplementary materials, Fig. S2). Thus,
 330 we advocate that the combination of molecular techniques with microscopic observations will help to obtain
 331 a more comprehensive and accurate analysis of eukaryotes species diversity in the environmental samples.

332

3.4 Comparison of bacterial and meiofaunal communities from abandoned site and from exploited site

The comparison of the bacterial and eukaryotic communities between the abandoned site and the exploited site by principal coordinate analysis (PCoA) showed a clear separation between the sites along the PCo1 axis explaining around 47% of the data distribution for the bacterial community (Fig. 5A) and 92% for the eukaryotic community (Fig. 5B). Noteworthy, the PCoA analysis showed a heterogeneity within the exploited site with a dispersion of the triplicates along the PCo2 axis explaining around 21% of the data distribution for bacteria (Fig. 5A) and around 5% for eukaryotes (Fig. 5B). However, cluster analysis based on Bray-Curtis distances confirmed that the bacterial and eukaryotic communities of the triplicates from the abandoned site were clustered together, clearly separated from the cluster formed by the bacterial and eukaryotic communities of the exploited site (Fig. 5C and 5D). The difference of the bacterial and eukaryotic communities between both sites was further confirmed by SIMPER analysis showing overall dissimilarity for bacterial community compositions (SIMPER: 50%) as well as for eukaryotic community compositions (SIMPER: 72%).

Accordingly, LEfSe analyses revealed taxa, bacterial and eukaryotic, differentially abundant between both sites (Fig. 6), which represent site-specific biomarkers as previously suggested in studies identifying environmental and contaminant microbial biomarkers (Segata *et al.* 2011; Jeanbille *et al.* 2016b). The abandoned site was characterized by 11 bacterial and 5 eukaryotic biomarkers (Fig. 6). Interestingly, ASVs related to *Loktanella* (Alphaproteobacteria) and Desulfobacterales, which are usually found in abandoned salterns and tidal flats (Lee *et al.* 2020), were identified among the bacterial biomarkers (Fig. 6A). The Desulfobacterales, sulfate reducing bacteria generally predominant in sulfate-rich sediment (Ruff *et al.* 2015), have been shown to influence the microbial re-colonization of abandoned salterns (Lee *et al.* 2020). Similarly, the Alphaproteobacteria *Loktanella*, has been described as a pioneer bacterium in marine biofilm formation (Lee *et al.* 2008). The identification of such biomarkers suggested that the bacterial community at the abandoned site was undergoing a re-organization. Regarding the eukaryotic biomarkers, ASVs related to Arthropoda, Nematoda and Platyhelminthes belonging both to Metazoa as specific members of abandoned site (Fig. 6B). It is known that Nematoda dominate in marine sediments, followed by Platyhelminthes, Arthropoda and a random assemblage of Gastrotricha, Annelida, Mollusca... (Fonseca *et al.* 2014; López-Escardó *et al.* 2018). This observation was in accordance with our hypothesis that the microbial composition in the abandoned site would tend to be similar to that found on the coast.

According to the LEfSe analyses, the exploited site was characterized by 9 bacterial and 14 eukaryotic biomarkers (Fig. 6). Consistent with the high salinity prevailing at the exploited site, ASVs related to halophilic bacterial genera were identified as biomarkers (Fig. 6A) such as *Marinobacter*, *Salinivibrio* and *Rhodohalobacter* that are usually found in hypersaline environments (Gorshkova 2003; Duran 2010; Kim *et al.* 2017; Xia *et al.* 2017; López-Hermoso *et al.* 2018; de la Haba *et al.* 2019). Such information showed the salinity impact of the microbial mat organisation, further highlighting the differences the contrasted organization with the microbial mat from the abandoned salterns.

The specific eukaryotic ASVs identified in the exploited site were more diverse than that identified in the abandoned site. For example, ASVs related to the Ciliophora phylum were detected in both sites but the populations were different at the class and subclass levels (Fig. 6B). Choreotrichia, Haptoria and Hypotrichia sub-classes, and Phyllopharyngea class were found to be significantly more abundant in the exploited site, while ASVs related to Litostomatea class were significantly predominant in the abandoned site (Fig. 6B). Such observation was in accordance with previous report showing that members of the Ciliophora phylum (ciliates) were influenced by the salinity, their abundance and biomass increasing with salinity together with a modification in their composition (Nche-Fambo, Tirok and Scharler 2016). Additionally, members of the Choreotrichia subclass have been reported dominant in saline environments with salinities reaching up to 64 psu (Nche-Fambo, Tirok and Scharler 2016). The other ASVs found significantly dominant in the exploited site were related to eukaryotic phototrophs related to the Chlorophyta phylum (*Dunaliella*), Dinophyceae class (Dinoflagellates), Haptista phylum (*Isochrysis* and *Pleurochrysis*), and Diatomea (Fig. 6B), indicating that the exploited site was dominated by eukaryotic phototrophs in comparison to the abandoned site. In contrast, the abandoned site was characterized by the presence of ASVs related to Nematoda, Arthropoda, Platyhelminthes and Ciliophora (Litostomatea) phyla, and Craspedida order, which were found significantly abundant (Fig. 6B). Such differences in eukaryotic composition between both sites suggested an important influence of the salinity in the organization of eukaryotic community that in turn might provoke changes in the food web as previously reported along a salinity gradient (Pedrós-Alió *et al.*, 2000). It is likely that the low diversity of preys on the abandoned site (bacteria and ciliates) favoured the development of Metazoa, explaining that Metazoa related ASVs (Nematoda, Arthropoda, and Platyhelminthes) were found significantly more abundant than in the exploited site. Indeed, the Metazoa related ASVs (Nematoda, Arthropoda, and Platyhelminthes) have been shown to be able to feed on bacteria (De Mesel *et al.* 2004;

391 Feazel *et al.* 2008; Hubas *et al.* 2010), diatoms (Montagna 1984), phytobenthos (algae and diatoms; Cowles
392 *et al.*, 1988), ciliates (Berk *et al.* 1977), and dinoflagellates (Cowles, Olson and Chisholm 1988; Turner
393 2004). In contrast, in the exploited site, the higher diversity of Ciliophora (ciliates) related ASVs together
394 with large diversity of eukaryotic phototrophs (Diatoms, Dinoflagellata and Chlorophyta; (Cupp 1943; Lewis
395 and McCourt 2004; Jordan 2012) indicated a larger diversity of preys than in the abandoned site. The ciliates
396 are well known as major predator for bacteria (Sherr and Sherr 1987; Parry 2004) but they can also feed
397 flagellates and other ciliates (Bernard and Rassoulzadegan 1990; Dolan and Coats 1991). It is likely that the
398 exploitation of the saltern resulted in more extreme conditions (higher salinity) reducing the bacterial
399 diversity with concomitant modification in food web structure.

400

401 **4. Conclusion**

402 After stopping the exploitation of the salterns, the abandoned site was characterized by lower salinity with
403 limited variations in comparison to the exploited site. Such conditions favoured the installation of a more
404 diverse bacterial community in the abandoned site while the eukaryotic diversity was lower with the
405 dominance of Nematoda. The presence of site specific ASVs, being significantly more abundant according to
406 the site (biomarkers), suggested a re-organisation of the microbial mat in the abandoned site with a structure
407 tending to get closer to that found in coastal areas. In contrast, the more extreme conditions at the exploited
408 site, characterized by salinity variations, resulted in reduced bacterial diversity with a modification of the
409 food web structure. The exploited site presented halophilic bacteria and a larger diversity of eukaryotic
410 metabolisms. It is likely that the human exploitation of salterns constrains the microbial composition of the
411 microbial mats, which undergo perceptible modifications several years after the abandonment of the salterns.
412 A monitoring of the microbial mats during several years (at least 15 years) on the abandoned salterns is
413 necessary to understand precisely how this structure could be resilient face to human activities.

414

415 **FUNDING**

416

417 C.M. was supported by a PhD grant from E2S-UPPA program and the Région Nouvelle-Aquitaine. We
418 thank the funding support from the ACI U. La Rochelle and the European programme ERANETMED
419 AQUASALT (NMED-0003-01).

420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454

ACKNOWLEDGEMENTS

The authors are grateful to Corentin Hervé, Nicolas Lachaussée, Philippe Pineau and the salterns owners Hélène and Tanguy Jouannet and the society Algorythm. We acknowledge the Molecular Core Facilities at LIENSs laboratory.

This work was performed in collaboration with the GeT core facility, Toulouse, France (<http://get.genotoul.fr>) supported by France Génomique National infrastructure, funded as part of “Investissement d’avenir” program managed by Agence Nationale pour la Recherche (contract ANR-10-INBS-09).

Conflicts of interest

None declared.

References

- Austen MC, Widdicombe S. Comparison of the response of meio- and macrobenthos to disturbance and organic enrichment. *J Exp Mar Biol Ecol* 2006;330:96–104.
- Baguley JG, Montagna PA, Hyde LJ et al. Metazoan meiofauna abundance in relation to environmental variables in the northern Gulf of Mexico deep sea. *Deep Sea Res Part Oceanogr Res Pap* 2006;53:1344–62.
- Bahram, M., Anslan, S., Hildebrand, F., Bork, P., Tedersoo, L., 2019. Newly designed 16S rRNA metabarcoding primers amplify diverse and novel archaeal taxa from the environment. *Environmental Microbiology Reports* 11(4), 487-494.
- Balvočiūtė, M., Huson, D.H., 2017. SILVA, RDP, Greengenes, NCBI and OTT — how do these taxonomies compare? *BMC Genomics* 18(2), 114.
- Ben Salem F, Ben Said O, Cravo-Laureau C et al. Bacterial community assemblages in sediments under high anthropogenic pressure at Ichkeul Lake/Bizerte Lagoon hydrological system, Tunisia. *Environ Pollut* 2019;252:644–56.
- Benlloch S, Lopez-Lopez A, Casamayor EO et al. Prokaryotic genetic diversity throughout the salinity gradient of a coastal solar saltern. *Environ Microbiol* 2002;4:349–60.
- Berk SG, Brownlee DC, Heinle DR et al. Ciliates as a food source for marine planktonic copepods. *Microb Ecol* 1977;4:27–40.
- Bernard C, Rassoulzadegan F. (PDF) Bacteria or microflagellates as a major food source for marine ciliates: Possible implications for the microzooplankton. ResearchGate 1990, DOI: <http://dx.doi.org/10.3354/meps064147>.

455 Bernhard AE, Marshall D, Yiannos L. Increased Variability of Microbial Communities in Restored
456 Salt Marshes nearly 30 Years After Tidal Flow Restoration. *Estuaries Coasts* 2012;35:1049–
457 59.

458 Boldina I, Beninger P, Coz M. Effect of long-term mechanical perturbation on intertidal soft-bottom
459 meiofaunal community spatial structure. *J Sea Res* 2014;85:85–91.

460 Bolhuis H, Cretoiu MS, Stal LJ. Molecular ecology of microbial mats. *FEMS Microbiol Ecol*
461 2014;90:335–350.

462 Bolhuis H, Stal LJ. Analysis of bacterial and archaeal diversity in coastal microbial mats using
463 massive parallel 16S rRNA gene tag sequencing. *ISME J* 2011;5:1701–12.

464 Bolyen E, Rideout JR, Dillon MR et al. Reproducible, interactive, scalable and extensible
465 microbiome data science using QIIME 2. *Nat Biotechnol* 2019;37:852–7.

466 Bordenave S, Fourçans A, Blanchard S et al. Structure and functional analyses of bacterial
467 communities changes in microbial mats following petroleum exposure. *Ophelia*
468 2004;58:195–203.

469 Boujelben I, Gomariz M, Martínez-García M et al. Spatial and seasonal prokaryotic community
470 dynamics in ponds of increasing salinity of Sfax solar saltern in Tunisia. *Antonie Van*
471 *Leeuwenhoek* 2012;101:845–57.

472 Callahan BJ, McMurdie PJ, Rosen MJ et al. DADA2: High-resolution sample inference from
473 Illumina amplicon data. *Nat Methods* 2016;13:581–3.

474 Caporaso JG, Lauber CL, Walters WA et al. Global patterns of 16S rRNA diversity at a depth of
475 millions of sequences per sample. *Proc Natl Acad Sci* 2011;108:4516–22.

476 Cardoso DC, Cretoiu MS, Stal LJ et al. Seasonal development of a coastal microbial mat. *Sci Rep*
477 2019;9:9035.

478 Caumette P, Matheron R, Raymond N et al. Microbial mats in the hypersaline ponds of
479 Mediterranean salterns (Salins-de-Giraud, France). *FEMS Microbiol Ecol* 1994;13:273–86.

480 Chronopoulou P-M, Fahy A, Coulon F et al. Impact of a simulated oil spill on benthic phototrophs
481 and nitrogen-fixing bacteria in mudflat mesocosms. *Environ Microbiol* 2013;15:242–52.

482 Coull BC. Role of meiofauna in estuarine soft-bottom habitats* - Coull - 1999 - Australian Journal
483 of Ecology - Wiley Online Library. *Aust J Ecol* 1999;24:327–343.

484 Cowles TJ, Olson RJ, Chisholm SW. Food selection by copepods: discrimination on the basis of
485 food quality. *Mar Biol* 1988;100:41–9.

486 Cupp EE. *Marine Plankton Diatoms of the West Coast of North America*. 1943.

487 Cvitković I, Despalatović M, Grubelić I et al. Spatio-temporal variability of meiofauna community
488 structure in abandoned salina of velike soline and adjacent area (Mediterranean, Adriatic
489 sea). *Fresenius Environ Bull* 2011;20:645–55.

490 Decho, A.W., Gutierrez, T., 2017. Microbial Extracellular Polymeric Substances (EPSs) in Ocean
491 Systems. *Frontiers in Microbiology* 8(922).

- 492 De Mesel I, Derycke S, Moens T et al. Top-down impact of bacterivorous nematodes on the
493 bacterial community structure: a microcosm study. *Environ Microbiol* 2004;6:733–44.
- 494 Dillon J, Carlin M, Gutierrez A et al. Patterns of microbial diversity along a salinity gradient in the
495 Guerrero Negro solar saltern, Baja CA Sur, Mexico. *Front Microbiol* 2013;4:399.
- 496 Dobretsov S, Abed RMM, Al Maskari SMS et al. Cyanobacterial mats from hot springs produce
497 antimicrobial compounds and quorum-sensing inhibitors under natural conditions. *J Appl*
498 *Phycol* 2011;23:983–93.
- 499 Dolan JR, Coats DW. A study of feeding in predacious ciliates using prey ciliates labeled with
500 fluorescent microspheres. *J Plankton Res* 1991;13:609–27.
- 501 Duran R. *Marinobacter*. Volume 3: Microbes and Communities Utilizing Hydrocarbons, Oils and
502 Lipids - Chapter 5. In: Timmis KN (ed.). *Handbook of Hydrocarbon and Lipid*
503 *Microbiology*. Vol 3. Berlin, Heidelberg: Springer, 2010, 1725–35.
- 504 Duran R, Bielen A, Paradžik T et al. Exploring Actinobacteria assemblages in coastal marine
505 sediments under contrasted Human influences in the West Istria Sea, Croatia. *Environ Sci*
506 *Pollut Res* 2015;22:15215–29.
- 507 Edgcomb, V.P., Bernhard, J.M., Summons, R.E., Orsi, W., Beaudoin, D., Visscher, P.T., 2014.
508 Active eukaryotes in microbialites from Highborne Cay, Bahamas, and Hamelin Pool (Shark
509 Bay), Australia. *The ISME Journal* 8(2), 418-429.
- 510 Elloumi J, Guermazi W, Ayadi H et al. Abundance and biomass of prokaryotic and eukaryotic
511 microorganisms coupled with environmental factors in an arid multi-pond solar saltern
512 (Sfax, Tunisia). *J Mar Biol Assoc U K* 2009;89:243–53.
- 513 Estrada M, Henriksen P, Gasol JM et al. Diversity of planktonic photoautotrophic microorganisms
514 along a salinity gradient as depicted by microscopy, flow cytometry, pigment analysis and
515 DNA-based methods. *FEMS Microbiol Ecol* 2004;49:281–93.
- 516 Falcón, L.I., Valdespino-Castillo, P.M., Alcántara-Hernández, R.J., Gómez-Acata, E.S., Yanez-
517 Montalvo, A., Águila, B., 2020. Stromatolites in Crater-Lake Alchichica and Bacalar
518 Lagoon, Astrobiology and Cuatro Ciénegas Basin as an Analog of Early Earth. Springer, pp.
519 183-201.
- 520 Feazel LM, Spear JR, Berger AB et al. Eucaryotic Diversity in a Hypersaline Microbial Mat. *Appl*
521 *Environ Microbiol* 2008;74:329–32.
- 522 Fei Xi X, Wang L, Jun Hu J et al. Salinity influence on soil microbial respiration rate of wetland in
523 the Yangtze River estuary through changing microbial community. *J Environ Sci China*
524 2014;26:2562–70.
- 525 Fleming, E.D., Prufert - Bebout, L., 2010. Characterization of cyanobacterial communities from
526 high - elevation lakes in the Bolivian Andes. *Journal of Geophysical Research:*
527 *Biogeosciences* 115(G2).
- 528 Fonseca VG, Carvalho GR, Nichols B et al. Metagenetic analysis of patterns of distribution and
529 diversity of marine meiobenthic eukaryotes. *Glob Ecol Biogeogr* 2014;23:1293–302.

- 530 Fourçans A, Oteyza TG de, Wieland A et al. Characterization of functional bacterial groups in a
531 hypersaline microbial mat community (Salins-de-Giraud, Camargue, France). FEMS
532 Microbiol Ecol 2004;51:55–70.
- 533 Fourçans A, Ranchou-Peyruse A, Caumette P et al. Molecular Analysis of the Spatio-temporal
534 Distribution of Sulfate-reducing Bacteria (SRB) in Camargue (France) Hypersaline
535 Microbial Mat. Microb Ecol 2008;56:90–100.
- 536 Fourçans A, Solé A, Diestra E et al. Vertical migration of phototrophic bacterial populations in a
537 hypersaline microbial mat from Salins-de-Giraud (Camargue, France). FEMS Microbiol
538 Ecol 2006;57:367–77.
- 539 van Gernerden H. Microbial mats: A joint venture. Mar Geol 1993;113:3–25.
- 540 Giani D, Seeler J, Giani L et al. Microbial mats and physicochemistry in a saltern in the Bretagne
541 (France) and in a laboratory scale saltern model. FEMS Microbiol Lett 1989;62:151–61.
- 542 Gómez-Villegas P, Vígara J, León R. Characterization of fungi from hypersaline environments of
543 solar salterns using morphological and molecular techniques. Mar Drugs 2018;16, DOI:
544 10.3390/md16090332.
- 545 Gorshkova NM. *Marinobacter excellens* sp. nov., isolated from sediments of the Sea of Japan. Int J
546 Syst Evol Microbiol 2003;53:2073–8.
- 547 Graham EB, Crump AR, Resch CT et al. Coupling Spatiotemporal Community Assembly Processes
548 to Changes in Microbial Metabolism. Front Microbiol 2016;7, DOI:
549 10.3389/fmicb.2016.01949.
- 550 de la Haba RR, López-Hermoso C, Sánchez-Porro C et al. Comparative Genomics and
551 Phylogenomic Analysis of the Genus *Salinivibrio*. Front Microbiol 2019;10, DOI:
552 10.3389/fmicb.2019.02104.
- 553 Heidelberg KB, Nelson WC, Holm JB et al. Characterization of eukaryotic microbial diversity in
554 hypersaline Lake Tyrrell, Australia. Front Microbiol 2013;4, DOI:
555 10.3389/fmicb.2013.00115.
- 556 Hubas C, Sachidhanandam C, Rybarczyk H et al. Bacterivorous nematodes stimulate microbial
557 growth and exopolymer production in marine sediment microcosms. Mar Ecol Prog Ser
558 2010;419:85–94.
- 559 Hugoni M, Domaizon I, Taib N, Biderre-Peti, C, Agogué H, Galand PE, Debroas D, Mary I.
560 Temporal dynamics of active Archaea in oxygen-depleted zones of two deep lakes. Env
561 Microbiol Rep 2015; 7: 321-329.
- 562 Jeanbille M, Gury J, Duran R et al. Response of Core Microbial Consortia to Chronic Hydrocarbon
563 Contaminations in Coastal Sediment Habitats. Front Microbiol 2016a;7, DOI:
564 10.3389/fmicb.2016.01637.
- 565 Jeanbille M, Gury J, Duran R et al. Chronic Polyaromatic Hydrocarbon (PAH) Contamination Is a
566 Marginal Driver for Community Diversity and Prokaryotic Predicted Functioning in Coastal
567 Sediments. Front Microbiol 2016b;7, DOI: 10.3389/fmicb.2016.01303.
- 568 Jordan RW. Haptophyta. ELS. American Cancer Society, 2012.

569 Jørgensen BB, Cohen Y. Solar Lake (Sinai). 5. The sulfur cycle of the benthic cyanobacterial mats.
570 Limnol Oceanogr 1977;22:657–66.

571 Kahle D, Wickham H. ggmap: Spatial Visualization with ggplot2. R J 2013;5:144.

572 Kapusta SC, Würdig NL, Bemvenuti CE et al. Meiofauna structure in Tramandaí-Armazém estuary
573 (South of Brazil). Acta Limnol Bras 2005;17:349–59.

574 Kim J-O, Lee H-J, Han S-I et al. *Marinobacter halotolerans* sp. nov., a halophilic bacterium isolated
575 from a saltern crystallizing pond. Int J Syst Evol Microbiol 2017;67:460–5.

576 Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, Glöckner FO. Evaluation of
577 general 16S ribosomal RNA gene PCR primers for classical and next-generation
578 sequencing-based diversity studies. Nucleic acids research 2013; 41: e1-e1.

579 Kolesnikov, A.V., Danelian, T., Gommeaux, M., Maslov, A.V., Grazhdankin, D.V., 2017.
580 Arumberiamorph structure in modern microbial mats: implications for Ediacaran
581 palaeobiology. Bulletin de la Société géologique de France 188(1-2), 5.

582 Lavergne C. Rôle (structure et fonction) des communautés procaryotes (bactéries et archées) dans le
583 cycle de l’azote d’une vasière littorale du Pertuis Charentais: influence des facteurs
584 biotiques et abiotiques par une approche multi-échelle. 2014:501.

585 Lee H, Heo YM, Kwon SL et al. Recovery of the benthic bacterial community in coastal abandoned
586 saltern requires over 35 years: A comparative case study in the Yellow Sea. Environ Int
587 2020;135:105412.

588 Lee J-W, Nam J-H, Kim Y-H et al. Bacterial communities in the initial stage of marine biofilm
589 formation on artificial surfaces. J Microbiol Seoul Korea 2008;46:174–82.

590 Lepère C, Domaizon I, Hugoni M et al. Diversity and Dynamics of Active Small Microbial
591 Eukaryotes in the Anoxic Zone of a Freshwater Meromictic Lake (Pavin, France). Front
592 Microbiol 2016;7, DOI: 10.3389/fmicb.2016.00130.

593 Lewis LA, McCourt RM. Green algae and the origin of land plants. Am J Bot 2004;91:1535–56.

594 López-Escardó D, Paps J, de Vargas C et al. Metabarcoding analysis on European coastal samples
595 reveals new molecular metazoan diversity. Sci Rep 2018;8:9106.

596 López-Hermoso C, de la Haba RR, Sánchez-Porro C et al. *Salinivibrio kushneri* sp. nov., a
597 moderately halophilic bacterium isolated from salterns. Syst Appl Microbiol 2018;41:159–
598 66.

599 McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis and
600 Graphics of Microbiome Census Data. Watson M (ed.). PLoS ONE 2013;8:e61217.

601 Martinez, J.N., Nishihara, A., Lichtenberg, M., Trampe, E., Kawai, S., Tank, M., Köhl, M., Hanada,
602 S., Thiel, V., 2019. Vertical distribution and diversity of phototrophic bacteria within a hot
603 spring microbial Mat (Nakabusa hot springs, Japan). Microbes and Environments 34(4),
604 374-387.

605 Mir J, MartÃ-nez-Alonso M, Esteve I et al. Vertical stratification and microbial assemblage of a
606 microbial mat in the Ebro Delta (Spain). FEMS Microbiol Lett 1991;86:59–68.

607 Montagna PA. In situ measurement of meiobenthic grazing rates on sediment bacteria and edaphic
608 diatoms. *Mar Ecol Prog Ser* 1984;18:119–30.

609 Nche-Fambo FA, Tirok K, Scharler UM. Hypersaline conditions cause distinct ciliate community
610 structure in a South African estuarine lake system. *J Plankton Res* 2016;38:878–87.

611 Nubel U, Garcia-Pichel F, Clavero E et al. Matching molecular diversity and ecophysiology of
612 benthic cyanobacteria and diatoms in communities along a salinity gradient. *Environ*
613 *Microbiol* 2000;2:217–26.

614 Oren A. Saltern evaporation ponds as model systems for the study of primary production processes
615 under hypersaline conditions. *Aquat Microb Ecol* 2009;56:193–204.

616 Pal S, Biswas R, Misra A, Sar A, Banerjee S, Mukherjee P, Dam B. Poorly known microbial taxa
617 dominate the microbiome of hypersaline Sambhar Lake salterns in India. *Extremophiles*
618 2020;24:875-885.

619 Parry JD. Protozoan Grazing of Freshwater Biofilms. *Advances in Applied Microbiology*. Vol 54.
620 Elsevier, 2004, 167–96.

621 Pedrós-Alió C, Calderón-Paz JI, MacLean MH et al. The microbial food web along salinity
622 gradients. *FEMS Microbiol Ecol* 2000;32:143–55.

623 Pillay D, Perissinotto R. Community structure of epibenthic meiofauna in the St. Lucia Estuarine
624 Lake (South Africa) during a drought phase. *Estuar Coast Shelf Sci* 2009;81:94–104.

625 Plominsky AM, Henríquez-Castillo C, Delherbe N et al. Distinctive Archaeal Composition of an
626 Artisanal Crystallizer Pond and Functional Insights Into Salt-Saturated Hypersaline
627 Environment Adaptation. *Front Microbiol* 2018;9, DOI: 10.3389/fmicb.2018.01800.

628 Prieto-Barajas, C.M., Valencia-Cantero, E., Santoyo, G., 2018. Microbial mat ecosystems: Structure
629 types, functional diversity, and biotechnological application. *Electronic Journal of*
630 *Biotechnology* 31, 48-56.

631 Pringault O, Duran R, Jacquet S et al. Temporal Variations of Microbial Activity and Diversity in
632 Marine Tropical Sediments (New Caledonia Lagoon). *Microb Ecol* 2008;55:247–58.

633 Quast C, Pruesse E, Yilmaz P et al. The SILVA ribosomal RNA gene database project: improved
634 data processing and web-based tools. *Nucleic Acids Res* 2012;41:D590–6.

635 R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical
636 Computing, Vienna, Austria. URL <https://www.R-project.org/>.<https://www.R-project.org/>.
637 Published 2020.

638 Rodrigues CM, Bio A, Amat F et al. Artisanal salt production in Aveiro/Portugal - an ecofriendly
639 process. *Aqua. Biosyst* 2011;7:3-14.

640 Ruff SE, Biddle JF, Teske AP et al. Global dispersion and local diversification of the methane seep
641 microbiome. *Proc Natl Acad Sci* 2015;112:4015–20.

642 Rzeznik-Orignac J, Fichet D, Boucher G. Spatio-temporal structure of the nematode assemblages of
643 the Brouage mudflat (Marennes Oléron, France). *Estuar Coast Shelf Sci* 2003;58:77–88.

644 Sánchez-Baracaldo, P., Bianchini, G., Wilson, J.D., Knoll, A.H., 2021. Cyanobacteria and
645 biogeochemical cycles through Earth history. *Trends in Microbiology*.

646 Schmidt, S., Darcy, J., Sommers, P., Gunawan, E., Knelman, J., Yager, K., 2017. Freeze–thaw
647 revival of rotifers and algae in a desiccated, high-elevation (5500 meters) microbial mat,
648 high Andes, Perú. *Extremophiles* 21(3), 573–580.

649 Segata N, Izard J, Waldron L et al. Metagenomic biomarker discovery and explanation. *Genome*
650 *Biol* 2011;12:R60.

651 Sherr EB, Sherr BF. High rates of consumption of bacteria by pelagic ciliates. *Nature*
652 1987;325:710–1.

653 Sierra, M.A., Li, Q., Pushalkar, S., Paul, B., Sandoval, T.A., Kamer, A.R., Corby, P., Guo, Y., Ruff,
654 R.R., Alekseyenko, A.V., Li, X., Saxena, D., 2020. The Influences of Bioinformatics Tools
655 and Reference Databases in Analyzing the Human Oral Microbial Community. *Genes* 11(8),
656 878.

657 Stal LJ, Bolhuis H, Cretoiu MS. Phototrophic marine benthic microbiomes: the ecophysiology of
658 these biological entities. *Environ Microbiol* 2019;21:1529–1551.

659 Stock A, Breiner H-W, Pachiadaki M et al. Microbial eukaryote life in the new hypersaline deep-sea
660 basin Thetis. *Extremophiles* 2012;16:21–34.

661 Tedersoo L, Tooming-Klunderud A, Anslan S. PacBio metabarcoding of Fungi and other
662 eukaryotes: errors, biases and perspectives. *New Phytol* 2018;217:1370–85.

663 Tran H-T, Wang H-C, Hsu T-W et al. Revegetation on abandoned salt ponds relieves the seasonal
664 fluctuation of soil microbiomes. *BMC Genomics* 2019;20, DOI: 10.1186/s12864-019-5875-
665 y.

666 Turner JT. The Importance of Small Planktonic Copepods and Their Roles in Pelagic Marine Food
667 Webs. *Zool Stud* 2004:13.

668 Valdespino-Castillo, P.M., Cerqueda-García, D., Espinosa, A.C., Batista, S., Merino-Ibarra, M., Taş,
669 N., Alcántara-Hernández, R.J., Falcón, L.I., 2018. Microbial distribution and turnover in
670 Antarctic microbial mats highlight the relevance of heterotrophic bacteria in low-nutrient
671 environments. *FEMS microbiology ecology* 94(9), fyy129.

672 Vogt, J.C., Abed, R.M.M., Albach, D.C., Palinska, K.A., 2018. Bacterial and Archaeal Diversity in
673 Hypersaline Cyanobacterial Mats Along a Transect in the Intertidal Flats of the Sultanate of
674 Oman. *Microbial Ecology* 75(2), 331–347.

675 Walmsley JG. THE ECOLOGICAL IMPORTANCE OF MEDITERRANEAN SALINAS. :15.

676 White, R.A., Visscher, P.T., Burns, B.P., 2021. Between a rock and a soft place: the role of viruses
677 in lithification of modern microbial mats. *Trends in Microbiology* 29(3), 204–213.

678 White, R.A., Wong, H.L., Ruvindy, R., Neilan, B.A., Burns, B.P., 2018. Viral Communities of Shark
679 Bay Modern Stromatolites. *Frontiers in Microbiology* 9(1223).

680 Wieland A, Kühl M, McGowan L et al. Microbial Mats on the Orkney Islands Revisited:
681 Microenvironment and Microbial Community Composition. *Microb Ecol* 2003;46:371–90.

682 Wong HL, Smith D-L, Visscher PT et al. Niche differentiation of bacterial communities at a
683 millimeter scale in Shark Bay microbial mats. *Sci Rep* 2015;5:15607.

684 Wong, H.L., Visscher, P.T., White Iii, R.A., Smith, D.-L., Patterson, M.M., Burns, B.P., 2017.
685 Dynamics of archaea at fine spatial scales in Shark Bay mat microbiomes. *Scientific Reports*
686 7(1), 46160.

687 Xia J, Xie Z-H, Dunlap CA et al. *Rhodohalobacter halophilus* gen. nov., sp. nov., a moderately
688 halophilic member of the family Balneolaceae. *Int J Syst Evol Microbiol* 2017;67:1281–7.

689 Yanez-Montalvo, A., Gómez-Acata, S., Águila, B., Hernández-Arana, H., Falcón, L.I., 2020. The
690 microbiome of modern microbialites in Bacalar Lagoon, Mexico. *PloS one* 15(3), e0230071.

691 Yilmaz P, Wegener Parfrey L, Yarza P et al. The SILVA and “All-species living tree project (LTP)”
692 taxonomic frameworks. *Nucleic Acids Res* 2013;42, DOI: 10.1093/nar/gkt1209.

693 Zhao J, Meng Y, Drewer J, Skiba UM, Prosser JI, Gubry-Rangin C. Differential ecosystem function
694 stability of ammonia-oxidizing archaea and bacteria following short-term environmental
695 perturbation. *mSystems* 2020, 5, e00309-20.

696

697 **Figure caption**

698 **Figure 1: Localisation of the sampling sites at the Ré Island (France).** Microbial mats were sampled in
699 abandoned (red point) and exploited (green point) sites in independent triplicates corresponding to three
700 ponds in each site.

701
702 **Figure 2: Diversity indexes for bacterial (A, B) and eukaryotic (C, D) communities in the abandoned**
703 **(red) and exploited (green) sites.** The boxplots show observed richness (A, C) and Shannon (B, D) indexes
704 determined at the ASV level from 16S (Bacteria) and 18S (Eukarya) rRNA gene sequences. Different letters
705 indicate significant differences based on Student t-test at $p < 0.05$ ($n = 3$).

706
707 **Figure 3: Bacterial community compositions at the phylum level in the abandoned and exploited sites.**
708 The bar plots represent each replicate, presenting the phyla with a relative abundance above 1%. The
709 class level within Proteobacteria phylum is shown.

710
711 **Figure 4: Eukaryotic community compositions at the phylum level in the abandoned and exploited**
712 **sites.** The bar plots represent each replicate, presenting the phyla with a relative abundance above 1%.

713
714 **Figure 5: Comparison of bacterial and eukaryotic communities inhabiting the abandoned (red) and**
715 **exploited (green) sites.** (A, B) Principal coordinate analysis (PCoA) comparing bacterial (A) and eukaryotic
716 (B) communities. (C, D) Clustering analysis based on Bray-Curtis distances.

717
718 **Figure 6: Comparison of bacterial (A) and eukaryotic (B) communities inhabiting the abandoned (red)**
719 **and exploited (green) sites by linear discriminant analysis effect size (LEfSe).** The analysis was
720 performed with the 50 more abundant bacterial and eukaryotic ASVs of each site.

721

722

727

728

730

731

732 Figure 4

733

734

735

737

738

740

741

743

744

745

746