1 Genomic and fossil windows into the secret lives of the most ancient fungi

2
3 Mary L. Berbee^{1†}, Christine Strullu-Derrien^{2,3}, Pierre-Marc Delaux⁴, Paul K. Strother⁵, Paul
4 Kenrick², Marc-André Selosse^{3,6}, John W. Taylor⁷

5
6 ¹Department of Botany, University of British Columbia, Vancouver, BC, Canada

7 ²Department of Earth Sciences, The Natural History Museum, London, UK

8 ³Institut Systématique Evolution Biodiversité, Muséum national d'Histoire naturelle, CNRS,
9 Sorbonne Université, Paris, France

10 ⁴LRSV, Université de Toulouse, CNRS, UPS, Castanet-Tolosan, France.

11 ⁵Department of Earth and Environmental Sciences, Boston College, Weston, MA, USA

12 ⁶Department of Plant Taxonomy and Nature Conservation, University of Gdańsk, Gdańsk,
13 Poland

14 ⁷Department of Plant and Microbial Biology, University of California, Berkeley, CA, USA

15 [†]email: mary.berbee@botany.ubc.ca

18 **Abstract**

19 Fungi have crucial roles in modern ecosystems as decomposers and pathogens, and they
20 engage in a variety of mutualistic associations with other organisms, especially plants. They
21 have a lengthy geological history, and there is an emerging understanding of their impact on
22 the evolution of Earth systems on a large scale. In this Review, we focus on the roles of fungi
23 in the establishment and early evolution of land and freshwater ecosystems. Today, questions
24 of evolution over deep time are informed by discoveries of new fossils and evolutionary
25 analysis of new genomes. Inferences can be drawn from evolutionary analysis by comparing
26 the genes and genomes of fungi with the biochemistry and development of their plant and
27 algal hosts. We then contrast this emerging picture against evidence from the fossil record to
28 develop a new, integrated perspective on the origin and early evolution of fungi.

30 **[H1] Introduction**

31 Fungi are crucial for thriving land plant communities as mycorrhizal symbionts^{1,2} and
32 decomposers³. For perhaps 500 million years^{4,5}, fungi have been supplying land plants with
33 phosphorus and other minerals, thereby speeding up photosynthesis and contributing to the
34 drawdown of atmospheric CO₂⁶⁻⁸. Yet phylogenomics tells us that fungi originated far earlier,

deep in the Precambrian⁹, perhaps a billion years ago^{2,10}. This information raises two important questions: When did fungi emerge, and what were their early lifestyles?

How and when fungi diversified and adapted to their present roles in Earth systems has long been hidden. Fossils representing simple filaments or spores of ancient fungi are uncommon in the Precambrian fossil record, and difficult to recognize or to distinguish from other organisms¹¹. Due largely to a lack of reliable fossil calibrations, comparisons of ages depend on dated fungal and plant phylogenies^{2,12}, providing a stimulating framework for thought, but these estimates are too imprecise to eliminate alternative hypotheses about ancient ecological relationships between kingdoms.

Here, we review recent landmark studies that open new windows on the understanding of the earliest fungi. Comparative analysis of genomes provides the most direct available evidence of ancient interdependence between fungi and algae or land plants. Fungi survive by secreting digestive enzymes and taking up the freed nutrients^{13,14}. The record — at least the remaining record — of the origin of enzymes targeting algal or plant cell walls is embedded in fungal genomes. As green algae evolved into land plants, and as the polysaccharides in their cell walls diversified¹⁵, the evolution of fungal enzymes surely kept pace^{3,16,17}. Comparative analysis of genomes of land plants¹⁸ and their closest algal relatives^{19,20} reveal the evolutionary origins of constituents of plant immune systems that are fundamental both for defense against fungal parasites and for the evolution of mycorrhizae²¹. Green algae, and by extension, the fungi that they nourished may have begun to evolve in freshwater environments, likely in the Precambrian, a billion years ago (1Ga), as highlighted by the geology^{22,23} and recent analysis of the phylogenies of algae^{24,25}.

We address the puzzle of what could have nourished early fungi prior to the origin of land plants by reviewing geological evidence of other life forms in shallow marine and terrestrial environments^{22,23,26}. We go on to consider specific fossils and compare their dates and traits to fungal phylogenies based on genomic analyses, asking which sorts of geological and fossil data can be used to resolve conflicts between rocks and clocks. Recently, Loron et al.²⁷ published evidence of 1 billion-year-old fungal fossils from Arctic Canada that they named *Ourasphaira giraldae*. A 1 billion-year-old fungus is twice the age of land plants, based on either fossils or molecular dating^{2,4}. This and other surprising fossil results invite challenge. We review the evidence for interpreting Precambrian fossils as 'fungi'^{28,29}, discuss the early occurrences of Paleozoic fungi, and highlight the potential contributions of new tools toward resolving their identities, for example, confocal scanning laser microscopy³⁰, synchrotron analyses^{27,29,31}, and chemical analyses^{27,29}. We document the tremendous strides

that have been made in characterizing fossils microscopically and chemically, but worry that even greater ones will be needed when the specimens are hundreds of millions to billions of years old and have endured pressures and temperatures far beyond those experienced by living organisms.

[H1] The first fungi: nutrition and ecology

[H2] Modern fungi evolved from phagotrophic, aerobic ancestors. Within the eukaryotic tree of life, Fungi, together with the nucleariid amoebas are the closest relatives of Holozoa³². Over a billion years ago^{2,10}, the fungal lineage evolved as a grade of motile, unicellular, amoeboid protists (FIG 1). The first amoeboid fungi may have been free living and, much like modern nucleariids³³ (FIG 1) and aphelids³⁴, equipped with filose pseudopodia and a phagotrophic mode of nutrition that involved engulfing and digesting organisms smaller than themselves^{35,36}. Phagotrophs diversified, giving rise to obligate parasites such as *Rozella*, which engulfs and digests the cytoplasm within a single host cell^{37,38}, and microsporidia, which lost the capacity for phagotrophy while refining their adaptations as intracellular parasites³⁹. In contrast to the ancestral phagotrophs, the vast majority of modern fungi, from aquatic or semi-aquatic Chytridiomycota to the land-inhabiting yeasts, molds and mushrooms of Dikarya, are osmotrophic^{13,14} (FIG 1). This mode of feeding involves the cytoplasmic uptake of nutrients that diffuse across cell walls and often is coupled with the secretion by the fungus of extracellular enzymes that release nutrients from substrates.

Fungi are almost all obligate aerobes or facultative anaerobes that require oxygen to complete their life cycles. Oxygen is needed for oxidative phosphorylation and for sterol biosynthesis. Additionally, genomes of early diverging fungi, including *Rozella allomyces*, Chytridiomycota¹⁶, Zoopagomycota and Dikarya¹⁰ encode lytic polysaccharide monooxygenases, enzymes that require oxygen to facilitate breakdown of polysaccharides⁴⁰. Just two groups of obligate anaerobic fungi are known and both independently lost the ancestral aerobic lifestyle. The microsporidia are one group; they live inside animal cells and compensate for the loss of their mitochondria and absence of oxidative respiration by absorbing nutrients and even ATP from their surrounding hosts⁴¹. The other group, the rumen fungi (Neocallimastigomycotina in the Chytridiomycota) adapted to the anaerobic, cellulose-rich environment in the digestive systems of herbivores. This adaptation has involved gain from rumen bacteria, by horizontal transfer, of multiple genes in several pathways needed for

anaerobiosis⁴². Given the rarity of obligately anaerobic fungi, it seems safe to state that the primary evolutionary radiation of their kingdom occurred in oxygenated environments.

[H2] *Plants are key to evolutionary success of osmotrophic fungi.* Modern land plants provide fungi with a superb source of reliable energy. Much of the carbon from net annual land-based productivity resulting from photosynthesis, currently about 56×10^{12} kg C/year⁴³, passes through fungi. Mycorrhizal fungi symbiotic with roots take ~13% of plant primary productivity⁴⁴. Saprotrophic fungi consume plant cell wall polymers, including the polysaccharides cellulose and hemicellulose, and can degrade even lignin, a recalcitrant, cross-linked, phenolic polymer. Annual production of cellulose, the most common organic polymer, accounts for $\sim 23 \times 10^{12}$ kg C/year, or ~40% of the total net annual plant productivity^{45,46}, and lignin, with $\sim 11 \times 10^{12}$ kg C/year, accounts for another 20% of the total productivity⁴⁷. Wood-rotting fungi degrade lignin not as a direct source of nutrition, as far as is known, but rather to expose the wood's cellulose and hemicellulose, which the fungus then digests for energy^{46,48,49}.

Land plants and their closest green algal relatives, together constitute the streptophytes (TABLE 1; FIG 1). The facts that follow lead us to infer that early osmotrophic fungi adapted first to symbiosis or saprotrophy on freshwater streptophyte algae before gaining access to carbon provided by land plants.

[H2] *Fungal enzyme systems tracked plant evolution from freshwater to land.* Genome-based phylogenetic analysis of expansions of fungal enzyme-encoding genes offers insights into the compounds that early phagotrophic, saprotrophic and symbiotic fungi could digest. Cellulases are encoded in genomes of the vast majority of fungi from phagotrophic aphelids³⁶ to osmotrophs¹⁶, indicating that cellulose was an ancient source of nutrition for fungi. Cellulose evolved before the earliest fungi; enzymes for the synthesis of cellulose appear in cyanobacteria and are widely distributed across eukaryotic algae, protists and animals⁵⁰. Cyanobacteria secrete a complex assortment of extracellular polysaccharides that may include cellulose⁵¹ but this assortment can vary even within a species⁵². This variation, which has been shown experimentally to contribute to defense against viruses⁵³ and fungi⁵⁴, could have presented a barrier to digestion of cyanobacteria by early fungi. Cell walls of red and green algae also vary in composition but they share cellulose as an important structural component⁵⁰. Thus, cellulases retained in fungal genomes may represent a signature of successful adaptation to digestion of eukaryotic algae.

Osmotrophic fungi share a rich array of additional enzymes that target other plant cell wall components: hemicelluloses that include xyloglucans¹⁶, pectins¹⁷ and lignin⁴⁸.

Fortunately for evolutionary inferences, these components, and by extension, the fungal enzymes needed to degrade them seem to have arisen much more recently than cellulose and cellulases. Pectins and xyloglucans are known only in streptophytes^{15,55} and early diverging streptophyte algae have few if any copies of genes for synthesis or modification of these polysaccharides⁵⁶.

Phylogenetic reconstructions show that multiple gene copies of diverse pectinases are retained in plant-saprotrophic species of fungi ranging from Chytridiomycota to Dikarya¹⁷. In phylogenies, copies of the genes predicted to encode pectinases fall into diverse classes of enzymes: glycoside hydrolases, polysaccharide lyases, and carbohydrate esterases. The pectinase gene copies are inferred to have multiplied in number through five gene duplications in the common ancestor of osmotrophic fungi¹⁷. Pectinase gene copies were lost convergently from some lineages that adapted to nutrition sources other than streptophytes and for example, few pectinase genes remain in *Batrachochytrium*, a genus that adapted to parasitism on amphibians (Table 1). However, most osmotrophic fungi retained pectinases throughout their evolution, implying that they have been under continuing selective pressure to take their nutrition from enzymatic attack on the pectins of cell walls of living or dead streptophyte algae and land plants¹⁷.

Although not analyzed in formal phylogenies, Chytridiomycota and Dikarya share predicted xyloglucanases^{57,58}, further evidence that early osmotrophs digested the streptophyte cell walls. Lignin is restricted to land plants, and an expansion of peroxidase enzymes needed for lignin decomposition is credited with contributing to the diversification of Basidiomycota³.

Fungal enzymes presumably diversified after their substrates became available. The age of the first xyloglucan- and pectin-containing streptophyte is difficult to estimate. Because few fossils are available, dates are poorly constrained leading to wide confidence intervals within molecular clock studies, and to incongruence across studies. Assuming that the pectins and xyloglucans evolved in streptophytes after the divergence of Mesostigmatophyceae from other algal species (FIG 1, FIG 2), this would indicate an origin sometime between 650 and 1326 million years ago (Ma; *i.e.*, either during the Neoproterozoic (891–629 Ma⁴; ~850–650 Ma⁵⁹ or Mesoproterozoic (1326–920 Ma)² (FIG 2). Based on the fossil record, wood evolved in early Devonian plants (407–397 Ma)^{60,61}, consistent with estimates that Basidiomycota diversified beginning at 295 Ma (95% highest posterior density interval (HPD) 399–195 Ma)³. Therefore, the age of pectin-and hemicellulose- containing

streptophytes provides a maximum age for the evolution of the osmotrophs, and the age of lignin-containing plants provides a maximum age for the evolution of the Basidiomycota. [H2] *Osmotrophic fungi evolved in freshwater habitats.* Streptophyte algae are inferred to have diversified in freshwater settings^{24,25}, and the fungi that digested them must have shared their habitat. Phylogenetic analysis points to a freshwater origin even for red algae, green algae, and their shared ancestors²⁴, beginning at perhaps ~1.5 Ga⁵⁹ (billion years ago; Mesoproterozoic) (FIG 2), followed by later radiations in the oceans. If these evolutionary inferences are correct, then even the earliest phagotrophic fungi also had the opportunity to establish nutritional relationships in a freshwater setting.

Phylogenies further support the evolutionary origin of osmotrophic fungi in freshwater, including the modern fungi that now live in seawater. Freshwater habitats predominate across early-diverging lineages including modern nucleariids and aquatic fungi (FIG 1)⁶². Most of the extant fungi found in seawater appear in phylogenies to be nested among isolates from land, a pattern that is consistent with many independent adaptations to the marine environment^{32,63-65}. The few fungi currently considered to be novel marine species may have close relatives on land that are as yet undiscovered. Strikingly, the fungi most commonly recovered from deep ocean environments are also recovered from tap water, ice cubes, and swimming pools⁶⁶. An estimated 95% or more of modern fungal species diversity is still undocumented by science^{67,68}. The undescribed species from land, freshwater and marine habitats, the ‘dark matter’ of the fungal world⁶⁹, may yet be the key to unlocking the most ancient secrets of fungal origins.

[H2] *Streptophyte algae recognized fungi before land plants evolved fungal mutualisms.* Documented alga-fungus interactions suggest that even before land plants arose, the genetic machinery to recognize contact with fungi had evolved in freshwater streptophyte algae²¹. In modern streptophyte algae, associations with multiple fungal species, including Chytridiomycota, Basidiomycota and Ascomycota, are inferred from metagenomic 28S and 18S rDNA sequence data⁷⁰. Land plants recognize fungi via LysM-RLK receptors localized to their plasma membranes, and streptophyte algae have proteins homologous to these receptors²¹. In land plants, the signaling cascades initiated by receptor stimulation are directed toward defensive, immunity-related pathways or to mutually beneficial mycorrhizal symbioses⁷¹. The genes responsible for the perception of fungal signals and for the downstream signaling pathway are inferred to have been present in the most recent common ancestor of streptophytes²¹. From this ancient origin, these genes show differential patterns of retention. Among streptophyte algae, the loss of the LysM-RLK receptor is inferred for

several species in Zygnematophyceae, whereas it is retained in Charales^{19,72}. Streptophyte algae are missing other gene modules that land plants require for mycorrhizal association with fungi, including those involved in the formation of arbuscules²¹. The presence of genes encoding receptors known to trigger immune responses⁷¹, combined with the absence of the downstream suite of symbiotic genes for mycorrhizal formation²¹, raises the possibility that streptophyte algae were defending themselves against fungal parasites. By contrast, the land plants including hornworts, liverworts and angiosperms share homologs of the necessary symbiotic genes, indicating that the immediate common ancestor of land plants could already form mycorrhizae^{20,21} (FIG 1). On balance, recent analysis is consistent with a hypothesis that ancestral and modern streptophyte algae could recognize fungi but that intracellular mycorrhizal symbiosis is a shared, derived character of land plants.

The abundance and broad phylogenetic distribution of arbuscular mycorrhizal fungi in living land plants is consistent with ancient and enduring relationships between the kingdoms in land-based ecosystems⁷³. Today, arbuscular mycorrhizal fungi are exclusively found within the Mucoromycota, and the origin of (or perhaps predisposition for) modern mutualisms was probably established in the last common ancestor of this fungal clade (FIG 1)⁷⁴. Glomeromycotina has been proposed as the original clade of arbuscular mycorrhizal fungi (FIG 1)⁷⁴, because its species are extremely widespread among modern land plants and because they evolved obligate biotrophy early and retained it throughout their diversification. Recently, Bidartondo et al.⁷⁵ proposed that Endogonales (Mucoromycotina, FIG 1) could also have been the earliest endomycorrhizal fungi, based on observation, and on phylogenetic and physiological studies of liverworts, hornworts, lycophytes and ferns⁷⁵⁻⁸¹. Complicating reconstruction of the ancestral states, mycorrhizae in Mucoromycotina, unlike those of Glomeromycotina, evolved multiple times and their symbiotic networks do not form nested subsets of species⁷⁷. The proposals are not mutually exclusive and proponents of both could claim support from the evidence that Mucoromycotina and Glomeromycotina can simultaneously colonize some hornworts, lycophytes and liverworts^{76,82}.

[H1] Ecosystems of early fungi

[H2] [Au: Again, I've deleted the level two heading here.] Precambrian beginnings of terrestrial, land and water ecosystems. Our understanding of early fungi in freshwater or on land is severely hampered by the rarity of well-preserved terrestrial sediments during the Proterozoic (2600-541 Ma)^{22,24} and Early Paleozoic (541-419 Ma)^{83,84}. Due to the strong sedimentary bias, many of the important and recently discovered Proterozoic eukaryotic

fossils are from tidal or shallow marine sediments⁸⁵⁻⁸⁷. Other geological lines of inquiry indicate that primary production in terrestrial ecosystems underwent greater changes during these periods than at any other time in Earth history. Decreases in the $^{13}\text{C}/^{12}\text{C}$ ratio in Neoproterozoic carbonate rocks starting at 1 Ga have been interpreted as an indicator of a substantial influx of carbon from widespread photosynthesizing communities on land that preferentially incorporated $^{12}\text{CO}_2$ ²³. Although the earliest fossils strongly suggestive of chlorophyte green algae are from coastal marine sediments of the Early Neoproterozoic, from ~1.0 Ga⁸⁸, geochemical traces of green algal C₂₈ and C₂₉ sterols did not appear in marine rocks until ~ 0.65 Ga, suggesting that the biomass of marine algae remained low for hundreds of millions more years²⁶ (FIG 2). This scenario is consistent with the view that, in the early Neoproterozoic, nutrition for fungi may have been more readily available in freshwater than marine environments.

Although direct fossil evidence of eukaryotes living in freshwater or on land in the Neoproterozoic is extremely limited, two geological sequences that show strong, although not undisputed, characteristics of freshwater lake sequences, both approximately 1 Ga in age, stand as exceptions to this general rule and corroborate the data from carbon isotope ratios. Preliminary studies of microfossils from the Nonesuch Shale (Michigan, Wisconsin, USA) suggest that primary production was maintained by planktonic cyanobacteria and benthic cyanobacterial mats^{22,89,90}. Unicellular cyst-forming eukaryotes were present and might also have contributed to primary production, although in freshwater settings there is no direct evidence of morphologically specialized eukaryotic algal plankton beyond simple spherical cells and cell clusters^{22,90}. In the Torridonian Group (Scotland, UK) there is some additional evidence of minute (< 1 mm), possibly thalloid fossils that may have grown on emergent surfaces²². Fossil fungi have not been recorded from land or freshwater sedimentary sequences of this period of time. Patterned perforations in organic walled fossils look like signs of damage by osmotrophic organisms, possibly bacteria, but possibly also fungi (FIG 3 B).

[H2] *Continental carbon sources diversify: Cambrian–Silurian Periods.* During the Early Paleozoic, continental ecosystems underwent dramatic changes with the origin and diversification of land plants and several groups of arthropods (Myriapoda, Hexapoda, and Arachnida)⁸³. Recent calibrated phylogenies trace the divergence of the land plant lineage to freshwater antecedents in the Ediacaran Period (Late Neoproterozoic at ~600 Ma) but place the first diversification of the crown group to between 515 and 474 Ma (Mid Cambrian–Early Ordovician)⁴ (FIG 2). Similarly, phylogenetic analyses of arthropods date land colonization by Myriapoda probably to the Early Cambrian (~540 Ma) and slightly later for Hexapoda and Arachnida, between Ordovician and Silurian (~500–440 Ma)⁹¹. Fungi on land, including Mucoromycota, long predate these events, their first radiations taking place > 600 Ma as judged by dated phylogenies^{2,10,17}. This dating is consistent with a fungal mediated colonization of land by plants, perhaps as a mycorrhizal symbiosis^{21,72,92}.

As in the Neoproterozoic, terrestrial sediments are rare in Cambrian, Ordovician and Silurian rocks, limiting our access to fossil evidence for life in freshwater or on land^{83,84}. Terrestrially-derived organic remains are first found in the middle Cambrian (FIG 3, for example in estuarine muds of the Bright Angel Shale, ~505 Ma in the eastern Grand Canyon (Arizona, USA))⁹³. By the mid-Paleozoic Era (~449 Ma), evidence for abundant organic carbon that could be consumed by fungi is known from the coalified remains of microbial mats probably derived from exposed soil crusts⁹⁴. During the Ordovician through to the mid Silurian (485–419 Ma), evidence for plant life on land also comes indirectly from spores deposited in shallow estuarine marine sediments^{22,83,95}, and from shifts in the geological carbon isotope record possibly attributable to land plant expansion⁹⁶. Taken together, these data suggest that the Early Paleozoic starting at 541 Ma witnessed an extension and diversification of the types of plant-based carbon that are targeted by enzyme systems of osmotrophic fungi, including cellulose, hemicellulose, pectins, cutins, and sporopollenin.

[H1] First fungi come into sight

[H2] *Equivocal evidence of fossil Fungi from Proterozoic saline deposits.* Direct fossil evidence of fungi during the Proterozoic is questionable and limited to a few examples from deep-sea basalts or shallow marine sediments (FIG 2, TABLE 2). The oldest fossils attributed to fungi are filaments found within vesicles and fissures in basalts of the Ongeluk Formation (South Africa) that formed on a seafloor 2.4 billion years ago²⁸. If correctly attributed, these fossils would imply that fungi are twice as old as estimates derived from calibrations of the fungal tree of life (FIG 2)^{2,10,59}. A suitable source of nutrition for such ancient heterotrophs

would be difficult to infer. Furthermore, the deep biosphere is anaerobic, which is inconsistent with the metabolic requirement for oxygen by most fungi. Although intriguing, these filaments do not exhibit any features that would establish a compelling link to fungi, and in our view, even their biological origin remains questionable.

Much more convincing in terms of biogenicity are filamentous fossils recovered from 1.01 to 0.89 Ga (Early Neoproterozoic) shallow estuarine marine deposits of the Grassy Bay Formation Arctic Canada (Table 2)^{27,97}. Named *Ourasphaira giraldae*, these are branched, apparently septate, and they develop conspicuous spheres (33–80 µm diameter) interpreted as spores. By themselves, these features are not compelling evidence of an affinity with fungi as they are also found in algae. Analysis of the organics by Fourier-transform infrared (FTIR) spectroscopy yielded absorption bands that were interpreted as indicating chitin or chitosan typical of fungi (BOX 1)²⁷. We contend however that controls would be needed to show that FTIR could distinguish fossilized, partially degraded chitin from cellulose⁹⁸, and that lacking these controls, the affinities of *O. giraldae* with fungi in general and Dikarya in particular remain doubtful. Most recently, pseudo-septate filaments attributed to fungi were discovered in 0.80 to 0.72 Ga (Early-Mid Neoproterozoic) sediments recovered from deep drill cores made originally during the 1950s in the Mbuji-Mayi Supergroup (Democratic Republic of Congo)²⁹ (TABLE 2). It should be noted that an alternative perspective on the dating of this sequence would make these fossils very much older, perhaps as much as ~1Ga⁹⁹. The sediments are a marine influenced dolomite that is thought to have been subject to aerial exposure. Evidence of chitin was inferred using a combination of complementary analytical techniques (TABLE 2, BOX 1). The common theme among these recent reports is a filamentous structure with few morphological characteristics. Establishing a fungal affinity relies on ruling out filamentous bacteria or archaea and photosynthetic eukaryotes by establishing the presence of chitin residues in thermally altered and partly decomposed organics using various chemical analytical techniques. Even if one accepts the evidence of chitin, it is not possible to specify a more precise position for these fossils in the fungal tree of life.

[H2] *Fossil evidence of Fungi in the Paleozoic.* Direct evidence of fossil fungi from the Cambrian through Silurian Periods is sparse. Fossil hyphae and spores of *Palaeoglomus grayi* from the Ordovician Guttenberg Formation (Wisconsin, USA) are frequently cited as the earliest evidence of Glomeromycotina at ~460 Ma¹⁰⁰. Although resembling extant Glomeromycotina, they are not associated with any plant fossils. Furthermore, these remains

could be more recent contaminants of the porous dolomitic sediment in which they were found¹¹.

Tortotubus protuberans (also known as *Ornatifilum lornensis*), when first documented from Silurian sediments, was interpreted as an ascomycete with what appeared to be lateral, flask-shaped, spore-producing cells¹⁰¹. However, Smith¹⁰² compared material from several Silurian and Devonian sites, and he showed that the flask-shaped cells were broken bases of filaments instead of spore-producing structures, casting serious doubt on an affinity with Ascomycota. Fossils of *T. protuberans* are plausibly of terrestrial origin and their features include branched filaments with pseudo-septate pores. Smith's study illustrates the value of basing interpretation on careful observations and a rich set of collections.

The beginning of the Devonian Period (419 Ma) saw a substantial change in the nature of the geological record marked by the widespread development of terrestrially deposited sediments on a global scale. The shift from dominantly marine to terrestrial preservation resulted in the first appearance of many land-dwelling lineages in the fossil record^{4,83,103}. Many fossil sites from across Europe, Asia, the Americas and Australia testify to the diversity of life on land, but one site stands out because organisms and their interactions were remarkably preserved. The Rhynie chert (407 Ma, FIG 3) preserves elements of a geothermal wetland including seven species of small herbaceous vascular plants (~ 20 cm maximum height), representatives of eight groups of arthropods, one nematode, several groups of green algae including a charophycean streptophyte alga, oomycetes, testate amoebae, cyanobacteria and numerous varied fungi, including Chytridiomycota, Blastocladiomycota, Mucoromycota, and possibly Ascomycota^{11,104-108}. Furthermore, Rhynie chert plants and fungi demonstrated saprotrophic^{11,30,109} (FIG 4) as well as symbiotic (both parasitic¹⁰⁸⁻¹¹¹ and mutualistic^{76,112,113}) interactions (FIG 3). Mycorrhiza-like associations in the Rhynie chert provide the earliest evidence of a relationship that can be traced through the geological record up to the present time. Fossils document a Mucoromycotina mycorrhizal association⁷⁶, in addition to two well-known fossils of arbuscular mycorrhizal associations involving Glomeromycotina^{76,112,113}. In early fossil plants (lacking roots or rhizomes) the development of arbuscular mycorrhizae was in prostrate and aerial axes^{76,112,113}, a situation similar to that found in some modern bryophytes, ferns and orchids¹¹⁴. The diverse and well-established mycorrhizal interactions between fungi and plants in the Rhynie chert are consistent with molecular phylogenic and genomic inferences that these interactions predate the Devonian Period by over 100 Ma.

[H2] *Puzzling fungi of the Devonian Period.* As the sedimentary window on to early ecosystems opened during the Devonian Period, many new organisms came into view for the first time. When interpreting their affinities and the nature of their interactions, it is tempting to shoehorn them into modern groups or categories that might not have existed at the time. Yet many taxa of the Devonian may now be extinct without descendants.

A prime example of the problem of identifying fossil fungi based on the morphology of extant species is *Paleopyrenomycites devonicus* from the Rhynie chert. This important fossil is widely used in dating the fungal tree of life (TABLE 2)^{115,116} and an affinity with Ascomycota is generally accepted. However, its position within the clade—whether basal in Ascomycota or nested among Pezizomycotina—is contentious^{2,12,117,118}. It is worth noting that dispersed spores and other evidence of an Ascomycota radiation do not appear in the sedimentary record until much later, during the Mesozoic Era, >250 Ma^{119,120}, providing support for the idea that *Paleopyrenomycites devonicus* represents an extinct clade of early-diverging Ascomycota.

Among the most unexpected fossils allied to fungi is *Prototaxites*, which is encountered in Devonian rocks ~419-359 Ma¹²¹. The largest specimens resembled tree trunks or logs, reaching over 8 meters in length although small specimens also occurred. Based on its structure of interwoven filaments and tubes of various sizes, Hueber¹²² hypothesized that *Prototaxites logani* was a giant sporophore of a species in the Basidiomycota. Recently, an affinity with Ascomycota was suggested for the small *Prototaxites taiti* with the discovery of a 1 mm-thick layer of meiotic cells, 'asci', each with more than 8 spores¹²¹. The ecological role of *Prototaxites* remains enigmatic. Hobbie and Boyce¹²³ concluded that the carbon isotope ratios of *Prototaxites* indicated that it was a saprotrophic fungus. Selosse argued that the surrounding Devonian primary producers could not have supported such a large saprotroph, and suggested that lichenization, which would have enabled carbon acquisition from algal partners, was more likely¹²⁴.

Adding to the mysteries of *Prototaxites* is the recently described *Cyathinema digermulense*¹²⁵. This organically-preserved fossil from the Nyborg formation in Arctic Norway at 580 Ma resembles the charcoalfied body parts of *Prototaxites* sp.¹²¹, although *Cyathinema* predates *Prototaxites* by 150 million years. In particular, the cups-and-tubes of *Cyathinema* resemble the tubular threads in the hymeninal layer of *Prototaxites*. A more prosaic interpretation of the phylogenetic relationships of *Cyathinema* would place it on the red algal lineage, an interpretation consistent with its probable location in a shallow-marine habitat, rather than in a terrestrial habitat as occupied by *Prototaxites*. This example

illustrates the implicit challenges in distinguishing fossil fungi from algae, while highlighting the importance of finding more fossils to understand the diversity of early eukaryotes.

Lichens, mutualistic symbioses between fungi and cyanobacteria or chlorophyte algae, are polyphyletic, arising from multiple lineages of Ascomycota and Basidiomycota. Based on recent molecular clock analysis of multiple lichen fungi and their diverse, polyphyletic algal associates, Nelsen et al.¹² concluded that extant lichen lineages originated after, not before land plants. Lichenization probably enabled diversification of the partners after the rise of multicellular land plants, by adaptation to niches unsuitable for the latter⁷³. Cyanobacteria are found in ~ 15% of lichens and they have been available as potential photosynthetic partners for ~2.4 Ga, yet there is no evidence for their involvement in lichens until after the evolution of land plants¹².

The early fossil evidence for lichens is scant¹²⁶. The controversial interpretations of *Dickinsonia* fossils from the Ediacaran (635 to 541 Ma) as lichens¹²⁷ or fungi¹²⁸ have been widely criticized¹²⁶ and are further contradicted by recent geochemical evidence of cholestane, which is a residue of cholesterol, a sterol that is more consistent with an animal than fungal interpretation¹²⁹. Likewise, the lichen-like symbioses with cyanobacteria in the marine Doushantuo Formation (551 to 635 Ma)¹³⁰ are less than compelling and alternative hypotheses involving filamentous bacteria rather than fungi are possible. The lichen-like *Winfrenatia reticulata* described in the terrestrial Rhynie chert (407 Ma)¹³¹⁻¹³³ bears no morphological similarity to extant lichens, and, although it represents a loose consortium of fungi and cyanobacteria (TABLE 2), a mutualistic relationship cannot be inferred from the morphological clues that the fossil presents. The earliest, most compelling evidence of fossil lichens comes from Early Devonian (415 Ma) sediments of the Ditton Group (England)^{134,135}. Two types of thallus-like fossils preserved in charcoal (*Chlorolichenomycites salopensis* and *Cyanolichenomycites devonicus*) have an internally stratified structure. Fungal hyphae are exceptionally well-preserved; photobionts less so but plausibly represented by clusters of pyrite crystals. The fungus was interpreted as a Pezizomycotina but it would be older than estimates of the stem age of extant lichens (FIG 2)¹². Here again, the Devonian fossils might be extinct lineages of lichens that are unrepresented in molecular phylogenetic studies of living species.

[H1] Conclusions and future directions

We contend that fungi were early colonists of freshwater environments that were already occupied by biofilms of cyanobacteria and photosynthetic eukaryotes, the forerunners of

green and red algae and land plants. The earliest fungi were protist-like with phagotrophic nutrition. Osmotrophic nutrition and mycelial growth initially developed as responses to the availability of abundant carbon sources produced by photosynthetic streptophytes. The diversification of pectinase and xyloglucanase genes in fungi and the capacity for streptophyte algae to recognize and respond to fungi indicate that fungal adaptations to land plants as a source of carbon began with their freshwater algal ancestors. Another major step enabled the rise of the current terrestrial flora and fungi: the mycorrhizal association. The rise and spread of woody plants on land triggered a later diversification of oxidative fungal enzymes capable of depolymerizing lignin to make available for fungal nutrition the vast stores of cellulose and hemicellulose trapped in wood. Perhaps similar patterns could also be observed for other complex organic molecules that appeared early in the geological record (for example, sporopollenin and cutin).

Osmotrophic fungi that live in modern ocean sediments and in the deep biosphere are most probably derived from freshwater or land species. Rare Precambrian freshwater sediments do exist, and these perhaps hold the best prospects for the discovery of mutually corroborating fossils of fungi. It would be fascinating to find new, deeply diverging extant species from freshwater and marine habitats and then to put such fungi into a phylogenomic perspective. Fungal diversity is still poorly known and bringing the fungal dark matter to light will help to decipher the origins of their kingdom. A deeper understanding of the modern diversity of fungi in soils, freshwater and marine systems is needed to understand fungal roles in modern ecosystems and will be helpful to the interpretation of the fossils in the early geological record.

We have reviewed the profound insights that are resulting as a flood of new techniques are applied to the analysis of fungal genomes and to the analysis of fossil chemistry and morphology. At present, dated phylogenies and genomic evidence for interactions with streptophyte hosts indicate that fungi are older than the unambiguous evidence for them provided by their 407 Ma Rhynie chert fossils. Fascinating, newly discovered fossils backed by chemical analysis are far older, but their occurrences are separated by gaps of over a hundred million years, and their identity as fungi is equivocal and has yet to be corroborated by information about their habitats or ecology.

We anticipate that discoveries and interpretation of new fossils will close gaps to reveal patterns of fungal continuity, radiation and extinction through the ages. The sophistication of evolutionary inferences from genomes will be increased by taking into account increasingly robust patterns recorded in the fossils. We hope that new researchers

and their funding agencies take note of the opportunities that are arising for the critical challenge of fundamental ideas about ancient fungi and their ecology. Recognizing the fungi, correlating them with habitat, and placing them into the tree of life, although never straightforward, will pave the way to a better understanding of ancient ecosystems.

References

[Au: For references that are particularly worth reading (5-10% of the total), please provide a single bold sentence that indicates the importance of the work.]

Acknowledgements

The authors thank Victor Leshyk (<http://victorleshyk.com/paleo.html>) for the reconstruction of Rhynie landscape. [Au: Do you have any other acknowledgements?]

Author contributions

The authors contributed equally to all aspects of the article.

Competing interests

The authors declare no competing interests.

Peer review information

Nature Reviews Microbiology thanks Antonis Rokas and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Table 1. **Host organisms for fungi through geological time**

Host		Hosts and associated fungi through geological time
Cyanobacteria	NRMICRO-19-159-Berbee-t1.1	Cyanobacteria made up a large fraction of aquatic biomass through the Precambrian, but there is no convincing fossil evidence that early fungi fed upon them. The diversity of fungi associated with modern cyanobacteria is low.
Streptophyte	NRMICRO-19-	Before land plants, early saprotrophic or symbiotic

green algae	159-Berbee-t1.2	fungi may have used freshwater streptophyte algae as a source of nutrition. Genomes of both algae and fungi point to a relationship going back ~750 Ma. Ancestral streptophytes gave rise to some species such as <i>Nitella hyalina</i> ¹³⁶ , shown here, which remained aquatic; other descendants evolved into land plants, possibly aided by mutually beneficial relationships with fungi.
Land plants	NRMICRO-19-159-Berbee-t1.3	Since their earliest proposed origin ~500 Ma ⁴ , land plants including bryophytes and vascular plants have been the most important source of carbon for the vast majority of saprotrophic and symbiotic fungi. Land plants are also the main beneficiaries of fungal ecosystem services including mycorrhizal partnerships and decomposition.
Green algae or cyanobacteria in lichens	NRMICRO-19-159-Berbee-t1.4	. Lichens are stable, mutualistic symbiotic associations where eukaryotic algae and/or cyanobacteria provide carbon to their fungal partners. Dated phylogenies show that lichens arose convergently, and that the clades of algae, cyanobacteria, and fungi that gave rise to extant lichens did not radiate until ~400 Ma, after land plants had begun to diversify ¹² .
Animals	NRMICRO-19-159-Berbee-t1.5	. Modern fungi include the predominantly aquatic Chytridiomycota, some of which cause diseases (such as chytridiomycosis in amphibians). Fungal species that infect animals are derived from more general saprotrophs, for example ref ¹³⁷ ; fungal evolution did not depend primarily on animals.

487 Ma, million years ago. Images provided by (top to bottom): NASA, Kenneth Karol (New
488 York Botanical Garden), Daniel Mosquin (University of British Columbia Botanical Garden;
489 both liverwort and lichen) and REF.¹³⁸.

490

491
492
493
494

Table 2. Interpretation of fossils: fungi or equivocal?

Fossil		Age and morphology (techniques)	Environmental conditions	Interpretation	Reason for the interpretation
Unnamed filament ²⁸	NRMICR O-19-159-Berbee-t2.1	2.4 Ga; filament (Syn, LM and SEM)	Not organic, anaerobic, marine environment	Uncertain biogenicity; probably not fungal	Inconsistent with other data. Too old; molecular clock dates eukaryotes 1.9-1.7 Ga ⁵⁹ ; first complex walled (eukaryotic) acritarchs ~1.7 Ga ¹³⁹
<i>Ourasphaira giralda</i> ^{27,97}	NRMICR O-19-159-Berbee-t2.2;	1.0-0.89 Ga; filaments and spores (LM, SEM, TEM, μ FTIR and Raman	Organic from estuarine environment	Possible fungal affinity, might be protist or alga	Unexpected but of great interest. No other bona fide fungal fossils are of similar age; fungal phylogeny predicts

		spectroscopy)			non-filamentous fungi this early ¹⁷ . Possible source of nutrition: consistent with fossil evidence of marine algae ^{88,140} ; predates geochemical evidence of expansion of marine eukaryotic algae ²⁶
Unnamed mycelium-like structures ²⁹³	NRMICR O-19-159-Berbee-t2.3	810 to 715 Ma; filaments (LM, SEM, TEM, μ FTIR, XANES, Raman spectroscopy and chitin staining)	Organic from coastal, lagoon, perennial lacustrine pond	Uncertain fungal affinity, might be alga	Consistent with fungal phylogeny ¹⁷ ; possible source of nutrition consistent with fossil evidence of marine algae ^{88,140} ; predates geochemical

					l evidence of expansion of marine eukaryotic algae ²⁶
<i>Cyathinema digermulense</i> ¹²⁵	NRMICR O-19-159-Berbee-t2.4;	580 Ma; filaments (LM, SEM and Raman spectroscopy)	Organic from shallow marine environment, a deltaic environment cannot be excluded	Uncertain affinity: share some resemblance with <i>Prototaxites</i> ; might be alga	Consistent with fungal phylogeny ¹⁷ ; possible source of nutrition consistent with fossil evidence of marine algae ^{88,140}
<i>Palaeoglomus grayi</i> ^{100,141}	NRMICR O-19-159-Berbee-t2.5	460 Ma; filaments and spores (LM)	Organic from shallow marine environment	Fungal (Glomeromycotina); possibly contaminant, found in dolomitic rock	Consistent with fungal and plant phylogenies ^{5,17} ; early plants, a possible source of nutrition, would have been available ¹⁴²
<i>Tortotubus</i> ^{102,143}	NRMICR O-19-159-Berbee-	419-359 Ma; filaments and spores	Organic from shallow marine or	Possibly fungal but unlike any living taxon, leaving open a	Consistent with fungal and plant phylogeny ⁵ ;

	t2.6	(LM and SEM)	terrestrial environment	possible protist or algal affinity	early land plants, a possible source of nutrition were present ¹⁴² .
<i>Paleopyrenomycites</i> 115,116	NRMICR O-19- 159- Berbee- t2.7	407 Ma; filaments, sporophor es and spores (LM)	Organic from terrestrial environment ; sporulating structure in land plant axis	Fungal, likely Dikarya; possibly extinct clade of early- diverging Ascomycota	Unexpected : corroboratin g Ascomycot a fossils appear ~250 Ma; consistent with expectation s for Dikarya, the fossil is immersed in a plant stem that it used for nutrition; fossil widely used to calibrate fungal molecular clocks
<i>Winfrenatia</i> ^{131,133}	NRMICR	407 Ma;	Organic	Affinity	Consistent

	O-19-159-Berbee-t2.8	filaments (LM)	from terrestrial environment ; cyanobacteria associated with fungal filaments	unresolved; could be a lichen in the broad sense, but not in the narrower sense of a stable, spatially correlated arrangement of a fungus and a photobiont	with fungal phylogeny ¹⁷ , and with fossil evidence of fungi and cyanobacteria ¹²
<i>Prototaxites</i> ^{121,122}	NRMICR O-19-159-Berbee-t2.9	419-370 Ma; filaments, sporophores and spores (LM, SEM and CLSM)	Organic from terrestrial environments	Affinity unresolved; possibly fungal, but unlike any living taxon	Available sources of nutrition included algae ²⁴ , microbial mats ⁹⁴ , and early land plants ¹⁴² .

Syn: Synchrotron; LM: light microscopy; SEM: scanning electron microscopy; TEM: transmission electron microscopy; μ FTIR: Fourier transform infrared microspectroscopy; CLSM: confocal laser scanning laser microscopy; XANES: X-ray absorption near edge structure. Images provided by (top to bottom): image reproduced with permission from REF.²⁸; REF.⁹⁷; REF.²⁹; REF.¹²⁵; Dirk Redecker (Université de Bourgogne); Paul Strother; Christine Strullu-Derrien; Hans Kerp (University of Münster); and Christine Strullu-Derrien.

Fig. 1. Evolution of early fungi. Fungi and animals evolved in the Precambrian from phagotrophic, amoeba-like ancestors that engulf nutrients into food vacuoles within their cells and left no known fossil record. The dated phylogeny (top) shows that modern, multicellular

fungi descended from unicellular phagotrophs that evolved cell walls and became osmotrophic, secreting digestive enzymes into the surrounding environment. Fungi originated in freshwater and undertook several evolutionary transitions (bottom). (1) Predating fungi, fossils of cyanobacteria are known from freshwater⁸⁹ and marine habitats (a). Stem Viridiplantae are ancestors to all green algae²⁵(b). (2) Phagotrophic amoebae (c) gave rise to fungi^{35,62}. (3) Unicellular streptophyte alga (d) arose. Early osmotrophic fungi (e) parasitized a multicellular streptophyte alga and saprotrophic fungi (f) digested detritus including streptophyte cell walls. (4) Cells of early vascular plants contain symbiotic fungal arbuscules¹¹² and hyphal coils⁷⁶ (g). (5) Wood, substrate for radiation of decay fungi³, evolved in the Devonian (407-397 Ma)^{60,61}. Dates of divergence of plants and chlorophyte algae are from Morris⁴, of most fungi from Chang et al.^{17,19}, of microsporidia, aphelids and *Nuclearia* are arbitrary, and all others from Parfrey et al.⁵⁹ Image (c) is adapted with permission from REF.³³

Figure 2. Geological ages of fungi, host plants, and pivotal events in eukaryote

diversification. Estimates from molecular clocks (dashed lines) are compared with direct fossil or biomarker evidence (solid lines). Perplexing 'fungal' fossils are shown below the geological timeline. These and other recently discovered fossils delight by inviting speculation while confounding expectations. Permission to reproduce images is from: (a) REF.²⁸; (b) REF.⁹⁷; (c) REF.²⁹; (d) REF.¹²⁵; (e) Dirk Redecker (Université de Bourgogne); (f) Paul Strother; (g) Christine Strullu-Derrien; (h) Hans Kerp (University of Münster); (i) Christine Strullu-Derrien..

Figure 3. Early Paleozoic fungi: landscapes, hosts and fossils. The top row shows early terrestrial life. Fossilized cyanobacteria (~530 Ma) accumulated around the edges of a former vernal pool, sitting on oxidized Cambrian Tapeats Sandstone in the Grand Canyon (a). Patterned perforations (arrows) in the vesicle walls indicate microbial, possibly actinomycete or fungal damage to unicellular microfossil from 1 Ga lake sediments, from the Torridonian Sequence in Scotland (b). Landscapes of cryptogamic crusts of fungi, algae, and bryophytes in Neoproterozoic to Cambrian landscapes (850-485 Ma) may have looked like the basalt fields in Iceland (c). Middle and bottom rows show Early Devonian 407 Ma Rhynie environment and Rhynie chert fossils. Leafless plants surrounded silica-rich hot springs in Rhynie, Scotland (d). Fossilized axes of early land plants were preserved (e). Bottom row shows microscopic images of fossils from thin sections. Arrows indicate mycorrhizal fungal arbuscules, fungal hyphae that branch into successively finer filaments, within cells of extinct land plant *Aglaophyton major* (f). The streptophyte alga *Palaeonitella cranii* consists of an

axis with whorls of filaments (bars 100 μm) (g, h) Asterisks indicate successive internodes in a healthy alga (left) (g); asterisks indicate successive internode cells enlarged in response to infection by a parasitic fungus (right) (h), The zoosporangium of a Chytridiomycota fungus is attached to a wavy-walled rounded structure by a rhizoid (arrow) (i).

Permission to reproduce images is from: Paul Strother (a, b); Anthony Ives (University of Wisconsin-Madison) (c); Victor Leshyk (Victor Leshyk Illustration) (d); Christine Strullu-Derrien (f); REF¹¹ (g, h); REF¹⁰⁸ (i).

Figure 4. Rhynie chert sporangium. Confocal laser scanning microscopy reveals internal structural detail of a sporangium of *Retesporangicus lyonii* from the 407 Ma Rhynie chert. Light microscopy (a), confocal laser microscopy (b-d), image processed (c, d³⁰ to show outer surface of the sporangium (c) and closely packed interior spores (d). Bottom two images reproduced with permission from REF.³⁰.

Box 1. New tools for old fossils

Molecular approaches

- Molecular clocks estimate geological ages of nodes using fossil calibrations, and by assuming that modeled molecular change is proportional to time. Reliable clocks would be based on accurate phylogenies, correct interpretation and dating of fossils¹⁴⁴, and on correct modeling of molecular evolutionary processes.
- Criteria for reliable clocks are hard to meet. The temptation is strong to constrain clade ages closely by over-interpreting the few available fossil calibration points, ignoring alternative interpretations that would more realistically reveal uncertainty.

Microscopy

- Confocal laser scanning microscopy (CLSM) yields 3D images of minute objects in exquisite detail with a resolution of $<1\ \mu\text{m}$ ¹⁴⁵. This method resolved spores encased in fungal sporangia and embedded in Rhynie chert^{108,146}. The CLSM digital tomographic datasets (sets of optical sections) are converted computationally into 3D visualizations or animations of fungal structures³⁰. Similar to other techniques listed below, CLSM is largely non-destructive.
- Scanning electron microscopic analysis of weathering patterns in mineral particles, together with detection of high levels of iron by energy-dispersive X-ray spectroscopy can help distinguish fungal fossils in paleosols³¹.
- Fluorescent stains characterize cell wall components, although without high specificity. Wheat-germ agglutinin complexed with a fluorophore labels the N-acetylglucosamine residues in chitin. However, it also binds other sugars, and N-

acetylglucosamine occurs in other organisms, for example, sheaths of some cyanobacteria¹⁴⁷. Calcofluor white fluorescence labels chitin, cellulose and other polysaccharides.

Spectroscopic methods

- Raman microspectroscopy can reveal the geological history of exposure to heat and pressure of an organic fossil^{29,148,149}. This data can help distinguish *bona fide* fossils from recent contaminants.
- Fourier transform-infrared (FITR) microspectroscopy involves infrared irradiation of samples and it detects organic functional groups including the amino groups that distinguish chitin from cellulose^{27,98}.
- Synchrotron X-ray absorption near edge structure (XANES) uses high energy irradiation to quantify speciation and bonding in inorganic-rich fossils^{29,98}.

In summary, although new genomic, microscopic and spectroscopic techniques cannot roll back the passage of time, they can, in combination, provide tantalizing new clues that challenge our concepts of the age, morphology and ecology of the most ancient fungi.

Glossary

Mycorrhiza: Soil fungi make close contact with plant roots or other organs to enter a mutualistic symbiosis, providing mineral nutrients to the plant, and gaining carbon energy in return.

Symbionts/symbiosis: Symbionts are unlike organisms that live together in a symbiosis, a close relationship; used here in the historical sense of De Bary 1879, the relationship can be mutualistic, benefiting both partners, or parasitic where one organism benefits at the expense of another.

Precambrian: The geological interval beginning with the origin of the Earth at 4.6 Ga and ending at 541 Ma; encompassing the Proterozoic Eon (2500-541 Ma).

Terrestrial: Non-marine geological environments; includes freshwater as well as land.

Paleozoic: The geological era extending from 541 to 252 Ma, after the Precambrian.

Nucleariid amoebas: Free-living, unicellular amoeboid protists on the fungal lineage that entrap their prey with thread-like filose pseudopodia.

Holozoa: Multicellular animals (Metazoa) and their closest (less well known) unicellular relatives, including *Sphaeroforma* and collar flagellates.

Aphelids: Unicellular parasites counted among fungi, the amoeba-like aphelids feed by injecting themselves into an algal cell, and then engulfing and ingesting its contents.

Arbuscules: Symbiotic, tree-like, branched fungal filaments within a plant cell that function as sites of plant-fungus nutrient exchange in arbuscular mycorrhizae.

Biotrophy: Nutrition from other living organisms; may refer to mutualists or parasites that are not killing their host.

Thalloid: Resembling a plant-like body but lacking roots, stems, leaves.

Testate amoebae: Unicellular protists, polyphyletic, with an organic, mineral or agglutinated shell (test) that partially encloses the cell.

Prostrate axes: Creeping stem-like branches of early fossil plants that can be surficial or subterranean and that do not have all the characters of a root or a rhizome of living plants.

Aerial axes: Erect stem-like branches of early fossil plants that do not have all the characters of a stem of living plants.

Sporophore: Spore-producing structure in fungi; a mushroom is a modern example.

Acritarchs: Organic-walled microfossils of biological affinity, some eukaryotic.

ToC blurb

Fungi originated in a freshwater environment and their evolution accompanied the rise of algae and land plants. In this Review, Berbee and colleagues examine the fossil and genomic record of ancient fungi and the inferences we can make about their lifestyle.

- 1 Strullu-Derrien, C., Selosse, M. A., Kenrick, P. & Martin, F. M. The origin and evolution of mycorrhizal symbioses: from palaeomycology to phylogenomics. *New Phytologist* **220**, 1012-1030, doi:10.1111/nph.15076 (2018).
- Thorough review of fossil and phylogenetic evidence for the geological first appearances and diversification of mycorrhizae.**
- 2 Lutzoni, F. *et al.* Contemporaneous radiations of fungi and plants linked to symbiosis. *Nature Communications* **9**, doi:10.1038/s41467-018-07849-9 (2018).
- 3 Floudas, D. *et al.* The Paleozoic origin of enzymatic lignin decomposition reconstructed from 31 fungal genomes. *Science* **336**, 1715-1719, doi:10.1126/science.1221748 (2012).

- 4 Morris, J. L. *et al.* The timescale of early land plant evolution. *Proceedings of the National Academy of Sciences of the United States of America* **115**, E2274-E2283, doi:10.1073/pnas.1719588115 (2018).
- Timescale of evolution of green algae and land plants based on thoughtful calibrations and genome-scale analysis.**
- 5 Puttick, M. N. *et al.* The interrelationships of land plants and the nature of the ancestral embryophyte. *Current Biology* **28**, 733-+, doi:10.1016/j.cub.2018.01.063 (2018).
- 6 Lenton, T. M. *et al.* Earliest land plants created modern levels of atmospheric oxygen. *Proceedings of the National Academy of Sciences of the United States of America* **113**, 9704-9709, doi:10.1073/pnas.1604787113 (2016).
- 7 Andrews, M. Y., Leake, J. R., Palmer, B. G., Banwart, S. A. & Beerling, D. J. Plant and mycorrhizal driven silicate weathering: Quantifying carbon flux and mineral weathering processes at the laboratory mesocosm scale. *Applied Geochemistry* **26**, S314-S316, doi:10.1016/j.apgeochem.2011.03.072 (2011).
- 8 Quirk, J. *et al.* Constraining the role of early land plants in Palaeozoic weathering and global cooling. *Proceedings of the Royal Society B-Biological Sciences* **282**, doi:10.1098/rspb.2015.1115 (2015).
- 9 Cohen, K., Finney, S., Gibbard, P. & Fan, J. The ICS International Chronostratigraphic Chart. *Episodes* **36**, 199-204, doi:10.18814/epiiugs/2013/v36i3/002 (2013).
- 10 Chang, Y. *et al.* Phylogenomics of Endogonaceae and evolution of mycorrhizas within Mucoromycota. *New Phytologist* **222**, 511-525, doi:10.1111/nph.15613 (2019).
- 11 Taylor, T., Krings, M. & Taylor, E. *Fossil Fungi*. (Elsevier/Academic Press, 2015).
- 12 Nelsen, M. P., Lücking, R., Boyce, C. K., Lumbsch, H. T. & Ree, R. H. No support for the emergence of lichens prior to the evolution of vascular plants. *Geobiology* **18**, 3-13, doi:10.1111/gbi.12369 (2020).
- Time-calibrated phylogenies show that the fungal and algal lineages that form lichens have evolved after vascular plants; lichens in the modern sense could not have been the earliest terrestrial organisms.**
- 13 Richards, T. A., Leonard, G. & Wideman, J. G. What Defines the "Kingdom" Fungi? *Microbiology Spectrum* **5**, doi:10.1128/microbiolspec.FUNK-0044-2017 (2017).
- 14 Berbee, M. L., James, T. Y. & Strullu-Derrien, C. Early Diverging Fungi: Diversity and Impact at the Dawn of Terrestrial Life. *Annu. Rev. Microbiol.* **71**, 41-60, doi:10.1146/annurev-micro-030117-020324 (2017).
- 15 Mikkelsen, M. D. *et al.* Evidence for land plant cell wall biosynthetic mechanisms in charophyte green algae. *Annals of Botany* **114**, 1217-1236, doi:10.1093/aob/mcu171 (2014).
- 16 Lange, L. *et al.* Origin of fungal biomass degrading enzymes: Evolution, diversity and function of enzymes of early lineage fungi. *Fungal Biology Reviews* **33**, 82-97, doi:10.1016/j.fbr.2018.09.001 (2019).
- 17 Chang, Y. *et al.* Phylogenomic analyses indicate that early fungi evolved digesting cell walls of algal ancestors of land plants. *Genome Biology and Evolution* **7**, 1590-1601, doi:10.1093/gbe/evv090 (2015).

- 18 Liang, Y. *et al.* Lipochitooligosaccharide recognition: an ancient story. *New Phytologist* **204**, 289-296, doi:10.1111/nph.12898 (2014).
 - 19 Cheng, S. *et al.* Genomes of subaerial Zygnematophyceae provide insights into land plant evolution. *Cell* **179**, 1057-1067.e1014, doi:<https://doi.org/10.1016/j.cell.2019.10.019> (2019).
 - 20 Li, F.-W. *et al.* *Anthoceros* genomes illuminate the origin of land plants and the unique biology of hornworts. *Nature Plants* **6**, 259-272, doi:10.1038/s41477-020-0618-2 (2020).
 - 21 Delaux, P. M. *et al.* Algal ancestor of land plants was preadapted for symbiosis. *Proceedings of the National Academy of Sciences of the United States of America* **112**, 13390-13395, doi:10.1073/pnas.1515426112 (2015).
- Comparative analysis of plant genes required for mycorrhiza formation shows that green algae and land plants share orthologous proteins for recognition of fungi, while proteins for mutualistic interactions arose in the common ancestor of the land plants.**
- 22 Wellman, C. H. & Strother, P. K. The terrestrial biota prior to the origin of land plants (embryophytes): a review of the evidence. *Palaeontology* **58**, 601-627, doi:10.1111/pala.12172 (2015).
- Precambrian terrestrial fossils are rare but this review outlines evidence from carbon isotopes, weathering patterns of clay, and from fossil diversity pointing to early Precambrian diversification of a complex terrestrial biota.**
- 23 Knauth, L. P. & Kennedy, M. J. The late Precambrian greening of the Earth. *Nature* **460**, 728-732, doi:10.1038/nature08213 (2009).
- Geological data on decreasing ratios of ^{13}C to ^{12}C in marine carbonate rocks is interpreted to represent expanding, photosynthetic, terrestrial biomass, beginning in the Neoproterozoic, ~850 Ma.**
- 24 Sánchez-Baracaldo, P., Raven, J. A., Pisani, D. & Knoll, A. H. Early photosynthetic eukaryotes inhabited low-salinity habitats. *Proceedings of the National Academy of Sciences of the United States of America* **114**, E7737-E7745, doi:10.1073/pnas.1620089114 (2017).
- Surprisingly, ancestral habitats of cyanobacteria, red and green algae first are inferred phylogenetically to be non-saline, even though the earliest fossils of these taxa are marine. The lack of early terrestrial fossils may reflect missing data due to a bias against terrestrial fossilization.**
- 25 Del Cortona, A. *et al.* Neoproterozoic origin and multiple transitions to macroscopic growth in green seaweeds. *Proceedings of the National Academy of Sciences of the United States of America* **117**, 2551-2559, doi:10.1073/pnas.1910060117 (2020).
 - 26 Brocks, J. J. *et al.* The rise of algae in Cryogenian oceans and the emergence of animals. *Nature* **548**, 578-+, doi:10.1038/nature23457 (2017).
- Biomarkers, steranes preserved in marine rocks, indicate that green algae first became abundant in the seas some ~650 Ma.**

- 27 Loron, C. C. *et al.* Early fungi from the Proterozoic era in Arctic Canada. *Nature*, doi:10.1038/s41586-019-1217-0 (2019).
- A fossil >900 Ma has morphological characters and a chemical profile consistent with identification as a fungus.**
- 28 Bengtson, S. *et al.* Fungus-like mycelial fossils in 2.4-billion-year-old vesicular basalt. *Nature Ecology & Evolution* **1**, doi:10.1038/s41559-017-0141 (2017).
- 29 Bonneville, S. C. *et al.* Molecular identification of fungi microfossils in a Neoproterozoic shale rock. *Science Advances* **6**, doi:10.1126/sciadv.aax7599 (2020).
- 30 Strullu-Derrien, C. *et al.* New insights into the evolutionary history of Fungi from a 407 Ma Blastocladiomycota fossil showing a complex hyphal thallus. *Philosophical Transactions of the Royal Society B: Biological Sciences* **373** (2018).
- 31 Mitchell, R. L., Strullu-Derrien, C. & Kenrick, P. Biologically mediated weathering in modern cryptogamic ground covers and the early Paleozoic fossil record. *Journal of the Geological Society* **176**, 430-439, doi:10.1144/jgs2018-191 (2019).
- 32 Baldauf, S., Romeralo, M. & Carr, M. in *Evolution from the Galapagos, Two Centuries after Darwin* (eds Gabriel Trueba & Carlos Montúfar) 73-106 (Springer, 2013).
- 33 Yoshida, M., Nakayama, T. & Inouye, I. *Nuclearia thermophila* sp nov (Nucleariidae), a new nucleariid species isolated from Yunoko Lake in Nikko (Japan). *European Journal of Protistology* **45**, 147-155, doi:10.1016/j.ejop.2008.09.004 (2009).
- 34 Letcher, P. M. & Powell, M. J. A taxonomic summary of Aphelidiaceae. *Ima Fungus* **10**, doi:10.1186/s43008-019-0005-7 (2019).
- 35 Cavalier-Smith, T. Megaphylogeny, cell body plans, adaptive zones: causes and timing of eukaryote basal radiations. *Journal of Eukaryotic Microbiology* **56**, 26-33, doi:10.1111/j.1550-7408.2008.00373.x (2009).
- 36 Torruella, G. *et al.* Global transcriptome analysis of the aphelid *Paraphelidium tribonemae* supports the phagotrophic origin of fungi. *Communications Biology* **1**, doi:10.1038/s42003-018-0235-z (2018).
- Aphelids are on the fungal lineage and they exemplify how an ancestral species may have transitioned from phagotrophy to osmotrophy because they retain genes related to both of these alternative nutritional modes.**
- 37 Held, A. A. Development of *Rozella* in *Allomyces* - a single zoospore produced numerous zoosporangia and resistant sporangia. *Canadian Journal of Botany* **58**, 959-979 (1980).
- 38 Powell, M. J. & Letcher, P. M. Ultrastructure of early stages of *Rozella allomycis* (Cryptomycota) infection of its host, *Allomyces macrogynus* (Blastocladiomycota). *Fungal Biology* **123**, 109-116, doi:10.1016/j.funbio.2018.11.009 (2019).
- 39 Quandt, C. A. *et al.* The genome of an intranuclear parasite, *Paramicrosporidium saccamoebae*, reveals alternative adaptations to obligate intracellular parasitism. *Elife* **6**, doi:10.7554/eLife.29594 (2017).
- 40 Hedegard, E. D. & Ryde, U. Molecular mechanism of lytic polysaccharide monooxygenases. *Chemical Science* **9**, 3866-3880, doi:10.1039/c8sc00426a (2018).
- 41 James, T. Y. *et al.* Shared signatures of parasitism and phylogenomics unite cryptomycota and microsporidia. *Current Biology* **23**, 1548-1553, doi:10.1016/j.cub.2013.06.057 (2013).

- 42 Murphy, C. L. *et al.* Horizontal gene transfer as an indispensable driver for evolution of Neocallimastigomycota into a distinct gut-dwelling fungal lineage. *Applied and Environmental Microbiology* **85**, doi:10.1128/aem.00988-19 (2019).
- 43 Zhao, M. S., Heinsch, F. A., Nemani, R. R. & Running, S. W. Improvements of the MODIS terrestrial gross and net primary production global data set. *Remote Sensing of Environment* **95**, 164-176, doi:10.1016/j.rse.2004.12.011 (2005).
- 44 Fatichi, S., Manzoni, S., Or, D. & Paschalis, A. A mechanistic model of microbially mediated soil biogeochemical processes: a reality check. *Global Biogeochemical Cycles* **33**, 620-648, doi:10.1029/2018gb006077 (2019).
- 45 Duchesne, L. C. & Larson, D. W. Cellulose and the evolution of plant life. *Bioscience* **39**, 238-241, doi:10.2307/1311160 (1989).
- 46 Janusz, G. *et al.* Lignin degradation: microorganisms, enzymes involved, genomes analysis and evolution. *Fems Microbiology Reviews* **41**, 941-962, doi:10.1093/femsre/fux049 (2017).
- 47 Wyman, C. E. Ethanol from lignocellulosic biomass - technology, economics, and opportunities. *Bioresource Technology* **50**, 3-16, doi:10.1016/0960-8524(94)90214-3 (1994).
- 48 Jeffries, T. W. Biodegradation of lignin-carbohydrate complexes. *Biodegradation* **1**, 163-176, doi:10.1007/bf00058834 (1990).
- 49 Leisola, M., Pastinen, O. & Axe, D. D. Lignin—designed randomness. *BIO-Complexity* **12**, 1-11, doi:10.5048/BIO-C.2012.3.c (2012).
- 50 Popper, Z. A. *et al.* in *Annual Review of Plant Biology, Vol 62* Vol. 62 *Annual Review of Plant Biology* (eds S. S. Merchant, W. R. Briggs, & D. Ort) 567-+ (Annual Reviews, 2011).
- 51 Nobles, D. R. & Brown, R. M. The pivotal role of cyanobacteria in the evolution of cellulose synthases and cellulose synthase-like proteins. *Cellulose* **11**, 437-448, doi:10.1023/B:CELL.0000046339.48003.0e (2004).
- 52 Uhliarikova, I. *et al.* Extracellular biopolymers produced by freshwater cyanobacteria: a screening study. *Chemical Papers* **73**, 771-776, doi:10.1007/s11696-018-0625-1 (2019).
- 53 Avrani, S., Wurtzel, O., Sharon, I., Sorek, R. & Lindell, D. Genomic island variability facilitates *Prochlorococcus*-virus coexistence. *Nature* **474**, 604-608, doi:10.1038/nature10172 (2011).
- 54 Agha, R., Gross, A., Rohrlack, T. & Wolinska, J. Adaptation of a chytrid parasite to its cyanobacterial host is hampered by host intraspecific diversity. *Frontiers in Microbiology* **9**, doi:10.3389/fmicb.2018.00921 (2018).
- 55 Del-Bem, L. E. Xyloglucan evolution and the terrestrialization of green plants. *New Phytologist* **219**, 1150-1153, doi:10.1111/nph.15191 (2018).
- 56 Wang, S. *et al.* Genomes of early-diverging streptophyte algae shed light on plant terrestrialization. *Nature Plants* **6**, 95-106, doi:10.1038/s41477-019-0560-3 (2020).
- 57 Alvarez-Cervantes, J., Diaz-Godinez, G., Mercado-Flores, Y., Gupta, V. K. & Anducho-Reyes, M. A. Phylogenetic analysis of beta-xylanase SRXL1 of *Sporisorium reilianum* and its relationship with families (GH10 and GH11) of Ascomycetes and Basidiomycetes. *Scientific Reports* **6**, doi:10.1038/srep24010 (2016).
- 58 Lange, L., Barrett, K., Pilgaard, B., Gleason, F. & Tsang, A. Enzymes of early-diverging, zoosporic fungi. *Applied Microbiology and Biotechnology* **103**, 6885-6902, doi:10.1007/s00253-019-09983-w (2019).
- 59 Parfrey, L. W., Lahr, D. J. G., Knoll, A. H. & Katz, L. A. Estimating the timing of early eukaryotic diversification with multigene molecular clocks. *Proceedings of the*

- National Academy of Sciences of the United States of America* **108**, 13624-13629, doi:10.1073/pnas.1110633108 (2011).
- 60 Gerrienne, P. *et al.* A Simple Type of Wood in Two Early Devonian Plants. *Science* **333**, 837-837, doi:10.1126/science.1208882 (2011).
- 61 Strullu-Derrien, C. *et al.* The earliest wood and its hydraulic properties documented in c. 407-million-year-old fossils using synchrotron microtomography. *Botanical Journal of the Linnean Society* **175**, 423-437, doi:10.1111/boj.12175 (2014).
- 62 Galindo, L. J. *et al.* Combined cultivation and single-cell approaches to the phylogenomics of nuclearioid amoebae, close relatives of fungi. *Philosophical Transactions of the Royal Society B-Biological Sciences* **374**, doi:10.1098/rstb.2019.0094 (2019).
- 63 Rama, T., Hassett, B. T. & Bubnova, E. Arctic marine fungi: from filaments and flagella to operational taxonomic units and beyond. *Botanica Marina* **60**, 433-452, doi:10.1515/bot-2016-0104 (2017).
- 64 Richards, T. A. *et al.* Molecular diversity and distribution of marine fungi across 130 European environmental samples. *Proceedings of the Royal Society B-Biological Sciences* **282**, doi:10.1098/rspb.2015.2243 (2015).
- 65 Amend, A. *et al.* Fungi in the marine environment: open questions and unsolved problems. *Mbio* **10**, doi:10.1128/mBio.01189-18 (2019).
- Reviews current understanding of marine fungi while raising tantalizing, unanswered questions about their diversity and ecosystem function.**
- 66 Grossart, H. P. *et al.* Fungi in aquatic ecosystems. *Nature Reviews Microbiology* **17**, 339-354, doi:10.1038/s41579-019-0175-8 (2019).
- 67 Hawksworth, D. L. Presidential address 1990 - The fungal dimension of biodiversity: magnitude, significance, and conservation. *Mycological Research* **95**, 641-655 (1991).
- 68 Taylor, D. L. *et al.* A first comprehensive census of fungi in soil reveals both hyperdiversity and fine-scale niche partitioning. *Ecological Monographs* **84**, 3-20, doi:10.1890/12-1693.1 (2014).
- 69 Grossart, H. P., Wurzbacher, C., James, T. Y. & Kagami, M. Discovery of dark matter fungi in aquatic ecosystems demands a reappraisal of the phylogeny and ecology of zoospore fungi. *Fungal Ecology* **19**, 28-38, doi:10.1016/j.funeco.2015.06.004 (2016).
- 70 Knack, J. J. *et al.* Microbiomes of streptophyte algae and bryophytes suggest that a functional suite of microbiota fostered plant colonization of land. *International Journal of Plant Sciences* **176**, 405-420, doi:10.1086/681161 (2015).
- 71 Limpens, E., van Zeijl, A. & Geurts, R. in *Annual Review of Phytopathology, Vol 53* Vol. 53 *Annual Review of Phytopathology* (ed N. K. VanAlfen) 311-334 (2015).
- 72 Nishiyama, T. *et al.* The *Chara* genome: Secondary complexity and implications for plant terrestrialization. *Cell* **174**, 448-464, doi:10.1016/j.cell.2018.06.033 (2018).
- 73 Selosse, M. A. & Le Tacon, F. The land flora: a phototroph-fungus partnership? *Trends in Ecology & Evolution* **13**, 15-20, doi:10.1016/s0169-5347(97)01230-5 (1998).
- 74 Spatafora, J. W. *et al.* A phylum-level classification of zygomycete fungi based on genome-scale data. *Mycologia* **108**, 1028-1046, doi:DOI: 10.3852/ 16-042 (2016).
- 75 Bidartondo, M. I. *et al.* The dawn of symbiosis between plants and fungi. *Biology Letters* **7**, 574-577, doi:10.1098/rsbl.2010.1203 (2011).
- 76 Strullu-Derrien, C. *et al.* Fungal associations in *Horneophyton ligneri* from the Rhynie Chert (c. 407 million year old) closely resemble those in extant lower land

plants: novel insights into ancestral plant-fungus symbioses. *New Phytologist* **203**, 964-979, doi:10.1111/nph.12805 (2014).

Rhynie chert fossils provide the earliest undisputed evidence of extant fungal phyla, as in this paper that describes a 407-year-old land plant colonized by fungi representing two lineages that are still engaged in mycorrhizal symbiosis.

- 77 Rimington, W. R., Pressel, S., Duckett, J. G., Field, K. J. & Bidartondo, M. I. Evolution and networks in ancient and widespread symbioses between Mucoromycotina and liverworts. *Mycorrhiza* **29**, 551-565, doi:<https://doi.org/10.1007/s00572-019-00918-x> (2019).
- 78 Desiro, A., Duckett, J. G., Pressel, S., Villarreal, J. C. & Bidartondo, M. I. Fungal symbioses in hornworts: a chequered history. *Proceedings of the Royal Society B-Biological Sciences* **280**, doi:10.1098/rspb.2013.0207 (2013).
- 79 Field, K. J. *et al.* First evidence of mutualism between ancient plant lineages (Haplomitriopsida liverworts) and Mucoromycotina fungi and its response to simulated Palaeozoic changes in atmospheric CO₂. *New Phytologist* **205**, 743-756, doi:10.1111/nph.13024 (2015).
- 80 Feijen, F. A. A., Vos, R. A., Nuytinck, J. & Merckx, V. Evolutionary dynamics of mycorrhizal symbiosis in land plant diversification. *Scientific Reports* **8**, doi:10.1038/s41598-018-28920-x (2018).
- 81 Hoysted, G. A. *et al.* Mucoromycotina fine root endophyte fungi form nutritional mutualisms with vascular plants. *Plant Physiology* **181**, 565-577, doi:10.1104/pp.19.00729 (2019).
- 82 Field, K. J. *et al.* Functional analysis of liverworts in dual symbiosis with Glomeromycota and Mucoromycotina fungi under a simulated Palaeozoic CO₂ decline. *Isme Journal* **10**, 1514-1526, doi:10.1038/ismej.2015.204 (2016).
- 83 Kenrick, P., Wellman, C. H., Schneider, H. & Edgecombe, G. D. A timeline for terrestrialization: consequences for the carbon cycle in the Palaeozoic. *Philosophical Transactions of the Royal Society B-Biological Sciences* **367**, 519-536, doi:10.1098/rstb.2011.0271 (2012).
- 84 Tomescu, A., Klymiuk, A., Matsunaga, K., Bippus, A. & Shelton, G. in *Their World: A Diversity of Microbial Environments* Vol. 1 (ed Christon J. Hurst) 69-169 (Springer, 2016).
- 85 Knoll, A. in *Fundamentals of Geobiology* (eds AH Knoll, DE Canfield, & K Konhauser) 297-314 (Wiley-Blackwell, 2012).
- 86 Knoll, A. H. Paleobiological perspectives on early eukaryotic evolution. *Cold Spring Harbor Perspectives in Biology* **6**, doi:10.1101/cshperspect.a016121 (2014).
- 87 Knoll, A. H., Javaux, E. J., Hewitt, D. & Cohen, P. Eukaryotic organisms in Proterozoic oceans. *Philosophical Transactions of the Royal Society B-Biological Sciences* **361**, 1023-1038, doi:10.1098/rstb.2006.1843 (2006).
- 88 Tang, Q., Pang, K., Yuan, X. & Xiao, S. A one-billion-year-old multicellular chlorophyte. *Nature Ecology & Evolution*, doi:10.1038/s41559-020-1122-9 (2020).
- 89 Strother, P. K. & Wellman, C. H. Palaeoecology of a billion-year-old non-marine cyanobacterium from the Torridon Group and Nonesuch Formation. *Palaeontology* **59**, 89-108, doi:10.1111/pala.12212 (2016).
- 90 Strother, P. K., Battison, L., Brasier, M. D. & Wellman, C. H. Earth's earliest non-marine eukaryotes. *Nature* **473**, 505-509, doi:10.1038/nature09943 (2011).

- 948 91 Lozano-Fernandez, J. *et al.* A molecular palaeobiological exploration of arthropod
949 terrestrialization. *Philosophical Transactions of the Royal Society B-Biological*
950 *Sciences* **371**, doi:10.1098/rstb.2015.0133 (2016).
- 951 92 Pirozynski, K. A. Interactions between fungi and plants through the ages. *Canadian*
952 *Journal of Botany-Revue Canadienne De Botanique* **59**, 1824-1827, doi:10.1139/b81-
953 243 (1981).
- 954 93 Baldwin, C. T., Strother, P. K., Beck, J. H. & Rose, E. Palaeoecology of the Bright
955 Angel Shale in the eastern Grand Canyon, Arizona, USA, incorporating
956 sedimentological, ichnological and palynological data. *Geological Society, London,*
957 *Special Publications* **228**, 213, doi:10.1144/GSL.SP.2004.228.01.11 (2004).
- 958 94 Tomescu, A. M. F., Pratt, L. M., Rothwell, G. W., Strother, P. K. & Nadon, G. C.
959 Carbon isotopes support the presence of extensive land floras pre-dating the origin of
960 vascular plants. *Palaeogeography Palaeoclimatology Palaeoecology* **283**, 46-59,
961 doi:10.1016/j.palaeo.2009.09.002 (2009).
- 962 95 Edwards, D., Morris, J. L., Richardson, J. B. & Kenrick, P. Cryptospores and
963 cryptophytes reveal hidden diversity in early land floras. *New Phytologist* **202**, 50-78,
964 doi:10.1111/nph.12645 (2014).
- 965 96 Krause, A. J. *et al.* Stepwise oxygenation of the Paleozoic atmosphere. *Nature*
966 *Communications* **9**, doi:10.1038/s41467-018-06383-y (2018).
- 967 97 Loron, C. C., Rainbird, R. H., Turner, E. C., Greenman, J. W. & Javaux, E. J.
968 Organic-walled microfossils from the late Mesoproterozoic to early Neoproterozoic
969 lower Shaler Supergroup (Arctic Canada): Diversity and biostratigraphic significance.
970 *Precambrian Research* **321**, 349-374, doi:10.1016/j.precamres.2018.12.024 (2019).
- 971 98 Ehrlich, H. *et al.* Discovery of 505-million-year old chitin in the basal demosponge
972 *Vauxia gracilenta*. *Scientific Reports* **3**, doi:10.1038/srep03497 (2013).
- 973 99 Francois, C. *et al.* Contributions of U-Th-Pb dating on the diagenesis and sediment
974 sources of the lower group (BI) of the Mbuji-Mayi Supergroup (Democratic Republic
975 of Congo). *Precambrian Research* **298**, 202-219,
976 doi:10.1016/j.precamres.2017.06.012 (2017).
- 977 100 Redecker, D., Kodner, R. & Graham, L. E. Glomalean fungi from the Ordovician.
978 *Science* **289**, 1920-1921 (2000).
- 979 101 Sherwood-Pike, M. A. & Gray, J. Silurian fungal remains: probable records of the
980 Class Ascomycetes. *Lethaia* **18**, 1-20 (1985).
- 981 102 Smith, M. R. Cord-forming Palaeozoic fungi in terrestrial assemblages. *Botanical*
982 *Journal of the Linnean Society* **180**, 452-460, doi:10.1111/boj.12389 (2016).
- 983 103 Kenrick, P. & Crane, P. R. The origin and early evolution of plants on land. *Nature*
984 **389**, 33-39, doi:10.1038/37918 (1997).
- 985 104 Edwards, D., Kenrick, P. & Dolan, L. History and contemporary significance of the
986 Rhynie cherts-our earliest preserved terrestrial ecosystem. *Philosophical Transactions*
987 *of the Royal Society B-Biological Sciences* **373**, doi:10.1098/rstb.2016.0489 (2018).
- 988 105 Strullu-Derrien, C., Kenrick, P., Goral, T. & Knoll, A. H. Testate Amoebae in the
989 407-Million-Year-Old Rhynie Chert. *Current Biology* **29**, 461-+,
990 doi:10.1016/j.cub.2018.12.009 (2019).
- 991 106 Krings, M., Harper, C. J. & Taylor, E. L. Fungi and fungal interactions in the Rhynie
992 chert: a review of the evidence, with the description of *Perexiflasca tayloriana* gen. et
993 sp nov. *Philosophical Transactions of the Royal Society B-Biological Sciences* **373**,
994 doi:10.1098/rstb.2016.0500 (2018).
- 995 107 Krings, M., Taylor, T. N. & Dotzler, N. Fossil evidence of the zygomycetous fungi.
996 *Persoonia* **30**, 1-10, doi:10.3767/003158513x664819 (2013).

997 108 Strullu-Derrien, C. *et al.* A new chytridiomycete fungus intermixed with crustacean
998 resting eggs in a 407-million-year-old continental freshwater environment. *PLoS One*
999 **11**, doi:10.1371/journal.pone.0167301 (2016).

1000 109 Taylor, T. N. *et al.* Fungi from the Rhynie chert: a view from the dark side.
1001 *Transactions of the Royal Society of Edinburgh - Earth Sciences* **94**, 457-473 (2004).

1002 110 Krings, M. *et al.* Fungal endophytes in a 400-million-yr-old land plant: infection
1003 pathways, spatial distribution, and host responses. *New Phytologist* **174**, 648-657,
1004 doi:10.1111/j.1469-8137.2007.02008.x (2007).

1005 111 Taylor, T. N., Hass, H. & Remy, W. Devonian Fungi: Interactions with the green
1006 Alga *Palaeonitella*. *Mycologia* **84**, 901-910 (1992).

1007 112 Remy, W., Taylor, T. N., Hass, H. & Kerp, H. Four hundred-million-year-old
1008 vesicular arbuscular mycorrhizae. *Proceedings of the National Academy of Sciences*
1009 *USA* **91**, 11841-11843 (1994).

1010 113 Taylor, T. N., Remy, W., Hass, H. & Kerp, H. Fossil arbuscular mycorrhizae from the
1011 early Devonian. *Mycologia* **87**, 560-573 (1995).

1012 114 Strullu-Derrien, C. & Strullu, D. G. Mycorrhization of fossil and living plants.
1013 *Comptes Rendus Palevol* **6**, 483-494, doi:10.1016/j.crpv.2007.09.006 (2007).

1014 115 Taylor, T. N., Hass, H. & Kerp, H. The oldest fossil ascomycetes. *Nature* **399**, 648,
1015 doi:10.1038/21349 (1999).

1016 116 Taylor, T. N., Hass, H., Kerp, H., Krings, M. & Hanlin, R. T. Perithecial ascomycetes
1017 from the 400 million year old Rhynie chert: an example of ancestral polymorphism
1018 (vol 96, pg 1403, 2004). *Mycologia* **97**, 269-285 (2005).

1019 117 Taylor, J. W. & Berbee, M. L. Dating divergences in the Fungal Tree of Life: review
1020 and new analyses. *Mycologia* **98**, 838-849 (2006).

1021 118 Lücking, R., Huhndorf, S., Pfister, D. H., Plata, E. R. & Lumbsch, H. T. Fungi
1022 evolved right on track. *Mycologia* **101**, 810-822, doi:10.3852/09-016 (2009).

1023 119 Kalgutkar, R. M. & Jansonius, J. Synopsis of fossil fungal spores, mycelia and
1024 fructifications. *American Association of Stratigraphic Palynologists Contributions*
1025 *Series* **39**, 1-423 (2000).

1026 120 Pirozynski, K. A. Fossil fungi. *Annual Review of Phytopathology* **14**, 237-246 (1976).

1027 121 Honegger, R., Edwards, D., Axe, L. & Strullu-Derrien, C. Fertile *Prototaxites taiti*: a
1028 basal ascomycete with inoperculate, polysporous asci lacking croziers. *Philosophical*
1029 *Transactions of the Royal Society B-Biological Sciences* **373**,
1030 doi:10.1098/rstb.2017.0146 (2018).

1031 122 Hueber, F. M. Rotted wood--alga--fungus: the history and life of *Prototaxites*
1032 Dawson 1859. *Review of Palaeobotany and Palynology* **116**, 123-158 (2001).

1033 123 Hobbie, E. A. & Boyce, C. K. Carbon sources for the Palaeozoic giant fungus
1034 *Prototaxites* inferred from modern analogues. *Proceedings of the Royal Society B-*
1035 *Biological Sciences* **277**, 2149-2156, doi:10.1098/rspb.2010.0201 (2010).

1036 124 Selosse, M. A. *Prototaxites*: A 400 Myr old giant fossil, a saprophytic
1037 holobasidiomycete, or a lichen? *Mycological Research* **106**, 642-644,
1038 doi:10.1017/s0953756202226313 (2002).

1039 125 Agic, H. *et al.* Organically-preserved multicellular eukaryote from the early Ediacaran
1040 Nyborg Formation, Arctic Norway. *Scientific Reports* **9**, doi:10.1038/s41598-019-
1041 50650-x (2019).

1042 126 Lücking, R. & Nelsen, M. in *Transformative Paleobotany* (eds Michael Krings, C. J.
1043 Harper, N. R. Cúneo, & G. W. Rothwell) 551-590 (Academic Press, 2018).

1044 127 Retallack, G. J. Ediacaran life on land. *Nature* **493**, 89-92, doi:10.1038/nature11777
1045 (2013).

1046 128 Peterson, K. J., Waggoner, B. & Hagadorn, J. W. A fungal analog for Newfoundland
1047 Ediacaran fossils? *Integrative and Comparative Biology* **43**, 127-136,
1048 doi:10.1093/icb/43.1.127 (2003).

1049 129 Bobrovskiy, I. *et al.* Ancient steroids establish the Ediacaran fossil *Dickinsonia* as one
1050 of the earliest animals. *Science* **361**, 1246-+, doi:10.1126/science.aat7228 (2018).

1051 130 Yuan, X. L., Xiao, S. H. & Taylor, T. N. Lichen-like symbiosis 600 million years ago.
1052 *Science* **308**, 1017-1020 (2005).

1053 131 Karatygin, I. V., Snigirevskaya, N. S. & Vikulin, S. V. The most ancient terrestrial
1054 lichen *Winfrenatia reticulata*: A new find and new interpretation. *Paleontological*
1055 *Journal* **43**, 107-114, doi:10.1134/s0031030109010110 (2009).

1056 132 Taylor, T. N., Hass, H. & Kerp, H. A cyanolichen from the Lower Devonian Rhynie
1057 chert. *American Journal of Botany* **84**, 992-1004 (1997).

1058 133 Taylor, T. N., Hass, H., Remy, W. & Kerp, H. The oldest fossil lichen. *Nature* **378**,
1059 244-244, doi:10.1038/378244a0 (1995).

1060 134 Honegger, R., Edwards, D. & Axe, L. The earliest records of internally stratified
1061 cyanobacterial and algal lichens from the Lower Devonian of the Welsh Borderland.
1062 *New Phytologist* **197**, 264-275, doi:10.1111/nph.12009 (2013).

1063 135 Honegger, R., Axe, L. & Edwards, D. Bacterial epibionts and endolichenic
1064 actinobacteria and fungi in the Lower Devonian lichen *Chlorolichenomycites*
1065 *salopensis*. *Fungal Biology* **117**, 512-518, doi:10.1016/j.funbio.2013.05.003 (2013).

1066 136 Lewis, L. A. & McCourt, R. M. Green algae and the origin of land plants. *American*
1067 *Journal of Botany* **91**, 1535-1556, doi:10.3732/ajb.91.10.1535 (2004).

1068 137 Joneson, S., Stajich, J. E., Shiu, S. H. & Rosenblum, E. B. Genomic transition to
1069 pathogenicity in chytrid fungi. *Plos Pathogens* **7**, doi:10.1371/journal.ppat.1002338
1070 (2011).

1071 138 Gewin, V. Riders of a modern-day Ark. *Plos Biology* **6**, 18-21,
1072 doi:10.1371/journal.pbio.0060024 (2008).

1073 139 Miao, L. Y., Moczyłowska, M., Zhu, S. X. & Zhu, M. Y. New record of organic-
1074 walled, morphologically distinct microfossils from the late Paleoproterozoic
1075 Changcheng Group in the Yanshan Range, North China. *Precambrian Research* **321**,
1076 172-198, doi:10.1016/j.precamres.2018.11.019 (2019).

1077 140 Butterfield, N. J. *Bangiomorpha pubescens* n. gen., n. sp.: implications for the
1078 evolution of sex, multicellularity, and the Mesoproterozoic/Neoproterozoic radiation
1079 of eukaryotes. *Paleobiology* **26**, 386-404, doi:10.1666/0094-
1080 8373(2000)026<0386:bpngns>2.0.co;2 (2000).

1081 141 Redecker, D. New views on fungal evolution based on DNA markers and the fossil
1082 record. *Research in Microbiology* **153**, 125-130 (2002).

1083 142 Rubinstein, C. V., Gerrienne, P., de la Puente, G. S., Astini, R. A. & Steemans, P.
1084 Early Middle Ordovician evidence for land plants in Argentina (eastern Gondwana).
1085 *New Phytol* **188**, 365-369, doi:10.1111/j.1469-8137.2010.03433.x (2010).

1086 143 Johnson, N. G. Early Silurian palynomorphs from the Tuscarora Formation in central
1087 Pennsylvania and their paleobotanical and geological significance. *Review of*
1088 *Palaeobotany and Palynology* **45**, 307-360, doi:10.1016/0034-6667(85)90006-5
1089 (1985).

1090 144 Wolfe, J. M., Daley, A. C., Legg, D. A. & Edgecombe, G. D. Fossil calibrations for
1091 the arthropod Tree of Life. *Earth-Science Reviews* **160**, 43-110,
1092 doi:10.1016/j.earscirev.2016.06.008 (2016).

1093 145 Peng, Y. B., Bao, H. M. & Yuan, X. L. New morphological observations for
1094 Paleoproterozoic acritarchs from the Chuanlinggou Formation, North China.
1095 *Precambrian Research* **168**, 223-232, doi:10.1016/j.precamres.2008.10.005 (2009).

- 1096 146 Pirie, M. D. & Doyle, J. A. Dating clades with fossils and molecules: the case of
1097 Annonaceae. *Botanical Journal of the Linnean Society* **169**, 84-116,
1098 doi:10.1111/j.1095-8339.2012.01234.x (2012).
- 1099 147 Shi, C. S., Schopf, J. W. & Kudryavtsev, A. B. Characterization of the stem anatomy
1100 of the eocene fern *Dennstaedtiopsis aerenchymata* (Dennstaedtiaceae) by use of
1101 confocal laser scanning microscopy. *American Journal of Botany* **100**, 1626-1640,
1102 doi:10.3732/ajb.1300027 (2013).
- 1103 148 Strullu-Derrien, C., Wawrzyniak, Z., Goral, T. & Kenrick, P. Fungal colonization of
1104 the rooting system of the early land plant *Asteroxylon mackiei* from the 407-Myr-old
1105 Rhynie chert (Scotland, UK). *Botanical Journal of the Linnean Society* **179**, 201-213,
1106 doi:10.1111/boj.12307 (2015).
- 1107 149 Saint Martin, J. P. & Saint Martin, S. Exquisite preservation of a widespread
1108 filamentous microorganism in French Cretaceous ambers: Crucial for revising a
1109 controversial fossil. *Comptes Rendus Palevol* **17**, 415-434,
1110 doi:10.1016/j.crpv.2017.05.003 (2018).
- 1111 150 Baludikay, B. K. *et al.* Raman microspectroscopy, bitumen reflectance and illite
1112 crystallinity scale: comparison of different geothermometry methods on fossiliferous
1113 Proterozoic sedimentary basins (DR Congo, Mauritania and Australia). *International*
1114 *Journal of Coal Geology* **191**, 80-94, doi:10.1016/j.coal.2018.03.007 (2018).
- 1115 151 Marshall, C. P., Javaux, E. J., Knoll, A. H. & Walter, M. R. Combined micro-Fourier
1116 transform infrared (FTIR) spectroscopy and micro-Raman spectroscopy of
1117 Proterozoic acritarchs: A new approach to Palaeobiology. *Precambrian Research* **138**,
1118 208-224, doi:10.1016/j.precamres.2005.05.006 (2005).

1119
1120