Lipidome variations of deep-sea vent shrimps according to acclimation pressure: A homeoviscous response?

B. Shillito, J. Ravaux, M. Zbinden, C. Desurmont, D. Barthélémy, D. Farabos, G. Després, A. Lamazière

► To cite this version:

B. Shillito, J. Ravaux, M. Zbinden, C. Desurmont, D. Barthélémy, et al.. Lipidome variations of deep-sea vent shrimps according to acclimation pressure: A homeoviscous response?. Deep Sea Research Part I: Oceanographic Research Papers, 2020, 161, pp.103285. 10.1016/j.dsr.2020.103285 . hal-03330586

HAL Id: hal-03330586

<https://hal.science/hal-03330586>

Submitted on 3 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Lipidome variations of deep-sea vent shrimps according to acclimation pressure : a homeoviscous response ?

Shillito, B. ^{1*}, Désurmont, C. ^{1,2}, Barthélémy, D. ³, Farabos, D. ², Despres, G. ², Ravaux, J. ¹,
Zbinden, M. ¹, Lamazière, A. ²

¹ Laboratoire de Biologie des Organismes et Ecosystèmes Aquatiques (BOREA), MNHN, CNRS-2030,
IRD-207, Sorbonne Université, UCN, UA

² Sorbonne Université, INSERM, Centre de Recherche St-Antoine, CRSA, Département de Métabolomique
Clinique, Hôpital Saint Antoine, AP-HP, F-75012 Paris France

³ Océanopolis, Port de Plaisance du Moulin Blanc BP 91039, 29210 Brest Cedex1, France

ABSTRACT

The present study questions the ability of marine fauna to modulate the molecular composition of their membrane lipids, as a function of environmental hydrostatic pressure conditions. Specific variations in the cell lipidome composition are presented for the deep-sea hydrothermal vent shrimp *Mirocaris fortunata*, acclimated at the laboratory for several months at both natural (18 MPa) and atmospheric pressures, but fed with the same diet. These animals display physiological capabilities of acclimation to pressure variations, possibly driven by a particular composition of their lipid species. In particular, it is shown that shrimps exposed to atmospheric pressure exhibit a significantly lower level of the mono-unsaturated fatty acid, vaccenic acid (C18:1n-7), compared to those maintained at natural pressure. The observed fatty acid variations are consistent with a homeoviscous response, i.e. a modulation of the lipidome composition, in response to physical constraints, in order to balance the effects of pressure on membrane order. Additionally, supervised multivariate data analysis was performed for complex lipids and revealed an increase in levels of sterols, sphingomyelin, and ether-phosphatidylethanolamine lipids for animals exposed to atmospheric pressure, with respect to natural pressure. These observations strongly suggest that lipidic domains are involved in the shrimp's response to experimental conditions, and their consistency with a homeoviscous response to pressure is discussed.

30 **KEY WORDS** : Hydrothermal vent shrimps, Pressure tolerance, Viscosity, Homeostasis,
31 Homeoviscous, Lipid signature, Lipidomes, AbyssBox
32

33 **(*) : Corresponding author**

34 e-mail : Bruce.Shillito@upmc.fr

35 postal address : B. Shillito, équipe AMEX-BOREA, Sorbonne Université,
36 7 Quai Saint-Bernard, Bâtiment A, 4ème étage, Paris 75005, FRANCE
37

38 **Competing interests statement** : The authors declare that they have no known competing financial
39 interests or personal relationships that could have appeared to influence the work reported in this
40 paper.
41

42 **Contributions of authors** : BS designed the acclimation experiments, BS CD and AL wrote the
43 manuscript, which was further reviewed by all authors. MZ JR DB supervised in situ sampling,
44 transport, and maintenance of deep-sea shrimps, JR and MZ undertook conditioning of samples at
45 the end of the acclimation period. DF GD AL undertook the determination of lipidomes by GC-MS,
46 and CD AL carried out statistical analyses of data.

INTRODUCTION

Marine depths exceeding 1000 m occupy about 75% of the global ocean volume. The deep sea is therefore probably the largest biotope on earth (Fang et al., 2010), and discoveries in the last decades have brought the picture of a high biodiversity environment (Ramirez-Llodra et al., 2010). Along with these findings, another emerging view is that the deep sea has been the subject of successive colonisation events by shallow water species, spread over geological times (reviews in Priede & Froese, 2013, Brown & Thatje, 2014), and the opposite phenomenon (from deep to shallow waters) is also likely (Barnes & Kuklinsky, 2010, Brown & Thatje, 2014). The pace at which such migratory events occurred remains unknown : is it a "slow" phenomenon submitted to evolutionary speciation, or a "fast" one, driven by the migration of individual specimens capable of acclimatisation to new environmental conditions ?

The existence of a depth zonation in the distribution of extant species, possibly explained by a genetically constrained physiological tolerance regarding hydrostatic pressure variations (Carney 2005, Brown & Thatje 2014) would support the first hypothesis. Hydrostatic pressure in the marine biosphere increases linearly with depth, ranging from 0.1 MPa at sea level (i.e. the so-called "atmospheric pressure"), to about 110 MPa in the deepest trenches, at a depth of about 11000m (Jamieson et al., 2010). Several studies have long documented severe impacts on behavioral and motor activity, and survival, either during pressurization experiments on shallow-water organisms (review in Pradillon 2012, Ammendolia et al., 2018), or by observations of decompressed deep-sea animals following their recovery at depth (reviews in MacDonald, 1997, Somero 1992). Furthermore, *in vitro* experiments have also established that pressure variations in the 10-50 MPa range impact on many aspects of biological systems, such as enzyme function, or trans-membrane signaling (reviews in Somero 1992, McDonald, 1997, Pradillon 2012). So, although other environmental parameters may vary between habitats at different depths (i.e. light intensity, temperature, food availability, oxygen levels, etc.. (Ramirez-Llodra et al., 2010)), and thereby possibly control the distribution of a species (Brown & Thatje, 2015), pressure alone would seem sufficient to restrict the occurrence of a species within a given depth range.

Some studies have opened the way to another (non-exclusive) hypothesis: direct colonisation by specimens capable of acclimatisation to new environmental conditions. Some *in situ* studies have reported spectacular bathymetric migrations, such as that of antarctic krill (*E. superba*, from the surface to a depth of beyond 3000 m, Clarke & Tyler, 2008), or Mediterranean red shrimp (*Aristeus antennatus*, depth from 80 m to 3300 m, Sarda et al., 2004). More recently, laboratory studies have shown that some shallow-water invertebrates could survive for several days at

81 pressures far beyond those encountered in the wild: the ditch shrimp *Palaemonetes varians*,
82 exclusively reported in depths of less than 1m, presented survival of 70% after 28 days at a
83 simulated depth of 1000 m (10 MPa, [Cottin et al., 2012](#)). Finally, the deep-sea hydrothermal vent
84 shrimp *Mirocaris fortunata*, the subject of the present study, not only displays an important
85 distribution of depth (from 850 to 4050 m, [Desbruyères et al., 2006](#), [Fouquet et al., 2008](#)), but has
86 also been successfully acclimated for several months at atmospheric pressure ([Shillito et al., 2015](#)),
87 far out of its reported natural bathymetric range. It has been argued that the potential to undertake
88 bathymetric migration may be species-specific, suggesting there may be "winners and losers" in the
89 face of near-future climate-driven migration patterns ([Ammendolia et al., 2018](#), [Brown & Thatje,](#)
90 [2015](#)). In other words, some species would be able to display physiological plasticity in response to
91 pressure variation (i.e. capable of acclimatisation), while others would not. Comparison between
92 deep and shallow species point at a number of biological features which respond to pressure
93 conditions, such as protein functioning ([Pradillon, 2012](#)) or osmolyte levels ([Yancey et al, 2013](#)),
94 but most importantly, the molecular composition of biological membranes has long been identified
95 as a highly pressure-sensitive feature (review in [Hazel & Williams, 1990](#)).

96 Biological membranes are a complex and dynamic assembly of lipids and proteins, and as
97 for temperature, variations in hydrostatic pressure can modify the conformation and packing of the
98 molecules composing these liquid-state lipidic bilayers, thereby impacting their molecular order.
99 However, pressure and temperature effects appear to be in opposition: an increase in pressure tends
100 to increase membrane order (i.e. "rigidifies" the membrane), while an increase in temperature
101 lowers membrane order (i.e. "fluidizes" the membrane). Living organisms respond to such
102 perturbations by modulating the molecular composition of their membranes, in order to oppose to
103 the deleterious (ordering or disordering) effects of the environmental change, thereby allowing to
104 maintain membrane order, and therefore cellular homeostasis ([Hazel, 1995](#), [Ernst et al., 2016](#)). This
105 response is termed "homeoviscous", and was first observed regarding thermal variations in bacteria
106 ([Sinensky, 1974](#)). Today it is widely accepted that all poikilothermic organisms (which do not
107 control their temperature) may undertake such a response in the face of thermal variations,
108 including within the lifespan of a given individual organism, i.e. as an acclimatory response ([Parent](#)
109 [et al., 2008](#), for example). The variations in membrane compositions of deep-sea animals submitted
110 to different pressures were first evidenced by comparing different fish species originating from
111 various depths ([Cossins & MacDonald, 1986](#), [Behan et al., 1992](#)). An important study also showed
112 that a given bacterial strain would modify its membrane composition within a few days, according
113 to incubation pressure ([Delong & Yayanos, 1985](#)). Since then, much of our knowledge on the
114 homeoviscous response to pressure has come from studies of bacteria ([Wirsén et al., 1987](#), [Allen et](#)
115 [al., 1999](#), [Grossi et al., 2010](#), [Fumiyoshi, 2013](#), [Wang et al., 2014](#), [Cario et al., 2015](#)), which are

116 more amenable to pressure incubation than metazoans, mainly because of a limitation in pressure
117 equipments suitable for large organisms. Therefore, our knowledge of the homeoviscous response
118 to pressure in animals remains limited. One of the major remaining questions is whether the above-
119 mentioned "winners", i.e. species displaying acclimatisation capacities in response to pressure
120 variation, are capable of a such a response.

121 Studies of the homeoviscous response to pressure have so far focused mainly on the study of
122 the relative proportions of saturated fatty acids (SFA), mono-unsaturated fatty acids (MUFA), and
123 poly-unsaturated fatty acids (PUFA) which compose the hydrophobic fraction of membrane lipids.
124 Schematically, high levels of SFA increase membrane order, while the presence of one or several
125 double bonds in MUFA and PUFA lead to disordered assemblies ([Guschina & Harwood, 2006](#)).
126 Therefore, organisms at high pressure are expected to display higher levels of unsaturated fatty
127 acids and lower levels of SFA, thus counteracting the ordering effect of pressure, and a reverse
128 situation is expected for exposure at low pressure. However, given the complexity of eucaryotic
129 biomembranes, composed of several hundreds of different molecules, an emerging view is that the
130 homeoviscous response is probably more complex than described up to now, and it is not only a
131 matter of acyl chain composition. This view is supported by the growing outcome of contemporary
132 quantitative lipidomics technology, which may allow to follow complex lipid changes during the
133 homeoviscous response (review in [Ernst et al., 2016](#)). Finally, our understanding of the genetic and
134 enzymatic regulation of the lipidomic response to pressure remains partial, and again owes much to
135 studies of bacteria or yeast model organisms ([Fernandes et al., 2004](#), [Freitas et al., 2012](#)). More
136 generally, the way by which membrane homeostasis is achieved remains unclear ([Ernst et al.,](#)
137 [2018](#)).

138 We here present results regarding the lipidomic compositions of the deep-sea shrimp
139 *Mirocaris fortunata*, successfully acclimated at the public aquarium of Océanopolis at different
140 pressure regimes (as opposed to "acclimated" when occurring in the wild), in the context of a
141 permanent public exhibition. Because of the importance in pressure variation between our
142 experimental groups (18 MPa vs. atmospheric pressure), and because of the duration of the
143 incubation in a controlled environment (several months, an exceptional situation regarding deep-sea
144 fauna), our hypothesis was that potential lipidomic signature responses would be detectable, and
145 that the additional use of high output lipidomics would help address the following questions :

- 146 1 - Are the observed successful acclimations accompanied by variations in lipidome composition,
147 and if so, do these variations compare with homeoviscous responses documented in the literature?
148 2 - Beyond revealing classic characteristics of a homeoviscous response, could the complex lipid
149 analysis bring original insights regarding the response to hydrostatic pressure?

150 MATERIAL AND METHODS

151 Biological model

152 This study focuses on the hydrothermal vent shrimp *Mirocaris fortunata* (Martin &
153 Christiansen, 1995). This species from the Alvinocarididae family is endemic to the Mid-Atlantic
154 Ridge (MAR), where it can be encountered in proximity to vent openings, at temperatures around
155 10 °C (Desbruyères et al., 2001, Desbruyères et al., 2006, Matabos et al., 2015). Believed to be an
156 omnivorous shrimp (Colaço et al., 2007), *M. fortunata* is found in great abundance on
157 *Bathymodiolus* mussel beds, making its collection relatively easy. This species has a wide
158 bathymetric distribution (850 m to 4050 m depth, Desbruyères et al., 2006, Fouquet et al., 2008),
159 thereby providing a good model to investigate pressure-driven homeoviscous adaptations of deep-
160 sea organisms (Shillito et al., 2006). Moreover, several specimens originating from depths of 1700-
161 1800 m have already been successfully acclimated in captivity for several years, both at *in situ* and
162 atmospheric pressure at the Océanopolis public aquarium (Brest, France), (Shillito et al., 2015).
163 These captive animals are the subjects of this study.

164

165 Specimen collection, transport and maintenance

166 Sampling of animals took place during two cruises, in July 2012 and 2013, respectively
167 (Momarsat, 2012; Biobaz, 2013), on board the oceanographic vessels "Thalassa" (Momarsat, 2012)
168 and "Pourquoi Pas?" (Biobaz, 2013) using the Remotely Operated Vehicle (ROV) Victor6000
169 (IFREMER) at the Lucky Strike vent field (MAR, 37°17'6N, ~ 1750 m depth). Specimens were
170 sampled with a suction device operated by the hydraulic arm of the ROV. Immediately after
171 recovery onboard the ship, the shrimps were stored at atmospheric pressure in tanks of
172 approximately 5 – 10 l of aerated seawater in the cold room (5 – 9 °C), in groups of few individuals
173 (< 5). At the end of the cruises, the shrimps were shipped to the Océanopolis public aquarium (air
174 freight by "Flying Sharks", Lisbon, Portugal).

175 On arrival at Océanopolis, husbandry of the shrimps was performed by staff members of the
176 aquarium. Most of the shrimps were acclimated at atmospheric pressure in a dark room (10 °C) in
177 groups of around 50 *M. fortunata* in ~ 80 l tanks each equipped with a 25W heating element
178 wrapped in PVC foam. This heater provides a localised “hotspot” with respect to the surrounding 10
179 °C environment, in the 13-28 °C range, in order to provide a thermal gradient mimicking their
180 natural environment. Lights were switched on only a few minutes per day for cleaning, feeding and
181 mortality counts. For the purposes of public exhibition, other individuals were maintained in a 16.5
182 l pressurized aquarium named “AbyssBox” (described in Shillito et al., 2015), at a pressure of 18

183 MPa (thereby simulating the pressure prevailing in the natural environment of these shrimps). The
184 background temperature was set to 10 °C by a temperature-regulation unit which cooled both the in-
185 coming water (flow rate of 10 l/hour) and the experimental vessel. As for atmospheric-pressure
186 maintenance, a localized hotspot was provided for the shrimps, consisting of a ring-shaped tube
187 heated to 27 °C, wrapped in PVC foam. Because the pressurized shrimps were exposed to the
188 public, they were submitted to about eleven hours of dim artificial light per day (see [Shillito et al.,](#)
189 [2015](#) and [Matabos et al., 2015](#) for further details on shrimp maintenance and behaviour).

190

191 **Experimental design**

192 A first group (the BOX group) of five *M. fortunata* was maintained in the AbyssBox at a
193 pressure of 18 MPa. The second group (the AP group) consisted of five *M. fortunata* maintained at
194 atmospheric pressure. During the experiment which lasted five months, both groups were
195 exclusively fed with Liptoqua food pellets (Liptosa, Madrid, Spain) every 4 – 5 days. These pellets
196 were inert food (as opposed to live diet), apparently obtained from dried organic extracts, and their
197 composition was not provided by the manufacturer. Shrimps in the AbyssBox were fed without
198 decompression by an isobaric feeding line (see [Shillito et al., 2015](#)). At the end of the experiment,
199 the shrimps (2 - 3 cm body length) were collected and dissected (immediately upon pressure release
200 for the AbyssBox group). Abdominal muscle tissues and food pellets were subsequently frozen in
201 liquid nitrogen.

202

203 **Lipid analysis**

204 ***Fatty Acid Methyl Ester Analysis***

205 The frozen samples (muscle tissues and food pellets) were first homogenized using a tissue
206 lyser apparatus (Quiagen, Les Ulis, France). A micro-method for total lipid extraction and
207 methylation was applied from 20 µL serum. Fatty acid methyl esters (FAME) were prepared and
208 assayed by GC-MS in the positive chemical ionization mode with ammonia as the reagent gas
209 (GC6890-MS5975; Agilent Technologies, Les Ulis, France) as described previously ([Wolf &](#)
210 [Quinn, 2008](#)). The response factors of fatty acids were calibrated with a weighed mixture (FAME
211 Mix Supelco® 37, Sigma-Aldrich Chimie, L'Isle d'Abeau Chesnes, Saint Quentin-Fallavier,
212 France). 25 fatty acids were detected and quantified.

213

214

Sterol Analysis

Sterols in abdominal muscle tissue homogenates were extracted with a solvent mixture containing chloroform/methanol 2/1 (v/v) spiked with internal standards as described extensively elsewhere ([Chevy et al., 2005](#)). Briefly, lipids were partitioned in chloroform after the addition of saline and saponified by methanol potassium hydroxide (0.5 N, 60 °C, 15 min). The fatty acids released were then methylated with BF₃-methanol (12%, 60 °C, 15 min) to prevent from interfering with the chromatography of sterols. Sterols were further re-extracted in hexane and silylated, and the resultant derivatives were separated by gas chromatography (GC) (Hewlett–Packard 6890 series) in a medium polarity capillary column RTX-65, (Restek, Evry, France). The mass spectrometer (Agilent 5975 inert XL) in series with the GC was set up for the detection of positive ions, which were produced in the electron impact mode at 70 eV. Sterols were identified by the fragmentogram in the scanning mode, and quantified by selective monitoring of the specific ions after normalization with the internal standards and calibration with weighed standards.

Glycerophospholipid Analysis

Total lipids were extracted from abdominal muscle tissues by the method of [Folch et al. \(1957\)](#). Individual phospholipid classes and their molecular species were quantified by a triple quadrupole Qtrap 6500, ABSciex (Les Ulis, France) in tandem with a Shimadzu Nexera XR liquid chromatography system (Shimadzu France, Marne la Vallée, France). A YMC-Pack PVA-Sil, bonded with a monomolecular polymer coating of vinyl alcohol (PVA) (Particle size : 5 µm Pore size : 120 Å, Usable pH range : 2.0-9.5 ; (YMC, Japan)) was used to separate the different lipid classes. Concentrations of the lipid compounds (including PE, PC, SM) were determined by comparing the peak area of each complex lipid with that of standards added with a known quantity. In total, 108 complex lipid compounds were quantified. The method is fully described in [Wolf & Quinn \(2008\)](#) and in [Lamaziere et al. \(2011\)](#).

Statistical analysis

In order to provide a broad view of the variations between the lipidomes of the shrimps in response to the two pressure conditions, Principal Component Analyses (PCA) were performed using RStudio software (1.1.4 version, packages ggplot2, factoextra, factoMineR, 2018 versions). The analysis was performed on both concentration and percentage data for the fatty acids and the complex lipids. This analysis also allowed to spot if any particular contamination occurred in some samples, by detecting outliers in the scores plot.

247 Following the PCAs, and in order to focus on the lipidome differences driven by pressure
248 changes only, Orthogonal Projections to Latent Structures Discriminant Analyses (OPLS-DA) were
249 then performed on the same data, using SIMCA 15.0 software (Umetrics, Umeå). The OPLS-DA,
250 introduced by Trygg & Wold (2002), is a multivariate analysis for omics datasets, which helps to
251 understand what distinguishes two groups of samples according to the factors that vary between
252 them. It is an extension to the supervised PLS regression method (Wold & Martens, 1983), which
253 removes variation from X (descriptor variables, here each lipid compound proportion or
254 concentration) that is not correlated to Y (the variable to explain, here the pressure condition) in the
255 first component of the model. Therefore, it is supposed to generate models easier to interpret
256 compared to those generated by PLS-DA (Checa et al., 2015, Trygg & Wold, 2002). The
257 importance of variables on projection for the predictive component values (VIPpred) for the
258 generated models were then calculated in order to detect which lipid compounds (fatty acids and
259 complex lipids) were most responsible for the separation of the two groups of shrimps. They
260 provide a score value for each independent variable X summarizing their importance in the
261 projection used by the PLS model to explain Y (Checa et al., 2015, Galindo-Prieto et al., 2014).
262 Hence, the higher the VIPpred score of a variable, the more important for the explanation of Y. VIP
263 values higher than one are generally admitted to consider a variable as significantly important
264 (Chong & Jun, 2005; Sun et al., 2012). Data were mean-centered and variance-scaled prior to PCA
265 and PLS-DA multivariate analyses.

266 We subsequently tested the difference in mean proportion between the groups of pressure
267 for the individual fatty acids (see Table 1). Data were tested with Shapiro–Wilks in order to
268 evaluate their distribution. If the distribution was not normal, a non-parametric Wilcoxon-Mann
269 Whitney test was applied. If the distribution was normal a t-test or a Welch test (if the variances of
270 the two groups were different) was performed. All statistics were performed using an α -level of
271 0.05. These tests were performed using RStudio software.

272 In the case of groups of variables (sum of the saturated, mono-unsaturated, and poly-unsaturated
273 fatty acids, shown in Figure 2), the two experimental groups were statistically compared, however
274 without using Bonferroni corrections. Although debated among statisticians (Nakagawa, 2004), the
275 use of such corrections accounts for the inflation of type I errors in multiple testing. The statistical
276 power of the corresponding results has therefore not passed the rigor of Bonferroni corrections,
277 however these data were not central in our approach, they were rather used as comparison points
278 with other studies.

279

280 RESULTS

281 Overview of *M. fortunata* lipidomic response to acclimation pressure

282 The two groups of shrimp samples form two separated clusters in the first two PCA
283 components performed on the concentration data for the fatty acids and the complex lipid contents
284 (Figures S1-S2 in supplementary material). The two groups were segregated in two clusters
285 opposed on a diagonal plan regarding their fatty acids concentrations, while they were opposed on
286 the first axis regarding their complex lipid concentrations. No strong outlier was detected. The
287 loading plots (not shown) and the raw data suggested that this clear segregation could be explained
288 by generally higher lipid concentrations in BOX shrimps, with respect to AP shrimps. Indeed, mean
289 concentration of fatty acids of BOX shrimps was 1317 +/- 283 ng/mg while that of AP shrimps was
290 590 +/- 380 ng/mg. Similarly, the muscle of BOX shrimps contained higher levels of PE (431 +/-
291 189 against 189 +/- 85 nanoM/mg) and PC (144 +/- 18 against 67 +/- 15 nanoM/mg). A similar
292 trend was observed for sterol levels (9.56 +/- 2.11 nanoM/mg for BOX against 7.60 +/- 3.57
293 nanoM/mg for AP shrimps), and also for SM levels (12.92 +/- 2.12 nanoM/mg for BOX, vs. 11.16
294 +/- 1.86 nanoM/mg for AP shrimps).

295

296 The two groups of shrimps also formed two distinct groups when the PCA analyses were
297 performed on data expressed as percentage of total lipid fatty acids, in order to avoid size effects
298 due to concentration differences (Figures S3-S4 in supplementary material). In the subsequent
299 OPLS-DA regarding fatty acid percentages, the two groups of shrimps segregated through the first
300 two components regarding their fatty acid compositions (Figure 1). In the score plot of the OPLS-
301 DA performed on complex lipid compositions, the two groups were also well separated (Figure S5
302 supplementary material). The two AP and BOX groups exhibited important phenotype
303 dissimilarities leading to a clear segregation of the observations in the variability plane (first two
304 components).

305

306

307

317

318

% weight	BOX	SD	AP	SD	trend	p-value	Food
C12:0	0,21	0,13	0,48	0,28			0,06
C14:0	0,46	0,22	1,29	0,44	UP	**	1,60
C14:1	0,02	0,01	0,06	0,02	UP	*	0,03
C16:0	15,04	2,02	16,93	1,11			6,53
C16:1n-9	5,04	1,70	4,60	0,85			3,02
C16:1n-7	0,08	0,01	0,12	0,03	UP	*	0,17
C16:2	0,03	0,01	0,06	0,02	UP	*	0,28
C17:0	0,12	0,04	0,23	0,05	UP	*	0,08
C18:0	1,69	0,45	2,72	0,47	UP	**	0,85
C18:1n-9	10,66	3,44	10,57	1,89			5,78
C18:1n-7	23,12	1,54	13,14	2,03	DOWN	**	5,94
C18:2n-6	1,73	0,33	1,96	0,75			2,93
C18:3n-6	0,13	0,01	0,23	0,04	UP	**	0,55
C18:3n-3	0,07	0,04	0,23	0,09	UP	**	0,00
C18:4	0,08	0,01	0,17	0,05	UP	**	1,94
C20:2	0,10	0,02	0,14	0,01	UP	*	0,02
C20:3n-9	0,07	0,02	0,12	0,05			0,02
C20:3n-6	0,04	0,00	0,08	0,02			0,00
C20:4n-6	0,68	0,16	1,01	0,29	UP	*	0,20
C20:5n-3	22,16	3,65	26,15	2,24			46,38
C22:5n-6	0,09	0,05	0,09	-			0,04
C22:5n-3	0,63	0,15	0,58	0,12			0,59
C22:6n-3	17,43	2,83	18,66	2,10			22,81
C24:0	0,11	0,09	0,30	0,13			0,05
C24:1	0,29	0,08	0,38	0,14			0,13

319

320

321 **Table 1:** Average fatty acid composition (% weight) of the *Mirocaris fortunata* and food lipid data
 322 set. Trends between BOX (n = 5 ind.) and AP (n = 5 ind.) shrimps are indicated (* symbol : p-value
 323 < 0.05, ** symbol : p-value < 0.01)

324

325

326 **Total fatty acid membrane composition**

327 25 fatty acids were detected in the analysis (**Fig. 1A**, and **Table 1**). The AP shrimps contain
328 a significantly higher proportion of saturated fatty acids than BOX shrimps (mean values : 21.95 %
329 and 17.58 %, respectively, p-value = 0.013), and a significantly lower proportion of mono-
330 unsaturated fatty acids (mean values : 28.86 % and 39.22 %, respectively, p-value = 0.011) (**Figure**
331 **2**). No difference was detected between BOX and AP shrimps levels of poly-unsaturated fatty acids
332 (p-value = 0.062).

333 Five predominant fatty acids each exceeded 10% of the composition, and accounted for 86.9
334 % in terms of proportion : C16:0 (15.98 % on average in all the shrimps), C18:1 n-7 (18.13 % on
335 average), C18:1 n-9 (10.61 % on average), C20:5 n-3 (known as “EPA”) (24.15 % on average) and
336 C22:6 n-3 (known as “DHA”) (18.4 % on average) (**Table 1**). No significant difference was found
337 between the levels of these predominant fatty acids within the two groups of shrimps, except for
338 C18:1 n-7, a MUFA. AP shrimps’ proportions of C18:1 n-7 were significantly lower than for BOX
339 shrimps (mean values : 13.14 % +/- 2.03, and 23.12 % +/- 1.54 respectively, p-value = 0.008). In
340 the OPLS-DA load plot (**Fig. 1B**), this fatty acid appeared to strongly discriminate the two
341 experimental groups. C18:1 n-7 also displayed the highest VIPpred score value (1.51) in the OPLS-
342 DA model performed on the fatty acid compositions (not shown). C18:1 n-7 alone is likely to
343 account for the higher proportion of MUFAs in BOX shrimps (**Table 1**, **Fig. 2**), as no other
344 predominant fatty acid seemed to vary between the two groups. The other most discriminating fatty
345 acids between the two groups of shrimps (according to their VIPpred values > 1, not shown), were
346 C18:3 n-6, C18:0, C14:0, C18:3 n-3, C17:0, C16:2, C14:1, C20:2, C18:4, C16:1 n-7, C20:4 n-6.
347 They were all present in proportions smaller than 3 % in the muscle of the shrimps, and these
348 proportions were all significantly higher in AP shrimps (**Table 1**). Finally, the food was analyzed
349 and two major components appeared to exceed 10% of the total composition : C20:5 n-3 (EPA,
350 46.38 %) and C22:6 n-3 (DHA, 22.82 %).

351 352 **Fatty acid metrics conventionally used to assess membrane order**

353 The average chain length (ACL) index, introduced by [Poynter et al. \(1989\)](#) was calculated as
354 follows for each shrimp sample: $ACL = (\sum [C_i] \times i) / \sum [C_i]$, where $[C_i]$ is the concentration of
355 the fatty acids containing i carbon atoms. Average ACL index was 18.65 +/- 0.19 for BOX
356 shrimps, and 18.68 +/- 0.13 for AP shrimps. The saturation ratio and unsaturation index were also
357 calculated according to [Cossins & MacDonald \(1986\)](#), i.e. the ratio of the weight % of saturated to
358 unsaturated fatty acids (for saturation ratio), and the sum of weight % multiplied by the number of
359 double bonds for each component (for unsaturation index) . The saturation ratio was 0.214 +/- 0.038

360 for BOX shrimps, and 0.282 ± 0.027 for AP shrimps. Finally, the unsaturation index was 265.7
 361 ± 24.7 for BOX shrimps, and 284.7 ± 15.9 for AP shrimps.
 362

363
 364
 365 **Figure 2:** Proportion (weight %) of saturated (SAT), mono-unsaturated (MUFAs) and poly-
 366 unsaturated (PUFAs) fatty acids in *Mirocaris fortunata* muscle after five months maintained at
 367 atmospheric pressure (AP, white bars, $n = 5$) or natural pressure (18MPa) in the Abyssbox (BOX ,
 368 black bars, $n = 5$). Mean values are shown and error bars correspond to the standard deviation.
 369

Complex lipid membrane composition

108 complex lipids were detected in the analysis, which targeted phosphatidylethanolamine (PE, 44 molecular species detected, including ether-PE (PE-O, 14 molecular species)), phosphatidylcholines (PC, 44 molecular species detected, including ether-PC (PC-O), 13 species), sphingomyelins (SM, 15 molecular species detected), and sterols (5 molecular species detected). For one sample (AP3), the sterol analysis could not be achieved for technical reasons, and therefore this individual observation was removed for the calculation of the following complex lipid proportions (molar %, **Figure 3**). Phosphatidylethanolamine (PE) was the dominant class in both groups of samples, accounting for 70.3 $\pm$ 7.6 % of complex lipid data composition in BOX shrimps, and 67.8 $\pm$ 11.7 % in AP shrimps (**Fig. 3**). Remarkably, these proportions were the addition of 33.0 % of ether-link PE (PE-O) and 37.3 % of ester-link PE for BOX shrimps, while they broke down in 42.3 % of ether-link PE (PE-O) and 25.5 % of ester-link PE, for AP shrimps. Phosphatidylcholines (PC) represented 25.6 $\pm$ 6.5 % of complex lipid data composition of the BOX shrimps and 24.3 $\pm$ 8.2 % of that of the AP shrimps. BOX shrimps contained 2.4 $\pm$ 0.9 % of sphingomyelin (SM), while AP ones contained 4.7 $\pm$ 2.3 %. Finally, sterols account for 1.7 $\pm$ 0.4 % in BOX shrimps and 3.2 $\pm$ 1.5 % in AP shrimps.

The VIPpred scoring of the OPLS-DA was performed on the complex lipids proportion dataset (**Figure 4**), in order to understand which lipid compounds separated the two groups (**Fig. S5 supplementary material**). **Figure 4** displays the 44 variables with VIPpred values higher than 1. Among the latter, 9 were sphingomyelins, all of which were dominant in the AP shrimps. All the detected sterols appeared in this figure (Cholesterol, 7-Lathosterol, Desmosterol, 7-DHC, 8-DHC), and were all present in higher proportions in the AP shrimps. Moreover, 23 phosphatidylethanolamines were contained in this list. Among them, the 8 that include an ether link (PE O) were more abundant in proportion in AP shrimps and the great majority of the others (14/15), which include an ester link (PE), were more abundant in proportion in BOX shrimps. For seven phosphatidylethanolamines, the ether (PE-O) and corresponding ester (PE) species were both present in the list. This list also included seven PC (all ester species), 3 of which were found in larger proportion in AP shrimps samples, while the 4 other ones displayed higher levels in BOX shrimps.

401

402

403

404

405 **Figure 3:** Complex lipids composition (molar %) of the data sets from *Mirocaris fortunata* muscle
406 after five months maintained at atmospheric pressure (AP, white bars, n = 4) or natural pressure
407 (18MPa) in the Abyssbox (BOX , black bars, n = 5). Mean values are shown and error bars
408 correspond to the standard deviation.

409

410

Figure 4: List of the variable importance for the projection (VIP) for the predictive components with scores higher than 1 (arbitrary unit). These VIPs refer to the OPLS-DA performed on % composition of complex lipids raw data. Each lipid molecular species is colored according to its relative abundance within specific experimental conditions: white/black bars indicate higher levels at atmospheric pressure (AP shrimps, n = 4)/18 MPa (BOX shrimps, n = 5) respectively.

422 DISCUSSION

423 The aim of the present work was to test whether the successful acclimation of a deep-sea
424 shrimp at two different pressures would allow to observe significant changes in membrane lipid
425 composition, as expected according to the concept of homeoviscous adaptation (review in [Ernst et](#)
426 [al., 2016](#)).

427 *The experimental context*

428 To our knowledge, it is the first time that lipidomic analyses have been conducted following
429 long-term acclimation (i.e. several months) of deep-sea animals in captivity. The shrimps studied
430 here had reached the Oceanopolis aquarium in July 2013 at the latest, and were sampled in
431 December 2014, meaning a minimum acclimation period of 17 months, with identical food during
432 the last five months.

433 While the AP group was maintained at atmospheric pressure, the BOX group was kept at its
434 natural hydrostatic pressure, corresponding to the depth of the site of origin (1700-1800 m, [Shillito](#)
435 [et al., 2015](#)). Except for hydrostatic pressure, our aim was to minimize environmental differences
436 which could influence lipid compositions of the studied animals. Water chemistry, temperature and
437 diet were the three main issues identified. Regarding nutrition, it is a well-known fact that lipid
438 composition of an organism is directly influenced by the fatty acid content of its food, justifying the
439 widespread use of fatty acids as tracers in trophic webs (review in [Kelly & Scheibling, 2012](#)).
440 Therefore identical food was provided to both BOX and AP shrimps during the last five months
441 prior to lipid analyses, in order to erase any differential diet effect on lipid composition. As to the
442 thermal environment, it is also well established that habitat temperature has a direct effect on lipid
443 composition of poikilothermic organisms ([Ernst et al., 2016](#)). In our experimental context, the
444 global temperature was 10°C, however for both groups a "hotspot" was provided (24-27 °C), and
445 AP and BOX shrimps were often seen aggregating on it, between periods of swimming or feeding
446 activity within the 10°C environment ([Matabos et al., Shillito et al., 2015](#)). Therefore a possible
447 effect of temperature on lipid composition cannot be ruled out, since we have no certainty on the
448 actual thermal regime experienced by the experimental animals. It should be noted however that
449 *Mirocaris fortunata* is a fairly eurythermal hydrothermal vent shrimp, with recorded habitat
450 temperature ranging from 4 to 24 °C *in situ* ([Desbruyères et al., 2001](#)). Finally, regarding water
451 chemistry, care was taken that both groups were supplied through the same seawater line.

452

453

454

455 *Higher levels of FA and complex lipids in pressurised samples (BOX)*

456 The concentration data (in ng/mg or nanomol/mg, as opposed to relative data, in %) showed
457 that the shrimps maintained at natural pressure (BOX shrimps, 18 MPa) were more concentrated in
458 lipid compounds than those maintained at atmospheric pressure (AP shrimps, 0.1 MPa). Overall,
459 mean concentration of fatty acids of BOX shrimps was two-fold of that of AP shrimps (1317 vs.
460 590 ng/mg), as for PE (431 vs. 189 nanoM/mg) and PC (144 vs. 67 nanoM/mg) levels. Such a trend
461 was already observed when comparing different animal species from various habitat depths
462 (shrimps : [Yerlikaya et al., 2013](#), or other animal species, [Bockus & Seibel, 2016](#)). The reasons for
463 such a phenomenon are unclear: although hydrostatic pressure was proposed as one among several
464 possible drivers of this phenomenon, ecological factors such as food limitation yielding increased
465 lipid storage were also proposed. In our case study, variations of the complex lipid species PE and
466 PC show that this response involves membrane contents, rather than lipid reserves. The decrease in
467 lipid levels in AP shrimps could then correspond to a limited microvilli or endomembrane system
468 synthesis, or lower cell divisions rates, at atmospheric pressure. Further structural studies (using
469 electron microscopy) would certainly help to test this hypothesis at the subcellular level.

471 *The major FA components*

472 Four of the five major FA found in our samples showed no significant variation between the
473 two experimental treatments. Among these, the two PUFA EPA (C20:5 n-3 / 24.15 %) and DHA
474 (C22:6 n-3 / 18.4 %) display levels which may appear unusually high for adult hydrothermal vent
475 fauna. The interpretation for lower levels in the wild (absence in several mid-Atlantic ridge animals,
476 [Colaço et al., 2007](#), [Zhu et al., 2015](#), or less than 6% for adult *Mirocaris* shrimp, [Pond et al., 1997](#),
477 and adult *Rimicaris* shrimp, [Streit et al., 2015](#)) is that EPA and DHA are considered as markers of a
478 photosynthetically-derived diet, of planktonic origin ([Kelly & Scheibling 2012](#), [Parrish, 2013](#)).
479 Therefore, they are rare in the hydrothermal vent food web, which is based on bacterial
480 chemosynthesis ([Colaço et al., 2007](#)). The higher levels observed in our experimental specimens
481 likely correspond to the EPA and DHA enrichment of the diet proposed in captivity (46 and 23%
482 respectively, see **Table 1**). The two other major FA that did not show significant variation between
483 our experiments are the SFA C16:0 (15.98 %), and the MUFA C18:1 n-9 (10.61 %). Such levels of
484 these FA are commonly observed in marine organisms ([Kelly & Scheibling 2012](#), [Parrish, 2013](#)),
485 including vent fauna ([Colaço et al., 2007](#)).

487 Finally, only one major FA showed a significant variation, the MUFA C18:1n-7 (vaccenic
488 acid), reaching 23.12% +/- 1.54 for natural pressure acclimation, and 13.14% +/- 2.03 for
489 atmospheric pressure acclimation (**Table 1**). In trophic studies, this FA is a common dietary tracer
490 (Kelly & Scheibling 2012, Parrish, 2013) and for example the ratio C18:1n-9/C18:1n-7 may be
491 used to discriminate carnivory from detritivory in deep-sea fauna (Drazen et al., 2008). At deep-sea
492 vents, vaccenic acid appears as a major component in several species (Streit et al., 2015, Colaço et
493 al., 2007), including Mirocaris (in the 6 to 21 % range, Pond et al., 1997). In the wild, these high
494 levels reflect the importance of food sources of bacterial origin, in the chemosynthetically-based
495 vent trophic web (Kelly & Scheibling 2012). In the present study, shrimps were in captivity and fed
496 with industrial food pellets, which contain a significant amount of C18:1n-7 (about 6%, **Table 1**).
497 Whether vaccenic acid in Mirocaris is here of endogenous or exogenous origin remains to be
498 determined. However, given that identical diet was provided in both experiments (AP and BOX),
499 the observed variation can not be explained by a diet effect.

500 In bacteria, C18:1n-7 is common and strongly involved in the homeoviscous response to
501 temperature, with an increased biosynthesis facing cold temperatures, resulting in lower membrane
502 order to counteract the ordering effect of cold temperatures (reviews in Zhang & Rock, 2008,
503 Hulbert et al., 2014, Ernst et al., 2016). Recent studies demonstrated that for *E. coli*, C18:1n-7
504 played a "disordering" role, and was the major MUFA involved in the regulation of membrane
505 order of this bacteria (Budin et al., 2018). Regarding a homeoviscous response to pressure, an
506 increased level of C18:1n-7 was also observed in response to increasing cultivation pressure, in the
507 case of a deep-sea bacteria (Allen et al., 1999). Accordingly, variations in the level of this
508 unsaturated FA impact membrane order, lower levels meaning higher order ("rigidifying effect"),
509 and vice-versa. The important variation of vaccenic acid observed in our experiment is consistent
510 with a homeoviscous response: exposed to atmospheric pressure, the shrimps displayed lower levels
511 of C18:1n-7, in order to counteract the disordering ("fluidizing") effect of lower pressure. We
512 therefore propose the existence of a homeoviscous response in the animals studied here, although
513 further measurements of membrane order and composition in our experimental conditions are
514 necessary for a complete demonstration.

515 Finally, it is interesting to note that C18:1n-7 (vaccenic acid) belongs to the same
516 biosynthetic pathway as the FA C16:1n-7 (Kelly & Scheibling 2012) observed at greater levels in
517 deep nematods (Van Campenhout et al., 2016) and fishes (Radnaeva et al., 2017), with respect to
518 shallower species. Whether this pathway (desaturation of C16:0 towards C16:1n-7, followed by
519 elongation towards C18:1n-7) was shut down following acclimation to atmospheric pressure (AP

520 samples), or whether C18:1n-7 was selectively accumulated through diet at natural pressure (BOX
521 samples), remains to be investigated.

522 *The global response in FA composition*

523 The decrease in vaccenic acid levels observed upon acclimation to atmospheric pressure is
524 likely responsible alone for the apparently lower levels of MUFA (**Fig. 2, Table 1**). This decrease
525 occurred along with increases of several other fatty acids: an increase in SFA levels likely relates to
526 the significant increases of C14:0, C18:0, and to a lesser extent C17:0 (**Table 1, Fig. 2**). These
527 significant MUFA and SFA variations probably contribute to the apparently higher saturation ratio
528 observed for atmospheric-pressure-acclimated shrimps (AP samples), and are also consistent with a
529 homeoviscous response. Inversely, the situation seems unclear regarding PUFA variations: while
530 several PUFA species clearly showed a marked up-regulation upon acclimation to atmospheric
531 pressure (namely C18:3n-6, C18:3n-3, C18:4, and to a lesser extent C16:2, C20:2, and C20:4n-6,
532 **Table 1**), these molecules nevertheless accounted for rather small proportions in the global
533 compositions (less than 1%), and displayed even smaller variations, as opposed to the ca. 10%
534 variation observed for vaccenic acid. At a global level furthermore (**Fig. 2**), there was no apparent
535 variation regarding either PUFA global levels (figure 3), or unsaturation index (265.7 +/- 24.7 vs.
536 284.7 +/- 15.9).

537 Most studies of lipid composition variations in response to increasing pressure and/or depth
538 have emphasized a modulation of MUFA vs. SFA levels (increase vs. decrease), while the
539 implication of PUFA variations is not clear. In the case of bacteria, some studies showed an
540 involvement of PUFA (enhanced biosynthesis under increasing pressure, [DeLong & Yayanos, 1986](#),
541 [Wang et al., 2014](#)), while others did not ([DeLong & Yayanos, 1985](#), [Allen et al., 1999](#), [Grossi et al.,](#)
542 [2010](#)). The situation also remains to be clarified regarding animals. In a study on antarctic copepods
543 sampled in the wild at increasing depths (from the surface down to approximately 1000m), [Pond et](#)
544 [al. \(2014\)](#) observed a significant decrease of SFA (consistent with our results) in the form of C16:0
545 and C18:0, but no specific variations of MUFA were mentioned (unlike our results). These studies
546 also found a significant increase of the PUFA DHA (C22:6 n-3) with depth (not observed in our
547 study), which they proposed to be consistent with homeoviscous adaptation theory. Another study
548 on different fish species sampled from the surface down to 1600 m depth in lake Baïkal lead to
549 decreased levels of SFA, and increased levels of MUFA (both consistent with our results, and with
550 a homeoviscous response, but here mainly due to C16:1n-7 and C18:1n-9 regarding MUFAs), yet
551 another trend for PUFA (decrease, including DHA), in deeper species ([Radnaeva et al., 2017](#)).
552 Another study comparing shallow-water and deep-sea nematod species (1300 m depth) lead to a
553 similar trend, regarding SFA (decrease), and MUFA (increase, C16:1n-7 in particular), but here

554 accompanied by an increase in PUFA levels (EPA and DHA in particular) for deeper species (Van
555 Campenhout et al., 2016). PUFA may help regulate membrane order, as suggested from *in vitro*
556 experiments (Guschina & Harwood 2006), however their role in that respect remains to be clarified
557 when dealing with whole animals sampled from the wild, with reduced knowledge on the
558 environmental and nutritional histories, as opposed to laboratory experiments. In our laboratory
559 study however, samples of the same species, fed for several months on identical diet, did not appear
560 to use PUFAs to adjust to experimental conditions.

561 Finally, the average chain length (ACL) calculated for each FA composition (BOX vs. AP
562 samples) remained unchanged (18.65 +/- 0.19 for BOX shrimps, and 18.68 +/- 0.13). Shifts in ACL
563 have been observed as a possible response to membrane order perturbation, following
564 environmental stress in bacteria (review in Yoon et al., 2015). Such a response does not seem to
565 occur in our study.

566 ***Variations in complex lipid composition***

567 As stated by Ernst et al. (2016), until now most studies have characterized the homeoviscous
568 response with a specific focus on the acyl chain composition, involving their degree of saturation
569 (relative compositions in terms of SFA MUFA PUFA). However, as a result of the development of
570 quantitative lipidomics technology, an increasing number of studies emphasize the importance of
571 complex lipidome variations of organisms facing changes in their thermal environment. To our
572 knowledge, the complex lipid composition of organisms experiencing pressure variation has little
573 been investigated so far, and hypotheses on lipid composition adjustments according to pressure are
574 scarce. In this study, we examined the compositions in PC, PE, SM and sterols of our two shrimp
575 pressure groups.

576 Cholesterol is known to regulate membrane order depending on its physical phase. It is
577 supposed to favor the order in the liquid phase (Urbina et al., 1995, Ayee & Levitan, 2016). But few
578 studies report a correlation between pressure and cholesterol content in metazoans. Vettier et al.
579 (2006) compared the cholesterol levels in the yellow and the silver stages of the European eel. The
580 yellow eel is the pre-migratory stage that lives in shallow fresh-waters. It never encounters high
581 pressure, contrary to the silver stage, which can be encountered at close to 1000 m depth during its
582 spawning migration (Righton et al., 2016). They suggested that the three-fold lower cholesterol
583 levels in silver eels could contribute in maintaining membrane order during their migration at depth.
584 Similarly, reduced proportions of cholesterol found in phocids were believed to help them cope
585 with high hydrostatic pressure while deep-diving (Williams et al., 2001). In our results, sterols (and
586 particularly cholesterol) were among the most discriminating complex lipids between the two

587 pressure groups (**Fig. 4**), representing on average 3.21 % of the complex lipids in the AP shrimps,
588 and 1.69 % in the BOX shrimps (**Fig. 3**).

589 As for sterols, we observed that AP shrimps tend to have higher proportions of
590 sphingomyelins than BOX shrimps on average (See **Figs 3-4**). Moreover, 9 out of the 15 detected
591 sphingomyelins were important to discriminate the two groups, and all of them were dominant in
592 AP shrimps (**Fig. 4**). This suggests that pressure also influences the SM levels. Finally, an
593 intriguing discovery is that 8 discriminating ether PE were dominant in AP shrimps, while most of
594 their corresponding ester versions (6) were dominant in BOX shrimps. It is interesting to note that
595 in another study on yeast ether lipid deficient mutants ([Shi et al., 2016](#)), a reduction in C18:1n-7
596 levels was observed, suggesting that production of these compounds were regulated by the same
597 mechanisms, in order to regulate membrane order. Also, the increase of ether lipids levels in
598 response to increasing temperature has long been reported, and their role in the homeoviscous
599 response was suggested to contribute to thermal compensation of membrane function ([Hazel &](#)
600 [Williams, 1990](#)).

601

602 Put together, cholesterol and sphingomyelins are known to be important components of
603 ordered lipid domains originally named "rafts" ([Simons & Ikonen, 1997](#), [Foster et al., 2003](#)). More
604 recently ether glycerophospholipids (PE species in particular), which also increase membrane order
605 ([Demediuk et al., 1983](#)), were also shown to be components of these domains and were believed to
606 stabilize these physical structures ([Wallner & Schmidz 2011](#), [Braverman & Moser, 2012](#);
607 [Tulodzieka et al., 2016](#), [Koivuniemi, 2017](#), [Dean & Lodhi, 2018](#)). Our data would thus be
608 consistent with the hypothesis of a modulation of the organization of these domains through specific
609 changes in the lipid signature of AP shrimps. This particular phenotype would possibly increase
610 membrane stability to maintain appropriate traffic and cell signaling, in response to the disordering
611 effect of lower pressure. Indeed, these domains were also known to be involved in many other
612 biological processes, such as signaling processes or vesicle traffic ([Foster et al., 2003](#)). More than
613 just contributing to structural stability, modifications in lipid fingerprints in AP shrimps might be
614 involved in regulating several cell functions impaired by the acclimation to atmospheric pressure. A
615 combined lipid/protein study, to further investigate the proteins, the regulation pathways commonly
616 associated with these domains and the variation of their relative gene expressions would allow to
617 test this hypothesis.

618

619 **Conclusion**

620 This study provided new insights on the lipidomic response of organisms following
621 acclimation to different pressures, and showed variations in lipid compositions, both at the fatty
622 acids and the complex lipids levels. It is the very first time that this response has been studied in a
623 deep-sea metazoan under long-term controlled conditions (pressure, temperature, food...), thanks to
624 the use of rearing equipments at different pressures, a unique opportunity provided by the
625 Océanopolis aquarium. It is also the first study reporting complex lipid variations in response to
626 pressure acclimation. One limitation of our study is that the lipidomes are not exhaustive since
627 several other complex lipids were not quantified (phosphatidylinositols, phosphatidylserines, ...),
628 meaning a possible bias in the observed variations. However, the clear response in terms of
629 vaccenic acid (C18:1n-7) variation, a molecule known to undoubtedly influence membrane order,
630 strongly suggests the existence of a homeoviscous response (i.e. a regulation of the membrane's
631 physical state) of *Mirocaris fortunata* following acclimation to atmospheric pressure. Whether the
632 accompanying complex lipid variations participate to a homeoviscous adaptation remains to be
633 determined. Further studies combining exhaustive lipidome determination and physical
634 measurements of membrane fluidity under different pressure conditions are needed to make the link
635 between these observations and membrane order. A complementary approach in order to better
636 understand the mechanisms underlying this lipidomic response would be to target the enzymes
637 involved in the production of the compounds which quantities differ, as well as follow the
638 expression levels of the corresponding genes.

639
640 Finally, the study presented here compares the lipidomes of shrimps that have been
641 maintained in captivity at different pressures, for several months. The observed lipidomic response
642 undoubtedly illustrates an acclimation process, thereby demonstrating the phenotypic plasticity of
643 these organisms regarding environmental variations. Further investigations should aim at evaluating
644 the kinetics of this response, by measuring lipidome variations throughout the acclimation period.
645 Bearing in mind that *Mirocaris fortunata* is found from 850 m to 4050 m depth, this would surely
646 provide new knowledge on the capacity of these shrimps to distribute over such a large bathymetric
647 range in the wild, and on their potential to undertake successful migrations towards yet unoccupied
648 depths, in response to near-future climate changes.

649
650

651 **ACKNOWLEDGEMENTS**

652 We wish to warmly thank Océanopolis staff members for their important contributions in shrimp
653 acclimation. We acknowledge the captains and crews of the oceanographic ships Thalassa
654 (Momarsat 2012 cruise) and Pourquoi Pas? (Biobaz 2013 cruises), and the pilots of Victor 6000
655 Remotely Operated Vehicle, for their dedicated assistance during sampling. We are also grateful to
656 G. Hamel, F. Lallier, M. Cannat, J. Blandin, P.-M. Sarradin, J. Sarrazin, P. Shillito, A. Colaço, and
657 R. Santos for scientific advice and support. We thank the two anonymous reviewers for their
658 important contribution to this manuscript. The AbyssBox project was funded by Océanopolis
659 Brest'Aim S.E.M., and this work also benefited from the support of the MIDAS European
660 Community project (European Union Seventh Framework Programme FP7/2007–2013) under the
661 grant agreement no. 603418. We also acknowledge initial financial support of the G.D.R. ECCHIS
662 (Ifremer, CNRS, UPMC).

663 BIBLIOGRAPHY

- 664 Allen, E. E., Facciotti, D., Bartlett, D. H. (1999). Monounsaturated but not polyunsaturated fatty acids are
 665 required for growth of the deep-sea bacterium *Photobacterium profundum* SS9 at high pressure and low
 666 temperature. *Applied and Environmental Microbiology*, 65(4), 1710–1720.
- 667
- 668 Ammendolia, J., Hamel, J.-F., Mercier, A. (2018) Behavioural responses to hydrostatic pressure in selected
 669 echinoderms suggest hyperbaric constraint of bathymetric range. *Marine Biology* , 165: 145.
 670 <https://doi.org/10.1007/s00227-018-3399-7>
- 671
- 672 Ayee, M.A., Levitan, I. (2016) Paradoxical impact of cholesterol on lipid packing and cell stiffness.
 673 *Frontiers in Bioscience*, Landmark, 21, 1245-1259. doi: 10.2741/4454
- 674
- 675 Barnes, D.K., Kuklinski, P. (2010) Bryozoans of the Weddell Sea continental shelf, slope and abyss: did
 676 marine life colonize the Antarctic shelf from deep water, outlying islands or in situ refugia following
 677 glaciations ? *J. Biogeogr.*, 37, 1648–1656. doi:10.1111/j.1365-2699.2010.02320.x
- 678
- 679 Behan, M.K., Macdonald, A.G., Jones, G.R., Cossins, A.R. (1992) Homeoviscous adaptation under
 680 pressure: the pressure dependence of membrane order in brain myelin membranes of deep-sea fish,
 681 *Biochimica et Biophysica Acta*. 1103, 317-323
- 682
- 683 Bligh, E. G., Dyer, W. J. (1959). A rapid method of total lipid extraction and purification. *Canadian*
 684 *Journal of Biochemistry and Physiology*, 37(8), 911–917. <https://doi.org/10.1139/o59-099>
- 685
- 686 Bockus A., B., Seibel, B., A. (2016) Trimethylamine oxide accumulation as a function of depth in Hawaiian
 687 mid-water fishes. *Deep-Sea Research I*, 112 : 37–44
 688 <http://dx.doi.org/10.1016/j.dsr.2016.03.005>
- 689
- 690 Braverman, N. E., Moser, A. B. (2012). Functions of plasmalogen lipids in health and disease. *Biochimica*
 691 *et Biophysica Acta (BBA) - Molecular Basis of Disease*, 1822(9), 1442–1452.
 692 <https://doi.org/10.1016/j.bbadis.2012.05.008>
- 693
- 694 Brown, A., Thatje, S. (2014) Explaining bathymetric diversity patterns in marine benthic invertebrates and
 695 demersal fishes: physiological contributions to adaptation of life at depth. *Biol. Rev.*, 89, pp. 406–426. doi:
 696 10.1111/brv.12061
- 697
- 698 Brown, A., Thatje, S. (2015). The effects of changing climate on faunal depth distributions determine
 699 winners and losers. *Global Change Biology*, 21(1), 173–180. <https://doi.org/10.1111/gcb.12680>

700

701 Budin, I., De Rond, T., Chen, Y., Chan, L.J.G., Petzold, C.J., Keasling, J.D. (2018) Viscous control of
702 cellular respiration by membrane lipid composition. *Science*, 362, 1186-1189. DOI: 10.1126/
703 science.aat7925

704

705 Cario, A., Grossi, V., Schaeffer, P., Oger, P. M. (2015). Membrane homeoviscous adaptation in the piezo-
706 hyperthermophilic archaeon *Thermococcus barophilus*. *Frontiers in Microbiology*, 6, art. 1152.
707 <https://doi.org/10.3389/fmicb.2015.01152>

708

709 Carney, R. (2005). Zonation of Deep Biota on Continental Margins. In R. Gibson, J. Gordon, & R.
710 Atkinson (Eds.), *Oceanography and Marine Biology* (Vol. 20051650, pp. 211–278). CRC Press.
711 <https://doi.org/10.1201/9781420037449.ch6>

712

713 Checa, A., Bedia, C., Jaumot, J. (2015). Lipidomic data analysis: Tutorial, practical guidelines and
714 applications. *Analytica Chimica Acta*, 885, 1–16. <https://doi.org/10.1016/j.aca.2015.02.068>

715

716 Chevy, F., Humbert, L., Wolf, C. (2005). Sterol profiling of amniotic fluid: a routine method for the
717 detection of distal cholesterol synthesis deficit. *Prenatal diagnosis*, 25, 1000-1006. doi: 10.1002/pd.1254

718

719 Chong, I.-G., Jun, C.-H. (2005). Performance of some variable selection methods when multicollinearity is
720 present. *Chemometrics and Intelligent Laboratory Systems*, 78(1–2), 103–112.
721 <https://doi.org/10.1016/j.chemolab.2004.12.011>

722

723 Clarke, A., Tyler, P.A. (2008) Adult Antarctic Krill Feeding at Abyssal Depths. *Current Biology*, 18, 282–
724 285, DOI 10.1016/j.cub.2008.01.059

725

726 Colaço, A., Desbruyères, D., Guezennec, J. (2007). Polar lipid fatty acids as indicators of trophic
727 associations in a deep-sea vent system community. *Marine Ecology*, 28(1), 15–24.
728 <https://doi.org/10.1111/j.1439-0485.2006.00123.x>

729

730 Cossins, A. R., Macdonald, A. G. (1986) Homeoviscous adaptation under pressure. III. The fatty acid
731 composition of liver mitochondrial phospholipids of deep-sea fish. *Biochimica et Biophysica Acta*, 860,
732 325-335

733

734 Cottin, D., Brown, A., Oliphant, A., Mestre, N., Ravaux, J., Shillito, B., Thatje, S. (2012) Sustained
735 hydrostatic pressure tolerance of the shallow water shrimp *Palaemonetes varians* at different temperatures:
736 Insights into the colonisation of the deep sea. *Comp. Biochem. Physiol.*, A, 162, 357–363.
737 doi:10.1016/j.cbpa.2012.04.005

738

739 Dean, J.M., Lodhi, I.J. (2018) Structural and functional roles of ether lipids. *Protein and Cell*, 9(2):196–
740 206. doi : 10.1007/s13238-017-0423-5

741

742 DeLong, E. F., Yayanos, A. A. (1985). Adaptation of the Membrane Lipids of a Deep-Sea Bacterium to
743 Changes in Hydrostatic Pressure. *Science*, New Series, 228(4703), 1101–1103.

744

745 DeLong, E. F., Yayanos, A. A. (1986). Biochemical Function and Ecological Significance of Novel
746 Bacterial Lipids in Deep-Sea Procaryotes. *Applied Environmental Microbiology*, 51, 4, 734-737.

747

748 Demediuk, P., Cowan, D.L., Moscatelli, E (1983) Effects Of Plasmenylethanolamine On The dynamic
749 Properties Of The Hydrocarbon Region Of Mixed Phosphatidylcholine Phosphatidylethanolamine Aqueous
750 dispersions. A spin Label Study. *Biochimica et Biophysica Acta*, 730, 263-270

751

752 Desbruyères, D., Biscoito, M., Caprais, J.-C., Colaço, A., Comtet, T., Crassous, P., ... Vangriesheim, A.
753 (2001). Variations in deep-sea hydrothermal vent communities on the Mid-Atlantic Ridge near the Azores
754 plateau. *Deep Sea Research Part I: Oceanographic Research Papers*, 48(5), 1325–1346.
755 [https://doi.org/10.1016/S0967-0637\(00\)00083-2](https://doi.org/10.1016/S0967-0637(00)00083-2)

756

757 Desbruyères, D., Segonzac, M., & Bright, M. (2006). *Handbook of Deep-Sea Hydrothermal Vent Fauna*
758 (Second compl. rev. ed). Linz: Art & Publishing.

759

760 Drazen, J.C., Phleger, C.F., Guest, M.A., Nichols, P.D. (2008) Lipid, sterols and fatty acids of abyssal
761 polychaetes, crustaceans, and a cnidarian from the northeast Pacific Ocean: food web implications. *Marine*
762 *Ecology Progress Series*, 372: 157–167. doi: 10.3354/meps07707

763

764 Ernst, R., Ejsing, C. S., & Antonny, B. (2016). Homeoviscous Adaptation and the Regulation of Membrane
765 Lipids. *Journal of Molecular Biology*, 428(24), 4776–4791. <https://doi.org/10.1016/j.jmb.2016.08.013>

766

767 Ernst, R., Ballweg, S. and Levental, I. (2018) Cellular mechanisms of physicochemical membrane
768 homeostasis. *Current Opinion in Cell Biology* 2018, 53:44–51. doi.org/10.1016/j.ceb.2018.04.013

769

770 Fang, J., Zhang, L, Bazylnski D. A. (2010) Deep-sea piezosphere and piezophiles: geomicrobiology and
771 biogeochemistry. *Trends in Microbiology* 18, 413-422.

772

773 Fernandes, P. M. B., Domitrovic, T., Kao, C. M., Kurtenbach, E. (2004). Genomic expression pattern in
774 *Saccharomyces cerevisiae* cells in response to high hydrostatic pressure. *FEBS Letters*, 556(1–3), 153–160.
775 [https://doi.org/10.1016/S0014-5793\(03\)01396-6](https://doi.org/10.1016/S0014-5793(03)01396-6)

776

777 Folch, J., Lees, M., Sloane Stanley, G. H. (1957). A simple method for the isolation and purification of total
778 lipides from animal tissues. *The Journal of Biological Chemistry*, 226(1), 497–509.

779

780 Foster, L. J., de Hoog, C. L., Mann, M. (2003). Unbiased quantitative proteomics of lipid rafts reveals high
781 specificity for signaling factors. *Proceedings of the National Academy of Sciences*, 100(10), 5813–5818.
782 <https://doi.org/10.1073/pnas.0631608100>

783

784 Fouquet, Y., Cherkashov, G., Charlou, J.-L., Ondreas, H., Cannat, M., Bortnikov, N., Silantiev, S.,
785 Etoubleau, J. (2008). Serpentine cruise - ultramafic hosted hydrothermal deposits on the Mid- Atlantic
786 Ridge: First submersible studies on Ashadze 1 and 2, Logatchev 2 and Krasnov vent fields. *InterRidge*
787 *News*, 2008, Vol. 17, 15-20.

788

789 Freitas, J.M., Bravim, F., Buss, D.S., Lemos, E.M., Fernandes, A.A.R., Fernandes, P.M.B. (2012) Influence
790 of cellular fatty acid composition on the response of *Saccharomyces cerevisiae* to hydrostatic pressure
791 stress. *FEMS Yeast Research*, 12, 871–878 DOI: 10.1111/j.1567-1364.2012.00836.x

792

793 Fumiyoshi, A. (2013) Dynamic structural changes in microbial membranes in response to high hydrostatic
794 pressure analyzed using time-resolved fluorescence anisotropy measurement. *Biophysical Chemistry*, 183,
795 3–8 . doi.org/10.1016/j.bpc.2013.05.005

796

797 Galindo-Prieto, B., Eriksson, L., Trygg, J. (2014). Variable influence on projection (VIP) for orthogonal
798 projections to latent structures (OPLS): Variable influence on projection for OPLS. *Journal of*
799 *Chemometrics*, 28(8), 623–632. <https://doi.org/10.1002/cem.2627>

800

801 Grossi, V., Yakimov, M. M., Al Ali, B., Tapilatu, Y., Cuny, P., Goutx, M., ... Tamburini, C. (2010).
802 Hydrostatic pressure affects membrane and storage lipid compositions of the piezotolerant hydrocarbon-
803 degrading *Marinobacter hydrocarbonoclasticus* strain #5: Pressure dependent bacterial storage lipid
804 formation. *Environmental Microbiology*, 12(7), 2020–2033. [https://doi.org/10.1111/j.1462-](https://doi.org/10.1111/j.1462-2920.2010.02213.x)
805 [2920.2010.02213.x](https://doi.org/10.1111/j.1462-2920.2010.02213.x)

806

807 Guschina, I. A., Harwood, J. L. (2006). Mechanisms of temperature adaptation in poikilotherms. *FEBS*
808 *Letters*, 580(23), 5477–5483. <https://doi.org/10.1016/j.febslet.2006.06.066>

809

810 Hazel, J. R. (1995). Thermal Adaptation in Biological Membranes: Is Homeoviscous Adaptation the
811 Explanation? *Annual Review of Physiology*, 57(1), 19–42.
812 <https://doi.org/10.1146/annurev.ph.57.030195.000315>

813

814 Hazel, J. R., Williams, E. E. (1990). The role of alterations in membrane lipid composition in enabling
815 physiological adaptation of organisms to their physical environment. *Progress in Lipid Research*, 29(3),
816 167–227. [https://doi.org/10.1016/0163-7827\(90\)90002-3](https://doi.org/10.1016/0163-7827(90)90002-3)

817

818 Hulbert, A.J., Kelly, M.A., Abbott, S.K. (2014) Polyunsaturated fats, membrane lipids and animal
819 longevity. *J Comp Physiol B* , 184:149–166 DOI 10.1007/s00360-013-0786-8

820

821 Jamieson, A.J., Fujii, T., Mayor, D.J., Solan, M. and Priede, I.G. (2010) Hadal trenches: the ecology of the
822 deepest places on Earth. *Trends in Ecology and Evolution* , Vol.25 No.3, 190-197.
823 doi:10.1016/j.tree.2009.09.009

824

825 Kelly, J.R., Scheibling, R.E. (2012) Fatty acids as dietary tracers in benthic food webs. *Marine Ecology*
826 *Progress Series*, 446: 1–22. doi: 10.3354/meps09559

827

828 Koivuniemi, A. (2017) The biophysical properties of plasmalogens originating from their unique molecular
829 architecture. *FEBS Letters* , 591, 2700–2713. doi:10.1002/1873-3468.12754

830

831 Lamaziere, A., Richard, D., Barbe, U., Kefi, K., Bausero, P., Wolf, C., et al., (2011) Differential
832 distribution of DHA-phospholipids in rat brain after feeding: A lipidomic approach. *Prostaglandins Leukot*
833 *Essent Fatty Acids*, 84, 7-11

834

835 Macdonald, A.G. (1997) Hydrostatic pressure as an environmental factor in life processes. *Comparative*
836 *Biochemistry and Physiology* 116A, 291-297.

837

838 Martin, J. W., Christiansen, J. C. (1995). A new species of the shrimp genus *Chorocaris* Martin (Crustacea:
839 Decapoda: Bresiliidae) from hydrothermal vent fields along the Mid-Atlantic Ridge. *Proceedings of the*
840 *Biological Society of Washington*, 108, 220–227.

841

842 Matabos, M., Cuvelier, D., Brouard, J., Shillito, B., Ravaux, J., Zbinden, M., ... Sarrazin, J. (2015).
843 Behavioural study of two hydrothermal crustacean decapods: *Mirocaris fortunata* and *Segonzacia*
844 *mesatlantica* , from the Lucky Strike vent field (Mid-Atlantic Ridge). *Deep Sea Research Part II: Topical*
845 *Studies in Oceanography*, 121, 146–158. <https://doi.org/10.1016/j.dsr2.2015.04.008>

846

847 Nakagawa, S. (2004) A farewell to Bonferroni: the problems of low statistical power and publication bias.
848 *Behavioral Ecology*, Vol. 15, No. 6: 1044–1045. doi:10.1093/beheco/arh107

849

850 Parent, G.J., Pernet, F., Tremblay, R., Sévigny, J.-M., Ouellette, M. (2008) Remodeling of membrane lipids
851 in gills of adult hard clam *Mercenaria mercenaria* during declining temperature. *Aquatic Biology*, 3: 101–
852 109. doi: 10.3354/ab00073

853

854 Parrish, C.R. (2013) Lipids in Marine Ecosystems. *ISRN Oceanography*, 2013, 16. doi :
855 10.5402/2013/604045

856

857 Pond, D.W., Segonzac, M., Bell, M.V., Dixon, D.R., Fallick, A.E., John, R., Sargent, J.R. (1997) Lipid and
858 lipid carbon stable isotope composition of the hydrothermal vent shrimp *Mirocaris fortunata*: evidence for
859 nutritional dependence on photosynthetically fixed carbon. *Marine Ecology Progress Series*, 157, 221-231.

860

861 Pond, D. W., Tarling, G. A., Mayor, D. J. (2014). Hydrostatic Pressure and Temperature Effects on the
862 Membranes of a Seasonally Migrating Marine Copepod. *PLoS ONE*, 9(10), e111043.
863 <https://doi.org/10.1371/journal.pone.0111043>

864

865 Poynter, J. G., Farrimond, P., Robinson, N., Eglinton, G. (1989). Aeolian-Derived Higher Plant Lipids in
866 the Marine Sedimentary Record: Links with Palaeoclimate. In M. Leinen & M. Sarnthein (Eds.),
867 *Paleoclimatology and Paleometeorology: Modern and Past Patterns of Global Atmospheric Transport* (pp.
868 435–462). Dordrecht: Springer Netherlands. https://doi.org/10.1007/978-94-009-0995-3_18

869

870 Pradillon, F. (2012). High hydrostatic pressure environments. In E. M. Bell (Ed.), *Life at extremes:*
871 *environments, organisms and strategies for survival* (pp. 271–295). Wallingford: CABI.
872 <https://doi.org/10.1079/9781845938147.0271>

873

874 Priede, I. G., Froese, R. (2013). Colonization of the deep sea by fishes: colonization of the deep sea.
875 *Journal of Fish Biology*, 83(6), 1528–1550. <https://doi.org/10.1111/jfb.12265>

876

877 Radnaeva, L. D., Popov, D. V., Grahl-Nielsen, O., Khanaev, I. V., Bazarsadueva, S. V., Käkälä, R. (2017).
878 Fatty acid composition in the white muscle of Cottoidei fishes of Lake Baikal reflects their habitat depth.
879 *Environmental Biology of Fishes*, 100(12), 1623–1641. <https://doi.org/10.1007/s10641-017-0670-6>

880

881 Ramirez-Llodra, E., Brandt, A., Danovaro, R., De Mol, B., Escobar, E., German, C.R., Levin, L.A.,
882 Martinez Arbizu, P., Menot, L., Buhl-Mortensen, P., Narayanaswamy, B.E., Smith, C.R., Tittensor, D.P.,
883 Tyler, P.A., Vanreusel, A., Vecchione, M. (2010) Deep, diverse and definitely different: unique attributes
884 of the world’s largest ecosystem. *Biogeosciences*, 7, 2851–2899, doi:10.5194/bg-7-2851-2010

885

886 Righton, D., Westerberg, H., Feunteun, E., Okland, F., Gargan, P., Amilhat, E., ... Aarestrup, K. (2016).
887 Empirical observations of the spawning migration of European eels: The long and dangerous road to the
888 Sargasso Sea. *Science Advances*, 2(10), e1501694–e1501694. <https://doi.org/10.1126/sciadv.1501694>
889
890 Sarda, F., D’Onghia, G.-F., Politou, C.Y., Company, J.B., Maiorano, P., Kapiris, K. (2004) Deep-sea
891 distribution, biological and ecological aspects of *Aristeus antennatus* (Risso, 1816) in the western and
892 central Mediterranean Sea. *Scientia. Marina.*, 68 (Suppl. 3): 117-127
893
894 Shi, X., Tarazona, P., Brock, T. , Browse, J., Feussner, I., Watts, J.L. (2016) A *Caenorhabditis*
895 *elegans* model for ether lipid biosynthesis and function. *J. Lipid Res.* 57: 265–275. DOI
896 10.1194/jlr.M064808
897
898 Shillito, B., Le bris, N., Hourdez, S., Ravaux, J., Cottin, D., Caprais, J.-C., Jollivet, D., Gaill, F. (2006).
899 Temperature resistance studies on the deep-sea vent shrimp *Mirocaris fortunata*. *Journal of Experimental*
900 *Biology*, 209(5), 945–955. <https://doi.org/10.1242/jeb.02102>
901
902 Shillito, B., Ravaux, J., Sarrazin, J., Zbinden, M., Sarradin, P.-M., Barthelemy, D. (2015). Long-term
903 maintenance and public exhibition of deep-sea hydrothermal fauna: The AbyssBox project. *Deep Sea*
904 *Research Part II: Topical Studies in Oceanography*, 121, 137–145.
905 <https://doi.org/10.1016/j.dsr2.2015.05.002>
906
907 Simons, K., Ikonen, E. (1997). Functional rafts in cell membranes. *Nature*, 387(6633), 569–572.
908 <https://doi.org/10.1038/42408>
909
910 Sinensky, M. (1974). Homeoviscous Adaptation - A Homeostatic Process that Regulates the Viscosity of
911 Membrane Lipids in *Escherichia coli*. *Proceedings of the National Academy of Sciences*, 71(2), 522–525.
912 <https://doi.org/10.1073/pnas.71.2.522>
913
914 Somero, G. N. (1992). Adaptations to high hydrostatic pressure. *Annual Review of Physiology*, 54, 557–
915 577. <https://doi.org/10.1146/annurev.ph.54.030192.003013>
916
917 Streit, K., Bennet, S.A., VanDover, C.L., Coleman, M. (2015) Sources of organic carbon for *Rimicaris*
918 *hybisae*: Tracing individual fatty acids at two hydrothermal vent fields in the Mid-Cayman rise. *Deep-Sea*
919 *Research Part I*, 100, 13–20. <http://dx.doi.org/10.1016/j.dsr.2015.02.003>
920
921 Sun, X.-M., Yu, X.-P., Liu, Y., Xu, L., Di, D.-L. (2012). Combining bootstrap and uninformative variable
922 elimination: Chemometric identification of metabonomic biomarkers by nonparametric analysis of

discriminant partial least squares. *Chemometrics and Intelligent Laboratory Systems*, 115, 37–43.
<https://doi.org/10.1016/j.chemolab.2012.04.006>

Trygg, J., Wold, S. (2002). Orthogonal projections to latent structures (O-PLS). *Journal of Chemometrics*, 16(3), 119–128. <https://doi.org/10.1002/cem.695>

Tulodziecka, K., Diaz-Rohrer, B.B., Farley, M.M., Chan, R.B., Di Paolo, G., Levental, K.R., Waxham, M.N., Levental, I. (2016) Remodeling of the postsynaptic plasma membrane during neural development. *Molecular Biology of the Cell*, 27, 3480–3489. doi/10.1091/mbc.E16-06-0420

Urbina, J. A., Pekerar, S., Le, H., Patterson, J., Montez, B., Oldfield, E. (1995). Molecular order and dynamics of phosphatidylcholine bilayer membranes in the presence of cholesterol, ergosterol and lanosterol: a comparative study using ²H-, ¹³C- and ³¹P-NMR spectroscopy. *Biochimica et Biophysica Acta (BBA) - Biomembranes*, 1238(2), 163–176. [https://doi.org/10.1016/0005-2736\(95\)00117-L](https://doi.org/10.1016/0005-2736(95)00117-L)

Van Campenhout, J., Vanreusel, A. (2016). Closely related intertidal and deep-sea Halomonhystera species have distinct fatty acid compositions. *Helgoland Marine Research*, 70(1). <https://doi.org/10.1186/s10152-016-0467-6>

Vettier, A., Labbe, C., Amerand, A., Da Costa, G., Le Rumeur, E., Moisan, C., Sebert, P. (2006). Hydrostatic pressure effects on eel mitochondrial functioning and membrane fluidity. *Undersea & Hyperbaric Medicine: Journal of the Undersea and Hyperbaric Medical Society, Inc*, 33(3), 149–156.

Wallner, S., Schmitz, G. (2011) Plasmalogens the neglected regulatory and scavenging lipid species. *Chemistry and Physics of Lipids*, 164, 573–589 doi: 10.1016/j.chemphyslip.2011.06.008

Wang, J., Li, J., Dasgupta, S., Zhang, L., Golovko, M.Y., Golovko, S.A., Fang, J. (2014) Alterations in Membrane Phospholipid Fatty Acids of Gram-Positive Piezotolerant Bacterium *Sporosarcina* sp. DSK25 in Response to Growth Pressure. *Lipids*, 49:347–356. DOI 10.1007/s11745-014-3878-7

Williams, E. E., Stewart, B. S., Beuchat, C. A., Somero, G. N., Hazel, J. R. (2001). Hydrostatic-pressure and temperature effects on the molecular order of erythrocyte membranes from deep-, shallow-, and non-diving mammals. *Canadian Journal of Zoology*, 79(5), 888–894. <https://doi.org/10.1139/cjz-79-5-888>

Wirsén, C.O., Jannasch, H.W., Wakeham, S.G., Canuel, E.A. (1987) Membrane Lipids of a Psychrophilic and Barophilic Deep-sea Bacterium. *Current Microbiology*, 14, pp. 319–322

960 Wold, S., Martens, H., Wold, H. (1983). The multivariate calibration problem in chemistry solved by the
 961 PLS method. In B. Kågström & A. Ruhe (Eds.), *Matrix Pencils* (Vol. 973, pp. 286–293). Berlin,
 962 Heidelberg: Springer Berlin Heidelberg. <https://doi.org/10.1007/BFb0062108>
 963

964 Wolf, C., Quinn, P.J. (2008) Lipidomics: practical aspects and applications, *Prog Lipid Res*, 47, 15-36
 965

966 Yancey, P.H., Geringer, M.E., Drazen, J.C., Rowden, A.A., Jamieson, A. (2014) Marine fish may be
 967 biochemically constrained from inhabiting the deepest ocean depths, *Proc. Nat. Acad. Sci.* vol. 111, 12,
 968 4461–4465. doi/10.1073/pnas.1322003111
 969

970 Yerlikaya, P., Topuz, O. K., Buyukbenli, H. A., Gokoglu, N. (2013). Fatty Acid Profiles of Different
 971 Shrimp Species: Effects of Depth of Catching. *Journal of Aquatic Food Product Technology*, 22(3), 290–
 972 297. <https://doi.org/10.1080/10498850.2011.646388>
 973

974 Yoon Y., Lee, H., Lee, S., Kima, S., Choi, K.-H. (2015) Membrane fluidity-related adaptive response
 975 mechanisms of foodborne bacterial pathogens under environmental stresses. *Food Research International*,
 976 72 25–36. <http://dx.doi.org/10.1016/j.foodres.2015.03.016>
 977

978 Zhang, Y.M., Rock, C.O. (2008) Membrane lipid homeostasis in bacteria. *Nature reviews microbiology*, 6,
 979 222-233. doi:10.1038/nrmicro1839
 980

981 Zhu, S., Ye, M., Yan, X., Zhou, Y., Wang, C., Xu, J. (2015). Lipid Adaptation of Shrimp *Rimicaris*
 982 *exoculata* in Hydrothermal Vent. *Lipids*, 50(12), 1233–1242. <https://doi.org/10.1007/s11745-015-4081-1>
 983