Olfactory detection of trace amounts of plant volatiles is correlated with testosterone in a passerine bird

Jessica L. Graham, Thierry D Charlier, Francesco Bonadonna, Samuel Caro

► To cite this version:

Jessica L. Graham, Thierry D Charlier, Francesco Bonadonna, Samuel Caro. Olfactory detection of trace amounts of plant volatiles is correlated with testosterone in a passerine bird. *Hormones and Behavior*, 2021, 136, pp.105045. 10.1016/j.yhbeh.2021.105045 . hal-03330902

HAL Id: hal-03330902

<https://hal.science/hal-03330902>

Submitted on 1 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Olfactory detection of trace amounts of plant volatiles is correlated with testosterone in a passerine bird

Jessica L. Graham^{1,†}, Thierry D. Charlier², Francesco Bonadonna¹, Samuel P. Caro^{1,*}

*Corresponding author: Samuel.caro@cefe.cnrs.fr; 33.(0)4.67.61.33.08 (P)

¹CEFE, Univ Montpellier, CNRS, EPHE, IRD, Univ Paul Valéry Montpellier 3, Montpellier, France

Email: Francesco.bonadonna@cefe.cnrs.fr

²Univ Rennes, Inserm, EHESP, Irset (Institut de Recherche en Santé, Environnement et Travail), UMR_S 1085, F-35000 Rennes, France

Email: Thierry.charlier@univ-rennes1.fr

[†]Current Address: Department of Biological Sciences, North Dakota State University, Fargo, ND 58102, USA

Email: Jessica.l.graham@ndsu.edu

Abstract

In response to damage by insects, plants release herbivore-induced plant volatiles (HIPVs) into the air. Insectivorous birds exploit these cues and, consequently, reduce the damages inflicted to the plants. However, little is known about whether they solely use HIPVs as foraging cues, or if they also use them to modulate traits linked to reproduction. As caterpillars are the primary food source required for insectivorous birds to raise offspring, their ability to locate and predict future peaks in caterpillar biomass using olfaction is likely to be advantageous. Therefore, we tested whether an insectivorous songbird that naturally inhabits oak dominated forests can be trained to detect early spring infestation by hatchling caterpillars, at a time when oaks begin bursting, and birds prepare to breed. Tree buds were either infested with caterpillars or left as a control and visually obscured in a Y-Maze choice test. Additionally, we measured testosterone and 17 β -estradiol as they influence olfactory perception in mammals and are linked to reproduction in vertebrates. After being trained to associate the presence of HIPVs with that of food, blue tits spent more time with, were more active around, and more frequently chose to first visit the infested trees, showing that blue tits can smell caterpillar activity. Males with higher testosterone spent more time around infested trees, suggesting that foraging behavior during the pre-breeding season is linked with a major reproductive signal. There was no relationship between foraging and estradiol in females. These results are an important foundation for further investigation of the role of hormones in avian olfaction and how smell may be useful for making breeding decisions that could improve reproductive success.

Keywords: seasonal reproduction, olfaction, alarm HIPVs, testosterone, estradiol

Introduction

Plants often emit chemical cues that protect them from damage caused by herbivores. In the case of herbivorous insects, volatile alarm signals released into the air attract carnivorous arthropods that protect the plants from damage (Turlings and Benrey, 1998; Van Poecke et al., 2001) by significantly reducing the number of herbivorous insects present on the plant (Kessler and Baldwin, 2001). In addition, some carnivorous arthropods appear to exploit these olfactory cues to prepare for reproduction. For example, ladybird beetles (*Harmonia axyridis* and *Oenopia conglobata*) experiencing a prey shortage will invest more in reproduction when presented with olfactory cues from aphid-infested plants (Rondoni et al., 2017). While most work on the influence of plant signals on animal behavior and physiology has focused on insects (Turlings and Benrey, 1998; Van Poecke et al., 2001), less is known about the abilities of vertebrates to detect and exploit these plant cues (Mrazova et al., 2019).

The olfactory ability of birds has become a popular area of study over the past few decades. Research has shown birds are capable of using smell in recognition of individuals (Bonadonna and Nevitt, 2004; Bonadonna and Sanz-Aguilar, 2012; Krause et al., 2012; Whittaker et al., 2011), selecting nest building materials (Clark and Mason, 1987; Petit et al., 2002), navigation (Bonadonna and Gagliardo, 2021; Gagliardo et al., 2013), and even modifying behavior to avoid predators (Amo et al., 2008; Roth II et al., 2008). There is growing evidence that insectivorous birds can also use their sense of smell in a foraging context, to detect infestation of plants by herbivorous insects (Amo et al., 2013; Mennerat et al., 2005). For example, the experimental infestation of birch trees (*Betula pubescens*) increases avian predation rates (Mäntylä et al., 2008). Similarly, birds are attracted to branches from infested pine trees (*Pinus sylvestris*) more than branches from uninfested trees (Mäntylä et al., 2017). However, in both these studies birds were able to see the infested trees and infestation can cause visual changes to trees (Amo et al., 2013). To our knowledge, only one published study exclusively tested the olfactory ability of birds to detect infested plants (Amo et al., 2013).

Among insects, caterpillars are particularly important for insectivorous birds, as they represent the primary food source required to successfully raise offspring (Naef-Daenzer and Keller, 1999). Thus, it may be expected that terrestrial birds may also use odors associated with food to detect it (Amo et al., 2013) and thereby constitute one of the most important selection factors in those bird species (Blondel et al., 1999). One such odor potentially associated with future food availability during the breeding season is herbivore-induced plant volatiles (HIPVs) released by plants in response to caterpillar herbivory. Volatile signals of caterpillar presence could modulate various behavioral and physiological processes in birds. If insectivorous birds like great tits use alarm signals from plants to forage for themselves and for feeding their chicks at a time when caterpillar biomass is large (i.e., late instar stages) and tree leaves are fully developed (Amo et al, 2013), birds may also use alarm signals from plants earlier in the season, when caterpillar biomass is still low because they just hatched and are even too small to be considered food. However, the olfactory detection of plants infested with young caterpillars has never been studied.

It has previously been shown that when young leaves are infested, monoterpene emissions are produced, but in significantly smaller amounts compared to mature leaves (Brilli et al., 2009). Cotton buds infested with *Helicoverpa zea* (Röse and Tumlinson, 2004) and oak buds infested with first instar green oak tortrix (*Tortrix viridana*) and winter moth (*Operophtera brumata*) caterpillars (about 1mm long) (Graham et al. unpublished results) already emit alarm volatile compounds, but in minute amounts compared to fully developed leaves. At this early stage, caterpillars live as miners in the buds, and are therefore invisible to potential predators (Du Merle and Mazet, 1983). Detecting such minute amounts of volatile signals could be challenging for birds, but there could nevertheless be several potential benefits to be gained from using such early signals of caterpillar presence in the environment. Birds might, for example, use the timing and intensity of the odorous signal to assess the quality of territories (Girling et al., 2011; Nyström, 1997). They could also use those signals to adjust clutch size in preparation for high or low insect

abundance (Gols et al., 2003; Hussell and Quinney, 1987; Marciniak et al., 2007), or potentially even to adjust laying date in response to year-to-year variation in the date of emergence of caterpillars. In order to determine if birds do make use of these early signals, we must first determine whether birds are capable of detecting the trace amounts of HIPVs emitted from developing buds, independently of visual cues. This is the main purpose of the present study.

If birds are detecting trace amounts of plant HIPVs and using those cues to make breeding decisions, any modification of reproductive behavior is likely to be primed by reproductive hormones (Ball and Balthazart, 2009, 2004; Wingfield et al., 1987). The relationship between chemosignals and their influence on the reproductive system is well documented in mammals, fish, and amphibians (Kawai et al., 2009; Nikonov et al., 2017; Petrulis, 2013), but less is known in birds (Caro and Balthazart, 2010), let alone for olfactory signals other than bodily odors, like plant or food odors for example. Work in several avian species suggest sex-steroids, like testosterone, stimulate more persistent food searching behavior (Andrew, 1972; Daisley et al., 2005; Németh et al., 2015), which is likely to be beneficial in early spring for acquiring resources for successful reproduction (e.g., territory, mate, food) (Lack, 1968; Wingfield, 1984). Additionally, olfactory sensitivity in rats appears to be upregulated by local sex-steroid production in the olfactory epithelium (Horie et al., 2017; Lupo et al., 1986), though whether this is the case in birds is unknown. Alternatively, it may be the exposure to some relevant odor signals that regulates sex-steroid dynamics, similarly to the stimulatory effect of pheromones on sex-steroid hormones (Miller and Maner, 2010). In any case, the relationship between behavior in response to an odor associated with food and pre-breeding reproductive hormones has not been studied.

The purpose of this study was to determine if insectivorous songbirds can detect the emissions of caterpillar infested trees early in the spring by smell only, and whether this is linked to their endocrine status. As blue tits (*Cyanistes caeruleus*) primarily inhabit oak dominated forests, we infested young oak trees with two common species of freshly hatched caterpillars that, later in the season, represent a major part of their natural diet in temperate zones: winter moths

(*Operophtera brumata*) and green oak tortrix (*Tortrix viridana*). We hypothesized that after training, birds would be able to differentiate between infested and uninfested oak trees by smell during the early stages of bud development, and that individuals that are more reproductively ready would spend more time near the infestation.

Methods

Ethics

Blue tits were trapped and maintained under licenses 2015615-147 issued by the Direction Régionale de l'Environnement, de l'Aménagement et du Logement Languedoc-Rousillon; and 15-XIX-116 issued by the Direction Départementale de la Protection des Populations de l'Hérault. Experiments were run under the license 2017-XIX-075 from the Direction Départementale de la Protection des Populations de l'Hérault, and approved by the Ethical committee N°036 with reference APAFIS#8608-2017012011062214 v4.

Housing of blue tits (*Cyanistes caeruleus*)

This experiment took place at the Centre National de la Recherche Scientifique (CEFE-CNRS) in Montpellier, France (43°37'56"N, 3°52'E) from 15 April to 2 May 2018. We used 42 hand-raised blue tits (females: $n = 20$, males: $n = 22$). Birds were moved indoors between 28 February and 14 March 2018 and were housed individually in cages (0.8 x 0.4 x 0.35m). Males and females were housed in separate rooms and females were never observed laying eggs for the duration of the study. Individuals were provided water and food *ad libitum*. Food consisted in a cake made of sunflower seed grease, eggs, sugar, wheat flour and high-protein pellets (Show 1+2 crumble, Versele-Laga, Deinze, Belgium), supplemented with minerals, vitamins, amino-acids and carotenoids (Nutribird A21, Versele-Laga, Deinze, Belgium; Nekton-S, Günter Enderle, Pforzheim, Germany; Yel-Lux, Versele-Laga, Deinze, Belgium). Mealworms were provided 3 times per week. Windows in the rooms allowed natural light to filter in while light timers were set

to turn on high-frequency TL tubes (Phillips TL5 HO 90 De Luxe 49W 965, Phillips, Eindhoven, The Netherlands) 15 minutes post-sunrise and turn off 15 minutes prior to sunset, providing natural increases and decreases in light intensity to occur (Fleissner and Fleissner, 2002). Sunrise and sunset times were determined using the United States Naval Observatory data base (www.usno.navy.mil/USNO). When moved, birds' feathers were checked to confirm they were in good condition, so flight capability was not inhibited.

Oak Trees

In the spring of 2018, 80 downy oak trees (*Quercus pubescens* Willd.; approx. 2 m high, and 0.8 m of diameter) grown in pots were maintained on the north side of a building ($n = 20$), south side of a building ($n = 20$), or at a higher elevation to delay bud burst ($n = 40$). All trees were treated with copper in a Bordeaux mixture (AMM 9500302) and a rapeseed oil insecticide (Naturen Eradibug, AMM 2110150) in the winter preceding each experiment to reduce the chance any fungus and wild insects infested the trees and caused other HIPV emissions than the ones induced by our caterpillar infestation. Once the oak buds began to elongate and lose their scales, they were considered ready to use in the experiment as caterpillars are unable to perforate the protective scales to enter the bud prior to this stage of development (Du Merle and Mazet, 1983). Of the 80 total trees, 24 were used in this experiment and 33 were used in a separate study specifically measuring HIPV production of oak buds experimentally infested with the same caterpillar species as here (Graham et al. unpublished results).

Caterpillars and bud infestation

We utilized 2 species of caterpillars for this project: winter moth (*Operophtera brumata*) and green oak tortrix (*Tortrix viridana*). Eggs of green oak tortrix were collected from tree branches the previous summer from Corsica, France while winter moth eggs were collected in the lab on

paper strips in The Netherlands. Eggs were kept in an unheated building overwinter and moved outdoors in spring so caterpillars and trees were experiencing the same weather conditions.

Once caterpillars began to hatch, oak trees with buds at an appropriate stage were either infested with caterpillars (using a small paintbrush to gently place caterpillars on the bud) or left as a control, 72 hrs prior to being used in a Y-maze (table 1). We chose 72 hrs post-infestation because data collected from experimentally infested oak buds found HIPVs significantly increase 48 – 72 hrs post-infestation when compared to uninfested, control buds (Graham et al. unpublished data). A single oak tree was placed at the end of each active arm (4 arms total, see below), so only 4 trees were used at one time. To infest a tree, 20 caterpillars were placed across 5 branches with stage 4 (bud elongated, swollen and green) – 6 (bud bursting, leaf shoots can be distinguished but are still intricate) buds (Du Merle and Mazet, 1983). *T. viridana* were given priority as they are native to the area, however, when not enough were available, *O. brumata* were used to keep the number of caterpillars on infested trees consistent. We recently found there is no difference in the volatile compounds emitted by oak buds when they were infested by either one of these two caterpillar species (Graham et al, unpublished results). A small wire cage was placed around the infested bud(s) and covered by a mesh bag to ensure caterpillars remained on the bud. The cage kept the bag from coming into contact with the bud(s), to prevent alarm emission due to mechanical stress (Kesselmeier and Staudt, 1999). Mean individual caterpillar mass for this study was 7.799 ± 0.990 mg. Control trees remained uninfested but had cages and bags on 5 branches to control for any emissions the tree may have produced in response to the presence of the cages. An individual bird was never exposed to the same pair of trees more than once and trees used on the day of the experiment had never been used during any of the training (see below).

Y-Mazes

Two large outdoor y-mazes (2.5x2m arms, figure S1) were set up facing southeast. Two arms were designated as active arms (one control and one treatment arm) and a neutral arm containing no stimuli where the bird entered the maze. Each active arm of the maze had five food dishes and one perch, and one additional perch was placed in the neutral arm of the maze. Food dishes were the same as those used to feed birds when housed in outdoor aviaries, so they were already familiar to the birds. A small box was mounted on the door in the neutral arm to introduce birds into the maze. Mazes were visually isolated from each other by a hedge. A wind break was set up around the maze that additionally reduced visibility beyond the arms of the maze. Stands placed at the far end of the two active arms of the maze (outside the maze, against the mesh) contained the trees. Those stands were covered with a white fabric on three sides (those visible from the inside of the maze) to block any changes in color infested trees may display compared to uninfested trees (Amo et al., 2013). These enclosures also had small fans (92x92x25mm, 36m³/h, EbmPapst, Chelmsford, UK) to blow air from the trees toward the center of the maze (placed at 80 cm, 130 cm, and 180 cm from the ground). Behavioral observations always started as soon as a bird left the entrance box. Habituation, training, and testing began as early as 08:40 and were completed by 16:45 each day (For full experimental design, see table 1). A water bath was also available in each arm of the maze, so the birds always had access to fresh water.

Habituation

We were unable to observe 42 birds in a day, so we ran the experiment twice. Group 1 consisted of 20 birds (10 males, 10 females) sampled from 15 April to 21 April 2018. Group 2 consisted of 22 birds (12 males, 10 females) sampled from 26 April to 2 May 2018. To habituate birds to the mazes, flocks of 3 – 4 birds were released into the maze for a minimum of 2 hours. Food was removed from the birds' housing cages 60 – 90 minutes before a bird was introduced to the maze, to stimulate food searching. A mixture of food rewards (cake, mealworms) were

placed in all 10 dishes at the end of both arms of the maze to encourage exploration. No trees or caterpillars were used during habituation. However, the empty tree enclosures and fans were in place so birds could habituate to their visual and auditory presence.

Training

Over the course of 5 consecutive days, individual birds were released into the mazes for 30 min with 1 food item placed in each dish only in the side of the maze containing an infested tree (Amo et al., 2013). Empty food dishes remained in the side of the maze with the control tree. The type of food reward was changed each day to keep birds from learning the scent of the food reward instead of the oak emissions (see table 1). The quantity of the food reward was recorded so we could keep track of whether birds were consuming the reward, and therefore whether they had a chance to associate the presence of oak emission with food. Additionally, the birds experienced the infested tree approximately equally on each side of the maze over the successive training sessions, so they would not become familiar with finding food on one side of the maze. Due to an odd number of trainings, half of the birds experienced the infested tree and food reward three times on the right side of the maze, while the other half of the birds experienced the infested tree and food reward three times on the left side of the maze. The side the infested tree was on was alternated between the left and right side daily for every individual. Order of individuals was randomly assigned, but one male and one female were always being tested in parallel in the two mazes with the exception of 2 males being tested at the same time during the second round of the experiment.

During the first training session, we observed that birds were less motivated to search for the food reward in the afternoon compared to the morning. To overcome this potential confounder, we alternated birds from the second group between morning and afternoon training sessions, so every bird experienced the maze 2 – 3 times in the morning. This reduced the number of birds that did not find the reward in at least half of the trainings from 40.0% of birds in the first round to

22.7% in the second round. In addition, inclement weather during the second round of trainings resulted in 12 birds only undergoing 4 rounds of training as opposed to 5 rounds.

Trials

Trials were performed once per individual and conducted in similar way to the training. A bird was released into the maze through the small box mounted on the door. The trial started as soon as a bird left the box and each bird was given 30 minutes to explore the maze. Food dishes were kept in the maze, but clean and empty of any food reward. Placement of the infested and control tree was randomized for each side of the maze and swapped after half of the birds had undergone a trial. Additionally, while 2 new trees were used in each maze on days 4, 7, and 9 (table 1), the trees were exchanged between mazes on days 5, 6, and 8 so that an individual bird experienced all 4 trees in up to 3 different combinations (e.g., day 1 – Trees 1 (infested) & 2 (control), day 2 – trees 3 (infested) & 4 (control), day 3 – trees 2 (control) & 3 (infested)). After trees were moved, the maze was left empty for 1 hour to allow any emissions caused by moving the tree to dissipate. Two Go Pro video cameras (GoPro, Inc., USA) were attached to the neutral perch, one facing into each active arm.

Behavioral Observations

Behavior was recorded by an observer blind to the treatment, using the free event logging software: Behavioral Observation Research Interactive Software (BORIS version 6.3.5, Friard and Gamba 2016). Videos in the left and right arm of the mazes could be watched at the same time and predetermined behavioral events were time stamped as they were observed. If a bird could not be seen in either video, they were presumed to be in the neutral arm of the maze. We measured (i) first arm of the maze visited after a bird entered the maze, (ii) time spent in each arm of the maze, and (iii) activity in each arm of the maze over the 30 min trial. Activity is the sum

of all movements where a bird was no longer in contact with its current perch (e.g., flying or hopping to a new or the original location).

Blood Sampling

Immediately after being removed from the maze on the trial day, a blood sample was collected from the wing vein. Samples were stored on ice until centrifugation to collect plasma. Plasma was stored at -20°C until assayed for plasma 17 β -estradiol and plasma testosterone in females and males, respectively.

Testosterone Assay

Plasma testosterone was measured using a commercially available enzyme immunoassay kit on a single plate (Enzo Life Sciences, ADI-900-065). Testosterone was extracted from 35 μ L of plasma using solid phase extraction with C18 columns (100 mg C18 material, Sep-Pak Vac 1cc, Waters, Milford, MA, USA), dried under nitrogen gas at 40°C, and reconstituted overnight with 250 μ L assay buffer (Caro et al., 2019). Concentration was adjusted for samples that did not have 35 μ L of plasma available ($n = 9$ of 22). Reconstituted samples were plated in duplicate (100 μ L per well) and concentrations determined using a freely available online five-parameter logistic curve-fitting program (MyAssays Ltd., 2019). Samples below detection limit ($n = 1$) were set at sensitivity of the assay (5.67 pg/mL). Intra-plate variation was calculated using pooled blue tit plasma from non-breeding individuals. To be certain the pool would produce results near the middle of the curve, 43.6 pg of exogenous testosterone for every 100 μ L of reconstituted sample was added prior to extraction. The standard was placed in three random locations on the plate and intra-assay variation was calculated at 6.52%.

We validated the assay in blue tits following the methods of (Caro et al., 2019). Briefly, pooled plasma samples were spiked with exogenous testosterone and serially diluted. The assay

was determined to work correctly as percentages of tracer bound (B/B0) from the serial dilutions were parallel to the standard curve (see supplementary material, figure S2).

Estradiol Assay

Plasma 17 β -estradiol (E₂) was measured using a commercially available double-antibody ¹²⁵I-E₂ radioimmunoassay (DSL-4800, Ultra-sensitive Estradiol RIA, Beckman Coulter, Brea, CA, USA) that was modified by dilution of the tracer and the antibody to increase the sensitivity of the assay, following (Charlier et al., 2010) and (Caro et al., 2019). Like for testosterone, steroids were first extracted from 35 μ L of plasma using solid phase extraction with C18 columns (100 mg C18 material, Sep-Pak Vac 1cc, Waters, Milford, MA, USA), dried under nitrogen gas at 40°C, and reconstituted overnight with PBSg (PBS with 0.1% gelatin) containing 0.7% ethanol (Caro et al., 2019). Concentration was adjusted for samples that did not have 35 μ L of plasma available (n = 11 of 20). Resuspended samples were then assayed in duplicate (300 μ l per tube) and assay tubes were counted on a gamma counter (Automatic Gamma Counter, Perkin Elmer, Waltham, MA, USA). Concentrations of E₂ were obtained using a linear regression with the log-transformed concentrations of the standards provided in the assay kit. All samples were run in a single assay, and the intra-assay coefficient of variation, as estimated by assaying one high and one low concentration E₂ standard in duplicate, was 5.9%. Assay sensitivity was 0.66 pg/ml. It was defined as the highest point on the standard curve whose standard deviation did not overlap that of the blank standard (Wingfield and Farner, 1975). No sample was found to be below the detection limit. The assay procedure has previously been validated for blue and great tits (Caro et al., 2019).

Statistical Analyses

All statistical analyses were run in R version 4.0.3 (R Core Team, 2020). As a preliminary analysis we used an exact binomial test to determine that individuals were equally as likely to go to the right side of the maze first (26 of 42) as they were to go to the left side when they entered

the maze (independently of which side the infested tree was on). Our results showed that birds were not lateralized ($p = 0.16$) and thus we excluded this variable from future analyses. We additionally ran preliminary analyses on behavior during training sessions. Briefly, we found that (i) birds did not become more successful in locating the food reward as the number of trainings increased ($z = -0.09$, $p = 0.93$, table S1); that (ii) training success was independent of plasma testosterone ($z = -0.96$, $p = 0.34$) and estradiol ($z = 0.14$, $p = 0.89$); and (iii) the ability to successfully locate the food reward was significantly repeatable within individuals at 0.49 ($C/\sigma = 0.21$, 0.66 , $p < 0.001$, see supplementary materials for more details on these analyses).

After the preliminary analyses, we first ran a simple binomial test that compared the number of birds that chose to visit the infested side when they entered the maze, with the number of birds that first visited the control side. We then used a logistic regression to test the effect of several explanatory variables on first side chosen. We tested whether the first side chosen was influenced by the maze in which an individual was tested (Maze 1 or 2), the sex of the birds (M or F), the period the birds were tested (group 1 = habituation, training, and testing from 15 to 21 April, group 2 = habituation, training, and testing from 26 April to 2 May), the order of testing within a group (from 1 to 11, as a continuous variable), and the number of times no food reward was consumed during training (from 0 to 5, continuous variable, listed as 'Reward Left' in tables 2 and 3). Statistical output for these variables is reported in table 2 and significant values are reported in text. Effect sizes were calculated using package effectsize (Ben-Shachar et al., 2020). We used standardized regression coefficients (β), which are reported in table 2 along with 95% confidence intervals.

To determine if time and activity level in each side of the maze differed between infested and control, paired t-tests (one for time, and one for activity) were initially used. To test additional variables that could potentially affect time and activity level in each side of the maze, two linear models were then run (one for each response variable, like above). The response variable for time was calculated as number of seconds spent in the infested arm of the maze minus the

number of seconds spent in the control arm of the maze. More positive values indicate more time spent in the infested arm, while more negative values indicate more time spent in the control arm. Similarly, the response variable for activity was calculated as the number of movements made in the infested arm minus the number of movements made in the control arm. More positive values indicate higher activity levels in the infested arm and more negative values indicate higher activity levels in the control arm. Explanatory variables included in both models were maze, sex, test group, testing order, and number of times the food reward was not consumed during training.

In a final set of analyses, we tested whether plasma steroid levels played an additional role on the time and the activity levels of the birds tested. Because we measured a different hormone in each sex, we ran separate models for males and females. Like in the models above, maze, test group, test order, and remaining food reward were also added, together with testosterone (in males) and estradiol (in females) concentrations (both $\ln$ transformed to achieve normality). Only significant values are reported in text for variables other than hormones, but all values are reported in table 3. A single male had undetectable testosterone levels and was subsequently excluded from final hormone analyses. However, assigning the lowest detectable level of the kit (5.67 pg/mL) and including the male in statistical models did not change the results. Graphs were made using package ggplot2 (Wickham, 2016). We set $\alpha = 0.05$ and report mean $\pm$ SEM.

Results

Individuals were more likely to visit the side of the maze with the infested tree first (29 of 42) compared to the control tree ($p = 0.02$). On average, birds spent 250.70 s longer in the side of the maze with the infested tree compared to the side with the control tree ($t_{41} = 2.13$, $p = 0.04$, figure 1). Birds were also nearly twice as active in the infested side of the maze compared to the control side ($t_{41} = 3.31$, $p = 0.002$, figure 1). When additional variables are introduced into a more

complex model, they have no influence on first side chosen (all $p > 0.1$), time (all $p > 0.09$), or activity (all $p > 0.12$, values reported in table 2).

Male-restricted analyses showed that birds with higher levels of testosterone spent more time in the infested arm of the maze ($F_{1,15} = 8.01$, $p = 0.01$, $R^2 = 0.44$, figure 2A). Testosterone was however not related to activity levels ($F_{1,15} = 3.35$, $p = 0.09$, $R^2 = 0.43$, table 3). Mean testosterone levels were significantly higher in maze 2 males (386.49 ± 122.42 pg/mL) compared to maze 1 males (259.26 ± 33.2 pg/mL; $F_{1,15} = 9.352$, $p = 0.008$), but this pattern was primarily driven by two males in maze 2 with the highest testosterone levels of all 21 individuals (maze 2 mean with highest two males removed: 214.25 ± 32.60 pg/mL). In pre-breeding females, estradiol levels showed no relationship with time ($F_{1,14} = 0.01$, $p = 0.94$, figure 2B) or activity ($F_{1,14} = 2.56$, $p = 0.13$). Other variables were also all non-significant (table 3).

Discussion

Blue tits are able to detect minute amounts of plant volatiles

We found that blue tits are capable of detecting HIPVs emitted by buds of one single small oak infested by young caterpillars. After training, individuals were not only more likely to visit the infested side of the maze first, but they also spent more time and were more active in the infested side of the maze. Our results are supported by other studies showing passerine birds select infested plants over uninfested plants (Amo et al., 2013; Mäntylä et al., 2017, 2008). In addition to the fact that visual cues were excluded in our experiment, emissions from infested buds appear to vary in both composition and quantity of HIPVs when compared to the mature foliage that older caterpillars feed on (Brilli et al., 2009; Röse and Tumlinson, 2004). This is the first study showing birds can detect the trace amounts of HIPVs from trees infested by young caterpillars during early stages of bud development, when caterpillars are not large enough to be considered a food source.

While our birds were originally naïve, having been in captivity from 10d of age, they were trained to associate the presence of food in small dishes with the smell of one infested tree. Our results clearly indicate that the detection of HIPVs from early bud infestation can be a learned behavior. While one study found that naïve great tits do not discriminate between infested and uninfested apple trees, suggesting use of HIPVs from fully developed leaves for locating food is not innate (Amo et al., 2016), new findings in blue tits show that detection of HIPVs from developing buds is innate in this species. When presented with an artificial odor bouquet mimicking the HIPVs released from developing buds, naïve blue tits are more likely to choose to first visit vials diffusing HIPVs as opposed to a control (Delaitre et al. unpublished data). Similarly, blue petrel fledglings (*Halobaena caerulea*) appear to show an innate preference for dimethyl sulfide, an olfactory cue important for locating patchily distributed prey in the ocean (Bonadonna et al., 2006; Cunningham et al., 2003). Further studies are needed to determine which HIPVs are innately detected, which ones need to be learned, and which ones may influence reproduction in blue tits.

Males with higher testosterone levels spent more time in the infested arm

We found that spending more time in the infested arm of the maze compared to the control arm was correlated with higher testosterone levels in males. Although plasma testosterone and time spent in the infested arm were correlated, it is unclear which of the two variables may affect the other. One potential explanation for this relationship is that male blue tits with higher levels of testosterone are more sensitive to cues that could indicate territory that will have higher food abundance. European starlings (*Sturnus vulgaris*) for example, only exhibit an olfactory response to the odors of a preferred nesting material (De Groof et al., 2010) and a conditioned odor stimulus during the breeding season (Clark and Smeraski, 1990). While artificially increasing plasma testosterone levels during the non-breeding season does not seem to change olfactory sensitivity to preferred nesting material in European starlings (De Groof et al., 2010) or food odor cues in

female goldfish (*Carassius auratus*) (Ghosal and Sorensen, 2016), the lack of receptors to receive the increased signal might result in non-activation of seasonal changes in olfactory sensitivity. Olfactory processing regions in many other vertebrate species contain gonadotropin releasing hormone (GnRH) and sex steroid receptors (Kawai et al., 2009). Steroid receptor density appears to change with reproductive status in fish (Maruska and Fernald, 2010), while olfactory sensitivity and preference appear to be highest during reproduction in rats (Moffatt, 2003). As the breeding season approaches, males may upregulate receptor density in olfactory processing regions so higher levels of testosterone have a greater influence on olfactory sensitivity during the breeding season. It should also be noted that work in mammals show that exposure to steroid hormones will modulate the integration of chemosensory signals, including olfactory cues. This was particularly well studied in the context of hormonal-dependent onset of maternal behavior in mammals. For example, females begin to show a preference for bedding soiled by pups over clean bedding, around the peripartum period (Kinsley and Bridges, 1990) and this is known to be associated with a general rewiring of the maternal brain (Keller et al., 2019; Lévy and Keller, 2009). Steroid-dependent changes in the neural pathways underlying chemosensory integration was also highlighted in copulatory context (Fiber and Swann, 1996; Paredes et al., 1998). Additional work should be performed to define whether sensitivity to HIPVs is modulated at the level of the periphery or the central nervous system, or both.

An alternative explanation for the correlation between testosterone and time spent in the infested arm of the maze is the rapid response of testosterone production to certain stimuli. For example, males of some species can quickly upregulate hormone production in response to male-male competition (Archer, 2006; McGlothlin et al., 2008; Wingfield and Wada, 1989), presence of a female (Coquelin and Bronson, 1979; Ronay and Hippel, 2010; Stacey, 2003), and even after exposure to a nest box (Gwinner et al., 2002). Male song sparrows showed a significant increase from baseline testosterone levels just 10 min after the start of a simulated territorial intrusion (Wingfield and Wada, 1989) and the introduction of green vegetation to captive-housed canaries

(*Serinus canaria*) results in an increase in testosterone (Voigt et al., 2007). Our birds were in the maze and able to interact with the infested trees for up to 30 min, providing ample time for the HIPV cues experienced in the maze to produce measurable differences in testosterone based on time spent near the stimulus (i.e., infested trees).

A third potential hypothesis also considers the fact that we trained individuals to associate food with HIPVs and testosterone's apparent role in olfactory sensitivity in rats and mice. For example, in gonadectomized male rats, testosterone injections can restore odor discrimination capabilities and improve task performance (Kunkhyen et al., 2018). However, our results suggest this may not be the case in blue tits. While those individuals with higher levels of testosterone may have spent more time in the infested arm of the maze, testosterone level was not correlated with increased success in locating the food reward during training. Instead, we found that the ability to locate the food reward was repeatable within an individual. Therefore, we think it is unlikely that testosterone played a role in developing a preference for HIPVs, but further work is needed to clarify this relationship.

Another consideration in the role of testosterone in detecting HIPVs is the subsequent reduction in testosterone levels during chick rearing (Hegner and Wingfield, 1987; Hunt et al., 1999), which seems counterintuitive when the ability to locate prey during this period is critical for reproductive success (Visser and Gienapp, 2019). However, fully developed vegetation produces HIPVs in much higher quantities than the early-spring buds we studied here (Brilli et al., 2009; Röse and Tumlinson, 2004), thus, increased olfactory sensitivity may not need to be maintained to successfully forage once HIPVs are emitted in higher quantities in response to herbivory by large caterpillars.

Estradiol is not correlated with behavior in females

We found no correlation between estradiol and time spent in the infested arm in female blue tits. Estradiol does respond to GnRH within 30 min in female great tits (*Parus major*) and

1
2
3
4 472 dark-eyed juncos (*Junco hyemalis*) (Caro et al., 2019; Needham et al., 2019), suggesting we
5
6 473 would see similar results to testosterone in male blue tits, if perception of a reproductively-relevant
7
8 474 stimulus like HIPVs can indeed induce a surge in sex-steroid hormones. Yet female birds control
9
10 475 timing of reproduction more closely than males (Caro et al., 2009). While male birds initiate
11
12 476 reproductive activity at the beginning of the breeding season and can maintain their status for
13
14 477 weeks (Ball and Ketterson, 2008; Caro et al., 2006), increases in plasma estradiol in females is
15
16 478 tightly correlated with rapid yolk development (which occurs just prior to egg laying) and
17
18 479 decreases to pre-breeding values before the final yolky follicle is ovulated (Williams et al., 2004).
19
20 480 Our female blue tits did not lay eggs while housed indoors for this experiment. Similarly, in
21
22 481 European starlings, social stimuli via the presence of a male is considered to be important for
23
24 482 stimulating final maturation of ovarian follicles (Perfito et al., 2015) and we housed male and
25
26 483 female blue tits separately during this experiment. Thus, female blue tits may not produce
27
28 484 estradiol in response to herbivore induced HIPVs occurring early in the season because they
29
30 485 were not ready to begin egg laying yet. Instead, females may be able to indirectly benefit from
31
32 486 increased testosterone in males. Physiological and behavioral changes associated with
33
34 487 reproduction are typically observed earlier in males than females (Ball and Ketterson, 2008; Caro
35
36 488 et al., 2006). The ability to detect HIPVs may be important for males in locating a high-quality
37
38 489 territory.

39
40 490 Testosterone on the other hand, may also be important in detecting environmental cues
41
42 491 in females. Interestingly, androgen levels in female European starlings appear to be upregulated
43
44 492 prior to rapid yolk development (Williams et al., 2004). For example, testosterone is highest during
45
46 493 territory establishment and pre-laying stages in female tree swallows (*Tachycineta bicolor*); which
47
48 494 corresponds with timing of female competition for resources and mates (George and Rosvall,
49
50 495 2018). Testosterone patterns in female blue tits could also upregulate in response to HIPVs and/or
51
52 496 influence olfactory sensitivity like in males, but future studies examining the effects of HIPV
53
54
55
56
57
58
59
60
61
62
63
64
65

exposure on female reproductive physiology will be needed to unravel the pathway linking HIPV perception to behavioral decisions.

Conclusions

Our study is novel in showing that a passerine bird is able to detect trace HIPVs produced in response to caterpillar infestation of developing buds. As freshly hatched caterpillars are too small and cryptic to be used as food, individuals may use these cues when making other decisions that would improve future reproductive success. Selecting mates based on the quality of the territory they occupy (Alatalo et al., 1986), adjusting clutch size based on habitat quality (Siikamäki, 1995), and timing offspring rearing to occur during peak food availability (Post and Forchhammer, 2008; Reed et al., 2013) are a few possibilities that need to be explored. Our finding that males with higher testosterone levels spend more time with infested trees compared to controls also indicates a possible physiological interaction with HIPVs. Future studies should now focus on how insectivorous vertebrates may use these cues early in the season and how HIPVs affect vertebrate physiology.

Acknowledgements

The authors thank M. Staudt for sharing his expertise in HIPV production; D. Degueldre for building the Y-mazes; M.E. Visser and B. van Lith for providing the winter moth eggs; C. de Francheschi for providing the tortrix eggs; B. Buatois and N. Barthes for assistance setting up the testosterone assay; M.M. Gueguen and M. Simean for assistance with the estradiol assay; S. Bencheikh for watching videos; and S. Ben-Chehida for assisting with tree infestation and bird care during the experiment.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

521 **Funding**

522 This work was supported by the Agence Nationale de la Recherche (awarded to SPC, grant
523 number ANR-15-CE02-0005-01) and a PRESTIGE Postdoctoral Research Fellowship (awarded
524 to JLG, grant number PRESTIGE-2018-2-0007).

525

References

- Alatalo, R.V., Lundberg, A., Glynn, C., 1986. Female pied flycatchers choose territory quality and not male characteristics. *Nature* 323, 152–153.
- Amo, L., Dicke, M., Visser, M.E., 2016. Are naïve birds attracted to herbivore-induced plant defences? *Behaviour* 153, 353–366.
- Amo, L., Galván, I., Tomás, G., Sanz, J.J., 2008. Predator odour recognition and avoidance in a songbird. *Funct. Ecol.* 22, 289–293.
- Amo, L., Jansen, J.J., van Dam, N.M., Dicke, M., Visser, M.E., 2013. Birds exploit herbivore-induced plant volatiles to locate herbivorous prey. *Ecol. Lett.* 16, 1348–1355.
<https://doi.org/10.1111/ele.12177>
- Andrew, R.J., 1972. Changes in search behaviour in male and female chicks, following different doses of testosterone. *Anim. Behav.* 20, 741–750. [https://doi.org/10.1016/S0003-3472\(72\)80146-5](https://doi.org/10.1016/S0003-3472(72)80146-5)
- Archer, J., 2006. Testosterone and human aggression: an evaluation of the challenge hypothesis. *Neurosci. Biobehav. Rev.* 30, 319–345.
- Ball, G.F., Balthazart, J., 2009. Neuroendocrine regulation of reproductive behavior in birds.
- Ball, G.F., Balthazart, J., 2004. Hormonal regulation of brain circuits mediating male sexual behavior in birds. *Physiol. Behav., Male Sexual Function* 83, 329–346.
<https://doi.org/10.1016/j.physbeh.2004.08.020>
- Ball, G.F., Ketterson, E.D., 2008. Sex differences in the response to environmental cues regulating seasonal reproduction in birds. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 363, 231–246.
<https://doi.org/10.1098/rstb.2007.2137>
- Bates, D., Maechler, M., Bolker, B., Walker, S., 2015. Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* 67, 1–48. <https://doi.org/10.18637/jss.v067.i01>
- Ben-Shachar, M., Lüdtke, D., Makowski, D., 2020. effectsize: Estimation of Effect Size Indices and Standardized Parameters. *J. Open Source Softw.* 5, 2518. <https://doi.org/doi:10.21105/joss.02815>
- Blondel, J., Dias, P.C., Perret, P., Maistre, M., Lambrechts, M.M., 1999. Selection-based biodiversity at a small spatial scale in a low-dispersing insular bird. *Science* 285, 1399–1402.
- Bonadonna, F., Caro, S.P., Jouventin, P., Nevitt, G.A., 2006. Evidence that blue petrel, *Halobaena caerulea*, fledglings can detect and orient to dimethyl sulfide. *J. Exp. Biol.* 209, 2165–2169.
<https://doi.org/10.1242/jeb.02252>
- Bonadonna, F., Gagliardo, A., 2021. Not only pigeons: avian olfactory navigation studied by satellite telemetry. *Ethol. Ecol. Evol.* 33. <https://doi.org/10.1080/03949370.2021.1871967>
- Bonadonna, F., Nevitt, G.A., 2004. Partner-Specific Odor Recognition in an Antarctic Seabird. *Science* 306, 835–835. <https://doi.org/10.1126/science.1103001>
- Bonadonna, F., Sanz-Aguilar, A., 2012. Kin recognition and inbreeding avoidance in wild birds: the first evidence for individual kin-related odour recognition. *Anim. Behav.* 84, 509–513.
- Brilli, F., Ciccio, P., Frattoni, M., Prestininzi, M., Spanedda, A.F., Loreto, F., 2009. Constitutive and herbivore-induced monoterpenes emitted by *Populus euroamericana* leaves are key volatiles that orient *Chrysomela populi* beetles. *Plant Cell Environ.* 32, 542–552.
- Caro, S.P., Balthazart, J., 2010. Pheromones in birds: myth or reality? *J. Comp. Physiol. A* 196, 751–766.
- Caro, S.P., Charmantier, A., Lambrechts, M.M., Blondel, J., Balthazart, J., Williams, T.D., 2009. Local adaptation of timing of reproduction: females are in the driver's seat. *Funct. Ecol.* 23, 172–179.
- Caro, S.P., Cornil, C.A., van Oers, K., Visser, M.E., 2019. Personality and gonadal development as sources of individual variation in response to GnRH challenge in female great tits. *Proc. R. Soc. B Biol. Sci.* 286, 20190142. <https://doi.org/10.1098/rspb.2019.0142>

- Caro, S.P., Lambrechts, M.M., Chastel, O., Sharp, P.J., Thomas, D.W., Balthazart, J., 2006. Simultaneous pituitary–gonadal recrudescence in two Corsican populations of male blue tits with asynchronous breeding dates. *Horm. Behav.* 50, 347–360. <https://doi.org/10.1016/j.yhbeh.2006.03.001>
- Charlier, T.D., Po, K.W., Newman, A.E., Shah, A.H., Saldanha, C.J., Soma, K.K., 2010. 17 β -Estradiol levels in male zebra finch brain: combining Palkovits punch and an ultrasensitive radioimmunoassay. *Gen. Comp. Endocrinol.* 167, 18–26.
- Clark, L., Mason, J.R., 1987. Olfactory discrimination of plant volatiles by the European starling. *Anim. Behav.* 35, 227–235. [https://doi.org/10.1016/S0003-3472\(87\)80228-2](https://doi.org/10.1016/S0003-3472(87)80228-2)
- Clark, L., Smeraski, C.A., 1990. Seasonal shifts in odor acuity by starlings. *J. Exp. Zool.* 255, 22–29.
- Coquelin, A., Bronson, F.H., 1979. Release of luteinizing hormone in male mice during exposure to females: habituation of the response. *Science* 206, 1099–1101.
- Cunningham, G.B., Van Buskirk, R.W., Bonadonna, F., Weimerskirch, H., Nevitt, G.A., 2003. A comparison of the olfactory abilities of three species of procellariiform chicks. *J. Exp. Biol.* 206, 1615–1620.
- Daisley, J.N., Bromundt, V., Möstl, E., Kotrschal, K., 2005. Enhanced yolk testosterone influences behavioral phenotype independent of sex in Japanese quail chicks *Coturnix japonica*. *Horm. Behav.* 47, 185–194.
- De Groof, G., Gwinner, H., Steiger, S., Kempenaers, B., Van der Linden, A., 2010. Neural correlates of behavioural olfactory sensitivity changes seasonally in European starlings. *PLoS One* 5, e14337.
- Du Merle, P., Mazet, R., 1983. Stades phénologiques et infestation par *Tortrix viridana* L. (Lep., Tortricidae) des bourgeons du chêne pubescent et du chêne vert. *Acta Oecologica* 4, 47–53.
- Fiber, J.M., Swann, J.M., 1996. Testosterone differentially influences sex-specific pheromone-stimulated Fos expression in limbic regions of Syrian hamsters. *Horm. Behav.* 30, 455–473.
- Fleissner, G., Fleissner, G., 2002. Perception of Natural Zeitgeber Signals, in: Kumar, V. (Ed.), *Biological Rhythms*. Springer Berlin Heidelberg, pp. 83–93. https://doi.org/10.1007/978-3-662-06085-8_8
- Friard, O., Gamba, M., 2016. BORIS: a free, versatile open-source event-logging software for video/audio coding and live observations. *Methods Ecol. Evol.* 7, 1325–1330. <https://doi.org/10.1111/2041-210X.12584>
- Gagliardo, A., Bried, J., Lambardi, P., Luschi, P., Wikelski, M., Bonadonna, F., 2013. Olfactory oceanic navigation in an Atlantic seabird. *J Exp Biol* 216, 2798–2805.
- George, E.M., Rosvall, K.A., 2018. Testosterone production and social environment vary with breeding stage in a competitive female songbird. *Horm. Behav.* 103, 28–35.
- Ghosal, R., Sorensen, P.W., 2016. Male-typical courtship, spawning behavior, and olfactory sensitivity are induced to different extents by androgens in the goldfish suggesting they are controlled by different neuroendocrine mechanisms. *Gen. Comp. Endocrinol.* 232, 160–173.
- Girling, R.D., Stewart-Jones, A., Dherbecourt, J., Staley, J.T., Wright, D.J., Poppy, G.M., 2011. Parasitoids select plants more heavily infested with their caterpillar hosts: a new approach to aid interpretation of plant headspace volatiles. *Proc. R. Soc. B Biol. Sci.* 278, 2646–2653. <https://doi.org/10.1098/rspb.2010.2725>
- Gols, R., Roosjen, M., Dijkman, H., Dicke, M., 2003. Induction of Direct and Indirect Plant Responses by Jasmonic Acid, Low Spider Mite Densities, or a Combination of Jasmonic Acid Treatment and Spider Mite Infestation. *J. Chem. Ecol.* 29, 2651–2666. <https://doi.org/10.1023/B:JOEC.0000008010.40606.b0>
- Gwinner, H., Van't Hof, T., Zeman, M., 2002. Hormonal and behavioral responses of starlings during a confrontation with males or females at nest boxes during the reproductive season. *Horm. Behav.* 42, 21–31.
- Hegner, R.E., Wingfield, J.C., 1987. Effects of experimental manipulation of testosterone levels on parental investment and breeding success in male house sparrows. *The Auk* 104, 462–469.

- Horie, S., Yamaki, A., Takami, S., 2017. Presence of Sex Steroid-Metabolizing Enzymes in the Olfactory Mucosa of Rats. *Anat. Rec.* 300, 402–414.
- Hunt, K.E., Hahn, T.P., Wingfield, J.C., 1999. Endocrine influences on parental care during a short breeding season: testosterone and male parental care in Lapland longspurs (*Calcarius lapponicus*). *Behav. Ecol. Sociobiol.* 45, 360–369.
- Hussell, D.J.T., Quinney, T.E., 1987. Food abundance and clutch size of Tree Swallows *Tachycineta bicolor*. *Ibis* 129, 243–258.
- Kawai, T., Oka, Y., Eisthen, H., 2009. The role of the terminal nerve and GnRH in olfactory system neuromodulation. *Zoolog. Sci.* 26, 669–680.
- Keller, M., Vandenberg, L.N., Charlier, T.D., 2019. The parental brain and behavior: A target for endocrine disruption. *Front. Neuroendocrinol.* 54, 100765.
- Kesselmeier, J., Staudt, M., 1999. Biogenic volatile organic compounds (VOC): an overview on emission, physiology and ecology. *J. Atmospheric Chem.* 33, 23–88.
- Kessler, A., Baldwin, I.T., 2001. Defensive Function of Herbivore-Induced Plant Volatile Emissions in Nature. *Science* 291, 2141–2144. <https://doi.org/10.1126/science.291.5511.2141>
- Kinsley, C.H., Bridges, R.S., 1990. Morphine treatment and reproductive condition alter olfactory preferences for pup and adult male odors in female rats. *Dev. Psychobiol. J. Int. Soc. Dev. Psychobiol.* 23, 331–347.
- Krause, E.T., Krüger, O., Kohlmeier, P., Caspers, B.A., 2012. Olfactory kin recognition in a songbird. *Biol. Lett.* 8, 327–329.
- Kunkhyen, T., Perez, E., Bass, M., Coyne, A., Baum, M.J., Cherry, J.A., 2018. Gonadal hormones, but not sex, affect the acquisition and maintenance of a Go/No-Go odor discrimination task in mice. *Horm. Behav.* 100, 12–19.
- Lack, D.L., 1968. Ecological adaptations for breeding in birds.
- Lévy, F., Keller, M., 2009. Olfactory mediation of maternal behavior in selected mammalian species. *Behav. Brain Res.* 200, 336–345.
- Lupo, C., Lodi, L., Canonaco, M., Valenti, A., Dessì-Fulgheri, F., 1986. Testosterone metabolism in the olfactory epithelium of intact and castrated male rats. *Neurosci. Lett.* 69, 259–262. [https://doi.org/10.1016/0304-3940\(86\)90490-8](https://doi.org/10.1016/0304-3940(86)90490-8)
- Mäntylä, E., Alessio, G.A., Blande, J.D., Heijari, J., Holopainen, J.K., Laaksonen, T., Piirtola, P., Klemola, T., 2008. From plants to birds: higher avian predation rates in trees responding to insect herbivory. *PLoS One* 3, e2832.
- Mäntylä, E., Kleier, S., Kipper, S., Hilker, M., 2017. The attraction of insectivorous tit species to herbivore-damaged Scots pines. *J. Ornithol.* 158, 479–491.
- Marciniak, B., Nadolski, J., Nowakowska, M., Loga, B., Bańbura, J., 2007. Habitat and annual variation in arthropod abundance affects Blue Tit *Cyanistes caeruleus* reproduction. *Acta Ornithol.* 42, 53–62.
- Maruska, K.P., Fernald, R.D., 2010. Reproductive Status Regulates Expression of Sex Steroid and GnRH Receptors in the Olfactory Bulb. *Behav. Brain Res.* 213, 208–217. <https://doi.org/10.1016/j.bbr.2010.04.058>
- Mcglathlin, J.W., Jawor, J.M., Greives, T.J., Casto, J.M., Phillips, J.L., Ketterson, E.D., 2008. Hormones and honest signals: males with larger ornaments elevate testosterone more when challenged. *J. Evol. Biol.* 21, 39–48. <https://doi.org/10.1111/j.1420-9101.2007.01471.x>
- Mennerat, A., Bonadonna, F., Perret, P., Lambrechts, M.M., 2005. Olfactory conditioning experiments in a food-searching passerine bird in semi-natural conditions. *Behav. Processes* 70, 264–270. <https://doi.org/10.1016/j.beproc.2005.07.005>
- Miller, S.L., Maner, J.K., 2010. Scent of a Woman: Men's Testosterone Responses to Olfactory Ovulation Cues. *Psychol. Sci.* 21, 276–283. <https://doi.org/10.1177/0956797609357733>

- Moffatt, C.A., 2003. Steroid hormone modulation of olfactory processing in the context of socio-sexual behaviors in rodents and humans. *Brain Res. Rev.* 43, 192–206.
- Mrazova, A., Sam, K., Amo, L., 2019. What do we know about birds' use of plant volatile cues in tritrophic interactions? *Curr. Opin. Insect Sci.*
- MyAssays Ltd., 2019. Five Parameter Logistic Curve online data analysis tool. MyAssays Ltd.
- Naef-Daenzer, B., Keller, L.F., 1999. The foraging performance of great and blue tits (*Parus major* and *P. caeruleus*) in relation to caterpillar development, and its consequences for nestling growth and fledging weight. *J. Anim. Ecol.* 68, 708–718. <https://doi.org/10.1046/j.1365-2656.1999.00318.x>
- Needham, K.B., Bergeon Burns, C.M., Graham, J.L., Bauer, C.M., Kittilson, J.D., Ketterson, E.D., Hahn, T.P., Greives, T.J., 2019. Changes in processes downstream of the hypothalamus are associated with seasonal follicle development in a songbird, the dark-eyed junco (*Junco hyemalis*). *Gen. Comp. Endocrinol.* 270, 103–112. <https://doi.org/10.1016/j.ygcen.2018.10.012>
- Németh, Z., Adams, R.H., Ramenofsky, M., 2015. Androgens Increase Persistence But Do Not Affect Neophobia in a Problem-Solving Context in a Songbird. *Ethology* 121, 428–433.
- Nikonov, A.A., Butler, J.M., Field, K.E., Caprio, J., Maruska, K.P., 2017. Reproductive and metabolic state differences in olfactory responses to amino acids in a mouth brooding African cichlid fish. *J. Exp. Biol.* 220, 2980–2992.
- Nyström, K.G.K., 1997. Food density, song rate, and body condition in territory-establishing willow warblers (*Phylloscopus trochilus*). *Can. J. Zool.* 75, 47–58. <https://doi.org/10.1139/z97-006>
- Paredes, R.G., Lopez, M.E., Baum, M.J., 1998. Testosterone augments neuronal Fos responses to estrous odors throughout the vomeronasal projection pathway of gonadectomized male and female rats. *Horm. Behav.* 33, 48–57.
- Perfito, N., Guardado, D., Williams, T.D., Bentley, G.E., 2015. Social cues regulate reciprocal switching of hypothalamic Dio2/Dio3 and the transition into final follicle maturation in European starlings (*Sturnus vulgaris*). *Endocrinology* 156, 694–706.
- Petit, C., Hossaert-McKey, M., Perret, P., Blondel, J., Lambrechts, M.M., 2002. Blue tits use selected plants and olfaction to maintain an aromatic environment for nestlings. *Ecol. Lett.* 5, 585–589. <https://doi.org/10.1046/j.1461-0248.2002.00361.x>
- Petrulis, A., 2013. Chemosignals, hormones and mammalian reproduction. *Horm. Behav.* 63, 723–741. <https://doi.org/10.1016/j.yhbeh.2013.03.011>
- Post, E., Forchhammer, M.C., 2008. Climate change reduces reproductive success of an Arctic herbivore through trophic mismatch. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 363, 2367–2373. <https://doi.org/10.1098/rstb.2007.2207>
- R Core Team, 2020. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org>.
- Reed, T.E., Jenouvrier, S., Visser, M.E., 2013. Phenological mismatch strongly affects individual fitness but not population demography in a woodland passerine. *J. Anim. Ecol.* 82, 131–144.
- Ronay, R., Hippel, W. von, 2010. The presence of an attractive woman elevates testosterone and physical risk taking in young men. *Soc. Psychol. Personal. Sci.* 1, 57–64.
- Rondoni, G., Ielo, F., Ricci, C., Conti, E., 2017. Behavioural and physiological responses to prey-related cues reflect higher competitiveness of invasive vs. native ladybirds. *Sci. Rep.* 7, 1–9.
- Röse, U.S., Tumlinson, J.H., 2004. Volatiles released from cotton plants in response to *Helicoverpa zea* feeding damage on cotton flower buds. *Planta* 218, 824–832.
- Roth II, T.C., Cox, J.G., Lima, S.L., 2008. Can foraging birds assess predation risk by scent? *Anim. Behav.* 76, 2021–2027.
- Siikamäki, P., 1995. Habitat Quality and Reproductive Traits in the Pied Flycatcher: An Experiment. *Ecology* 76, 308–312. <https://doi.org/10.2307/1940652>

- Stacey, N., 2003. Hormones, pheromones and reproductive behavior. *Fish Physiol. Biochem.* 28, 229–235.
- Stoffel, M., Nakagawa, S., Schielzeth, H., 2017. rptR: repeatability estimation and variance decomposition by generalized linear mixed-effects models. *Methods Ecol. Evol.* 8, 1639–1644.
- Turlings, T.C.J., Benrey, B., 1998. Effects of plant metabolites on the behavior and development of parasitic wasps. *Écoscience* 5, 321–333. <https://doi.org/10.1080/11956860.1998.11682472>
- Van Poecke, R.M., Posthumus, M.A., Dicke, M., 2001. Herbivore-induced volatile production by *Arabidopsis thaliana* leads to attraction of the parasitoid *Cotesia rubecula*: chemical, behavioral, and gene-expression analysis. *J. Chem. Ecol.* 27, 1911–1928.
- Visser, M.E., Gienapp, P., 2019. Evolutionary and demographic consequences of phenological mismatches. *Nat. Ecol. Evol.* 3, 879–885.
- Voigt, C., Goymann, W., Leitner, S., 2007. Green matters! Growing vegetation stimulates breeding under short-day conditions in wild canaries (*Serinus canaria*). *J. Biol. Rhythms* 22, 554–557.
- Whittaker, D.J., Richmond, K.M., Miller, A.K., Kiley, R., Bergeon Burns, C., Atwell, J.W., Ketterson, E.D., 2011. Intraspecific preen oil odor preferences in dark-eyed juncos (*Junco hyemalis*). *Behav. Ecol.* 22, 1256–1263. <https://doi.org/10.1093/beheco/arr122>
- Wickham, H., 2016. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York, New York, NY, USA.
- Williams, T.D., Kitaysky, A.S., Vézina, F., 2004. Individual variation in plasma estradiol-17 β and androgen levels during egg formation in the European starling *Sturnus vulgaris*: implications for regulation of yolk steroids. *Gen. Comp. Endocrinol.* 136, 346–352. <https://doi.org/10.1016/j.ygcen.2004.01.010>
- Wingfield, J.C., 1984. Androgens and mating systems: testosterone-induced polygyny in normally monogamous birds. *The Auk* 101, 665–671.
- Wingfield, J.C., Ball, G.F., Jr., A.M.D., Hegner, R.E., Ramenofsky, M., 1987. Testosterone and Aggression in Birds. *Am. Sci.* 75, 602–608.
- Wingfield, J.C., Farner, D.S., 1975. The determination of five steroids in avian plasma by radioimmunoassay and competitive protein-binding. *Steroids* 26, 311–327.
- Wingfield, J.C., Wada, M., 1989. Changes in plasma levels of testosterone during male-male interactions in the song sparrow, *Melospiza melodia*: time course and specificity of response. *J. Comp. Physiol. A* 166, 189–194.

Day	1	2	3	4	5	6	7	8	9
Caterpillars	Infest Trees (1 - 4)			Infest Trees (5 - 8)		Infest Trees (9 - 12)			
Birds	Habituation			Training 1	Training 2	Training 3	Training 4	Training 5	Trials
Trees Used	No Trees			1 – 4	1 – 4	1 – 4	5 – 8	5 – 8	9 – 12
Reward	Meal Worms, Cake			Meal Worms	Wax Moth Larvae	Crickets	Beetle Larvae	Wax Moth Larvae	None

Table 1. Timeline of caterpillar infestation, habituation, training, and trials conducted in captive blue tits. Only 22 birds could be used at a time; thus, the protocol was repeated twice.

Tables

Table 2. Analyses of the variables that influence the first side chosen, the time spent, and the level of activity close to one caterpillar-infested oak in blue tits ($n = 42$).

Response Variable	Explanatory Variable	B [95% CI]	Estimate	S.E.	Test Statistic	p-value
First Side Chosen	Maze	0.56 [-0.81, 2.03]	1.11	1.40	$Z = 0.80$	0.426
	Sex	-1.32 [-3.02, 0.15]	-1.31	0.79	$Z = -1.66$	0.096
	Test Group	0.09 [-0.70, 0.87]	0.18	0.77	$Z = 0.23$	0.820
	Order	-0.73 [-2.19, 0.63]	-0.12	0.12	$Z = -1.03$	0.302
	Reward Left	-0.45 [-1.29, 0.32]	-0.25	0.23	$Z = -1.11$	0.266
Time	Maze	-0.56 [-1.17, 0.05]	-842.70	454.72	$F_{1,36} = 2.93$	0.096
	Sex	-0.11 [-0.75, 0.54]	-81.69	242.19	$F_{1,36} = 0.17$	0.682
	Test Group	-0.11 [-0.44, 0.23]	-161.72	248.89	$F_{1,36} = 0.48$	0.493
	Order	0.34 [-0.28, 0.95]	43.25	38.90	$F_{1,36} = 1.26$	0.270
	Reward Left	0.05 [-0.29, 0.40]	22.36	74.43	$F_{1,36} = 0.09$	0.766
Activity	Maze	-0.26 [-0.58, 0.07]	-30.54	19.11	$F_{1,36} = 2.53$	0.120
	Sex	0.10 [-0.55, 0.76]	6.10	19.56	$F_{1,36} = 0.10$	0.748
	Test Group	0.17 [-0.17, 0.51]	19.98	20.10	$F_{1,36} = 0.96$	0.333
	Order	-0.04 [-0.36, 0.29]	-0.70	3.14	$F_{1,36} = 0.05$	0.828
	Reward Left	0.03 [-0.32, 0.38]	0.89	6.01	$F_{1,36} = 0.02$	0.883

Bolded values were statistically significant ($p < 0.05$). Effect sizes (β) are standardized beta coefficients followed by the 95% confidence intervals in brackets.

Table 3: Analyses of the variables that influence the time spent and the level of activity close to one caterpillar-infested oak in male ($n = 21$) and female ($n = 20$) blue tits.

Response Variable	Explanatory Variable	B [95% CI]	Estimate	S.E.	Test Statistic	p-value
Males	Testosterone (ln)	0.54 [0.13, 0.95]	567.19	203.46	$F_{1,15} = 8.01$	0.013
	Maze	-0.57 [-0.94, -0.20]	-853.77	261.06	$F_{1,15} = 9.35$	0.008
	Time					
	Test Group	0.19 [-0.24, 0.62]	288.00	309.04	$F_{1,15} = 0.63$	0.440
	Order	0.28 [-0.10, 0.67]	70.41	45.16	$F_{1,15} = 2.60$	0.127
	Reward Left	0.10 [-0.32, 0.52]	48.90	93.63	$F_{1,15} = 0.27$	0.609
	Testosterone (ln)	0.53 [0.12, 0.94]	39.23	14.23	$F_{1,15} = 3.35$	0.087
	Maze	-0.65 [-1.02, -0.28]	-68.43	18.26	$F_{1,15} = 14.12$	0.002
	Activity					
	Test Group	0.32 [-0.11, 0.76]	34.07	21.62	$F_{1,15} = 2.78$	0.116
Females	Order	-0.11 [-0.49, 0.28]	-1.87	3.16	$F_{1,15} = 0.35$	0.563
	Reward Left	-0.02 [-0.44, 0.41]	-0.56	6.55	$F_{1,15} < 0.01$	0.932
	Estradiol (ln)	-0.07 [-0.65, 0.51]	-165.44	643.38	$F_{1,14} = 0.01$	0.940
	Maze	-0.03 [-0.60, 0.53]	-47.78	394.39	$F_{1,14} = 0.01$	0.928
	Time					
	Test Group	-0.18 [-0.78, 0.42]	-270.04	419.53	$F_{1,14} = 0.56$	0.467
	Order	-0.04 [-0.62, 0.54]	-10.16	66.03	$F_{1,14} = 0.01$	0.930
	Reward Left	0.08 [-0.52, 0.68]	32.89	118.07	$F_{1,14} = 0.08$	0.785
	Estradiol (ln)	-0.37 [-0.91, 0.17]	-78.11	53.53	$F_{1,14} = 2.56$	0.132
	Maze	0.05 [-0.47, 0.58]	7.22	32.81	$F_{1,14} = 0.06$	0.807
Activity	Test Group	0.12 [-0.43, 0.68]	16.66	34.91	$F_{1,14} = 0.21$	0.657
	Order	-0.08 [-0.62, 0.46]	-1.77	5.49	$F_{1,14} = 0.09$	0.766
	Reward Left	0.03 [-0.53, 0.59]	1.14	9.82	$F_{1,14} = 0.01$	0.909

Bolded values were statistically significant ($p < 0.05$). Effect sizes (β) are standardized beta coefficients followed by the 95% confidence intervals in brackets.

Figures

Figure 1. Blue tits detect herbivore-induced plant volatiles (HIPV) emitted by the buds of one single, small oak. On average, individuals spent 250.7 seconds longer in the infested arm of the maze than the control arm and were significantly more active in the infested arm of the maze compared to the control arm. Activity is the sum of all movements where a bird was no longer in contact with its current perch (e.g., flying or hopping to a new or the original location). Circles represent individual data points while squares represent mean of all individuals $\pm$ SEM.

Figure 2. Testosterone correlates with HIPV detection. (A) Males with higher levels of testosterone spent significantly more time in the infested arm of the maze compared to the control arm of the maze, (B) but this was not the case for the relationship with estradiol in females.

Supplementary Material

Behavioral analyses during training

We ran preliminary analyses on bird behavior related with successfully locating food during training. To test whether individuals became more successful at finding the food reward during trainings using a generalized linear mixed-model with binomial distribution in package lme4 (Bates et al., 2015). The response variable was a repeated measure coded as 1 (consumed food reward) or 0 (did not consume food reward) for each of the 5 training sessions. Our fixed factor of interest in the model was the training session (1 – 5). Individuals were tested in two mazes, during two different time periods, thus, to determine if these variables affected the results, the maze an individual was tested in (Maze 1 or 2) and the test group (group 1 = habituation, training, and testing from 15 to 21 April, group 2 = habituation, training, and testing from 26 April to 2 May) were both included in the model. Testing order is included as a measure of when an individual was tested during the day. Those listed as 1 were the first to be tested on that day and those listed as 10 or 11 were last to be tested that day. Sex (M or F) was also included. Statistical output for these variables is reported in table S1 and significant values are reported in the main text. Given the evidence that testosterone can influence olfactory discrimination in rats and mice (Kunkhyen et al., 2018), we also decided to run 2 additional models comparing testosterone in males and estradiol in females to training success. Furthermore, we calculated the repeatability (logit link-scale approximation) of consuming the food reward across the five trainings using the package rptR (Stoffel et al., 2017).

Table S1. Analyses of the variables that influence training success. All individuals were included in the initial analysis while only males and females were included in the testosterone and estradiol analyses, respectively.

Training Success	Explanatory Variable	B [95% CI]	Estimate	S.E.	Test Statistic	p-value
All Individuals (<i>n</i> = 42)	Trial Number	-0.02 [-0.40, 0.37]	-0.01	0.14	<i>z</i> = -0.09	0.931
	Maze	-0.16 [-0.88, 0.56]	-0.32	0.73	<i>z</i> = -0.43	0.666
	Sex	1.15 [-0.31, 2.61]	1.15	0.74	<i>z</i> = 1.55	0.121
	Test Group	0.68 [-0.06, 1.42]	1.35	0.75	<i>z</i> = 1.80	0.072
	Order	-0.06 [-0.76, 0.65]	-0.02	0.12	<i>z</i> = -0.15	0.879
Males (<i>n</i> = 21)	Testosterone (ln)	-0.55 [-1.66, 0.57]	-0.74	0.77	<i>z</i> = -0.96	0.336
Females (<i>n</i> = 20)	Estradiol (ln)	0.07 [-0.92, 1.06]	0.23	1.59	<i>z</i> = 0.14	0.887

Bolded values were statistically significant (*p*<0.05). Effect sizes (*β*) are standardized beta coefficients followed by the 95% confidence intervals in brackets.

Figure S1. Diagram of the Y-mazes used in the behavioral experiment. To test the ability of insectivorous songbirds to detect caterpillar infestation of oak buds by smell, a Y-maze was set up so birds entered from a small box in the neutral arm. Two GoPro cameras were mounted on a center perch to film activity in each of the active arms. An infested oak tree was on one side of the maze and an uninfested tree on the other. Both were visually isolated from the bird to obscure any visual changes resulting from infestation, but small fans were in place to help direct any odors toward the center of the maze. Each trial lasted 30 minutes.

Figure S2. Testosterone assay validation for blue tit plasma. To validate the assay in blue tits, 2 plasma pools were created from samples collected from blue tits in January 2018. The first pool was spiked with 19.7 pg of exogenous testosterone per well (low spike) and the second pool was spiked with 39.5 pg of exogenous testosterone per well (high spike) to ensure the entire serial dilution of the pooled samples would fall on the curve. Each plasma pool was then serially diluted to create a 5-point curve. During validation, testosterone was extracted from 35 μ L of plasma using solid phase extraction with C18 columns (100 mg C18 material, Sep-Pak Vac 1cc, Waters, Milford, MA, USA), dried under nitrogen gas at 40°C, and reconstituted overnight with 250 μ L assay buffer (Caro et al., 2019). Recovery after extraction was 57.3% and 64.4% for low and high spiked curves, respectively.