

On a Pólya's inequality for planar convex sets. Ilias Ftouhi

▶ To cite this version:

Ilias Ftouhi. On a Pólya's inequality for planar convex sets.. 2021. hal-03334368v2

HAL Id: hal-03334368 https://hal.science/hal-03334368v2

Preprint submitted on 26 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON A PÓLYA'S INEQUALITY FOR PLANAR CONVEX SETS

ILIAS FTOUHI

ABSTRACT. In this short note, we prove that for every bounded, planar and convex set Ω , one has

$$\frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|} \leq \frac{\pi^2}{12} \cdot \left(1 + \sqrt{\pi} \frac{r(\Omega)}{\sqrt{|\Omega|}}\right)^2,$$

where λ_1 , T, r and $|\cdot|$ are the first Dirichlet eigenvalue, the torsion, the inradius and the volume. The inequality is sharp as the equality asymptotically holds for any family of thin collapsing rectangles.

As a byproduct, we obtain the following bound for planar convex sets

$$\frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|} \le \frac{\pi^2}{12} \left(1 + \frac{2\sqrt{2(6+\pi^2)} - \pi^2}{4+\pi^2} \right)^2 \approx 0.996613..$$

which improves Polyá's inequality $\frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|} < 1$ and is slightly better than the one provided in [13]. The novel ingredient of the proof is the sharp inequality

$$\lambda_1(\Omega) \leq \frac{\pi^2}{4} \cdot \left(\frac{1}{r(\Omega)} + \sqrt{\frac{\pi}{|\Omega|}}\right)^2,$$

recently proved in [4].

1. INTRODUCTION

Let Ω be an open set in \mathbb{R}^n with finite Lebesgue measure $|\Omega|$. We denote

$$T(\Omega) := \sup_{w \in H_0^1(\Omega) \setminus \{0\}} \frac{\left(\int_{\Omega} w dx\right)^2}{\int_{\Omega} |\nabla w|^2 dx}$$

the torsion of the set Ω , and

$$\lambda_1(\Omega) := \inf_{u \in H_0^1(\Omega) \setminus \{0\}} \frac{\int_{\Omega} |\nabla u|^2 dx}{\int_{\Omega} u^2 dx}$$

the fundamental frequency of Ω which corresponds to the first eigenvalue of the Laplace operator with Dirichlet boundary condition of the set Ω .

In 1951, G. Pólya [9] proved the following inequality:

$$\frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|} \le 1,$$

which was recently improved by M. van den Berg et al. in [13], where the authors prove the inequality

(1)
$$\frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|} \le 1 - \frac{2n\omega_n^{2/n}}{n+2}\frac{T(\Omega)}{|\Omega|^{1+\frac{2}{n}}},$$

where ω_n is the measure of the ball with radius 1 in \mathbb{R}^n . They also show, by using homogenization arguments, that the upper bound 1 is optimal in the class of open sets with finite measure, see [13, Theorem 1.2], but is not optimal for the class of bounded, <u>planar</u> and convex sets. Indeed, in this case they obtain the following improved estimate

(2)
$$\frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|} \le 1 - \frac{1}{11560} \approx 0.999913...$$

Date: October 26, 2021.

Key words and phrases. spectral inequalities, convex sets, Blaschke-Santaló diagrams.

see [13, Theorem 1.5]. The optimal upper bound is conjectured to be given by $\frac{\pi^2}{12}$, which is asymptotically attained by any sequence of thin collapsing rectangles.

Let us now state the main results of the present note.

Theorem 1. We have for every bounded planar convex set Ω :

(3)
$$\frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|} \le \frac{\pi^2}{12} \cdot \left(1 + \sqrt{\pi}\frac{r(\Omega)}{\sqrt{|\Omega|}}\right)^2$$

The inequality is sharp as it is asymptotically attained by any family of thin collapsing rectangles. Moreover, for every sequence (Ω_k) of planar convex sets such that $|\Omega_k| = 1$ and $\lim_{k \to +\infty} d(\Omega_k) = +\infty$, we have

$$\limsup_{k \to +\infty} \frac{\lambda_1(\Omega_k) T(\Omega_k)}{|\Omega_k|} \le \frac{\pi^2}{12}.$$

As a byproduct, we obtain the following slight improvement of inequality (2):

Corollary 1. We have for every bounded planar convex set Ω :

(4)
$$\frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|} \le \frac{\pi^2}{12} \left(1 + \frac{2\sqrt{2(6+\pi^2)} - \pi^2}{4+\pi^2}\right)^2 \approx 0.996613...$$

We provide the complete proofs of Theorem 1 and Corollary 1 in Section 2 and state some comments in Section 3.

2. Proofs of The main results

2.1. Proof of Theorem 1. Let Ω be a bounded, planar and convex set. We combine the following inequality proved in [4]:

(5)
$$\lambda_1(\Omega) < \frac{\pi^2}{4} \cdot \left(\frac{1}{r(\Omega)} + \sqrt{\frac{\pi}{|\Omega|}}\right)^2,$$

with the following Makai's inequality [7]:

$$T(\Omega) < \frac{1}{3} |\Omega| r(\Omega)^2.$$

We have

$$\frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|} < \frac{1}{|\Omega|} \times \frac{\pi^2}{4} \cdot \left(\frac{1}{r(\Omega)} + \sqrt{\frac{\pi}{|\Omega|}}\right)^2 \times \frac{1}{3} |\Omega| r(\Omega)^2 = \frac{\pi^2}{12} \cdot \left(1 + \sqrt{\pi} \frac{r(\Omega)}{\sqrt{|\Omega|}}\right)^2.$$

The inequality is sharp as it is an asymptotically an equality for any family of thin vanishing rectangles, see [12, Proposition 3.2].

Let us now consider (Ω_k) a sequence of planar convex sets such that $|\Omega_k| = 1$ and $\lim_{k \to +\infty} d(\Omega_k) = +\infty$. We have for every $k \in \mathbb{N}$:

$$0 < \frac{r(\Omega_k)}{\sqrt{|\Omega_k|}} \le \frac{1}{\sqrt{|\Omega_k|}} \frac{2|\Omega_k|}{P(\Omega_k)} \le \frac{2\sqrt{|\Omega_k|}}{2d(\Omega_k)} = \frac{1}{d(\Omega_k)} \underset{k \to +\infty}{\longrightarrow} 0,$$

where we used the classical inequalities $r(\Omega_k) \leq \frac{2|\Omega_k|}{P(\Omega_k)}$ (see [1]) and $2d(\Omega_k) \leq P(\Omega_k)$.

We then conclude by using inequality (3) that

$$\limsup_{k \to +\infty} \frac{\lambda_1(\Omega_k) T(\Omega_k)}{|\Omega_k|} \le \frac{\pi^2}{12}.$$

2.2. Proof of Corollary 1. Let us prove the inequality

(6)
$$\frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|} < 1 - \frac{\pi}{\frac{\pi^2}{4} \left(\frac{\sqrt{|\Omega|}}{r(\Omega)} + \sqrt{\pi}\right)^2 + \pi}$$

We have by inequality (1) (see [13, Theorem 1.1]):

$$\frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|} \le 1 - \pi \frac{T(\Omega)}{|\Omega|^2},$$

which is equivalent to the inequality:

$$\frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|} \leq 1 - \frac{\pi}{|\Omega|\lambda_1(\Omega) + \pi}.$$

Indeed:

$$\begin{aligned} \frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|} &\leq 1 - \pi \frac{T(\Omega)}{|\Omega|^2} \Leftrightarrow \frac{T(\Omega)}{|\Omega|} \left(\lambda_1(\Omega) + \frac{\pi}{|\Omega|}\right) < 1 \\ &\Leftrightarrow \frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|} < \frac{\lambda_1(\Omega)}{\lambda_1(\Omega) + \frac{\pi}{|\Omega|}} = 1 - \frac{\pi}{|\Omega|\lambda_1(\Omega) + \pi}. \end{aligned}$$

We then combine the inequalities (5) and (6) to conclude:

$$\begin{aligned} \frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|} &\leq \min\left(\frac{\pi^2}{12} \cdot \left(1 + \sqrt{\pi}\frac{r(\Omega)}{\sqrt{|\Omega|}}\right)^2, 1 - \frac{\pi}{\frac{\pi^2}{4}\left(\frac{\sqrt{|\Omega|}}{r(\Omega)} + \sqrt{\pi}\right)^2 + \pi}\right) \\ &\leq \max_{x \in \left(0, \frac{1}{\sqrt{\pi}}\right]} \min\left(\frac{\pi^2}{12} \cdot \left(1 + \sqrt{\pi}x\right)^2, 1 - \frac{\pi}{\frac{\pi^2}{4}\left(\frac{1}{x} + \sqrt{\pi}\right)^2 + \pi}\right) \\ &= \frac{\pi^2}{12} \left(1 + \frac{2\sqrt{2(6 + \pi^2)} - \pi^2}{4 + \pi^2}\right)^2.\end{aligned}$$

The second inequality is a consequence of the estimate $0 < \frac{r(\Omega)}{\sqrt{|\Omega|}} \leq \frac{r(B)}{\sqrt{|B|}} = \frac{1}{\sqrt{\pi}}$, where *B* is any ball of \mathbb{R}^2 . Let us now detail the computations leading to the last equality. We introduce the functions $f: x \mapsto \frac{\pi^2}{12} \cdot (1 + \sqrt{\pi}x)^2$ and $g: x \mapsto 1 - \frac{\pi}{\frac{\pi^2}{4}(\frac{1}{x} + \sqrt{\pi})^2 + \pi}$. The function *f* is increasing on $(0, \frac{1}{\sqrt{\pi}}]$ and *g* is decreasing on $(0, \frac{1}{\sqrt{\pi}}]$, moreover, $\lim_{x \to 0^+} f(x) = \frac{\pi^2}{12} < 1 = \lim_{x \to 0^+} g(x)$ and $f(\frac{1}{\sqrt{\pi}}) = \frac{\pi^2}{3} > \frac{1}{1+\pi^2} = g(\frac{1}{\sqrt{\pi}})$. Thus the function $x \mapsto \min(f(x), g(x))$ attains its maximum on $(0, \frac{1}{\sqrt{\pi}}]$ at the point x_0 such that $f(x_0) = g(x_0)$ (see Figure 1). It remains then to solve the equation f(x) = g(x) on $(0, \frac{1}{\sqrt{\pi}}]$. The equation f(x) = g(x) can be written in the algebraic form $ax^3 + bx^2 + cx + d = 0$, where a, b, c and d are explicit constants. We then notice that $-\frac{1}{\sqrt{\pi}}$ is a solution, which means that the equation is equivalent to $a(x + \frac{1}{\sqrt{\pi}})(x^2 + b'x + c') = 0$, where b' and c' are explicit constants. It remains then to solve the second order equation $x^2 + b'x + c' = 0$ that admits two solutions

$$x_1 = -\frac{\pi^{3/2}}{4 + \pi^2} - \frac{2\sqrt{\frac{2(6+\pi^2)}{\pi}}}{4 + \pi^2} < 0 \text{ and } x_0 = \frac{2\sqrt{\frac{2(6+\pi^2)}{\pi}}}{4 + \pi^2} - \frac{\pi^{3/2}}{4 + \pi^2} \in (0, \frac{1}{\sqrt{\pi}}]$$

Thus

$$\max_{x \in (0, \frac{1}{\sqrt{\pi}}]} \min(f(x), g(x)) = f(x_0) = g(x_0) = \frac{\pi^2}{12} \left(1 + \frac{2\sqrt{2(6 + \pi^2)} - \pi^2}{4 + \pi^2} \right)^2$$

This ends the proof.



FIGURE 1. Curves of the functions f and g.

3. Some comments

We close the paper by the following comments:

• The result of the paper is limited to the planar case because it mainly relies on the inequality

$$\lambda_1(\Omega) < \frac{\pi^2}{4} \left(\frac{1}{r(\Omega)} + \sqrt{\frac{\pi}{|\Omega|}} \right)^2,$$

which is obtained by combining the inequalities $\lambda_1(\Omega) < \frac{\pi^2}{4}h(\Omega)^2$ (proved in [8]), where *h* is the Cheeger constant, and $h(\Omega) \leq \frac{1}{r(\Omega)} + \sqrt{\frac{\pi}{|\Omega|}}$ (proved in [4]). Up to our knowledge, the last inequality is only known for the planar case and a generalization to higher dimensions seems to be difficult as the proof relies on the explicit characterization of the Cheeger constant of planar convex sets (see [5]).

• Let us denote $B_{r(\Omega)}$ a ball of radius $r(\Omega)$ inscribed in Ω . The inclusion $B_{r(\Omega)} \subset \Omega$ combined with the monotonicity of λ_1 for the inclusion yield the inequality

$$\lambda_1(\Omega) \le rac{j_{01}^2}{r(\Omega)^2}$$

where j_{01} denotes the first zero of the first Bessel function. We note that even if inequality (7) is better than (5) for sets that are close to the ball (we refer to [4, Section 5.1.1] for more details), it is not sufficient to improve the upper bound given in Theorem 1.

• We recall that the infimum of the functional $\frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|}$ on the class of open sets is zero (see [12, Remark 2.4]. As for the class of bounded convex subsets of \mathbb{R}^n (with $n \ge 2$), it is conjectured that the infimum is given by the constant $\frac{\pi^2}{12} \frac{6}{(n+1)(n+2)}$, see [11, Conjecture 4.2]. In [13], the authors prove the following estimates

$$\frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|} \ge \left(\frac{\pi}{2}\right)^2 \frac{1}{n^{n+2}(n+2)}, \quad \text{ for } n \ge 3,$$

and

(7)

$$\frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|} \ge \frac{\pi^2}{48}, \quad \text{for } n = 2,$$

that have been improved in [2, Remark 4.1], where the authors provide the lower bound

$$\frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|} \ge \left(\frac{\pi}{2}\right)^2 \frac{1}{n(n+2)}.$$

We also refer to [14, Theorems 1.4 & 1.5] for finer lower bounds in restricted classes of convex planar sets.

• An interesting tool to visualize the inequalities relating three functionals is the Blaschke–Santaló diagram. In our case, we are interested in the functionals: torsion T, fundamental frequency λ_1 and measure $|\cdot|$ in the case of planar convex sets, which leads to consider set of points

$$\mathcal{D} = \{ (\lambda_1(\Omega), T^{-1}(\Omega)) \mid \Omega \subset \mathbb{R}^2 \text{ is convex and } |\Omega| = 1 \}.$$

We note that this diagram has been theoretically studied in [6] and we refer to [3, 11] for results in the case of open sets. In Figure 2, we plot an approximation of the diagram \mathcal{D} obtained by randomly generating 10⁵ convex polygons (we used the algorithm presented in [10]) for which we compute the involved functionals, we also plot the curves corresponding to the best known inequalities relating the 3 functionals, namely:

– the Kohler-Jobin inequality

$$T(\Omega)\lambda_1(\Omega)^2 \ge T(B)\lambda_1(B)^2,$$

- the Faber-Krahn inequality

$$|\Omega|\lambda_1(\Omega) \ge |B|\lambda_1(B),$$

- the inequality (4)

$$\frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|} \le \frac{\pi^2}{12} \left(1 + \frac{2\sqrt{2(6+\pi^2)} - \pi^2}{4+\pi^2} \right)^2 \approx 0.996613...$$

We also plot in dashed line the curves corresponding to the conjectures

$$\frac{\pi^2}{24} < \frac{\lambda_1(\Omega)T(\Omega)}{|\Omega|} < \frac{\pi^2}{12}.$$



FIGURE 2. Blaschke-Santaló diagram \mathcal{D} and inequalities.

Acknowledgements:

We want to thank warmly the anonymous referee who allows us to improve the writing and clarity of this paper. This work was partially supported by the project ANR-18-CE40-0013 SHAPO financed by the French Agence Nationale de la Recherche (ANR).

ON A PÓLYA'S INEQUALITY FOR PLANAR CONVEX SETS

References

- T. Bonnesen and W. Fenchel. Theorie der konvexen Körper. Springer-Verlag, Berlin-New York, 1974. Berichtigter Reprint.
- [2] L. Brasco and D. Mazzoleni. On principal frequencies, volume and inradius in convex sets. NoDEA Nonlinear Differential Equations Appl., 27(2):Paper No. 12, 26, 2020.
- [3] G. Buttazzo and A. Pratelli. An application of the continuous Steiner symmetrization to Blaschke-Santaló diagrams. ESAIM Control Optim. Calc. Var., 27:Paper No. 36, 13, 2021.
- [4] I. Ftouhi. On the cheeger inequality for convex sets. Journal of Mathematical Analysis and Applications, 504(2):125443, 2021.
- [5] B. Kawohl and T. Lachand-Robert. Characterization of Cheeger sets for convex subsets of the plane. Pacific J. Math., 225(1):103-118, 2006.
- [6] I. Lucardesi and D. Zucco. On Blaschke-Santaló diagrams for the torsional rigidity and the first Dirichlet eigenvalue. Annali di Matematica Pura ed Applicata (1923 -), 2021.
- [7] E. Makai. On the principal frequency of a membrane and the torsional rigidity of a beam. In Studies in mathematical analysis and related topics, pages 227–231. Stanford Univ. Press, Stanford, Calif., 1962.
- [8] E. Parini. Reverse Cheeger inequality for planar convex sets. J. Convex Anal., 24(1):107–122, 2017.
- [9] G. Pólya and G. Szegö. Isoperimetric Inequalities in Mathematical Physics. Annals of Mathematics Studies, No. 27. Princeton University Press, Princeton, N. J., 1951.
- [10] V. Sander. http://cglab.ca/ sander/misc/ConvexGeneration/convex.html.
- [11] M. van den Berg, G. Buttazzo, and A. Pratelli. On relations between principal eigenvalue and torsional rigidity. Communications in Contemporary Mathematics, 0(0):2050093, 0.
- [12] M. van den Berg, G. Buttazzo, and B. Velichkov. Optimization problems involving the first Dirichlet eigenvalue and the torsional rigidity. In *New trends in shape optimization*, volume 166 of *Internat. Ser. Numer. Math.*, pages 19–41. Birkhäuser/Springer, Cham, 2015.
- [13] M. van den Berg, V. Ferone, C. Nitsch, and C. Trombetti. On Pólya's inequality for torsional rigidity and first Dirichlet eigenvalue. *Integral Equations Operator Theory*, 86(4):579–600, 2016.
- [14] M. van den Berg, V. Ferone, C. Nitsch, and C. Trombetti. On a Pólya functional for rhombi, isosceles triangles, and thinning convex sets. *Rev. Mat. Iberoam.*, 36(7):2091–2105, 2020.

(Ilias Ftouhi) FRIEDRICH-ALEXANDER-UNIVERSITÄT ERLANGEN-NÜRNBERG, DEPARTMENT OF MATHEMATICS, CHAIR OF APPLIED ANALYSIS (ALEXANDER VON HUMBOLDT PROFESSORSHIP), CAUERSTR. 11, 91058 ERLANGEN, GERMANY. Email address: ilias.ftouhi@fau.de