Do DOM optical parameters improve the prediction of copper availability in vineyard soils?

Frédéric Ouédraogo, Jean-Yves Cornu, Noémie Janot, Christophe Nguyen, Mahaut Sourzac, Edith Parlanti, Laurence Denaix

► To cite this version:

Frédéric Ouédraogo, Jean-Yves Cornu, Noémie Janot, Christophe Nguyen, Mahaut Sourzac, et al.. Do DOM optical parameters improve the prediction of copper availability in vineyard soils?. Environmental Science and Pollution Research, 2022, 10.1007/s11356-021-16361-5 . hal-03348974

HAL Id: hal-03348974

<https://hal.science/hal-03348974>

Submitted on 20 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Do DOM optical parameters improve the prediction of copper availability
in vineyard soils?**

Frédéric Ouedraogo^{1,2,*}, Jean-Yves Cornu¹, Noémie Janot¹, Christophe Nguyen¹, Mahaut
Sourzac², Edith Parlanti², Laurence Denaix¹

¹ ISPA, INRAE, Bordeaux Sciences Agro, 33140, Villenave d'Ornon, France

² University of Bordeaux, UMR CNRS 5805, EPOC, 33400, Talence, France

*Corresponding author: Frédéric Ouedraogo

Tel: 05 57 12 24 46

Fax : 05 57 12 25 10

E-mail : frederic.ouedraogo@inrae.fr

Abstract

Accumulation of copper (Cu) in soils due to the application of fungicides may be toxic for organisms and hence affect winegrowing sustainability. Soil parameters such as pH and dissolved organic matter (DOM) are known to affect the availability of Cu. In this study, we investigated the contribution of chromophoric and fluorescent DOM properties to the prediction of Cu availability in 18 organic vineyard soils in the Bordeaux winegrowing area (France). The DOM parameters, assessed through absorbance and fluorescence analyses, and proxies for Cu availability (total soluble Cu and free ionic Cu^{2+}) were measured in 0.01 M KCl extracts. Total soluble Cu (Cu_{KCl}) varied 23-fold while free ionic Cu^{2+} varied by a factor of 4,600 among the soils. DOC concentrations were similar among the soils, but the samples differed in the quality of DOM as assessed by optical spectroscopy. Multilinear regression models with and without DOM quality parameters were investigated to predict Cu availability. The best model for Cu_{KCl} successfully explained 83 % of variance and included pH, Cu_{T} and two DOM fluorescence quality indices, the FI fluorescence index, which distinguishes between microbial and higher plant origins, and the HIX humification index. For the prediction of Cu^{2+} , pH alone explained 88 % of variance and adding DOM parameters did not improve modelling. The two Cu availability proxies were related to pH. This study confirms the prominent role of pH in Cu availability and underlines the importance of DOM quality to better predict Cu solubility.

Keywords

Copper, organic matter, fluorescence, absorbance, vineyard soils, availability, modelling.

1. Introduction

Copper-based fungicides like Bordeaux mixture have been repeatedly applied in vineyards since the end of the 19th century to control downy mildew of grapevine, sometimes causing Cu to accumulate in the soils (Brun et al. 2001; Ballabio et al. 2018). In European soils, the median of total Cu concentrations is 13 mg kg⁻¹. In vineyards soils, this value reaches 49 and 91 mg kg⁻¹ in Europe and France, respectively (Ballabio et al. 2018). In old vineyards, Cu concentrations in the topsoil can reach several hundred mg Cu kg⁻¹, as widely reported in several European countries including France, Italy, Spain, Greece, Portugal, Czech Republic (Fernández-Calviño et al. 2009; Komárek et al. 2010; Fernández-Calviño et al. 2010; Flores-Álvarez et al. 2011; Kelepertzis et al. 2017). Despite the reduction of the authorized application rate to 4 kg Cu ha⁻¹ year⁻¹ (EFSA et al. 2018), Cu-based fungicides continue to be used, particularly in organic viticulture for which there is currently no alternative that offers equally effective protection (Andrivon et al. 2018).

Cu is an essential micronutrient for all living organisms. Microorganisms need trace levels of Cu for their growth because it is a cofactor in enzymatic reactions (Trevors and Cotter 1990). In plants, copper plays key roles in cell wall metabolism and in photosynthetic electron transport (Yruela 2009). A Cu deficiency can affect vital functions in plant metabolism. However, excess copper can cause phytotoxicity through the formation of oxygen radicals which damage cells and inactivate protein structure, or by disturbing photosynthetic activity due to the photoinhibition induced by excessive Cu (Yruela 2005). In calcareous soils, excess Cu can also lead to Fe deficiency in graminaceous species such as durum wheat (Michaud et al. 2007) since Cu(II) has sufficient affinity for phytosiderophores to divert them from their initial objective of mobilising Fe(III). All these problems may affect plant growth and reduce agricultural productivity. Cu phytotoxicity has been reported in durum wheat grown in former vineyard soils with total Cu concentrations beginning at 114 mg kg⁻¹ (Michaud et al. 2007). Exposure to

Cu at soil concentrations above 16 mg Cu kg⁻¹ in the soil has also been shown to have a serious impact on earthworm *Eisenia fetida* reproduction by reducing the production of cocoons (Helling et al. 2000). Fernández-Calviño et al. (2010) reported that microbial enzymatic activity (phosphatase) in Portuguese vineyard soils decreased drastically above 250 mg Cu kg⁻¹. A recent meta-analysis of Cu in vineyard soils showed that microbial respiration was reduced by half above a stock of 200 kg Cu ha⁻¹ (Karimi et al. 2021). Excess copper may thus cause ecotoxicity in vineyard soils, although the ecotoxic thresholds vary widely depending on certain soil parameters like organic matter (OM) content or pH. Total Cu inducing 50 % of biomass decrease (EC₅₀) is a widely used indicator in ecotoxicity studies. It has been shown that EC₅₀ of lettuce and tomatoes increases with increasing OM content or decreasing pH (Recatalá et al. 2012; Sacristán et al. 2016). Cu toxicity may therefore be linked not only to total Cu amounts but also to certain bioavailable forms of Cu in the soils.

The bioavailability of metals is defined as the fraction of its environmentally available compound an organism takes up through physiologically driven processes (ISO 2006). Environmental availability is defined as the availability of metals in the soil measured using chemical methods (Harmsen 2007). Sauvé et al. (1996) showed that Cu concentrations in the tissues of several plants were closely linked to Cu in solution, more specifically to free Cu²⁺ activity. The free ion Cu²⁺ is thought to be the bioavailable form, and this hypothesis is supported by other authors who reported that Cu toxicity for different organisms (algae, earthworms, zooplankton) in soils and aquatic systems depended strongly on free Cu²⁺ activity (Verweij et al. 1992; Qiu et al. 2013; Tait et al. 2016).

The concentration of Cu in soil solutions and Cu²⁺ activity are controlled by soil properties such as total Cu content, pH and soil organic matter (SOM) content (Sauvé et al. 1997; Sauvé et al. 2000). pH is known to affect Cu solubility (Temminghoff et al. 1994), but also copper speciation in solution (Bravin et al. 2009). Cu biogeochemistry is controlled to a large extent

by its binding to organic matter, since SOM contains acidic functional groups (mainly phenolic and carboxylic groups) which have a particular and strong affinity for Cu (Manceau and Matynia 2010). Organic matter supplied to soils in different forms (fresh materials, compost, manure, sludge) has been shown to significantly reduce Cu availability in its soluble form (Navel and Martins 2014; Ferreira et al. 2018; Gonzaga et al. 2020) and its ionic form (Oustriere et al. 2016). Dissolved organic matter (DOM) plays a key role in the fate of metals in the soil. While SOM immobilises Cu through sorption onto solid phases (Sauvé et al. 2000), DOM increases Cu solubility by forming Cu complexes in soil solution. However, the free Cu fraction may also decrease due to complexation with DOM. Cu mobilisation from the solid phase into solution has been reported to be exacerbated with increasing aromatic DOM (Amery et al. 2007; Amery et al. 2008; Araújo et al. 2019). What is more, prediction of Cu in soil leachates was improved when the DOM aromaticity was taken into account in geochemical modelling (Amery et al. 2008) rather than only considering a percentage of reactive DOM (Djae et al. 2017). Phenolic and O-functional groups of aromatic molecules have high affinity for Cu binding and are highly stable, which may facilitate its solubilisation by DOM (Fuentes et al. 2013). These findings clearly underline the importance of DOM characteristics in the control of Cu dynamics in soil solutions.

The capacity of DOM to complex Cu and to promote its mobilisation from the solid phase not only depends on DOM quantity, but also on DOM quality (Baken et al. 2011; He et al. 2016; Kikuchi et al. 2017). DOM quality can be described through its optical properties assessed by Ultraviolet visible (UV-Vis) absorption and excitation-emission matrix (EEM) fluorescence spectroscopy (Jaffé et al. 2014). In the soil literature, DOM is largely characterised by its specific UV-absorbance index at 254 nm ($SUVA_{254}$), which is related to its aromaticity (Weishaar et al. 2003). But the measurements of $SUVA_{254}$ may be biased due to the presence of colloids or iron at high concentrations (Weishaar et al. 2003). Other spectroscopic indices

are widely used in studies on aquatic environments to characterise DOM (Yang and Hur 2014; Derrien et al. 2017) and its interactions with Cu (He et al. 2016). Another UV-Vis absorption index is the spectral slope ratio S_R which is inversely proportional to molecular size (Helms et al. 2008). Fluorescence indices include the humification index HIX, which expresses the degree of DOM condensation (Zsolnay et al. 1999), the biological index BIX, used to assess the recent production of autochthonous components (Huguet et al. 2009), and the fluorescence index FI which makes it possible to distinguish microbial versus higher plant origin of DOM (McKnight et al. 2001). In aquatic media, concentrations of metals (Cu, Zn, Cd) in sediment solutions were shown to be related to DOM aromaticity (HIX) and to the autochthonous (BIX) or microbial (FI) origins of molecules and not to DOM molecular size (S_R) (He et al. 2016; Zhang et al. 2020).

Up to now, only a few studies combined all these tools to describe DOM in contrasted agricultural soils (Romero et al. 2017; Li et al. 2019; Sun et al. 2020) and their effects on metal speciation in soil solution (Guo et al. 2015) or sediments (Xu et al. 2019). To our knowledge, only one paper reported the use of these spectroscopic indices for vineyard soils exposed to herbicide glyphosate and Cu-based fungicides (Daouk et al. 2015). The main goal of the present study was thus to examine the contribution of quality of DOM in predicting Cu availability in organic vineyard soils. We hypothesized that taking the DOM optical indices into account would improve the prediction of Cu environmental availability, assessed with total soluble Cu or free Cu^{2+} .

2. Material and methods

2.1 Soil sampling and analyses

Eighteen vineyard topsoils were sampled in the vine inter-rows at a depth of between 0 and 15 cm in March 2019 in the region of Bordeaux (France), located in the AOC (i.e., controlled

141 designation of origin) zones of ‘*Bordeaux supérieur*’, ‘*Blaye Côtes de Bordeaux*’ and ‘*Graves*’.

142 All the vineyards sampled are organic and the only fungicide treatment they receive is Cu. The
143 fungicide applied was mainly Bordeaux mixture (mixture of copper sulphate and lime).

144 The 18 soils were contrasted in terms of organic matter and pH based on the pedological
145 bedrock and agronomic practices. After sampling, the fresh soils were sieved to 4 mm and kept
146 at 4 °C before analysis to minimise the effects of sampling on dissolved organic carbon (Jones
147 and Willett 2006).

148 Soil pH (pH_{water}), total copper (Cu_T), organic carbon (C_{org}) and total nitrogen contents were
149 determined on separate air-dried soils. pH_{water} was determined by potentiometry in distilled
150 water in a 1:2.5 (w/v) soil-to-solution ratio (NF ISO 10390). Cu_T was determined after *aqua*
151 *regia* microwave-assisted digestion (Anton Paar Multiwave PRO Microwave Reaction
152 System®). Aliquots (0.5 g) of crushed dried sample were placed in a microwave Teflon flask
153 with an acid mix $\text{HNO}_3\text{:HCl}$ (3.75:1.25, v:v) (Aristar® for trace analysis, VWR Chemicals).
154 Extracts were obtained after centrifugation at 1,300 W at 180 °C for 20 minutes and filtration
155 on a 2- μm Whatman cellulose acetate filter. The concentration of Cu in the extracts was
156 measured by ICP-OES (Agilent®). A certified reference material (BCR-143R, 98 % Cu
157 recovery rate) was used to validate extraction and quantification.

158 C_{org} and N contents in soils were assessed by dry combustion (950 °C) with a CN analyser
159 (Flash EA 1112 series®) after carbonates were removed with HCl (NF ISO 10694). An organic
160 soil (ISE 973) was used as control.

161 2.2 Extraction of soil solutions

162 Before the soil solution was extracted, the wet soils were all adjusted to 80 % water holding
163 capacity (WHC) and incubated at 20 °C for one week to avoid a possible microbial flush and
164 its effects on Cu solubility and speciation (Cornu et al. 2007).

Soil solutions were extracted as follows: 15 g of wet soil were shaken with 30 mL of 0.01 M KCl (purity 99.5 %, Merck®) with 1:2 w/v (soil-to-solution ratio) at 35 rev. mn^{-1} for 2 h. The soil suspensions were centrifuged at 4,600 g for 10 min. The supernatants were collected and filtered through 0.2 μm cellulose acetate filters. Extractions were performed in triplicate. The extracted soil solutions were stored at 4 °C in the dark before spectroscopic analysis, which was performed less than seven days after extraction.

The 0.01 M KCl extraction was chosen for two reasons: (i) to be in the range of common ionic strength for soil pore water and (ii) to avoid DOM coagulation which may occur in the case of CaCl_2 extraction (Römken and Dolfing 1998). The 1:2 w/v ratio was used because lower extraction ratios tend to overestimate the concentrations of DOC recovered (Jones and Willett 2006).

2.3 Analyses of soil extracts

2.3.1 Total dissolved metals and Cu^{2+} concentrations

The pH of the extracts (pH_{KCl}) was measured immediately after extraction by potentiometry using a microelectrode (E16M331, Radiometer Analytical®). The total concentrations of dissolved Ca, Cu, Fe, Mn and Zn were determined by ICP-OES (Agilent ®) in a 5-mL aliquot acidified with 2 % HNO_3 (Aristar® for trace analysis purity 69 %, VWR Chemicals®). Free Cu activity was measured in non-acidified 5-mL aliquots using a cupric ion-selective electrode (ISE, 9629BNWP, Thermo Scientific Orion®). The ISE was calibrated in Cu solutions buffered with iminodiacetic acid and potassium phthalate with pCu ($= -\log_{10} a_{\text{Cu}^{2+}}$) ranging from 5.5 to 13 as described in (Bravin et al. 2009). The calibration curve ($R^2=0.98$, $n=67$) showed 85 % slope efficiency compared to the theoretical Nernstian slope (**Fig. S1**). The concentrations of free Cu^{2+} were determined from pCu values and from the Cu^{2+} activity coefficient, calculated from the extended Debye-Hückel equation (Ritsema 1993).

2.3.2 Dissolved organic matter parameters

The concentrations of dissolved organic carbon (DOC) in the soil extracts were determined using a TOC analyser (TOC-VCPH, Shimadzu®, Japan). The DOC was calculated as the difference between total carbon (TC) and inorganic carbon (IC) (DOC=TC - IC). The analyser was calibrated with a solution prepared from sodium carbonate, sodium bicarbonate, potassium hydrogen phthalate, and the 0.01 M KCl was used as the zero point in the calibration.

Absorbance indices

Soil solution UV-visible absorbance was measured using a Jasco® V-560 spectrophotometer, scanning from 210 to 700 nm with a 0.5 cm path length quartz cuvette. A 0.01 M KCl solution was used as a blank. Several indices were determined from absorbance spectra to characterise the DOM in the extracts. The specific UV absorbance at 254 nm (SUVA₂₅₄, L gC⁻¹ cm⁻¹) was calculated as follows:

$$\text{SUVA}_{254} = \frac{A_{254} \cdot 1000}{b \cdot \text{DOC}}$$

where A₂₅₄ is the measured absorbance at 254 nm (dimensionless) (**Fig. S2**), b is the path length (cm) and DOC the organic carbon concentration (mg L⁻¹).

The S_R index was calculated as the spectral slope ratio over 275-295 nm to 350-400 nm (Helms et al. 2008).

Fluorescence indices

The excitation-emission matrix (EEM) fluorescence spectra were recorded in a 0.5 cm quartz cuvette, thermostatted at 20 °C in the cell holder, using an Aqualog spectrofluorometer (Horiba Scientific®). Excitation scans were performed using a double-grating monochromator, a 150-Watt Xenon lamp with a 5 nm bandpass, and 2-s integration time at wavelengths of 240–800 nm at 5-nm intervals. Emission spectra were collected with high CCD detector gain at

approximately 0.58 nm (1 pixel) intervals at wavelengths of 245-830 nm. A blank EEM was acquired daily from the 0.01 M KCl solution used for soil extraction and subtracted from EEM spectra of all the samples to eliminate scattering peaks. EEM spectra were corrected for instrumental biases. When the maximum intensity of the absorbance spectrum was higher than 0.1, the samples were diluted with 0.01 M KCl to avoid inner filter effects. Fluorescence data are expressed in Raman units (R.U) after normalization to the area under the Raman peak of an ultrapure water blank (Milli-Q, Millipore®) at an excitation wavelength of 350 nm. TreatEEM software (<https://sites.google.com/site/daromasoft/home/treateem>) was used to remove remaining scatter peaks in EEMs and to calculate the fluorescence indices.

Several indices were obtained from fluorescence analysis. As defined by Zsolnay et al. (1999), the humification index HIX was calculated as the ratio H/L of the area under the curve for two spectral regions (between emission wavelengths 435 nm and 480 nm for H and between 300 nm and 345 nm for L) from the emission spectrum scanned for excitation at 254 nm (**Fig. S3**). The higher the HIX, the higher the degree of condensation and aromaticity. The biological index BIX was determined as the ratio of fluorescence intensity at an emission wavelength of 380 nm to that at 430 nm, at an excitation wavelength of 310 nm (Huguet et al. 2009). High BIX values are related to predominantly recent production of DOM linked to biological activity (Huguet et al. 2009). The fluorescence index FI was calculated as the ratio of the fluorescence intensity at emission wavelength of 450 nm to that at 500 nm while excitation was set at 370 nm (McKnight et al. 2001). FI is an indicator of the source of DOM, low values are associated with vascular-plant-derived compounds, while high values indicate microbial origin or microbially processed terrestrial DOM (Cory and McKnight 2005; Broder et al. 2017).

2.4 Statistical analyses

All statistical analyses were performed with R software version 4.0.2 (R core Team 2020). First, principal component analysis (PCA) was used to visualise correlations between variables and

possibly groups of soils. PCA was performed using the ‘FactomineR’ package (Husson et al. 2008). For the PCA, all the variables were log-transformed except pH_{water} and pH_{KCl} . Second, a correlation matrix was performed to investigate the relationships between the same variables using the R ‘corrplot’ package (Wei and Simko 2021). The Pearson correlation coefficients were considered as statistically significant at $p < 0.01$.

Multi-linear log-log regressions were used to identify and rank the contributions of soil variables to the prediction of Cu availability proxies (Cu_{KCl} and Cu^{2+}). Multi-linear regressions were performed using the ‘car’ package (Fox and Weisberg 2019). Each replicate was considered as one sample. For Cu_{KCl} prediction, soil and solution variables were entered as predictors (Cu_{T} , pH_{water} , C:N, C_{org} , SUVA_{254} , S_{R} , DOC, HIX, BIX, FI). Only solution variables were used for Cu^{2+} prediction (Cu_{KCl} , pH_{KCl} , SUVA_{254} , S_{R} , DOC, HIX, BIX, FI). The variables (except pH_{water} and pH_{KCl}) were log-transformed before the computation. A total of 2,047 and 255 models with one to maximum four predictors (to avoid overparameterization) were exhaustively tested for Cu_{KCl} and Cu^{2+} predictions, respectively. The best models had the maximum R^2_{adj} (goodness of fit) with a minimum number of predictors (parsimony). The selected models also had a variation inflation factor (VIF) below 2.3 to avoid multicollinearity. The root mean square error (RMSE) was calculated to compare the accuracy of the models including or not DOM parameters.

2.5 Modelling Cu^{2+} activity

Cu speciation in solution was predicted with ORCHESTRA software (Meeussen 2003) using the DOM ion-binding model NICA-Donnan (Kinniburgh et al. 1996; Koopal et al. 2005). The objective was to compare simulations of amounts of free Cu^{2+} with the corresponding experimental measurements. The input parameters used for the calculations were pH_{KCl} , total DOM, Cu^{2+} , Ca^{2+} , Zn^{2+} , Fe^{3+} and Mn^{3+} concentrations. Cl^- and K^+ concentrations were set to 0.01 M, i.e., their concentration in the extraction solution. DOM concentrations were calculated

assuming that DOM contained 50 % of C. Activity coefficients were calculated using the Davies equation. Fe activity was calculated by assuming its equilibrium with $\text{Fe}(\text{OH})_3$ ($\log K_s = 3.19$). Equilibrium reactions were described taking the equilibrium with atmospheric CO_2 ($p\text{CO}_2 = 3.9 \times 10^{-4}$ atm) into account.

Several hypotheses regarding DOM composition were examined: (i) 100 % humic acids (HA), (ii) 100 % fulvic acids (FA), or an inert DOM pool and an active DOM pool comprised of (iii) 65 % FA, or (iv) 30 % HA and 30 % FA. In each case, proton and metal binding parameters were set at their generic value (Milne et al. 2001, 2003). Sixty-five percent is the approximation generally used to determine the amount of free metal cations in soil solutions (Weng et al. 2002, Djae et al. 2017). In the second step, free Cu was optimised by adjusting the reactive DOM pool (noted rDOM) in each sample to match the experimental measurements.

3. Results and discussion

3.1 Soil properties

The main soil parameters are listed in **Table 1**.

Total concentration of Cu in the soil (Cu_T) ranged from 28 to 238 mg kg^{-1} with a median of 107 mg kg^{-1} . In the Aquitaine region of France, large quantities of Cu-based fungicides are used to prevent downy mildew caused by the wet climate. Accordingly, El Hadri et al. (2012) found that Cu_T in soils in the Aquitaine region ranged from 0.5 to 491 mg kg^{-1} , with maximum values found in vineyard soils. All the vineyard soils in our study fell within this range. It is worth noting that Cu_T concentrations may be very different in the same location, mainly depending on the age of the vineyard (Fernández-Calviño et al. 2009). This may explain the wide range of Cu_T (i.e., 8-fold) observed in the present study.

The pH_{water} of the soils ranged from 6.3 to 8.3 (**Table 1**). The most alkaline soils were associated with the highest carbonate contents (**Table 1**). The organic carbon content (C_{org}) ranged from 5 to 31 g kg^{-1} . Vineyard soils are generally poor in organic carbon. Coll et al. (2011) reported C_{org} values ranging from 10 to 13 g kg^{-1} for vineyard soils in the Languedoc-Roussillon region (South of France). Our results are also similar to those obtained by Parat et al. (2002) ($11 < \text{C}_{\text{org}} < 45 \text{ g kg}^{-1}$) in a study conducted in Burgundy viticultural region (east-central France).

The C:N ratio varied between 10.3 and 17.4 (**Table 1**). The C:N ratio varied between 10.3 and 17.4 (**Table 1**). This ratio provides information on the rate of SOM mineralisation (high for values below 12) and also on the carbon storage capacity of the soil (Swift et al. 1979).

3.2 Elemental composition of the soil extracts and Cu availability

Mean Fe, Mn, Ca and Zn concentrations are listed in **Table S1** in supplementary information.

The pH of the soil extracts (pH_{KCl}) fluctuated between 5.9 and 7.6 (**Table S1**).

Mean concentrations of DOC in KCl extracts ranged only from 15 to 28 mg L^{-1} (median 18) (**Table S1**). This corresponds to 0.08 and 0.33 % of total C_{org} , illustrating the low solubility of SOM. Furthermore, Zsolnay (1996) reported that DOC is the most mobile and reactive fraction of C_{org} .

The mean total concentration of Cu in the extracts (Cu_{KCl}) ranged from 0.11 to 2.64 μM , i.e., a factor of 23 (**Table S1**), with a median of 0.3 μM . Many studies in the literature used salt solutions - generally CaCl_2 - to extract Cu and considered it to be a correct surrogate for soil pore water Cu (Sauvé et al. 1997; Chaignon et al. 2003). Our Cu_{KCl} values were similar to those found by Djae et al. (2017) (3×10^{-2} - 3.7 μM Cu) with 2 mM KNO_3 / $\text{Ca}(\text{NO}_3)_2$ extractions. Similar ranges of total Cu assessed in CaCl_2 extracts were found in the literature: 0.11 – 2 μM (Sauvé et al. 1996), 0.01 – 5 μM (Sauvé et al. 1995).

309 The mean Cu^{2+} concentrations varied between 2.1×10^{-4} and $0.95 \mu\text{M}$, i.e., by a factor of 4,600
 310 (**Table S1**). This variability shows that our samples were more contrasted in terms of Cu^{2+} than
 311 Cu_{KCl} . The highest Cu^{2+} values were found in the most acidic soils. The percentage of free Cu,
 312 defined as the ratio of concentrations in Cu^{2+} and Cu_{KCl} (fCu^{2+}), ranged from 0.07 % to 50 %
 313 (**Table S1**). Our results are in agreement with Djae et al. (2017) who found Cu^{2+} concentrations
 314 between 3.10^{-7} and $2 \mu\text{M}$ and fCu^{2+} ranged from 0.01 % to 44 % for Cu-contaminated soils.
 315 But in moderately contaminated vineyard soils, Cornu et al. (2019) found Cu^{2+} of between 6×10^{-5} – $0.3 \mu\text{M}$ and fCu^{2+} of between 0.002 % - 10 %. These results reveal dramatic differences
 316 based on free Cu^{2+} , even within a collection of soils with the same source of Cu contamination.
 317
 318 Soil pore water Cu has been used as a proxy for Cu phytotoxicity (Michaud et al. 2008),
 319 assuming that few complex forms of Cu present in the solution play a role in Cu uptake by plant
 320 roots. The toxicity threshold found in the literature for various plants ($3.94\text{--}9.55 \mu\text{M}$) are all
 321 higher than the maximum Cu_{KCl} ($2.64 \mu\text{M}$) found in the present study (**Table 2**), suggesting
 322 that our 18 vineyard soils do not *a priori* represent a risk for grapevine and also possibly maize,
 323 grapevine and sunflower production (Ouzounidou 1995, Juang et al. 2012, Kolbas et al. 2014).
 324 Beyond total soluble Cu, the bioavailable form of Cu is generally considered to be its ionic
 325 species Cu^{2+} (Sauvé et al. 1998), so Cu^{2+} concentrations would be a far more reliable tool to
 326 assess Cu toxicity for organisms. In the present study, the Cu^{2+} measured in some KCl extracts
 327 was the same as that which caused a 50 % reduction ($\text{EC}_{50} = 6.3 \times 10^{-2} \mu\text{M}$) in root growth in
 328 durum wheat (Bravin et al. 2010) (**Table 2**). This EC_{50} value was found at $2.2 \mu\text{M}$ Cu^{2+} for
 329 cucumber (Kader et al. 2016). To our knowledge, no such toxicity indicator assessed via ionic
 330 Cu^{2+} has been established for grapevine. Four soils of our panel (**Table S1**) may be problematic
 331 if converted from grapevine to widely consumed crops like wheat. Qiu et al. (2013) reported
 332 that Cu^{2+} concentration was also an appropriate proxy to assess Cu toxicity for earthworms and
 333 found a EC_{50} of $1 \mu\text{M}$ Cu^{2+} for *Eisenia fetida* mortality (**Table 2**). In our panel of soils, this

suggests that one soil represents a potential toxic risk for this earthworm species (**Table 2, Table S1**).

Overall, the Cu^{2+} and Cu_{KCl} levels in the soils sampled in this study cannot be considered completely risk-free for various organisms, and may therefore affect the sustainability of the vineyards. The Cu^{2+} concentrations in about one-quarter of the soils were above the toxicity thresholds reported in the literature and may be toxic for organisms. It is thus important to examine and understand the factors that control Cu availability and potential toxicity.

3.3 Diversity of soils in terms of DOM properties

We calculated several spectroscopic indices to assess DOM quality (**Fig. S4**). SUVA_{254} values varied between 5 and 61 $\text{L g C}^{-1} \text{ cm}^{-1}$ (**Fig. 1, Table S1**). This range is similar to that reported in other studies on agricultural soils: 6–65 $\text{L g C}^{-1} \text{ cm}^{-1}$ (Amery et al. 2007); 27–46 $\text{L g C}^{-1} \text{ cm}^{-1}$ (Amery et al. 2008); 8–58 $\text{L g C}^{-1} \text{ cm}^{-1}$ (Supriatin et al. 2015). DOM is considered to be more aromatic when the SUVA_{254} index is higher (Weishaar et al. 2003). It has been reported that SUVA_{254} values below 30 $\text{L g C}^{-1} \text{ cm}^{-1}$ suggest that DOM is composed mainly of hydrophilic substances (Yeh et al. 2014), which is the case of about 80 % of our samples. S_R values varied three-fold between 0.42 and 1.28 (**Fig. 1, Table S1**), which is the same range as those reported by Tang et al. (2019) for Chinese agricultural soils (0.36–1.7). The S_R index is inversely related to molecular size, high S_R values indicate low molecular weight DOM (Helms et al. 2008) while low S_R values indicate high molecular weight and highly condensed organic molecules (Gao et al. 2017).

Fluorescence EEM spectra (**Fig. S4**) were acquired to describe heterogeneities in DOM in the samples more precisely by calculating indices such as FI, HIX and BIX. The FI values ranged from 1.10 to 1.34 (median 1.22) (**Fig. 1, Table S1**). FI values are usually used to distinguish

sources of DOM. McKnight et al. (2001) found that low FI values (~1.3) correspond to higher plant-derived molecules, while high values (~1.9) indicate microbial or microbially-processed DOM. This suggests that the majority of DOM collected from our samples originated from vascular plants, and that microbial contributions were limited. The BIX index is used to assess the relative production of recent DOM and usually ranges from 0.5 to 1. In the present study, about 75 % of the samples had BIX between 0.6 and 0.7, indicating a relatively low contribution of fresh autochthonous DOM substances (**Table S1**), while the other 25 % had BIX between 0.7 and 0.9, suggesting fresh DOM production by organisms (Huguet et al. 2009; He et al. 2016). In our samples the HIX values were contrasted, with almost 10-fold variation from 1.31 to 10.33 (median 5.75) (**Fig. 1, Table S1**). Two of the soils (D and L) were characterised by very low HIX values (**Table S1**) due to the presence of tyrosine-like fluorescence at an excitation wavelength of 240-260 nm and emission of 300-340 nm (**Fig. S4a**). This tyrosine-like fluorescence band was absent from all other EEM spectra that were similar to that of soil B (**Fig. S4b**). HIX stands for degree of humification or condensation of DOM (Zsolnay et al. 1999), but has also been used to characterise DOM aromaticity (Ohno et al. 2007). Our HIX results are in the same range as those reported by Daouk et al. (2015) for Swiss vineyard soils (0.1-7.0) and as those reported by Tang et al. (2019) (2 to 10). However, HIX values varying from 10 to 30 have been reported in several arable soils (Kalbitz et al. 2003). Compared to the latter range, our soils had low HIX values and can be characterized as weakly-humified.

3.4 Relationships between the variables

We performed a principal component analysis (PCA) of all 18 soil extracts to describe the relationships between the solution variables and to identify homogeneous groups. 64% of total variance is explained by the first two principal components (**Fig. 2a**). The first principal component revealed 44.5 % variance of data and was positively associated with Cu_{KCl} , Cu^{2+} , Fe concentrations, and SUVA_{254} , and was negatively associated with pH_{water} , pH_{KCl} and HIX.

The second principal component explained 19.2 % of variance and was positively associated with C_{org} , the C:N ratio, and negatively associated with FI and BIX. Two groups can be distinguished on the principal plan plot (**Fig. 2b**): group 1 is composed of the eight most alkaline soils ($7.8 < pH_{water} < 8.3$), while group 2 is composed of the 10 most acidic soils ($6.3 < pH_{water} < 7.5$). The differences between the groups are mainly explained by the pH_{water} values (PCA correlation coefficient with Component 1: -0.92; with Component 2: 0.03). On the first axis, the soils in group 1 share high values of pH and HIX, and low values of Cu_{KCl} , Fe, and Cu^{2+} . The opposite trend was observed in group 2. On the second axis, soils in group 2 have more contrasted FI and BIX than those in group 1. Indeed, in group 1, FI values ranged from 1.20 to 1.31; and BIX values from 0.67 to 0.75. Comparatively, these indices show a larger range in group 2, with 1.10 - 1.34 for FI and 0.59 - 0.86 for BIX. Group 1 is composed of alkaline soils with low Cu concentrations in solution and relatively highly-humified DOM, while group 2 is composed of more acidic soils with high concentrations of Cu_{KCl} and Cu^{2+} , and contrasted contributions of microbial DOM.

Pearson's correlation coefficients were analysed to describe the links between the variables measured in the soils and in the extracts (**Fig. 3**). pH_{water} and pH_{KCl} were strongly and positively correlated ($r=0.94$, $p < 0.01$). Total dissolved Fe and Cu concentrations were negatively correlated with pH (**Fig. 3**), with $r=0.77$, 0.73 ($p < 0.01$), respectively. **Fig. 4a** reports total Cu in extracts as a function of total Cu in the soils. We found no immediate relationship between Cu availability and the level of contamination. Cu_{KCl} concentrations globally increased with decreasing soil pH (**Fig. 4b**). This result confirms the fact that metals are more soluble in acidic soils, as reported by McBride et al. (1997). None of the Fe and Cu concentrations were significantly correlated with DOC (**Fig. 3**). Cu^{2+} was strongly linked to Cu_{KCl} ($r=0.82$, $p < 0.01$) (**Fig. 3, Fig. 4c**), showing that Cu solubilisation affects its concentration of free ions. A slightly positive relationship was found between pH_{KCl} and HIX (**Fig. 3**), suggesting that the most

humified DOM was present in the most alkaline samples, as confirmed by the results of the PCA.

Cu^{2+} concentrations and $f\text{Cu}^{2+}$ were strongly linked to the pH of the extracts, with correlation coefficients of -0.94 and -0.91 respectively (**Fig. 3, Fig. 4d**), suggesting that Cu^{2+} decreased with increasing pH. The strong relationship between pH and Cu^{2+} has already been reported (McBride and Blasiak 1979, Sauvé et al. 1995, Laurent et al. 2020), confirming that pH is the main factor controlling Cu speciation in solution. When pH increases, DOM is more negatively charged due to the deprotonation of its functional groups. This promotes its complexation with Cu, leading to a drastic decrease in Cu^{2+} activity in solution (Bravin et al. 2009; Ren et al. 2015). Conversely, at low pH, competition for DOM binding sites is high between Cu^{2+} and H^+ , which promotes increasing Cu^{2+} activity. A pH above 6.5 seems to be the safety threshold for the soils in our collection in terms of Cu toxicity when Cu^{2+} is used as a proxy (**Fig. 4d**).

No tangible relationship was found between C_{org} values and DOC concentrations in the extracts (**Fig. 3**), meaning that the solubilisation rates of C_{org} may differ with the soil conditions or properties. None of DOM optical indices was significantly linked to DOC concentrations in the KCl extracts, indicating that DOM quality was not related to the quantity of DOM in solution. No significant relationship was found between the S_R and the SUVA_{254} indices. SUVA_{254} and HIX indices were negatively linked in this study ($r = -0.71$, $p < 0.01$) (**Fig. 3**). This negative relationship is all the more surprising as these indices are both related to DOM aromaticity. In the literature, SUVA_{254} and HIX are generally positively correlated (He et al. 2016; Tang et al. 2019). However, such unexpected differences between these two indices have already been observed (Inamdar et al. 2011; Broder et al. 2017). Considering the differences in pH between soils, and given that fluorescence is pH-dependent, one may wonder if the variations observed in these two indices are not affected by the effect of pH on spectroscopic analyses. However, an impact on fluorescence analyses is unlikely in our range of soil pH, and would, in any case,

be less than differences due to the sources and composition of DOM (Hudson et al. 2007). The presence of protein-like fluorescence was observed in EEM spectra of only two soils (D and L) and cannot explain the negative correlation between HIX and SUVA₂₅₄. Moreover, it has been reported that SUVA₂₅₄ measurements may be subject to some bias since Fe, colloids, and other constituents also absorb at 254 nm (Weishaar et al. 2003; Yang and Hur 2014). Since there were huge differences in Fe concentrations among our samples (i.e., varying by a factor of 12, **Table S1**), the SUVA₂₅₄ results in this study should be interpreted with caution. In a collection of samples with such contrasted concentrations of dissolved Fe as those measured in the present study, it would therefore not be appropriate to use the SUVA₂₅₄ index to assess DOM aromaticity, and HIX appears to be a more reliable tool for that purpose. No significant relationship was found between HIX and S_R, suggesting that, in our soil extracts, DOM condensation may not be directly related to its molecular size. Moreover, the negative correlation between HIX and BIX (**Fig. 4b**) ($r = -0.82$, $p < 0.01$) indicates that the most aromatic DOM is associated with older molecules derived from higher plants (Tang et al. 2019), which is quite consistent. No significant link was found between HIX and FI. Nevertheless, the positive relationship between BIX and FI ($r = 0.63$, $p < 0.01$) was not surprising and suggested that the soils with DOM originating from a microbial source were related to the highest autochthonous contributions, as reported in other works (Tang et al. 2019; Musadji et al. 2020). Overall, these results indicate that the 18 soils sampled in our study differed in DOM quality even though they had similar concentrations of DOC. Given the significant affinity of Cu for DOM, these differences in DOM quality are expected to affect the geochemistry of Cu in solution.

3.5 Prediction of Cu availability using the properties of the soils and the extracts

The risk of Cu environmental toxicity could be better assessed if its availability could be predicted using the variables measured in the soil and in the soil solution. However, organisms

often use mechanisms to resist the stress caused by Cu toxicity. For instance, some plants take up and sequester Cu in cells where it cannot cause harm (Küpper et al. 2009). Nevertheless, as Cu environmental availability is linked to its total (Cu_{KCl}) and ionic (Cu^{2+}) concentrations in soil solutions, predicting these concentrations could enable more accurate evaluation of potential Cu toxicity to a certain extent (see **Section 3.2**). **Table 3** shows the best models calculated to predict Cu_{KCl} and Cu^{2+} among the 18 vineyard soils sampled in this study.

3.5.1. Prediction of total Cu in extracts Cu_{KCl}

Among the one-variable models tested to predict Cu_{KCl} , the model with pH_{water} (a) showed the best goodness of fit ($R^2_{\text{adj}} = 0.575$, **Table 3**). This suggests that pH is the main factor driving Cu solubilisation in the soils sampled in this study. However, this model showed marked dispersion of points around the 1:1 line (**Fig. 5a**), more visible in soils in group 1 than in soils in group 2. The prediction was better when Cu_T was added to pH (model (b)), the R^2_{adj} value raised to 0.71 and the RMSE decreased from 0.28 to 0.23 (**Table 3**). The points were then less dispersed than with the preceding model, except in soils in group 2 (**Fig. 5b**). The respective coefficients of pH and Cu_T are suitable for the presumed mechanisms. Indeed, it has been reported that metal desorption increases with a decrease in pH, which is why metals tend to be more soluble in acidic environments (McBride and Blasiak 1979; Degryse et al. 2009). Moreover, at higher pH, Cu oxyhydroxides or carbonates may form and precipitate thereby reducing the Cu in solution (McBride 1989). As expected, the model showed that the higher the Cu_T , the higher the concentration of soluble Cu in solution (Cu_{KCl}). Sauv et al. (1997) underlined the importance of pH and Cu_T in Cu solubilisation, but in the reverse hierarchy. These authors found that in 66 soils with diverse sources of Cu contamination, 90 % of variation in soluble Cu was explained by total Cu in the soils and the importance of pH was minimal.

This could be explained by the wider range of Cu_T in their study ($14\text{--}3000\text{ mg kg}^{-1}$) which reinforced the contribution of Cu_T in predicting soluble Cu.

The concentration of DOC did not appear in the best models, presumably because of its limited variability in our initial dataset. However, Cu_{KCl} predictability was dramatically improved when DOM quality parameters were added as predictors in addition to pH and Cu_T . This suggests that in the present study, Cu availability depended more on DOM characteristics than on its concentration. Among all the DOM indices tested in the regressions, only HIX and FI emerged significantly, and the best prediction was obtained by equation (c) (**Table 3, Fig. 5c**), which explained 83.2 % of variance, showing good accuracy of this model for the prediction of Cu_{KCl} . Compared to model (b), model (c) produced lower RMSE (0.17) and a slope closer to 1 (0.85) (**Fig. 5c**). The contributions of pH_{water} , Cu_T , FI and HIX were all significant in the model ($p < 0.001$) (**Table 3**). The variance inflation factor (VIF) did not identify correlations between the predictors ($\text{VIF} < 2.3$). HIX and FI were both expressed with negative coefficients in the model, suggesting that the solubilisation of Cu is enhanced with lowly-condensed plant-derived DOM molecules. This suggests that, in the vineyard soils studied here, molecules with low aromaticity have the most affinity for Cu for its solubilisation. This finding contradicts previous studies suggesting that the most aromatic DOM had higher affinity for Cu and promoted its solubilisation through complexation (Amery et al. 2007; Amery et al. 2008; Araújo et al. 2019). In the present study, DOM composed of more hydrophilic molecules may have higher Cu-binding capacity because of its higher proportion of O-containing functional groups (hydroxyl, carboxylic acid, sulfonic acid and amine) than aromatic and hydrophobic substances (Baham et al. 1978; Leenheer 1981). Moreover, the negative coefficient of FI in the model suggests a higher plant origin for the DOM components that bind Cu. Our results are consistent with those of He et al. (2016), who found higher concentrations of heavy metals in sediment solutions with lower aromatic DOM molecules, which the authors linked to amino acids. Moreover, it has

previously been reported that the ability of DOM hydrophilic and low aromatic components to transport Cu is greater than that of hydrophobic molecules (Han and Thompson 1999; 2003). These authors explained that low aromatic and hydrophilic molecules were more mobile, while aromatic and complex components tended to be adsorbed on soil particles.

3.5.2. Prediction of free ionic Cu^{2+}

Among the models with just one predictor, the one with pH_{KCl} (model (d)) appeared to be the most effective and explained 88.3 % of Cu^{2+} variance (**Table 3**). The RMSE value was low (0.4), implying good accuracy of the prediction (**Fig. 5d**). The high R^2_{adj} (0.88) of model (d) highlights the importance of pH in predicting Cu^{2+} concentrations.

We hypothesised that free Cu^{2+} may be linked to the solubility of Cu, as we found a positive relationship between Cu^{2+} and Cu_{KCl} (**Fig. 4c**). When adding Cu_{KCl} to pH_{KCl} as predictors of Cu^{2+} (model (e)), the R^2_{adj} increased from 0.88 to 0.91. This slight improvement can be observed in **Fig. 5e**. The corresponding VIF was the maximum receivable (2.3), certainly due to the relative effect of pH on Cu solubilisation. The contributions of pH_{KCl} and Cu_{KCl} were all significant ($p < 0.001$) in explaining Cu^{2+} variability. The coefficients in the model suggest that Cu^{2+} increases with Cu_{KCl} , and decreasing pH. Compared to model (d), model (e) was more accurate with a lower RMSE (0.34 instead of 0.4) (**Table 3**).

BIX was the DOM index that appeared in the best three-variables model (model (f), **Table 3**). Models with more than three variables did not present any statistical improvement (data not shown). In model (f), adding BIX to pH_{KCl} and Cu_{KCl} did not enhance the effectiveness of the prediction compared to model (e) (**Fig. 5f**), with similar R^2_{adj} and RMSE (0.92 and 0.32 respectively) (**Table 3**). Even though the contribution of BIX to the model was significant ($p < 0.05$), the addition of BIX was not needed to explain the variability of our collection of samples. Moreover, the max VIF of model (f) (2.44) was higher than the fixed threshold of 2.3,

illustrating multicollinearity between the predictors that probably weakens the stability of the prediction. Taken together, these results suggest that including DOM indices does not improve the modelling of Cu^{2+} and that only pH_{KCl} and Cu_{KCl} were needed.

In the literature, pH is widely reported to play an important role explaining free ionic Cu^{2+} activities and the log-log relationship between these two variables is usually linear (McBride and Blasiak 1979; Sauvé et al. 1995; McBride et al. 1997; Sauvé et al. 1997; Bravin et al. 2009). Sauvé et al. (1997) proposed the following equation to predict Cu^{2+} activity in CaCl_2 extracts of a collection of wide Cu-contaminated soils: $\text{pCu} = -1.7 \log_{10}\text{Cu}_T + 1.4 \text{pH} + 3.4$. In contrast to our study, the OM parameters (concentrations and quality) were not included in the model calculations and the authors underlined the importance of the degree of Cu contamination in predicting Cu^{2+} . However, interestingly, the pH coefficient in our study (model (e)) is consistent with the results of McBride (1989) who reported that slopes for $\log\text{Cu}^{2+}$ - pH plots are generally lower than 2. This suggests that pH plays a constant role in Cu^{2+} prediction. But the optical parameters describing the quality of DOM in soil solutions were of no use in explaining the concentrations of Cu^{2+} .

3.6. Influence of DOC composition on Cu speciation

We calculated Cu speciation in solution based on thermodynamic models and on several hypotheses concerning DOM composition. We found that considering DOM as 100 % HA led to underprediction of free Cu^{2+} compared to experimental observations in all samples (**Fig. S5**). Considering all DOM as FA produced better results, but even using only 65 % of active DOM as FA (a widespread assumption in similar studies) resulted in marked discrepancies between model and experimental results. This observation - made here using the NICA-Donnan model to describe ion binding to DOM - is in agreement with previous estimations made using the Windermere humic aqueous model (WHAM) on a large set of soil solutions (Djae et al. 2017).

For the model results to fit the experimental observations, the percentage of active DOM, rDOM, was adjusted separately for each sample. The values of rDOM obtained ranged between 0.10 and 1.68, reflecting the wide variability of the active pool within DOM. This is consistent with the physically meaningful range of 35-215 % estimated by Djae et al. (2017). The variations in rDOM were compared to the variations in the chemical and optical parameters, but no multilinear regression model could be established (data not shown).

The amount of Cu bound to carboxylic or phenolic groups of DOM can be estimated using thermodynamic modelling. Results show that binding to phenolic groups becomes predominant above pH 7.4 (**Fig. S6**). This suggests that these DOM binding groups, deprotonated at alkaline pH, have a very high affinity for Cu. Thus, in alkaline soils, the nature of DOM binding sites will be important in controlling Cu speciation.

4. Conclusions

The main goal of this work was to assess the contribution of DOM spectroscopic indices to predicting two Cu availability proxies in 18 organic vineyard soils. Our results showed that in the KCl extracts, the total concentration of Cu varied 23-fold while that of free ionic Cu^{2+} varied 4,600-fold among soils, pointing to wide diversity of soils in terms of Cu availability. Four soils had concentrations of Cu^{2+} higher than the toxicity levels for organisms reported in the literature. These samples were considered as potentially harmful for vineyard soil organisms.

Our results also showed that Cu environmental availability was weakly related to the level of Cu contamination in the soils. pH was the main factor controlling Cu environmental availability, as it was closely and positively correlated with the concentrations of total Cu and free ionic Cu^{2+} measured in KCl extracts. Thus, in theory, toxicity problems can occur even in moderately Cu-contaminated soils when the soils are particularly acidic. It would thus be safer to maintain the pH above 6.5. To this end, liming is possible but should be applied with caution,

580 so as not to jeopardise plant uptake of other nutrients like Fe, whose availability is low at
581 alkaline pH.

582 This work highlighted the fact that even though the DOC concentrations were similar among
583 the soils, noticeable differences were found in DOM quality, as assessed by optical
584 spectroscopy. Absorbance and fluorescence analyses are thus appropriate to identify these
585 differences in chromophoric and fluorescent DOM in soils.

586 The prediction of total Cu in KCl extracts was clearly improved when DOM spectral indices
587 (HIX and FI) were taken into account, in addition to pH and Cu_T. This study thus emphasizes
588 that the quality of DOM affects the availability of Cu in the soil more than its quantity. Although
589 this finding now needs to be tested on a wider dataset (with different sources of contamination),
590 it suggests that organic management in Cu-contaminated vineyards needs to be redesigned, to
591 better control quality of dissolved organic matter production as a function of the nature of
592 organic matter amendments.

593

594 **Declarations**

595 **Ethics approval and consent to participate**

596 Not applicable

597 **Consent for publication**

598 Not applicable

599 **Availability of data and materials**

600 All data generated or analysed during this study are included in this published article and its
601 supplementary information files.

602 **Competing interests**

603 The authors declare that they have no competing interests.

604 **Funding**

605 This work was financially supported by the CNRS-INSU EC2CO program under the
606 ‘INTRIGUE’ project, by Bordeaux Sciences Agro Institute of Agricultural Sciences through
607 the PhD scholarship of Frédéric Ouedraogo and by the French National Institute for Agriculture,
608 Food and Environment (INRAE).

609 **Authors' contributions**

610 Frédéric Ouedraogo: Conceptualization, Investigation, Formal analysis, Writing-original draft

611 Jean-Yves Cornu: Supervision, Writing-original draft

612 Noémie Janot: Formal analysis, Supervision, Writing-review & editing

613 Christophe Nguyen: Formal analysis, Writing-review & editing

614 Mahaut Sourzac: Investigation

615 Edith Parlanti: Conceptualization, Funding acquisition, Supervision, Writing-original draft

616 Laurence Denaix: Conceptualization, Funding acquisition, Supervision, Writing-original draft

617 All authors read and approved the final manuscript.

618

619 **Acknowledgments**

620 The authors wish to thank Thierry Robert for help in soil sampling, Sylvie Bussière for
621 determining the DOC concentration in soil extracts, Cécile Coriou for technical and human
622 support in ICP-OES measurements and Sylvie Millin for C_{org} and N determinations. The authors
623 would also like to thank Dario Omanović for his assistance in adapting the TreatEEM program
624 and Nicolas Fanin for his support throughout the PhD research of Frédéric Ouedraogo.

References

- Amery F, Degryse F, Cheyns K, De Troyer I, Mertens J, Merckx R, Smolders E (2008) The UV-absorbance of dissolved organic matter predicts the fivefold variation in its affinity for mobilizing Cu in an agricultural soil horizon. *European Journal of Soil Science* 59(6):1087–1095. <https://doi.org/10.1111/j.1365-2389.2008.01078.x>
- Amery F, Degryse F, Degeling W, Smolders E, Merckx R (2007) The Copper-Mobilizing-Potential of Dissolved Organic Matter in Soils Varies 10-Fold Depending on Soil Incubation and Extraction Procedures. *Environmental Science & Technology* 41(7):2277–2281. <https://doi.org/10.1021/es062166r>
- Andrison D, Bardin M, Bertrand C, Brun L, Daire X, Fabre F, Gary C, Montarry J, Nicot P, Reignault P, Tamm L, Savini I (2018) Can organic agriculture give up copper as a crop protection product? Synthesis of the scientific assessment report. Institut national de la recherche agronomique (INRA) :68. <https://doi.org/hal-02944872>
- Araújo E, Strawn DG, Morra M, Moore A, Ferracciú Alleoni LR (2019) Association between extracted copper and dissolved organic matter in dairy-manure amended soils. *Environmental Pollution* 246:1020–1026. <https://doi.org/10.1016/j.envpol.2018.12.070>
- Baham J, Ball NB, Sposito G (1978) Gel Filtration Studies of Trace Metal-Fulvic Acid Solutions Extracted From Sewage Sludges. *Journal of Environmental Quality* 7(2):181–188. <https://doi.org/10.2134/jeq1978.00472425000700020006x>
- Baken S, Degryse F, Verheyen L, Merckx R, Smolders E (2011) Metal Complexation Properties of Freshwater Dissolved Organic Matter Are Explained by Its Aromaticity and by Anthropogenic Ligands. *Environ Sci Technol* 45(7):2584–2590. <https://doi.org/10.1021/es103532a>

648 Ballabio C, Panagos P, Lugato E, Huang J-H, Orgiazzi A, Jones A, Fernández-Ugalde O,
 649 Borrelli P, Montanarella L (2018) Copper distribution in European topsoils: An
 650 assessment based on LUCAS soil survey. *Science of The Total Environment* 636:282–
 651 298. <https://doi.org/10.1016/j.scitotenv.2018.04.268>

652 Bravin MN, Le Merrer B, Denaix L, Schneider A, Hinsinger P (2010) Copper uptake kinetics
 653 in hydroponically-grown durum wheat (*Triticum turgidum durum* L.) as compared with
 654 soil's ability to supply copper. *Plant Soil* 331(1–2):91–104.
 655 <https://doi.org/10.1007/s11104-009-0235-3>

656 Bravin MN, Martí AL, Clairotte M, Hinsinger P (2009) Rhizosphere alkalisation — a major
 657 driver of copper bioavailability over a broad pH range in an acidic, copper-contaminated
 658 soil. *Plant Soil* 318(1):257–268. <https://doi.org/10.1007/s11104-008-9835-6>

659 Broder T, Knorr K-H, Biester H (2017) Changes in dissolved organic matter quality in a
 660 peatland and forest headwater stream as a function of seasonality and hydrologic
 661 conditions. *Hydrology and Earth System Sciences* 21(4):2035–2051.
 662 <https://doi.org/10.5194/hess-21-2035-2017>

663 Brun LA, Maillet J, Hinsinger P, Pépin M (2001) Evaluation of copper availability to plants in
 664 copper-contaminated vineyard soils. *Environmental Pollution* 111(2):293–302.
 665 [https://doi.org/10.1016/S0269-7491\(00\)00067-1](https://doi.org/10.1016/S0269-7491(00)00067-1)

666 Chaignon V, Sanchez-Neira I, Herrmann P, Jaillard B, Hinsinger P (2003) Copper
 667 bioavailability and extractability as related to chemical properties of contaminated soils
 668 from a vine-growing area. *Environmental Pollution* 123(2):229–238.
 669 [https://doi.org/10.1016/S0269-7491\(02\)00374-3](https://doi.org/10.1016/S0269-7491(02)00374-3)

670 Coll P, Le Cadre E, Blanchart E, Hinsinger P, Villenave C (2011) Organic viticulture and soil
671 quality: A long-term study in Southern France. *Applied Soil Ecology*
672 :S0929139311001570. <https://doi.org/10.1016/j.apsoil.2011.07.013>

673 Cornu JY, Denaix L, Schneider A, Pellerin S (2007) Temporal evolution of redox processes
674 and free Cd dynamics in a metal-contaminated soil after rewetting. *Chemosphere*
675 70(2):306–314. <https://doi.org/10.1016/j.chemosphere.2007.06.003>

676 Cornu JY, Randriamamonjy S, Gutierrez M, Rocco K, Gaudin P, Ouerdane L, Lebeau T (2019)
677 Copper phytoavailability in vineyard topsoils as affected by pyoverdine supply.
678 *Chemosphere* 236:124347. <https://doi.org/10.1016/j.chemosphere.2019.124347>

679 Cory R, Mcknight D (2005) Fluorescence Spectroscopy Reveals Ubiquitous Presence of
680 Oxidized and Reduced Quinones in Dissolved Organic Matter. *Environmental science*
681 & technology 39:8142–9. <https://doi.org/10.1021/es0506962>

682 Daouk S, Frege C, Blanc N, Mounier S, Redon R, Merdy P, Lucas Y, Pfeifer H-R (2015)
683 Fluorescence spectroscopy to study dissolved organic matter interactions with
684 agrochemicals applied in Swiss vineyards. *Environ Sci Pollut Res* 22(12):9284–9292.
685 <https://doi.org/10.1007/s11356-015-4086-6>

686 Degryse F, Smolders E, Parker DR (2009) Partitioning of metals (Cd, Co, Cu, Ni, Pb, Zn) in
687 soils: concepts, methodologies, prediction and applications – a review. *European*
688 *Journal of Soil Science* 60(4):590–612. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2389.2009.01142.x)
689 [2389.2009.01142.x](https://doi.org/10.1111/j.1365-2389.2009.01142.x)

690 Derrien M, Yang L, Hur J (2017) Lipid biomarkers and spectroscopic indices for identifying
691 organic matter sources in aquatic environments: A review. *Water Research* 112:58–71.
692 <https://doi.org/10.1016/j.watres.2017.01.023>

693 Djae T, Bravin MN, Garnier C, Doelsch E (2017) Parameterizing the binding properties of
 694 dissolved organic matter with default values skews the prediction of copper solution
 695 speciation and ecotoxicity in soil: Heterogeneity in reactive DOM impacts soil Cu
 696 ecotoxicity. *Environmental Toxicology and Chemistry* 36(4):898–905.
 697 <https://doi.org/10.1002/etc.3622>

698 El Hadri H, Chéry P, Jalabert S, Lee A, Potin-Gautier M, Lespes G (2012) Assessment of
 699 diffuse contamination of agricultural soil by copper in Aquitaine region by using French
 700 national databases. *Science of The Total Environment* 441:239–247.
 701 <https://doi.org/10.1016/j.scitotenv.2012.09.070>

702 European Food Safety Authority (EFSA), Arena M, Auteri D, Barmaz S, Bellisai G, Brancato
 703 A, Brocca D, Bura L, Byers H, Chiusolo A, Court Marques D, Crivellente F, De
 704 Lentdecker C, Egsmose M, Erdos Z, Fait G, Ferreira L, Goumenou M, Greco L, Ippolito
 705 A, Istace F, Jarrah S, Kardassi D, Leuschner R, Lythgo C, Magrans JO, Medina P, Miron
 706 I, Molnar T, Nougadere A, Padovani L, Parra Morte JM, Pedersen R, Reich H, Sacchi
 707 A, Santos M, Serafimova R, Sharp R, Stanek A, Streissl F, Sturma J, Szentes C,
 708 Tarazona J, Terron A, Theobald A, Vagenende B, Verani A, Villamar-Bouza L (2018)
 709 Peer review of the pesticide risk assessment of the active substance copper compounds
 710 copper(I), copper(II) variants namely copper hydroxide, copper oxychloride, tribasic
 711 copper sulfate, copper(I) oxide, Bordeaux mixture. *EFSA Journal* 16(1).
 712 <https://doi.org/10.2903/j.efsa.2018.5152>

713 Fernández-Calviño D, Nóvoa-Muñoz JC, Díaz-Raviña M, Arias-Estévez M (2009) Copper
 714 accumulation and fractionation in vineyard soils from temperate humid zone (NW
 715 Iberian Peninsula). *Geoderma* 153(1–2):119–129.
 716 <https://doi.org/10.1016/j.geoderma.2009.07.024>

717 Fernández-Calviño D, Soler-Rovira P, Polo A, Díaz-Raviña M, Arias-Estévez M, Plaza C
 718 (2010) Enzyme activities in vineyard soils long-term treated with copper-based
 719 fungicides. *Soil Biology and Biochemistry* 42(12):2119–2127.
 720 <https://doi.org/10.1016/j.soilbio.2010.08.007>

721 Ferreira PAA, Marchezan C, Ceretta CA, Tarouco CP, Lourenzi CR, Silva LS, Soriani HH,
 722 Nicoloso FT, Cesco S, Mimmo T, Brunetto G (2018) Soil amendment as a strategy for
 723 the growth of young vines when replanting vineyards in soils with high copper content.
 724 *Plant Physiology and Biochemistry* 126:152–162.
 725 <https://doi.org/10.1016/j.plaphy.2018.03.003>

726 Flores-Álvarez JM, González I, García-de la Rosa LA, Mancilla-Peña F, Reyes-Gómez J (2011)
 727 Carbon paste electrodes modified with biosolids, soils and biocomposites utilized to
 728 study the interaction between organic matter and copper. *Journal of Environmental*
 729 *Management* 92(3):448–456. <https://doi.org/10.1016/j.jenvman.2010.08.033>

730 Fox J, Weisberg S (2019) *An R companion to applied regression*, Third Edition. Sage
 731 Publications

732 Fuentes M, Olaetxea M, Baigorri R, Zamarreño AM, Etienne P, Laine P, Ourry A, Yvin J-C,
 733 Garcia-Mina JM (2013) Main binding sites involved in Fe(III) and Cu(II) complexation
 734 in humic-based structures. *Journal of Geochemical Exploration* 129:14–17.
 735 <https://doi.org/10.1016/j.gexplo.2012.12.015>

736 Gonzaga MIS, Matias MI de AS, Andrade KR, Jesus AN de, Cunha G da C, Andrade RS de,
 737 Santos JC de J (2020) Aged biochar changed copper availability and distribution among
 738 soil fractions and influenced corn seed germination in a copper-contaminated soil.
 739 *Chemosphere* 240:124828. <https://doi.org/10.1016/j.chemosphere.2019.124828>

740 Guo X, Zhu N, Chen L, Yuan D, He L (2015) Characterizing the fluorescent properties and
 741 copper complexation of dissolved organic matter in saline-alkali soils using
 742 fluorescence excitation-emission matrix and parallel factor analysis. *J Soils Sediments*
 743 15(7):1473–1482. <https://doi.org/10.1007/s11368-015-1113-7>

744 Han N, Thompson ML (1999) Copper-Binding Ability of Dissolved Organic Matter Derived
 745 from Anaerobically Digested Biosolids. *Journal of Environmental Quality* 28(3):939–
 746 944. <https://doi.org/10.2134/jeq1999.00472425002800030026x>

747 Han N, Thompson ML (2003) Impact of Dissolved Organic Matter on Copper Mobility in
 748 Aquifer Material. *Journal of Environmental Quality* 32(5):1829–1836.
 749 <https://doi.org/10.2134/jeq2003.1829>

750 Harmsen J (2007) Measuring Bioavailability: From a Scientific Approach to Standard Methods.
 751 *Journal of environmental quality* 36:1420–8. <https://doi.org/10.2134/jeq2006.0492>

752 He W, Lee J-H, Hur J (2016) Anthropogenic signature of sediment organic matter probed by
 753 UV–Visible and fluorescence spectroscopy and the association with heavy metal
 754 enrichment. *Chemosphere* 150:184–193.
 755 <https://doi.org/10.1016/j.chemosphere.2016.01.116>

756 Helling B, Reinecke SA, Reinecke AJ (2000) Effects of the Fungicide Copper Oxychloride on
 757 the Growth and Reproduction of *Eisenia fetida* (Oligochaeta). *Ecotoxicology and*
 758 *Environmental Safety* 46(1):108–116. <https://doi.org/10.1006/eesa.1999.1880>

759 Helms JR, Stubbins A, Ritchie JD, Minor EC, Kieber DJ, Mopper K (2008) Absorption spectral
 760 slopes and slope ratios as indicators of molecular weight, source, and photobleaching of
 761 chromophoric dissolved organic matter. *Limnology and Oceanography* 53(3):955–969.
 762 <https://doi.org/10.4319/lo.2008.53.3.0955>

763 Hudson N, Baker A, Reynolds D (2007) Fluorescence analysis of dissolved organic matter in
 764 natural, waste and polluted waters—a review. *River Research and Applications*
 765 23(6):631–649. <https://doi.org/10.1002/rra.1005>

766 Huguet A, Vacher L, Relexans S, Saubusse S, Froidefond JM, Parlanti E (2009) Properties of
 767 fluorescent dissolved organic matter in the Gironde Estuary. *Organic Geochemistry*
 768 40(6):706–719. <https://doi.org/10.1016/j.orggeochem.2009.03.002>

769 Husson F, Josse J, Lê S (2008) FactoMineR: An R Package for Multivariate Analysis. *Journal*
 770 *of Statistical Software* 25. <https://doi.org/10.18637/jss.v025.i01>

771 Inamdar S, Singh S, Dutta S, Levia D, Mitchell M, Scott D, Bais H, McHale P (2011)
 772 Fluorescence characteristics and sources of dissolved organic matter for stream water
 773 during storm events in a forested mid-Atlantic watershed. *Journal of Geophysical*
 774 *Research: Biogeosciences* 116(G3). <https://doi.org/10.1029/2011JG001735>

775 ISO 2006 Soil quality — Requirements and guidance for the selection and application of
 776 methods for the assessment of bioavailability of contaminants in soil and soil materials

777 Jaffé R, Cawley KM, Yamashita Y (2014) Applications of Excitation Emission Matrix
 778 Fluorescence with Parallel Factor Analysis (EEM-PARAFAC) in Assessing
 779 Environmental Dynamics of Natural Dissolved Organic Matter (DOM) in Aquatic
 780 Environments: A Review. In: Rosario-Ortiz F (ed) ACS Symposium Series. American
 781 Chemical Society, Washington, DC, pp 27–73

782 Jones D, Willett V (2006) Experimental evaluation of methods to quantify dissolved organic
 783 nitrogen (DON) and dissolved organic carbon (DOC) in soil. *Soil Biology and*
 784 *Biochemistry* 38(5):991–999. <https://doi.org/10.1016/j.soilbio.2005.08.012>

785 Juang K-W, Lee Y-I, Lai H-Y, Wang C-H, Chen B-C (2012) Copper accumulation,
 786 translocation, and toxic effects in grapevine cuttings. *Environ Sci Pollut Res Int*
 787 19(4):1315–1322. <https://doi.org/10.1007/s11356-011-0657-3>

788 Kader M, Lamb DT, Wang L, Megharaj M, Naidu R (2016) Predicting copper phytotoxicity
 789 based on pore-water pCu. *Ecotoxicology* 25(3):481–490.
 790 <https://doi.org/10.1007/s10646-015-1605-7>

791 Kalbitz K, Schmerwitz J, Schwesig D, Matzner E (2003) Biodegradation of soil-derived
 792 dissolved organic matter as related to its properties. *Geoderma* 113(3):273–291.
 793 [https://doi.org/10.1016/S0016-7061\(02\)00365-8](https://doi.org/10.1016/S0016-7061(02)00365-8)

794 Karimi B, Masson V, Guillaud C, Leroy E, Pellegrinelli S, Giboulot E, Maron P-A, Ranjard L
 795 (2021) Ecotoxicity of copper input and accumulation for soil biodiversity in vineyards.
 796 *Environ Chem Lett.* <https://doi.org/10.1007/s10311-020-01155-x>

797 Kelepertzis E, Massas I, Fligos G, Panagiotou M, Argyraki A (2017) Copper accumulation in
 798 vineyard soils from Nemea, Greece. *Bulletin of the Geological Society of Greece*
 799 50:2192. <https://doi.org/10.12681/bgsg.14275>

800 Kikuchi T, Fujii M, Terao K, Jiwei R, Lee YP, Yoshimura C (2017) Correlations between
 801 aromaticity of dissolved organic matter and trace metal concentrations in natural and
 802 effluent waters: A case study in the Sagami River Basin, Japan. *Science of The Total*
 803 *Environment* 576:36–45. <https://doi.org/10.1016/j.scitotenv.2016.10.068>

804 Kinniburgh DG, Milne CJ, Benedetti MF, Pinheiro JP, Filius J, Koopal LK, Van Riemsdijk
 805 WH (1996) Metal Ion Binding by Humic Acid: Application of the NICA-Donnan
 806 Model. *Environ Sci Technol* 30(5):1687–1698. <https://doi.org/10.1021/es950695h>

807 Kolbas A, Marchand L, Herzig R, Nehnevajova E, Mench M (2014) Phenotypic seedling
808 responses of a metal-tolerant mutant line of sunflower growing on a Cu-contaminated
809 soil series: potential uses for biomonitoring of Cu exposure and phytoremediation. *Plant*
810 *Soil* 376(1):377–397. <https://doi.org/10.1007/s11104-013-1974-8>

811 Komárek M, Čadková E, Chrastný V, Bordas F, Bollinger J-C (2010) Contamination of
812 vineyard soils with fungicides: A review of environmental and toxicological aspects.
813 *Environment International* 36(1):138–151.
814 <https://doi.org/10.1016/j.envint.2009.10.005>

815 Koopal LK, Saito T, Pinheiro JP, Riemsdijk WH van (2005) Ion binding to natural organic
816 matter: General considerations and the NICA–Donnan model. *Colloids and Surfaces A:*
817 *Physicochemical and Engineering Aspects* 265(1):40–54.
818 <https://doi.org/10.1016/j.colsurfa.2004.11.050>

819 Küpper H, Götz B, Mijovilovich A, Küpper FC, Meyer-Klaucke W (2009) Complexation and
820 Toxicity of Copper in Higher Plants. I. Characterization of Copper Accumulation,
821 Speciation, and Toxicity in *Crassula helmsii* as a New Copper Accumulator. *Plant*
822 *Physiology* 151(2):702–714. <https://doi.org/10.1104/pp.109.139717>

823 Leenheer JA (1981) Comprehensive approach to preparative isolation and fractionation of
824 dissolved organic carbon from natural waters and wastewaters. *Environ Sci Technol*
825 15(5):578–587. <https://doi.org/10.1021/es00087a010>

826 Li S, Li M, Wang G, Sun X, Xi B, Hu Z (2019) Compositional and chemical characteristics of
827 dissolved organic matter in various types of cropped and natural Chinese soils. *Chem*
828 *Biol Technol Agric* 6(1):20. <https://doi.org/10.1186/s40538-019-0158-z>

829

830 Manceau A, Matynia A (2010) The nature of Cu bonding to natural organic matter. *Geochimica*
831 *et Cosmochimica Acta* 74(9):2556–2580. <https://doi.org/10.1016/j.gca.2010.01.027>

832 Meeussen JCL (2003) ORCHESTRA: An Object-Oriented Framework for Implementing
833 Chemical Equilibrium Models. *Environ Sci Technol* 37(6):1175–1182.
834 <https://doi.org/10.1021/es025597s>

835 McBride M, Sauve S, Hendershot W (1997) Solubility control of Cu, Zn, Cd and Pb in
836 contaminated soils. *European Journal of Soil Science* 48(2):337–346.
837 <https://doi.org/10.1111/j.1365-2389.1997.tb00554.x>

838 McBride MB (1989) Reactions Controlling Heavy Metal Solubility in Soils. In: Stewart BA
839 (ed) *Advances in Soil Science: Volume 10*. Springer, New York, NY, pp 1–56

840 McBride MB, Blasiak JJ (1979) Zinc and Copper Solubility as a Function of pH in an Acid
841 Soil. *Soil Science Society of America Journal* 43(5):866.
842 <https://doi.org/10.2136/sssaj1979.03615995004300050009x>

843 McKnight DM, Boyer EW, Westerhoff PK, Doran PT, Kulbe T, Andersen DT (2001)
844 Spectrofluorometric characterization of dissolved organic matter for indication of
845 precursor organic material and aromaticity. *Limnology and Oceanography* 46(1):38–
846 48. <https://doi.org/10.4319/lo.2001.46.1.0038>

847 Michaud AM, Bravin MN, Galleguillos M, Hinsinger P (2007) Copper uptake and
848 phytotoxicity as assessed in situ for durum wheat (*Triticum turgidum durum* L.)
849 cultivated in Cu-contaminated, former vineyard soils. *Plant Soil* 298(1):99–111.
850 <https://doi.org/10.1007/s11104-007-9343-0>

851 Michaud AM, Chappellaz C, Hinsinger P (2008) Copper phytotoxicity affects root elongation
852 and iron nutrition in durum wheat (*Triticum turgidum durum* L.). *Plant Soil* 310(1):151–
853 165. <https://doi.org/10.1007/s11104-008-9642-0>

854 Milne CJ, Kinniburgh DG, Tipping E (2001) Generic NICA-Donnan Model Parameters for
855 Proton Binding by Humic Substances. *Environ Sci Technol* 35(10):2049–2059.
856 <https://doi.org/10.1021/es000123j>

857 Milne CJ, Kinniburgh DG, van Riemsdijk WH, Tipping E (2003) Generic NICA–Donnan
858 Model Parameters for Metal-Ion Binding by Humic Substances. *Environ Sci Technol*
859 37(5):958–971. <https://doi.org/10.1021/es0258879>

860 Musadji NY, Lemée L, Caner L, Porel G, Poinot P, Geffroy-Rodier C (2020) Spectral
861 characteristics of soil dissolved organic matter: Long-term effects of exogenous organic
862 matter on soil organic matter and spatial-temporal changes. *Chemosphere* 240:124808.
863 <https://doi.org/10.1016/j.chemosphere.2019.124808>

864 Navel A, Martins JMF (2014) Effect of long term organic amendments and vegetation of
865 vineyard soils on the microscale distribution and biogeochemistry of copper. *Science of*
866 *The Total Environment* 466–467:681–689.
867 <https://doi.org/10.1016/j.scitotenv.2013.07.064>

868 Ohno T, Fernandez IJ, Hiradate S, Sherman JF (2007) Effects of soil acidification and forest
869 type on water soluble soil organic matter properties. *Geoderma* 140(1):176–187.
870 <https://doi.org/10.1016/j.geoderma.2007.04.004>

871 Oustriere N, Marchand L, Galland W, Gabbon L, Lottier N, Motelica M, Mench M (2016)
872 Influence of biochars, compost and iron grit, alone and in combination, on copper
873 solubility and phytotoxicity in a Cu-contaminated soil from a wood preservation site.

874 Science of The Total Environment 566–567:816–825.
875 <https://doi.org/10.1016/j.scitotenv.2016.05.091>

876 Ouzounidou G (1995) Responses of maize (*Zea mays* L.) plants to copper stress—I. Growth,
877 mineral content and ultrastructure of roots. *Environmental and Experimental Botany*
878 35(2):167–176. [https://doi.org/10.1016/0098-8472\(94\)00049-B](https://doi.org/10.1016/0098-8472(94)00049-B)

879 Parat C, Chaussod R, Leveque J, Dousset S, Andreux F (2002) The relationship between copper
880 accumulated in vineyard calcareous soils and soil organic matter and iron. *Eur J Soil*
881 *Science* 53(4):663–670. <https://doi.org/10.1046/j.1365-2389.2002.00478.x>

882 Qiu H, Vijver MG, He E, Peijnenburg WJGM (2013) Predicting copper toxicity to different
883 earthworm species using a multicomponent Freundlich model. *Environ Sci Technol*
884 47(9):4796–4803. <https://doi.org/10.1021/es305240n>

885 R core Team (2020) R A Lang. *Environ. Stat. Comput. R Found. Stat. Comput*, Vienna, Austria
886 URL. <https://www.r-project.org/>

887 Recatalá L, Sacristán D, Arbelo C, Sánchez J (2012) Can a Single and Unique Cu Soil Quality
888 Standard be Valid for Different Mediterranean Agricultural Soils under an Accumulator
889 Crop? *Water, Air, & Soil Pollution* 223. <https://doi.org/10.1007/s11270-011-0960-0>

890 Ren Z-L, Tella M, Bravin MN, Comans RNJ, Dai J, Garnier J-M, Sivry Y, Doelsch E, Straathof
891 A, Benedetti MF (2015) Effect of dissolved organic matter composition on metal
892 speciation in soil solutions. *Chemical Geology* 398:61–69.
893 <https://doi.org/10.1016/j.chemgeo.2015.01.020>

894 Ritsema CJ (1993) Estimation of activity coefficients of individual ions in solutions with ionic
895 strengths up to 0.3 mol dm⁻³. *Journal of Soil Science* 44(2):307–315.
896 <https://doi.org/10.1111/j.1365-2389.1993.tb00454.x>

897 Romero CM, Engel RE, D’Andrilli J, Chen C, Zabinski C, Miller PR, Wallander R (2017) Bulk
898 optical characterization of dissolved organic matter from semiarid wheat-based
899 cropping systems. *Geoderma* 306:40–49.
900 <https://doi.org/10.1016/j.geoderma.2017.06.029>

901 Römken PFAM, Dolfing J (1998) Effect of Ca on the Solubility and Molecular Size
902 Distribution of DOC and Cu Binding in Soil Solution Samples. *Environmental Science*
903 *& Technology* 32(3):363–369. <https://doi.org/10.1021/es970437f>

904 Sacristán D, Viscarra Rossel RA, Recatalá L (2016) Proximal sensing of Cu in soil and lettuce
905 using portable X-ray fluorescence spectrometry. *Geoderma* 265:6–11.
906 <https://doi.org/10.1016/j.geoderma.2015.11.008>

907 Sauvé S, Cook N, Hendershot WH, McBride MB (1996) Linking plant tissue concentrations
908 and soil copper pools in urban contaminated soils. *Environmental Pollution* 94(2):153–
909 157. [https://doi.org/10.1016/S0269-7491\(96\)00081-4](https://doi.org/10.1016/S0269-7491(96)00081-4)

910 Sauvé S, Dumestre A, McBride M, Hendershot W (1998) Derivation of soil quality criteria
911 using predicted chemical speciation of Pb²⁺ and Cu²⁺. *Environmental Toxicology and*
912 *Chemistry* 17(8):1481–1489. <https://doi.org/10.1002/etc.5620170808>

913 Sauvé S, Hendershot W, Allen HE (2000) Solid-Solution Partitioning of Metals in
914 Contaminated Soils: Dependence on pH, Total Metal Burden, and Organic Matter.
915 *Environ Sci Technol* 34(7):1125–1131. <https://doi.org/10.1021/es9907764>

916 Sauv S, McBride MB, Hendershot WH (1995) Ion-selective electrode measurements of
 917 copper(II) activity in contaminated soils. *Archives of Environmental Contamination and*
 918 *Toxicology* 29(3):373–379. <https://doi.org/10.1007/BF00212503>

919 Sauv S, McBride MB, Norvell WA, Hendershot WH (1997) Copper Solubility and Speciation
 920 of In Situ Contaminated Soils: Effects of Copper Level, pH and Organic Matter. *Water,*
 921 *Air, & Soil Pollution* 100(1):133–149. <https://doi.org/10.1023/A:1018312109677>

922 Sun R, Song C, Sun Y, Wang Q, Liu H, Wang J (2020) Agricultural Soil Organic Matters and
 923 Microbiome Are Shaped by Management and Hedgerows. *Agronomy* 10(9):1316.
 924 <https://doi.org/10.3390/agronomy10091316>

925 Supriatin S, Terrones CA, Bussink W, Weng L (2015) Drying effects on selenium and copper
 926 in 0.01M calcium chloride soil extractions. *Geoderma* 255–256:104–114.
 927 <https://doi.org/10.1016/j.geoderma.2015.04.021>

928 Swift MJ, Heal OW, Anderson JM (1979) Decomposition in terrestrial ecosystems. *Studies in*
 929 *ecology (USA)*

930 Tait TN, Cooper CA, McGeer JC, Wood CM, Smith DS (2016) Influence of dissolved organic
 931 matter (DOM) source on copper speciation and toxicity to *Brachionus plicatilis*. *Environ*
 932 *Chem* 13(3):496. <https://doi.org/10.1071/EN15123>

933 Tang J, Wang W, Yang L, Cao C, Li X (2019) Variation in quantity and chemical composition
 934 of soil dissolved organic matter in a peri-urban critical zone observatory watershed in
 935 Eastern China. *Science of The Total Environment* 688:622–631.
 936 <https://doi.org/10.1016/j.scitotenv.2019.06.270>

937 Temminghoff EJM, Van Der Zee SE a. TM, Keizer MG (1994) The influence of pH on the
 938 desorption and speciation of copper in a sandy soil. *Soil Science* 158(6):398–408

 939 Trevors JT, Cotter CM (1990) Copper toxicity and uptake in microorganisms. *Journal of*
 940 *Industrial Microbiology* 6(2):77–84. <https://doi.org/10.1007/BF01576426>

 941 Verweij W, Glazewski R, Haan HD (1992) Speciation of copper in relation to its bioavailability.
 942 *Chemical Speciation & Bioavailability* 4(2):43–51.
 943 <https://doi.org/10.1080/09542299.1992.11083177>

 944 Wei T, Simko V (2021) R package “corrplot”: Visualization of a Correlation Matrix.
 945 <https://github.com/taiyun/corrplot>. Accessed 5 Jan 2021

 946 Weishaar JL, Aiken GR, Bergamaschi BA, Fram MS, Fujii R, Mopper K (2003) Evaluation of
 947 Specific Ultraviolet Absorbance as an Indicator of the Chemical Composition and
 948 Reactivity of Dissolved Organic Carbon. *Environ Sci Technol* 37(20):4702–4708.
 949 <https://doi.org/10.1021/es030360x>

 950 Weng L, Temminghoff EJM, Lofts S, Tipping E, Van Riemsdijk WH (2002) Complexation
 951 with Dissolved Organic Matter and Solubility Control of Heavy Metals in a Sandy Soil.
 952 *Environmental Science & Technology* 36(22):4804–
 953 4810.<https://doi.org/10.1021/es0200084>

 954 Xu H, Zou L, Guan D, Li W, Jiang H (2019) Molecular weight-dependent spectral and metal
 955 binding properties of sediment dissolved organic matter from different origins. *Science*
 956 *of The Total Environment* 665:828–835.
 957 <https://doi.org/10.1016/j.scitotenv.2019.02.186>

958 Yang L, Hur J (2014) Critical evaluation of spectroscopic indices for organic matter source
 959 tracing via end member mixing analysis based on two contrasting sources. *Water*
 960 *Research* 59:80–89. <https://doi.org/10.1016/j.watres.2014.04.018>

961 Yeh Y-L, Yeh K-J, Hsu L-F, Yu W-C, Lee M-H, Chen T-C (2014) Use of fluorescence
 962 quenching method to measure sorption constants of phenolic xenoestrogens onto humic
 963 fractions from sediment. *Journal of Hazardous Materials* 277:27–33.
 964 <https://doi.org/10.1016/j.jhazmat.2014.03.057>

965 Yruela I (2009) Copper in plants: acquisition, transport and interactions. *Functional Plant Biol*
 966 36(5):409. <https://doi.org/10.1071/FP08288>

967 Yruela I (2005) Copper in plants. *Brazilian Journal of Plant Physiology* 17(1):145–156.
 968 <https://doi.org/10.1590/S1677-04202005000100012>

969 Zhang S, Chen X-W, Ye Q, Zhang Z-T, Kong S-F, Cao C, Wang J-J (2020) Dissolved
 970 Metal(loid) Concentrations and Their Relations with Chromophoric and Fluorescent
 971 Dissolved Organic Matter in an Urban River in Shenzhen, South China. *Water*
 972 12(1):281. <https://doi.org/10.3390/w12010281>

973 Zsolnay A (1996) Dissolved Humus in Soil Waters. In: *Humic Substances in Terrestrial*
 974 *Ecosystems*. Elsevier, pp 171–223

975 Zsolnay A, Baigar E, Jimenez M, Steinweg B, Saccomandi F (1999) Differentiating with
 976 fluorescence spectroscopy the sources of dissolved organic matter in soils subjected to
 977 drying. *Chemosphere* 38(1):45–50. [https://doi.org/10.1016/S0045-6535\(98\)00166-0](https://doi.org/10.1016/S0045-6535(98)00166-0)

978

Figure 1

Figure 2

Figure 2

Figure 3

Figure 4

Figure 4

Figure 5

Figure 5

Figure captions

Fig. 1: Distribution of DOM optical parameters (mean of replicates), the red points represent the mean values of all the soils

Fig. 2: Principal component analysis (PCA) of soil solution chemistry and Cu availability proxies in the dataset. All the variables except pH_{water} and pH_{KCl} were log-transformed. The correlation plot (a) shows the contribution of each variable to the two first principal components. The soil variables are shown in black, the extract variables in blue. Soil samples (grouped in triplicate) are coloured according to their pH_{water} values (b)

Fig. 3: Correlation matrix between soil properties (pH_{water} , C_{org} , C:N, Cu_{T}), solution parameters: pH of the extracts (pH_{KCl}), free ionic concentrations (Cu^{2+}), free ionic fractions (fCu^{2+}), total dissolved metals (Cu_{KCl} , Fe_{KCl}) and DOM properties (DOC, HIX, BIX, FI, S_{R} , SUVA_{254}). The colored boxes represent significant correlations ($p < 0.01$)

Fig. 4: Total Cu concentrations in extracts (Cu_{KCl}) as a function of total Cu in soils Cu_{T} (a) and pH of the soils (pH_{water}) (b); free ionic Cu concentrations (Cu^{2+}) as a function of total Cu concentrations in extracts (c) and pH of the extracts (pH_{KCl}) (d). The log-log plot is shown in (c) and only the y-axis is log-transformed in (b) and (d)

Fig. 5: Comparison of the multilinear models predicting Cu_{KCl} (a, b, c) and Cu^{2+} concentrations (d, e, f) in the 0.01 M KCl extracts. The dashed red line represents the 1:1 line, the dashed black line represents the regression line

Tables

Table 1: Chemical properties of the 18 vineyard soils analysed in this study. C_{org} stands for organic carbon, Cu_T for total Cu. Carbonate values <1 were not quantifiable

Soil id	AOC Zone	Soil classification (WRB)	Cu _T	pH _{water}	CaCO ₃	C _{org}	C:N
			mg kg ⁻¹		g kg ⁻¹	g kg ⁻¹	
A	Blaye Côtes de Bordeaux	Cambisol	28	8.1	< 1	9.9	10.6
B	Blaye Côtes de Bordeaux	Cambisol	35	8.3	27.6	13.7	10.9
C	Blaye Côtes de Bordeaux	Luvic Camisol	35	6.9	< 1	10.2	11.4
D	Bordeaux supérieur	Luvisol	73	6.3	< 1	10.5	11.5
E	Blaye Côtes de Bordeaux	Cambisol	80	7.9	< 1	8.3	10.7
F	Graves	Podzosol	84	6.5	< 1	16.9	15.3
G	Bordeaux supérieur	Luvisol	89	6.9	< 1	19.2	11.7
H	Bordeaux supérieur	Luvisol	103	7.9	3.2	12.6	10.6
I	Graves	Cambisol	104	8.3	19.5	7.5	11.5
J	Blaye Côtes de Bordeaux	Luvisol	110	8	1.1	11.3	11.6
K	Blaye Côtes de Bordeaux	Podzosol	121	6.5	< 1	11.8	13.0
L	Graves	Leptosol	155	7.1	< 1	14.6	15.5
M	Blaye Côtes de Bordeaux	Luvisol	175	6.5	< 1	9.3	11.9
N	Blaye Côtes de Bordeaux	Luvisol	176	7.8	< 1	5.4	11.7
O	Blaye Côtes de Bordeaux	Luvisol	181	7.3	< 1	15.6	12.6
P	Graves	Leptosol	193	7.5	< 1	30.8	17.4

Q	Blaye Côtes de Bordeaux	Luvisol	195	6.3	< 1	13.3	12.6
R	Blaye Côtes de Bordeaux	Cambisol	238	8.2	30.4	10.6	10.3

Table 2: Values of total Cu in solution and Cu²⁺ reported as toxicity thresholds in the literature, compared with the maximum values found in the present study

	Toxicity thresholds	Values (μM)	References
	Max value in this study	2.64	
Total Cu in solution (Cu _{KCl} in this study)	IC ₅₀ grapevine	3.94	Juang et al. 2012
	IC ₅₀ maize	6.50	Ouzounidou 1995
	IC ₅₀ sunflower	9.55	Kolbas et al. 2014
	Max value in this study	0.94	
Free ionic Cu ²⁺	EC ₅₀ durum wheat	0.06	Bravin et al. 2010
	EC ₅₀ cucumber	2.20	Kader et al. 2016
	EC ₅₀ <i>Eisenia fetida</i>	1.00	Qiu et al. 2013

Table 3: Best models for predicting Cu_{KCl} and Cu^{2+} in KCl extracts and associated statistics.

R^2_{adj} : adjusted r-squared, RMSE: root mean square error, max VIF: maximum value of the variance inflation factor, N: number of observations

Statistical significance: *** for p-value < 0.001, ** for p-value < 0.01, * for p-value < 0.05

	Models	R^2_{adj}	max VIF	RMSE (μM)	N
$\log_{10}\text{Cu}_{\text{KCl}} =$	$-0.465^{***} \text{pH}_{\text{water}} + 3.066$ (a)	0.575	-	0.28	53
	$-0.434^{***} \text{pH}_{\text{water}} + 0.615^{***} \log_{10}\text{Cu}_{\text{T}} + 1.598$ (b)	0.707	1.02	0.23	53
	$-0.312^{***} \text{pH}_{\text{water}} + 0.443^{***} \log_{10}\text{Cu}_{\text{T}} - 7.362^{***} \log_{10}\text{FI}$	0.832	1.91	0.17	53
	$-0.664^{***} \log_{10}\text{HIX} + 2.148$ (c)				
$\log_{10}\text{Cu}^{2+} =$	$-1.751^{***} \text{pH}_{\text{KCl}} + 9.589$ (d)	0.883	-	0.40	53
	$-1.380^{***} \text{pH}_{\text{KCl}} + 0.731^{***} \log_{10}\text{Cu}_{\text{KCl}} - 7.307$ (e)	0.912	2.30	0.34	53
	$-1.316^{***} \text{pH}_{\text{KCl}} + 0.73^{***} \log_{10}\text{Cu}_{\text{KCl}} + 2.818^{*} \log_{10}\text{BIX} +$	0.919	2.44	0.32	53
	7.338 (f)				