1 Research Paper

2 Membrane interactions accelerate the self-aggregation of 3 huntingtin exon 1 fragments in a polyglutamine 4 length-dependent manner

5

6 Arnaud Marquette ¹, Christopher Aisenbrey ¹ and Burkhard Bechinger ^{1,2,*}

7

¹ University of Strasbourg/CNRS, Chemistry Institute UMR7177

8

² Institut Universitaire de France

9

* Correspondence: beching@unistra.fr

10

11

12

13

14

15

16

17

18

Abstract: The accumulation of aggregated protein is a typical hallmark of many human neurodegenerative disorders including polyglutamine-related diseases such as chorea Huntington. Misfolding of the amyloidogenic proteins gives rise to self-assembled complexes and fibres. The huntingtin protein is characterized by a segment of consecutive glutamines, which when exceeding ~ 37 residues results in the occurrence of the disease. Furthermore, it has also been demonstrated that the 17-residue amino-terminal domain of the protein (htt17), located upstream of this polyglutamine tract, strongly correlates with aggregate formation and pathology. Here we demonstrate that membrane interactions strongly accelerate the oligomerization and β -amyloid fibril formation of htt17-polyglutamine segments. By using a combination of biophysical approaches the kinetics of fibre formation has been investigated and found to be strongly dependent to the presence of lipids, the length of the polyQ expansion and the polypeptide-to-lipid ratio. Finally, the implications for therapeutic approaches are discussed.

19

20

Citation: Marquette, A.; Aisenbrey, C.; Bechinger, B. Membrane interactions accelerate the self-aggregation of huntingtin exon 1 fragments in a polyglutamine length-dependent manner. *Int. J. Mol. Sci.* **2021**, *22*, x. <https://doi.org/10.3390/xxxxx>

21

22

23

24

25

26

27

28

Keywords: circular dichroism; DLS; thioflavin T fluorescence; peptide-lipid interactions; huntingtin; Huntington's disease; amyloid; htt17; membrane-driven aggregation

Received: date

Accepted: date

Published: date

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Copyright: © 2021 by the author. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

At least nine different hereditary diseases that are related to the expansion of a polyglutamine (polyQ) domain are known to date [1]. These so-called CAG repeat pathologies are all related by the propensity of their associated polypeptide to form insoluble β -sheet rich amyloid fibrils. Polyglutamine expansion promotes the self-assembly of fibrils and other types of aggregates that accumulate in inclusions found for example in the brain tissues of patients. By using modern imaging techniques these have been described as dynamic phase separated gel-like structures or as coexisting liquid / solid condensates [2,3]. Chorea Huntington is one of the best studied polyglutamine-related diseases. In this case it has been shown that the age of development and the severity of the disease correlate with the length of the polyQ stretch which is located within the amino-terminal domain of huntingtin, a protein encompassing about 3500 amino acids [1,4]. Symptoms of the disease develop when the polyQ expansion exceeds a critical length of ~ 37 glutamines [5,6]. Although the genetics of Huntington's disease (HD) are well studied, the exact biological functions of huntingtin remain speculative and the exact mechanism of pathogenic peptide aggregation remains a controversial topic [7].

It has been suggested that polyglutamines perturb neuronal membranes, result in their disruption concomitant with calcium dysregulation [8,9] thereby causing Huntington's or other amyloidogenic diseases [10–12]. Huntingtin or fragments of the protein have been shown to be involved in intracellular vesicle trafficking [12–14]. They associate with the ER, the Golgi-apparatus and endosomal vesicles [8,13,15–17]. Notably endosomal vesicle rupture has been shown to be a common mechanism in the redistribution between cells of large assemblies of α -synuclein, tau protein and huntingtin [18]. Furthermore, the length of the polyglutamine tract of huntingtin affects the redistribution of these polypeptides between the cytoplasm and the nucleus [19–21]. Finally, mitochondrial malfunction has also been associated with the pathogenesis of Huntington's disease [22,23]. Indeed, when fusion proteins of GST with exon 1 of huntingtin encompassing either 20 or 51 glutamines have been studied in association with brain lipid membranes protein oligomerization in the presence of long polyglutamine tracts is observed [24].

Studies on the various factors influencing the rate of aggregation include polyQ length, flanking sequences, posttranslational modification, protease activities on huntingtin and the presence of chaperones [1]. On the one hand, the macromolecular assembly of huntingtin or its fragments through polyQ interactions is modulated by association of the protein with membranes [12,25,26]. On the other hand, the length of the polyQ has an influence on these lipid interactions and the resulting membrane disruption [12,25]. Furthermore, the polyproline segment downstream of the polyQ domain has the opposite effect by reducing both the kinetics of aggregation and the formation of β -sheets by the polyQ region [25,27].

Importantly, the first 17 amino acids of huntingtin exon 1 (htt17, also abbreviated N17 or htt^{N17} by other authors), directly preceding the polyQ tract, and posttranslational modifications within this region have been shown to have a strong effect on the cellular localization of huntingtin and the propensity of the protein to aggregate [8,27–32]. Thus, disease pathogenesis in transgenic mice can be inhibited by mutations of serine 13 and 16 within this htt17 domain [33]. Furthermore, the htt17 sequence carries phosphorylation, SUMOlation and nuclear export sequences [1,30,34–38]. Membrane interactions of huntingtin *in vivo* require this htt17 domain [8,15]. Furthermore, it has been shown that htt17 enhances polyglutamine oligomerization [8,15,26,39–41] and is involved in the seeding and fibre maturation processes [26,42].

Recent structural investigations reveal high conformational plasticity of htt17 and the subsequent polyQ domain where htt17 association [26], its interactions with the membrane [43–47] or with other polypeptide domains are associated with random coil-helix structural transitions [48,49]. A recent NMR study has shown that in solution the htt17 sequence associates in a dimer of dimers preaggregation state [50,51]. By using EPR labels positioned at different locations along the exon 1 sequence made up with 46 Qs it was shown that the aggregation process is initiated by the N-terminal domain forming helical structures, followed by the polyQ adopting β -sheet conformations [26]. Membranes or seeds accelerate the aggregation process of this exon 1-Q46 construct. Notably in solution oligomers of 7–11 subunits have been observed [26] while it has been shown that a transition from helical coiled-coil to β -sheet can be part of the aggregation process of polyQ proteins in general [52]. Molecular dynamics calculations indicate that β -hairpin containing conformers of polyQ probably act as templates for subsequent fibril formation [53,54]. The membrane-associated structure of the non-aggregating exon 1-Q25 variant has been investigated using EPR and ODNP approaches [44]. In the presence of membranes, the structuration reaches till residue 22, in agreement with a helical conformation covering much of the htt17 domain [55], while from residue 30 on the protein is water exposed and dynamic [44]. The htt17 domain insertion is shallow and reversible with the helix axis parallel to the membrane surface [43,44,55,56].

Interestingly, the htt17 and the polyQ domains mutually influence each other and their conformational properties are coupled [57,58]. More recent investigations using electron microscopy and molecular dynamics calculations reveal tadpole like structures of exon 1 where htt17 together with polyQ forms a globular head domain increasing in size as the number of glutamines increases and where the polyproline extends as the tail [42,59,60]. The htt17 domain plays an important role in the polyQ aggregation and *de novo*, seeded or membrane-driven mechanisms have been distinguished [26]. Depending on the conditions, different aggregate sizes of globular or fibrous morphologies have been described and correlated to toxicity [17,61,62].

Aggregation, oligomer and/or fibril formation which are the causative events for the development and progression of Huntington's disease [17,61,62] require that polyglutamines are brought in contact with each other. This can occur by protein-mediated interactions of polyglutamines [1,63,64], by seeding or by local accumulation of polyglutamines at bilayer surfaces [26,43,65]. Notably, the reversible membrane interactions of the amphipathic helical htt17 domain have been characterized in a quantitative and lipid-dependent manner [43] and the importance of this domain to enhance polyglutamine aggregation has been shown *in vitro* and *in vivo* [8,15,39–41]. Another huntingtin domain mediating membrane interactions has been identified within residues 171–371, a region bearing an overall high cationic character [12]. Furthermore, biochemical and cell biological assays demonstrate the potential role of these anchoring domains in disease development [12,24,45,46,56,66].

Membrane surface induced conformational changes in proteins play a critical role in the aggregation process, for example, by concentrating and aligning polyglutamines in such a manner to promote nucleation of amyloid formation [25,26,43,67,68]. Furthermore, membranes could alter aggregate morphology to specific toxic species or stabilize potentially toxic, transient aggregation intermediates [10,69–71]. Therefore, investigations of how amyloid fibrils as well as their intermediate and protofibrillar states interact with membranes is of considerable importance [25,72]. In the case of huntingtin and synuclein it has been shown that fibrils are toxic to the cells [73] and that the docking of extra-cellular aggregates to the cell membranes is a key step of the vicious propagation-amplification cycle [74]. However, hardly any of these publications provide quantitative structural and biophysical data about interactions of huntingtin domains with membranes.

This prompted us to investigate in more detail the role of the membrane in the polyQ association kinetics. To this end we prepared constructs involving the membrane-anchoring htt17 domain followed by polyglutamines of different length. The structural changes were characterized in a time-dependent manner using CD spectroscopy and the supramolecular complexes formed by dynamic light scattering and ThT fluorescence. By investigating htt17 in the presence of polyQ domains as short as 9 glutamines the slower aggregation kinetics allow for a more controlled evaluation of the processes involved in aggregation and fibril formation similar to the use of htt17-polyQ constructs used in previous biophysical studies (e.g. [45,46,51,59,66,75]). The polyglutamine was then successively extended in small steps to measure the effect of polyQs on the aggregation dynamics. The results reveal a pronounced dependence of the htt17-driven aggregation rates on the presence of lipid bilayers, the length of the polyglutamine tract, and polypeptide concentration.

2. Results

The aggregation of huntingtin exon 1-derived peptides was studied by a combination of biophysical assays. The peptides encompass the first 17-residue amphipathic sequence known to reversibly interact with membranes [43] and polyglutamine stretches of variable length (Table 1).

T	able	htt17-Q17	MATLEKLMKAFESLKSF QQQ QQQ QQQ QQQ QQQ QQ
		htt17-Q12	MATLEKLMKAFESLKSF QQQ QQQ QQQ QQQ

1:
Ami

no

acid sequences of huntingtin exon 1-related peptides.

htt17-Q9	MATLEKLMKAFESLKSFQQQ QQQ QQQ
----------	------------------------------

2.1. CD spectroscopy

To test if membranes can increase the speed of polyglutamine aggregation and at the same time to get insight into their secondary structure, we recorded CD spectra of the polyQ peptides htt17-Q9, htt17-Q12 and htt17-Q17 as a function of time. Their structural changes were monitored in solution in the absence (Figure 1A-C) and in the presence of phospholipid vesicles (Figure 1D-F) in 10 mM Tris-HCl, pH 7 at a concentration of 0.1 mg/mL (about 30 μ M). The spectra were recorded between 260 and 194 nm where the spectral line shape correlates with the secondary structure composition of the peptides. Measurements were performed every 24.5 minutes and more than 18 spectra were recorded for each polypeptide sequence.

In aqueous solution the three peptides all adopt predominantly random coil conformations, without any significant spectral changes over the time period of the experiment (Figure 1A-C). When fitting the data using the DicroProt analysis software [76] 64% of the signal were associated with random coil conformations.

Figure 1D-F exhibits the time evolution of the peptide spectra in the presence of SUVs made of POPC/POPS at a 3/1 molar ratio. The same mass of peptides (0.1 mg/mL) was mixed with a suspension of vesicles at a lipid concentration of 0.5 mg/mL in 10 mM Tris-HCl, pH 7, giving final peptide-to-lipid (P/L) molar ratios of 1/19.6, 1/22 and 1/26 for htt17-Q9, htt17-Q12 and htt17-Q17, respectively. Interestingly, the structure of htt17-Q9 remains unchanged over time since all the spectra overlap almost perfectly (Figure 1D). An estimate of the secondary structure of the peptide at the beginning of the experiment gives 33 % α -helix, 22% β -sheet and 45% random-coil structures [76]. In contrast, the CD spectra shown in Figure 1E exhibit a significant increase in ellipticity over time in particular between 194 and 220 nm. Whereas the initial structure of the htt17-Q12 peptide resembles that of htt17-Q9 the β -sheet content of the Q12 sequence increases gradually, at the expense of α -helix and random-coil contributions. This effect is even more pronounced for htt17-Q17 where the whole spectrum changes over time to ultimately converge to positive ellipticity values over the whole spectral range (Figure 1F).

Figure 1: Time-dependent structural changes measured by circular dichroism. CD spectra of htt17-Q9, htt17-Q12 and htt17-Q17 ($C = 9.1 \cdot 10^{-2}$ mg/mL) in 10 mM Tris-HCl, pH 7 (A to C), and in presence of SUVs made of POPC/POPS 3/1 mole/mole ($C = 0.45$ mg/mL) (D to F) were recorded every 24.5 minutes. The progress of the spectral changes with time is depicted by arrows in panels E and F. Thereby the peptide-to-lipid ratios are 1/19.6, 1/22 and 1/26 for htt17-Q9, htt17-Q12 and htt17-Q17, respectively.

In order to characterize the kinetics of the peptide structural changes, we quantitatively analysed the time-dependent changes of ellipticity at 208 nm (Figure 2). The intensity at this wavelength is related to the helix secondary structures [77] but also affected by vesicle aggregation processes [78]. A mono-exponential function of the form $A + B \cdot (1 - \exp(-t/\tau))$, where A and B are fitting parameters and τ is the exponential time constant, describes well the intensity increase over time. The resulting fits are displayed in Figure 2. Although it should be noted that the absolute values are influenced by a number of environmental effects such as the stirring efficiency and the rate of dichroism increase, the aggregation rate $1/\tau$ follows $\text{htt17-Q17} > \text{htt17-Q12} > \text{htt17-Q9}$ (Fig. 2). This is confirmed by observations from longer polyglutamine peptides which aggregate too fast to be investigated by the techniques employed in this paper (not shown). Furthermore, the peptides investigated here only aggregate in the presence of lipids within the time frame of the experiments.

Figure 2: Aggregation kinetics by circular dichroism. The time-dependent intensity of the CD signal measured at 208 nm is shown for htt17-Q9, htt17-Q12 and htt17-Q17 in the presence of 0.45 mg/mL SUVs made of POPC/POPS 3/1 mole/mole in 10 mM Tris-HCl, pH 7. The results of least square fits with a mono-exponential function are displayed as solid lines. An error of ± 1 was estimated from the signal-to-noise ratio of the spectra.

2.2. Thioflavin T fluorescence

To follow the kinetics of htt17-polyQ β -sheet formation changes in the Thioflavin T (ThT) fluorescence were monitored in the presence of phospholipid bilayers. ThT is a popular reporter of amyloid aggregation because it demonstrates a strong shift and enhanced intensity of fluorescence emission upon binding to β -sheet rich fibrils [79,80]. The dye has been used to visualize and quantify the presence of misfolded protein or peptide aggregates *in vitro* and *in vivo* [81].

Figure 3 displays three sets of measurements performed at the same ThT concentration where a spectrum has been recorded every 2 minutes over a time interval of up to about one hour. The intensity of the fluorescence emission spectra in 10 mM Tris-HCl buffer at pH 7 increases with time in the presence of both $\approx 14.5 \mu\text{M}$ htt17-Q17 and SUVs made of POPC/POPS 3/1 (Figure 3A). In contrast, the fluorescence measured in control experiments with peptides only (Fig. 3B) or with SUVs only (Fig. 3C) remains unchanged. In the presence of liposomes alone an increase in the ThT fluorescence background suggests an interaction of the fluorophore with membranes (Fig. 3C). The spikes observed in the presence of protein are suggestive of light diffraction when aggregates pass through the light beam.

Figure 3: Time-dependent amyloid formation of htt17-Q17. The thioflavin T fluorescence is shown in the presence of 14.5 μM htt17-Q17 and SUVs made of 320 μM POPC/POPS 3/1 mole/mole in 10 mM Tris-HCl buffer, pH 7 (A). The peptide-to-lipid molar ratio was 1/22 and a spectrum was recorded every 2 min. The control experiments show almost no changes in thioflavin T fluorescence when exposed to the same amount of htt17-Q17 only (B) or SUVs only (C). The thioflavin T concentration was 5 μM in all recordings.

In order to follow the β -sheet formation, we measured the time-dependent fluorescence intensity of ThT in the presence of the three peptides at the fixed wavelength of $\lambda_{\text{fluor}} = 485 \text{ nm}$, while the dye was continuously excited at $\lambda_{\text{exc}} = 440 \text{ nm}$. To quantify the effect of increasing peptide concentrations, for each htt17-polyQ sequence three measurements were performed, namely at P/L ratios of 1/88, 1/44 and 1/22. Under most experimental conditions, an increase of the signal was measured over time, indicating peptide aggregation. In some cases, the signal first increased and then decayed, which is due to sedimentation of the peptides, aggregated and/or associated with the vesicles [82]. Indeed, in these samples, precipitates were detected by visual inspection (Fig. S1A) and fibrous structures made of peptide and associated lipids were observed by electron microscopy (Fig. S1B-D).

As illustrated in Figure 4, the P/L ratio has a direct effect on the efficiency of the aggregation process as well as on its kinetics. For all three peptides, increasing the P/L ratio makes the aggregation process more efficient. Therefore, the fluorescence intensities measured at P/L=1/22 are higher than for P/L=1/44 or P/L=1/88, while the kinetics (i.e. the τ values of the corresponding exponentials) are surprisingly similar when htt17-polyQ is investigated. Whereas for P/L ratios 1/22 a mono-exponential signal increase describes the signal increase reasonably well, additional processes initially cause negative slopes in the ThT fluorescence when htt17-Q9 is investigated at P/L ratios of 1/44 or 1/88, and for htt17-Q12 at P/L=1/88.

In order to test the reproducibility the aggregation kinetics of htt17-Q17 in the presence of lipids (P/L 1/22) was tested 5 times by two different investigators and two different techniques. The τ values were $21 \pm 3 \text{ min}$ ($n=5$) when determined by ThT fluorescence (Figs. 3 and 4C) and 29 min when the CD spectral changes were analyzed (Figs.

1F and 2). For htt17-Q12 the time evolution was an order of magnitude slower (251 ± 83 min, $n=3$; Figs. 1E, 2 and 4B).

Figure 4: Time-dependent thioflavin T fluorescenc as a function of polypeptide concentration. The thioflavin T fluorescence was continuously measured at 485 nm in the presence of SUVs and htt17-Q9 (A), htt17-Q12 (B) or htt17-Q17 (C), in 10 mM Tris-HCl, pH7. The peptide-to-lipid molar ratios are 1/22, 1/44 and 1/88, displayed as solid, short-dashed and long-dashed lines, respectively. The lipid concentration was kept constant ($C \approx 320 \mu\text{M}$) while the amount of peptide was adjusted to obtain the P/L ratios indicated. The ThT concentration was $\approx 5 \mu\text{M}$ in all the recordings.

2.3. Dynamic light scattering

Dynamic light scattering measurements (DLS) were performed to obtain a more detailed view of the vesicle-vesicle interactions and the subsequent sedimentation processes that was suspected to occur in some of the fluorescence and CD spectroscopy experiments. In these experiments the three htt17-polyQ peptides were exposed to suspensions of SUVs made of POPC/POPS 3/1 mole/mole (0.45 mg/mL , i.e. $580 \mu\text{M}$) in 10 mM Tris-HCl, pH 7 buffer (Figure 5). To start with, the htt17-Q9 concentration was $29 \mu\text{M}$ and the hydrodynamic diameter of the molecular assemblies and their corresponding polydispersity indexes (PDI) were measured every 18.7 minutes for more than 7 hours. Related experiments were performed in the presence of $25.9 \mu\text{M}$ htt17-Q12 and $22.9 \mu\text{M}$ htt17-Q17, respectively, i.e. the same concentrations by weight. This corresponds to the same experimental conditions also used for the CD measurements. When vesicles and htt17-Q17 are mixed the DLS data indicate a rapid initial increase in polydispersity indicating fast aggregation and/or vesicle agglutination within 90 min. The changes of the PDI in the

presence of htt17-Q12 occur at about half the pace. At the same time the apparent hydrodynamic radius of the two systems also increases but it should be kept in mind that the absolute value of this parameter is unreliable when the polydispersity approaches 1 (not shown). While the PDI and apparent hydrodynamic radii of htt17-Q12 and htt17-Q17 increase, the htt17-Q9 peptide left the light scattering unaffected (Figure 5). Thereby, the apparent changes in ellipticity were assigned to light scattering processes induced by supra-wavelength sized systems like vesicle and/or peptide clusters.

Figure 5: Time-dependent changes of polydispersity. The polydispersity index (PDI) is as a function of time for SUVs in the presence of htt17-Q9 (circles), htt17-Q12 (squares) or htt17-Q17 (triangles) in 10 mM Tris-HCl, pH7. Measurements were performed by dynamic light scattering in the absence of mechanical stirring. The same concentrations of lipids and peptides were used as in the CD experiments presented in Figures 1 and 2. Data points were recorded every 19–20 min.

3. Discussion

Previously it has been shown that huntingtin carries domains before and after the poly-Q tract that promote its reversible membrane association and it has been suggested that this membrane-association helps in polyQ aggregation [12,26,29,43]. In particular, the N-terminal 17 amino acids preceding the polyQ tract of huntingtin (usually abbreviated htt17 or N17 or htt^{N17}) are involved in the regulation of the spatio-temporal distribution of huntingtin or fragments thereof. Their association with membrane components and with different cellular compartments has been shown important for the development of symptoms of Huntington's disease [8,12,15,19–23,29,33,39–41]. Here we show that htt17-polyQ membrane association indeed strongly accelerates polypeptide aggregation in a manner that is dependent on the number of glutamines. A series of sequences was investigated with htt17 as a membrane anchor and various polyQ extensions (Table 1).

In order to quantitatively measure the aggregation kinetics biophysical experiments were performed at polypeptide concentrations of 3 to 30 μ M. This relatively low concentration is still suitable for spectroscopic analysis, but longer polyQ constructs tend to aggregate at very short time scales (not shown). Therefore, quantitative studies of the speed of aggregation were performed with constructs carrying a limiting number of glutamines (see also [83]). It should be noted that in their natural context the solubility of the full-length protein or its exon 1 domain is increased by the polyproline flanking se-

quence following the polyQ domain [31,32,40,48]. Therefore, as an alternative, exon 1 fragments were investigated where the carboxy-terminal proline-rich domain slows down the aggregation process probably through interactions with polyQ and htt17 (e.g. [26,42,84]). Because such additional interactions add an additional layer of complexity to the system here we have chosen to investigate the role of htt17 membrane interactions in polyglutamine aggregation using shorter constructs that aggregate at a rate still accessible to biophysical investigation. Furthermore, in the cellular environment, interactions with other proteins, chaperones and proteases, as well as other domains of the full-length protein assure that healthy cells are protected from huntingtin aggregation [85,86]. Therefore, it takes many years to develop the disease even when much longer polyQ mutants are present [1].

When diluted in aqueous buffer CD spectra of the peptides with short polyQ additions are indicative of predominantly random coil structure and some features of α -helical/ β -sheet conformations. Upon addition of POPC/POPS 3/1 membranes the CD spectra of htt17-polyglutamines carrying 12 or more glutamines change in appearance over the next few hours (Fig 1). Notably, the CD spectra of htt17-Q17 exhibit similar features when compared to htt(1-40), a construct carrying htt17, 17 glutamines and 6 prolines [52]. This observation suggests related structural changes in solution and in the presence of membranes. Electron microscopic images indicate that liposomes agglutinate with htt17-Q17 proteinaceous fibres (Fig. S1B-D). The increase in PDI observed by DLS (Fig. 5), the thioflavin fluorescence (Figs. 3 and 4) and the EM pictures (Fig. S1) are indicative that the peptides aggregate in the presence of membranes in a manner that depends on the peptide-to-lipid ratio (Fig. 4). In the case of htt17-Q17 the changes in the CD spectra obtained at concentrations of about 30 μ M peptide and 650 μ M lipid reach saturation after about 1.5h (Figs. 1 and 2).

At longer incubation times CD- and ThT fluorescence spectra decrease in intensity (Figs. 2 and 4C) because the larger peptide-lipid aggregates sink to the ground and/or stick to the surface of the glass tubes (Fig. S1). The spectral changes are suggestive of an increase in β -sheet conformation (Fig. 1) in agreement with other structural data [44,49,52,75], when at the same time a continuous increase in hydrodynamic radius is observed (Fig 5). Concomitant with this aggregation a quantitative analysis of the resulting CD spectra is hampered by light diffraction artefacts (Figs. 1F, 2 and 5). Peptides carrying a shorter polyQ segment aggregate more slowly or do not show spectral changes at all (Fig. 1D,E). The degree of aggregation increases but also its kinetics accelerates with the number of glutamines.

Whereas under the conditions investigated the peptides studied in this work remain in solution over many hours, even days in the absence of POPC/POPS vesicles, here we show that the addition of membranes strongly catalyses the aggregation process. Indeed, whereas aggregate formation *de novo* or through seeding are well established pathways, membranes have been shown to provide a third aggregation mechanism [26,87]. The data presented agrees with previous investigations where GDT-exon1 constructs have been found associated with rat postsynaptic membranes and where brain lipid vesicles accelerate nucleation and thereby fibril formation upon trypsin cleavage of 3 μ M GST-exon 1 encompassing 51 glutamines [24]. In the same study the presence of zwitterionic DMPC or of DOPC/SM/cholesterol slowed down this effect thereby being in-line with studies where htt17 membrane interactions were weak or absent for zwitterionic and cholesterol-rich membranes [43,88]. In a related manner a recent paper has revealed the importance of electrostatic contributions to the membrane-association of htt17 within exon 1 and the possible interference of amino acid modifications introducing additional negative charges [44].

The N-terminal 17-residue sequence has been demonstrated to adopt a largely helical conformations when being membranes-associated [44-46,55,89], when part of a htt17-polyQ fiber [42,49,57,90] or in aggregation intermediates [62,83]. Here we have shown that the membrane interactions of the htt17 flanking region of the polyQ domain

promotes the polyglutamine aggregation process. In a related manner, Htt17 also plays a leading role in catalysing the nucleation process during polyQ aggregation in solution. The htt17 amphipathic helix has been shown to associate into small oligomeric structures which serve as nucleation sites [26,62,83] from which polyQ fibrils elongate [26,90,91]. Furthermore, the flanking sequences play important additional roles in regulating the proteolytic degradation of the protein and its aggregates where the toxicity depends in a complex manner on the conformational subpopulation rather than the polyQ aggregation propensity [30,32,62,92].

Flanking regions of other polyQ proteins seem equally important in the regulation of their aggregation and fiber formation [1]. In particular previous studies showed that posttranslational modifications of flanking regions such as phosphorylation, SUMOylation or ubiquitination including of htt17 have an effect on polyQ aggregation [1,30,34–37]. This has been attributed to the changes of htt17 oligomerization being a consequence of such modifications, as well as modifications of its interaction surface with chaperones. The work presented here suggest that also its interactions with membranes should be strongly modified when e.g. negative phosphates are attached to its serines [28], when lysines are made unavailable for protein-protein and protein-lipid interactions [55,89,93] or the overall positive charge of htt17 is neutralized or inverted by posttranslational modifications thus abolishing the electrostatic attraction of the flanking regions to anionic membranes [12,44,55]. The numerous possibilities to interfere with protein-protein and protein-lipid interactions and thereby fibre formation explain why not all polyQ-extension diseases follow the same pattern and why the age onset of Huntington disease shows significant variation even for the same number of glutamines [1].

We suggest that the amphipathic htt17 helix, which reversibly associates with the membrane interface [44,55,89], concentrates and aligns the polyQ chains in such a manner to facilitate intermolecular interactions and fiber association [12,43]. Once the interactions with the membrane are established the aggregation process is enhanced by high peptide density (i.e. P/L ratio) and by extended polyQ chains but slowed down by the presence of the polyproline stretch that follows the poly-Q domain [48,94]. The polyQ length dependence may arise from the need to adopt an intermediate coiled-coil structure before transition into a β -sheet amyloid [52] as well as from interactions with other proteins or polypeptide domains [1,25,27]. Thereby these and other biophysical observations provide a rationale for a number of biochemical and cell biological experiments where huntingtin has been shown associated with membranes of intracellular organelles [8,12,15,19–23,28] or where the membrane anchoring domain htt17 has been demonstrated to promote the development of the disease [8,15,29,33,39–41].

Such lipid interactions have been shown to be modulators of aggregation and fibril formation also for α -synuclein [67,72,74], islet amyloid polypeptide [11] and β -amyloid [68,95]. Within these studies the detailed membrane composition including the resulting physico-chemical properties such as negative charge density, fluidity, saturation, curvature or interactions with specific lipids all play important roles in the aggregation process [11,43,44,66,88,96]. Thereby, the membrane interactions of polyQ flanking regions and their modulation by posttranslational modifications provides a possible therapeutic intervention site which has to our knowledge not been explored in greater detail.

4. Materials and Methods

4.1. Materials

All lipids were purchased from Avanti Polar Lipids (Alabaster, AL, USA). Water (HPLC grade), acetonitrile (99.8% HPLC grade), hexafluoro-2-propanol (99.5%), trifluoroacetic acid (TFA) (99.5%) and thioflavin (ThT) were from Sigma (St. Quentin Fallavier, France).

4.2. Peptide synthesis and purification

The peptides with the sequences shown in Table 1 were prepared by solid-phase synthesis using a Millipore 9050 automated peptide synthesizer and its standard Fmoc

(9-fluorenylme- thyoxy carbonyl) chemistry. They were purified by semi-preparative reverse phase high-performance liquid chromatography (Gilson, Villiers-le-Bel, France) using a preparative C18 column (Luna, C18-100Å-5µm, Phenomenex, Le Pecq, France) and an acetonitrile/water gradient. Their identity and purity (> 90%) were verified by analytical HPLC and MALDI mass spectrometry (MALDI-TOF Autoflex, Bruker Daltonics, Bremen, Germany). The fractions of interest were lyophilized for storage at -20°C.

4.3. Small unilamellar vesicles

The lipids were dissolved in chloroform/methanol (2/1 v/v). The solvent was evaporated under a flow of nitrogen gas in such a manner to form a homogeneous film on the walls of glass test tubes. The remaining solvent was removed by exposure for about 12 hours to high vacuum ($P < 100$ mPa). Subsequently, the lipids were hydrated in 10 mM Tris-HCl, pH 7 for more than an hour and subjected to several freeze-thaw cycles. Small unilamellar vesicles (SUV) were obtained after less than 1 min of tip sonication (Bandelin Sonopuls HD 200, Berlin, Germany). The lipid composition was chosen to represent the overall charge and main lipid components of the inner leaflet of the plasma membrane [97,98].

4.4. Circular dichroism spectroscopy

Circular dichroism (CD) spectra were recorded from 260 to 194 nm (spectral resolution: 1 nm, data pitch: 1 nm, scan speed: 100 nm/min) using a J-810 spectropolarimeter (Jasco, Tokyo, Japan). 20 µL of peptide solution (at 1 mg/mL, in 10 mM Tris-HCl buffer pH 7) were transferred into a quartz cuvette of 1 mm path length and mixed with 200 µL of SUVs ($C = 0.5$ mg lipid/mL) to reach the final concentration of peptide $\approx 9.1 \cdot 10^{-2}$ mg/mL (corresponding to 29, 26 and 22 µM for htt17-Q9, htt17-Q12 and htt17-Q17, respectively). The mixture was vortexed for ≈ 15 s prior to spectral acquisition. The secondary structure composition of the peptides was calculated from the spectra using a linear least-squares method implemented in the DicroProt analysis software [76].

4.5. Measurements of thioflavin-T fluorescence

A 100 mM stock solution of Thioflavin-T (ThT) was prepared in 10 mM Tris-HCl, pH 7 and stored in the dark at -20°C. The solution was thawed and diluted to 1 mM in the same buffer on the day of analysis. To prepare samples for fluorescence spectroscopy, peptide powders were suspended in some mL of mixed HFIP and TFA (1/1), dried and dissolved in the same solvent at least three times in order to obtain a clear solution. Thereafter, for each sample preparation an aliquot containing 0.3 mg of peptide, was taken and dried under a flow of N₂ gas for more than 30 minutes. A few mL of SUV suspension (at 0.25 mg/mL) was added to the dried peptide to reach the appropriate peptide-to-lipid ratio together with some µL of ThT solution to reach a 5 µM final concentration. The solution was then vortexed for about one minute. 1 mL of sample was rapidly transferred into a quartz cuvette and immediately placed in the sample holder of a Fluorolog 3-22 spectrometer (Horiba Jobin-Yvon, Longjumeau, France). The sample was excited at $\lambda_{exc} = 440$ nm while the dispersed fluorescence intensity was either recorded from $\lambda_{fluo} = 455$ to 651 nm or at the fixed wavelength of $\lambda_{fluo} = 485$ nm at a constant temperature of 25 °C. A resolution of $\Delta\lambda = 4$ nm was chosen for excitation as well as for analysis in order to obtain a good signal-to-noise ratio. The sample was constantly stirred at low speed with a small size rotated Teflon coated magnetic bar.

4.6. Dynamic light scattering

Measurements were performed on a Zetasizer Nano-S system (Malvern Instruments, Worcestershire, UK) equipped with a 4 mW He-Ne laser. Samples containing the mixture of peptides and vesicles were placed in a low volume quartz cell equilibrated at 25°C, and the light scattered backward was collected at an angle of $\theta = 173^\circ$. Data analysis was performed with the DTS Malvern software implemented on a personal computer.

4.7. Data analysis

To analyse the kinetic data, *i.e.* fluorescence intensity of ThT at 485 nm or circular dichroism at 208 nm as a function of time, a standard least-square fit analysis was implemented numerically. A three parameter single exponential function, $I(t) = A + B \cdot (1 - \exp(-t/\tau))$ was found to describe well the intensity increase of the signals.

Supplementary Materials: The following are available online at www.mdpi.com/xxx/s1, Figure S1: Picture and TEM image of htt17-Q17 in the presence of 100 nm LUVs.

Author Contributions: Methodology, formal analysis, investigation and data curation, A.M. and C.A.; writing—original draft preparation, A.M. and B.B.; conceptualization, writing—review and editing, all authors.; project administration and funding acquisition, B.B. All authors have read and agreed to the published version of the manuscript.

Funding: The financial contributions of the Agence Nationale de la Recherche (projects MemPep-Syn 14-CE34-0001-01, InMembrane 15-CE11-0017-01, Biosupramol 17-CE18-0033-3 and the LabEx Chemistry of Complex Systems 10-LABX-0026_CSC), the University of Strasbourg, the CNRS, the Région Grand-Est and the RTRA International Center of Frontier Research in Chemistry, the American Foundation for Research on Huntington's Disease (CHDI) are gratefully acknowledged.

Data Availability Statement: Data are available on request

Acknowledgments: We are grateful to Nicole Harmouche, Rabia Sarroukh, Caroline Lonez, Erick Goormaghtigh and Jean-Marie Ruysschaert for providing data of unpublished experiments and for discussion as well as Delphine Hatay for technical help during peptide synthesis and purification.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Kuiper, E.F.; de Mattos, E.P.; Jardim, L.B.; Kampinga, H.H.; Bergink, S. Chaperones in Polyglutamine Aggregation: Beyond the Q-Stretch. *Front Neurosci* **2017**, *11*, 145, doi:10.3389/fnins.2017.00145.
- Aktar, F.; Burudpakdee, C.; Polanco, M.; Pei, S.; Swayne, T.C.; Lipke, P.N.; Emtage, L. The huntingtin inclusion is a dynamic phase-separated compartment. *Life Sci Alliance* **2019**, *2*, doi:10.26508/lsa.201900489.
- Peskett, T.R.; Rau, F.; O'Driscoll, J.; Patani, R.; Lowe, A.R.; Saibil, H.R. A Liquid to Solid Phase Transition Underlying Pathological Huntingtin Exon1 Aggregation. *Mol Cell* **2018**, *70*, 588–601 e586, doi:10.1016/j.molcel.2018.04.007.
- Pandey, M.; Rajamma, U. Huntington's disease: the coming of age. *J Genet* **2018**, *97*, 649–664.
- Rubinsztein, D.C.; Leggo, J.; Coles, R.; Almqvist, E.; Biancalana, V.; Cassiman, J.J.; Chotai, K.; Connarty, M.; Crauford, D.; Curtis, A.; et al. Phenotypic characterization of individuals with 30–40 CAG repeats in the Huntington disease (HD) gene reveals HD cases with 36 repeats and apparently normal elderly individuals with 36–39 repeats. *Am J Hum Genet* **1996**, *59*, 16–22.
- Andresen, J.M.; Gayan, J.; Djousse, L.; Roberts, S.; Brocklebank, D.; Cherny, S.S.; Cardon, L.R.; Gusella, J.F.; MacDonald, M.E.; Myers, R.H.; et al. The relationship between CAG repeat length and age of onset differs for Huntington's disease patients with juvenile onset or adult onset. *Annals of Human Genetics* **2007**, *71*, 295–301.
- Wanker, E.E.; Ast, A.; Schindler, F.; Trepte, P.; Schnoegl, S. The pathobiology of perturbed mutant huntingtin protein-protein interactions in Huntington's disease. *J Neurochem* **2019**, *151*, 507–519, doi:10.1111/jnc.14853.
- Rockabrand, E.; Slepko, N.; Pantalone, A.; Nukala, V.N.; Kazantsev, A.; Marsh, J.L.; Sullivan, P.G.; Steffan, J.S.; Sensi, S.L.; Thompson, L.M. The first 17 amino acids of Huntingtin modulate its sub-cellular localization, aggregation and effects on calcium homeostasis. *Hum Mol Genet* **2007**, *16*, 61–77, doi:10.1093/hmg/ddl440 [pii] 10.1093/hmg/ddl440.
- Monoi, H.; Futaki, S.; Kugimiya, S.; Minakata, H.; Yoshihara, K. Poly-L-glutamine forms cation channels: relevance to the pathogenesis of the polyglutamine diseases. *Biophys.J* **2000**, *78*, 2892–2899.
- Terakawa, M.S.; Lin, Y.; Kinoshita, M.; Kanemura, S.; Itoh, D.; Sugiki, T.; Okumura, M.; Ramamoorthy, A.; Lee, Y.H. Impact of membrane curvature on amyloid aggregation. *Biochim Biophys Acta* **2018**, doi:10.1016/j.bbame.2018.04.012.

11. Relini, A.; Marano, N.; Gliozzi, A. Probing the interplay between amyloidogenic proteins and membranes using lipid monolayers and bilayers. *Adv Colloid Interface Sci* **2014**, *207*, 81-92, doi:10.1016/j.cis.2013.10.015.
12. Kegel-Gleason, K.B. Huntingtin interactions with membrane phospholipids: strategic targets for therapeutic intervention? *J Huntingtons Dis* **2013**, *2*, 239-250, doi:10.3233/JHD-130068.
13. del Toro, D.; Alberch, J.; Lazaro-Dieguez, F.; Martin-Ibanez, R.; Xifro, X.; Egea, G.; Canals, J.M. Mutant huntingtin impairs post-Golgi trafficking to lysosomes by delocalizing optineurin/Rab8 complex from the Golgi apparatus. *Mol Biol Cell* **2009**, *20*, 1478-1492, doi:10.1091/mbc.E08-07-0726.
14. Caviston, J.P.; Holzbaur, E.L. Huntingtin as an essential integrator of intracellular vesicular trafficking. *Trends Cell Biol* **2009**, *19*, 147-155, doi:10.1016/j.tcb.2009.01.005.
15. Atwal, R.S.; Xia, J.; Pinchev, D.; Taylor, J.; Epand, R.M.; Truant, R. Huntingtin has a membrane association signal that can modulate huntingtin aggregation, nuclear entry and toxicity. *Hum Mol Genet* **2007**, *16*, 2600-2615, doi:ddm217 [pii] 10.1093/hmg/ddm217.
16. Atwal, R.S.; Truant, R. A stress sensitive ER membrane-association domain in Huntingtin protein defines a potential role for Huntingtin in the regulation of autophagy. *Autophagy* **2008**, *4*, 91-93, doi:5201 [pii].
17. Leitman, J.; Ulrich Hartl, F.; Lederkremer, G.Z. Soluble forms of polyQ-expanded huntingtin rather than large aggregates cause endoplasmic reticulum stress. *Nat Commun* **2013**, *4*, 2753, doi:10.1038/ncomms3753.
18. Flavin, W.P.; Bousset, L.; Green, Z.C.; Chu, Y.; Skarpathiotis, S.; Chaney, M.J.; Kordower, J.H.; Melki, R.; Campbell, E.M. Endocytic vesicle rupture is a conserved mechanism of cellular invasion by amyloid proteins. *Acta Neuropathol* **2017**, *134*, 629-653, doi:10.1007/s00401-017-1722-x.
19. Tanaka, Y.; Igarashi, S.; Nakamura, M.; Gafni, J.; Torcassi, C.; Schilling, G.; Crippen, D.; Wood, J.D.; Sawa, A.; Jenkins, N.A.; et al. Progressive phenotype and nuclear accumulation of an amino-terminal cleavage fragment in a transgenic mouse model with inducible expression of full-length mutant huntingtin. *Neurobiol Dis* **2006**, *21*, 381-391, doi:S0969-9961(05)00216-0 [pii]10.1016/j.nbd.2005.07.014.
20. Hackam, A.S.; Hodgson, J.G.; Singaraja, R.; Zhang, T.Q.; Gan, L.; Gutekunst, C.A.; Hersch, S.M.; Hayden, M.R. Evidence for both the nucleus and cytoplasm as subcellular sites of pathogenesis in Huntington's disease in cell culture and in transgenic mice expressing mutant huntingtin. *Philosophical Transactions of the Royal Society of London Series B-Biological Sciences* **1999**, *354*, 1047-1055.
21. Xia, J.; Lee, D.H.; Taylor, J.; Vandelft, M.; Truant, R. Huntingtin contains a highly conserved nuclear export signal. *Hum Mol Genet* **2003**, *12*, 1393-1403.
22. Panov, A.V.; Gutekunst, C.A.; Leavitt, B.R.; Hayden, M.R.; Burke, J.R.; Strittmatter, W.J.; Greenamyre, J.T. Early mitochondrial calcium defects in Huntington's disease are a direct effect of polyglutamines. *Nature neuroscience* **2002**, *5*, 731-736, doi:10.1038/nn884 [pii].
23. Choo, Y.S.; Johnson, G.V.; MacDonald, M.; Detloff, P.J.; Lesort, M. Mutant huntingtin directly increases susceptibility of mitochondria to the calcium-induced permeability transition and cytochrome c release. *Hum Mol Genet* **2004**, *13*, 1407-1420, doi:10.1093/hmg/ddh162[pii].
24. Suopanki, J.; Gotz, C.; Lutsch, G.; Schiller, J.; Harjes, P.; Herrmann, A.; Wanker, E.E. Interaction of huntingtin fragments with brain membranes--clues to early dysfunction in Huntington's disease. *J Neurochem* **2006**, *96*, 870-884, doi:10.1111/j.1471-4159.2005.03620.x.
25. Burke, K.A.; Yates, E.A.; Legleiter, J. Biophysical insights into how surfaces, including lipid membranes, modulate protein aggregation related to neurodegeneration. *Front Neurol* **2013**, *4*, 17, doi:10.3389/fneur.2013.00017.
26. Pandey, N.K.; Isas, J.M.; Rawat, A.; Lee, R.V.; Langen, J.; Pandey, P.; Langen, R. The 17-residue-long N terminus in huntingtin controls stepwise aggregation in solution and on membranes via different mechanisms. *J Biol Chem* **2018**, *293*, 2597-2605, doi:10.1074/jbc.M117.813667.
27. Nagarajan, A.; Jawahery, S.; Matysiak, S. The effects of flanking sequences in the interaction of polyglutamine peptides with a membrane bilayer. *J Phys Chem B* **2014**, *118*, 6368-6379, doi:10.1021/jp407900c.

28. Atwal, R.S.; Desmond, C.R.; Caron, N.; Maiuri, T.; Xia, J.R.; Sipione, S.; Truant, R. Kinase inhibitors modulate huntingtin cell localization and toxicity. *Nature Chemical Biology* **2011**, *7*, 453–460, doi:10.1038/Nchembio.582.
29. Arndt, J.R.; Chaibva, M.; Legleiter, J. The emerging role of the first 17 amino acids of huntingtin in Huntington's disease. *Biomol Concepts* **2015**, *6*, 33–46, doi:10.1515/bmc-2015-0001.
30. Chaibva, M.; Jawahery, S.; Pilkington, A.W.t.; Arndt, J.R.; Sarver, O.; Valentine, S.; Matysiak, S.; Legleiter, J. Acetylation within the First 17 Residues of Huntingtin Exon 1 Alters Aggregation and Lipid Binding. *Biophys J* **2016**, *111*, 349–362, doi:10.1016/j.bpj.2016.06.018.
31. Crick, S.L.; Ruff, K.M.; Garai, K.; Frieden, C.; Pappu, R.V. Unmasking the roles of N- and C-terminal flanking sequences from exon 1 of huntingtin as modulators of polyglutamine aggregation. *Proc Natl Acad Sci U S A* **2013**, *110*, 20075–20080, doi:10.1073/pnas.1320626110.
32. Shen, K.; Calamini, B.; Fauerbach, J.A.; Ma, B.; Shahmoradian, S.H.; Serrano Lachapel, I.L.; Chiu, W.; Lo, D.C.; Frydman, J. Control of the structural landscape and neuronal proteotoxicity of mutant Huntingtin by domains flanking the polyQ tract. *Elife* **2016**, *5*, doi:10.7554/eLife.18065.
33. Gu, X.F.; Greiner, E.R.; Mishra, R.; Kodali, R.; Osmand, A.; Finkbeiner, S.; Steffan, J.S.; Thompson, L.M.; Wetzel, R.; Yang, X.W. Serines 13 and 16 Are Critical Determinants of Full-Length Human Mutant Huntingtin Induced Disease Pathogenesis in HD Mice. *Neuron* **2009**, *64*, 828–840, doi:10.1016/J.Neuron.2009.11.020.
34. Thompson, L.M.; Aiken, C.T.; Kaltenbach, L.S.; Agrawal, N.; Illes, K.; Khoshnan, A.; Martinez-Vincente, M.; Arrasate, M.; O'Rourke, J.G.; Khashwji, H.; et al. IKK phosphorylates Huntingtin and targets it for degradation by the proteasome and lysosome. *Journal of Cell Biology* **2009**, *187*, 1083–1099, doi:10.1083/Jcb.200909067.
35. Aiken, C.T.; Steffan, J.S.; Guerrero, C.M.; Khashwji, H.; Lukacsovich, T.; Simmons, D.; Purcell, J.M.; Menhaji, K.; Zhu, Y.Z.; Green, K.; et al. Phosphorylation of threonine 3: implications for Huntingtin aggregation and neurotoxicity. *J Biol Chem* **2009**, *284*, 29427–29436, doi:10.1074/jbc.M109.013193.
36. Steffan, J.S.; Agrawal, N.; Pallos, J.; Rockabrand, E.; Trotman, L.C.; Slepko, N.; Illes, K.; Lukacsovich, T.; Zhu, Y.Z.; Cattaneo, E.; et al. SUMO modification of Huntingtin and Huntington's disease pathology. *Science* **2004**, *304*, 100–104, doi:10.1126/Science.1092194.
37. Zheng, Z.Q.; Li, A.M.; Holmes, B.B.; Marasa, J.C.; Diamond, M.I. An N-terminal Nuclear Export Signal Regulates Trafficking and Aggregation of Huntingtin (Htt) Protein Exon 1. *Journal of Biological Chemistry* **2013**, *288*, 6063–6071, doi:10.1074/Jbc.M112.413575.
38. Sedighi, F.; Adegboyiro, A.; Legleiter, J. SUMOylation Prevents Huntingtin Fibrillization and Localization onto Lipid Membranes. *ACS Chem Neurosci* **2020**, *11*, 328–343, doi:10.1021/acscchemneuro.9b00509.
39. Kelley, N.W.; Huang, X.; Tam, S.; Spiess, C.; Frydman, J.; Pande, V.S. The predicted structure of the headpiece of the Huntingtin protein and its implications on Huntingtin aggregation. *J Mol Biol* **2009**, *388*, 919–927, doi:10.1016/j.jmb.2009.01.032.
40. Thakur, A.K.; Jayaraman, M.; Mishra, R.; Thakur, M.; Chellgren, V.M.; Byeon, I.J.; Anjum, D.H.; Kodali, R.; Creamer, T.P.; Conway, J.F.; et al. Polyglutamine disruption of the huntingtin exon 1 N terminus triggers a complex aggregation mechanism. *Nat Struct Mol Biol* **2009**, *16*, 380–389, doi:10.1038/nsmb.1570.
41. Tam, S.; Spiess, C.; Auyeung, W.; Joachimiak, L.; Chen, B.; Poirier, M.A.; Frydman, J. The chaperonin TRiC blocks a huntingtin sequence element that promotes the conformational switch to aggregation. *Nat Struct Mol Biol* **2009**, *16*, 1279–1285, doi:10.1038/nsmb.1700.
42. Boatz, J.C.; Piretra, T.; Lasorsa, A.; Matlahov, I.; Conway, J.F.; van der Wel, P.C.A. Protofilament Structure and Supramolecular Polymorphism of Aggregated Mutant Huntingtin Exon 1. *J Mol Biol* **2020**, *432*, 4722–4744, doi:10.1016/j.jmb.2020.06.021.

43. Michalek, M.; Salnikov, E.S.; Werten, S.; Bechinger, B. Membrane interactions of the amphipathic amino-terminus of huntingtin. *Biochemistry* **2013**, *52*, 847–858.
44. Tao, M.; Pandey, N.K.; Barnes, R.; Han, S.; Langen, R. Structure of Membrane-Bound Huntingtin Exon 1 Reveals Membrane Interaction and Aggregation Mechanisms. *Structure* **2019**, *27*, 1570–1580 e1574, doi:10.1016/j.str.2019.08.003.
45. Levy, G.R.; Shen, K.; Gavrilov, Y.; Smith, P.E.S.; Levy, Y.; Chan, R.; Frydman, J.; Frydman, L. Huntingtin's N-Terminus Rearrangements in the Presence of Membranes: A Joint Spectroscopic and Computational Perspective. *ACS Chem Neurosci* **2019**, *10*, 472–481, doi:10.1021/acscchemneuro.8b00353.
46. Ceccon, A.; Clore, G.M.; Tugarinov, V. Decorrelating Kinetic and Relaxation Parameters in Exchange Saturation Transfer NMR: A Case Study of N-Terminal Huntingtin Peptides Binding to Unilamellar Lipid Vesicles. *J Phys Chem B* **2018**, *122*, 11271–11278, doi:10.1021/acs.jpcc.8b07112.
47. Karanji, A.K.; Beasley, M.; Sharif, D.; Ranjbaran, A.; Legleiter, J.; Valentine, S.J. Investigating the interactions of the first 17 amino acid residues of Huntingtin with lipid vesicles using mass spectrometry and molecular dynamics. *J Mass Spectrom* **2020**, *55*, e4470, doi:10.1002/jms.4470.
48. Arndt, J.R.; Chaibva, M.; Beasley, M.; Kiani Karanji, A.; Ghassabi Kondalaji, S.; Khakinejad, M.; Sarver, O.; Legleiter, J.; Valentine, S.J. Nucleation Inhibition of Huntingtin Protein (htt) by Polyproline PPII Helices: A Potential Interaction with the N-Terminal alpha-Helical Region of Htt. *Biochemistry* **2020**, *59*, 436–449, doi:10.1021/acs.biochem.9b00689.
49. Hoop, C.L.; Lin, H.K.; Kar, K.; Hou, Z.; Poirier, M.A.; Wetzel, R.; van der Wel, P.C. Polyglutamine amyloid core boundaries and flanking domain dynamics in huntingtin fragment fibrils determined by solid-state nuclear magnetic resonance. *Biochemistry* **2014**, *53*, 6653–6666, doi:10.1021/bi501010q.
50. Ceccon, A.; Tugarinov, V.; Clore, G.M. Kinetics of Fast Tetramerization of the Huntingtin Exon 1 Protein Probed by Concentration-Dependent On-Resonance R1rho Measurements. *J Phys Chem Lett* **2020**, *11*, 5643–5648, doi:10.1021/acs.jpclett.0c01636.
51. Kotler, S.A.; Tugarinov, V.; Schmidt, T.; Ceccon, A.; Libich, D.S.; Ghirlando, R.; Schwieters, C.D.; Clore, G.M. Probing initial transient oligomerization events facilitating Huntingtin fibril nucleation at atomic resolution by relaxation-based NMR. *Proc Natl Acad Sci U S A* **2019**, *116*, 3562–3571, doi:10.1073/pnas.1821216116.
52. Fiumara, F.; Fioriti, L.; Kandel, E.R.; Hendrickson, W.A. Essential role of coiled coils for aggregation and activity of Q/N-rich prions and PolyQ proteins. *Cell* **2010**, *143*, 1121–1135, doi:10.1016/j.cell.2010.11.042.
53. Miettinen, Markus S.; Monticelli, L.; Nedumpully-Govindan, P.; Knecht, V.; Ignatova, Z. Stable Polyglutamine Dimers Can Contain β -Hairpins with Interdigitated Side Chains—But Not α -Helices, β -Nanotubes, β -Pseudohelices, or Steric Zippers. *Biophysical Journal* **2014**, *106*, 1721–1728, doi:<https://doi.org/10.1016/j.bpj.2014.02.027>.
54. Miettinen, M.S.; Knecht, V.; Monticelli, L.; Ignatova, Z. Assessing Polyglutamine Conformation in the Nucleating Event by Molecular Dynamics Simulations. *The Journal of Physical Chemistry B* **2012**, *116*, 10259–10265, doi:10.1021/jp305065c.
55. Michalek, M.; Salnikov, E.; Werten, S.; Bechinger, B. Structure and topology of the huntingtin 1–17 membrane anchor by a combined solution and solid-state NMR approach. *Biophys. J.* **2013**, *105*, 699–710.
56. Michalek, M.; Aisenbrey, C.; Bechinger, B. Investigation of membrane penetration depth and interactions of the amino-terminal domain of huntingtin: refined analysis by tryptophan fluorescence measurement. *Eur Biophys J* **2014**, *43*, 347–360, doi:10.1007/s00249-014-0966-9.
57. Bravo-Arredondo, J.M.; Kegulian, N.C.; Schmidt, T.; Pandey, N.K.; Situ, A.J.; Ulmer, T.S.; Langen, R. The folding equilibrium of huntingtin exon 1 monomer depends on its polyglutamine tract. *J Biol Chem* **2018**, *293*, 19613–19623, doi:10.1074/jbc.RA118.004808.
58. Kokona, B.; Rosenthal, Z.P.; Fairman, R. Role of the coiled-coil structural motif in polyglutamine aggregation. *Biochemistry* **2014**, *53*, 6738–6746, doi:10.1021/bi500449a.

59. Warner, J.B.t.; Ruff, K.M.; Tan, P.S.; Lemke, E.A.; Pappu, R.V.; Lashuel, H.A. Monomeric Huntingtin Exon 1 Has Similar Overall Structural Features for Wild-Type and Pathological Polyglutamine Lengths. *J Am Chem Soc* **2017**, *139*, 14456-14469, doi:10.1021/jacs.7b06659.
60. Newcombe, E.A.; Ruff, K.M.; Sethi, A.; Ormsby, A.R.; Ramdhan, Y.M.; Fox, A.; Purcell, A.W.; Gooley, P.R.; Pappu, R.V.; Hatters, D.M. Tadpole-like Conformations of Huntingtin Exon 1 Are Characterized by Conformational Heterogeneity that Persists regardless of Polyglutamine Length. *J Mol Biol* **2018**, *430*, 1442-1458, doi:10.1016/j.jmb.2018.03.031.
61. Kokona, B.; Johnson, K.A.; Fairman, R. Effect of helical flanking sequences on the morphology of polyglutamine-containing fibrils. *Biochemistry* **2014**, *53*, 6747-6753, doi:10.1021/bi501066q.
62. Sahoo, B.; Arduini, I.; Drombosky, K.W.; Kodali, R.; Sanders, L.H.; Greenamyre, J.T.; Wetzel, R. Folding Landscape of Mutant Huntingtin Exon1: Diffusible Multimers, Oligomers and Fibrils, and No Detectable Monomer. *PLoS One* **2016**, *11*, e0155747, doi:10.1371/journal.pone.0155747.
63. Bulone, D.; Masino, L.; Thomas, D.J.; Biagio, P.L.S.; Pastore, A. The Interplay between PolyQ and Protein Context Delays Aggregation by Forming a Reservoir of Protofibrils. *PLoS One* **2006**, *1*, e111.
64. Ignatova, Z.; Thakur, A.K.; Wetzel, R.; Gierasch, L.M. In-cell aggregation of a polyglutamine-containing chimera is a multistep process initiated by the flanking sequence. *Journal of Biological Chemistry* **2007**, *282*, 36736-36743.
65. Beasley, M.; Stonebraker, A.R.; Hasan, I.; Kapp, K.L.; Liang, B.J.; Agarwal, G.; Groover, S.; Sedighi, F.; Legleiter, J. Lipid Membranes Influence the Ability of Small Molecules To Inhibit Huntingtin Fibrillization. *Biochemistry* **2019**, *58*, 4361-4373, doi:10.1021/acs.biochem.9b00739.
66. Chaibva, M.; Gao, X.; Jain, P.; Campbell, W.A.t.; Frey, S.L.; Legleiter, J. Sphingomyelin and GM1 Influence Huntingtin Binding to, Disruption of, and Aggregation on Lipid Membranes. *ACS Omega* **2018**, *3*, 273-285, doi:10.1021/acsomega.7b01472.
67. O'Leary, E.I.; Lee, J.C. Interplay between alpha-synuclein amyloid formation and membrane structure. *Biochim Biophys Acta Proteins Proteom* **2019**, *1867*, 483-491, doi:10.1016/j.bbapap.2018.09.012.
68. Okada, Y.; Okubo, K.; Ikeda, K.; Yano, Y.; Hoshino, M.; Hayashi, Y.; Kiso, Y.; Itoh-Watanabe, H.; Naito, A.; Matsuzaki, K. Toxic Amyloid Tape: A Novel Mixed Antiparallel/Parallel beta-Sheet Structure Formed by Amyloid beta-Protein on GM1 Clusters. *ACS Chem Neurosci* **2019**, *10*, 563-572, doi:10.1021/acchemneuro.8b00424.
69. Qiang, W.; Yau, W.M.; Lu, J.X.; Collinge, J.; Tycko, R. Structural variation in amyloid-beta fibrils from Alzheimer's disease clinical subtypes. *Nature* **2017**, *541*, 217-221, doi:10.1038/nature20814.
70. Zhu, D.; Bungart, B.L.; Yang, X.; Zhumadilov, Z.; Lee, J.C.; Askarova, S. Role of membrane biophysics in Alzheimer's-related cell pathways. *Front Neurosci* **2015**, *9*, 186, doi:10.3389/fnins.2015.00186.
71. Kotler, S.A.; Walsh, P.; Brender, J.R.; Ramamoorthy, A. Differences between amyloid-beta aggregation in solution and on the membrane: insights into elucidation of the mechanistic details of Alzheimer's disease. *Chemical Society Reviews* **2014**, *43*, 6692-6700, doi:10.1039/C3cs60431d.
72. Fusco, G.; De Simone, A.; Arosio, P.; Vendruscolo, M.; Veglia, G.; Dobson, C.M. Structural Ensembles of Membrane-bound alpha-Synuclein Reveal the Molecular Determinants of Synaptic Vesicle Affinity. *Sci Rep* **2016**, *6*, 27125, doi:10.1038/srep27125.
73. Pieri, L.; Madiona, K.; Bousset, L.; Melki, R. Fibrillar alpha-synuclein and huntingtin exon 1 assemblies are toxic to the cells. *Biophys J* **2012**, *102*, 2894-2905, doi:10.1016/j.bpj.2012.04.050.
74. Monsellier, E.; Bousset, L.; Melki, R. alpha-Synuclein and huntingtin exon 1 amyloid fibrils bind laterally to the cellular membrane. *Sci Rep* **2016**, *6*, 19180, doi:10.1038/srep19180.
75. Yushchenko, T.; Deuerling, E.; Hauser, K. Insights into the Aggregation Mechanism of PolyQ Proteins with Different Glutamine Repeat Lengths. *Biophys J* **2018**, *114*, 1847-1857, doi:10.1016/j.bpj.2018.02.037.
76. Deleage, G.; Geourjon, C. An interactive graphic program for calculating the secondary structure content of proteins from circular dichroism spectrum. *Comput. Appl. Biosci.* **1993**, *2*, 197-199.

77. Greenfield, N.J. Methods to estimate the conformation of proteins and polypeptides from circular dichroism data. *Anal Biochem* **1996**, *235*, 1-10, doi:10.1006/abio.1996.0084.
78. Vermeer, L.S.; Marquette, A.; Schoup, M.; Fenard, D.; Galy, A.; Bechinger, B. Simultaneous Analysis of Secondary Structure and Light Scattering from Circular Dichroism Titrations: Application to Vectofusin-1. *Sci Rep* **2016**, *6*, 39450, doi:10.1038/srep39450.
79. Streets, A.M.; Sourigues, Y.; Kopito, R. R.; Melki, R.; Quake, S. R. Simultaneous measurement of amyloid fibril formation by dynamic light scattering and fluorescence reveals complex aggregation kinetics. *PloS one* **2013**, *8*, e54541, doi:10.1371/journal.pone.0054541.
80. LeVine, H., 3rd. Quantification of beta-sheet amyloid fibril structures with thioflavin T. *Methods Enzymol* **1999**, *309*, 274-284, doi:10.1016/s0076-6879(99)09020-5.
81. Biancalana, M.; Koide, S. Molecular mechanism of Thioflavin-T binding to amyloid fibrils. *Biochim Biophys Acta* **2010**, *1804*, 1405-1412, doi:10.1016/j.bbapap.2010.04.001.
82. Marquette, A.; Lorber, B.; Bechinger, B. Reversible liposome association induced by LAH4: a peptide with potent antimicrobial and nucleic acid transfection activities. *Biophys J* **2010**, *98*, 2544-2553, doi:S0006-3495(10)00322-X [pii]10.1016/j.bpj.2010.02.042.
83. Jayaraman, M.; Kodali, R.; Sahoo, B.; Thakur, A.K.; Mayasundari, A.; Mishra, R.; Peterson, C.B.; Wetzel, R. Slow amyloid nucleation via alpha-helix-rich oligomeric intermediates in short polyglutamine-containing huntingtin fragments. *J Mol Biol* **2012**, *415*, 881-899, doi:10.1016/j.jmb.2011.12.010.
84. Ceccon, A.; Tugarinov, V.; Ghirlando, R.; Clore, G.M. Abrogation of prenucleation, transient oligomerization of the Huntingtin exon 1 protein by human profilin I. *Proc Natl Acad Sci U S A* **2020**, *117*, 5844-5852, doi:10.1073/pnas.1922264117.
85. Sethi, R.; Roy, I. Stabilization of elongated polyglutamine tracts by a helical peptide derived from N-terminal huntingtin. *IUBMB Life* **2020**, doi:10.1002/iub.2288.
86. Smith, D.M. Could a Common Mechanism of Protein Degradation Impairment Underlie Many Neurodegenerative Diseases? *J Exp Neurosci* **2018**, *12*, 1179069518794675, doi:10.1177/1179069518794675.
87. Juhl, D.W.; Glattard, E.; Lointier, M.; Bampilis, P.; Bechinger, B. The antimicrobial and synergistic activities of PGLa and magainin 2 fibrils. *Frontiers in Cellular and Infection Microbiology* **2020**, *10*, 526459.
88. Gao, X.; Campbell, W.A.t.; Chaibva, M.; Jain, P.; Leslie, A.E.; Frey, S.L.; Legleiter, J. Cholesterol Modifies Huntingtin Binding to, Disruption of, and Aggregation on Lipid Membranes. *Biochemistry* **2016**, *55*, 92-102, doi:10.1021/acs.biochem.5b00900.
89. Cote, S.; Binette, V.; Salnikov, E.S.; Bechinger, B.; Mousseau, N. Probing the Huntingtin 1-17 membrane anchor on a phospholipid bilayer by using all-atom simulations. *Biophys J* **2015**, *108*, 1187-1198, doi:S0006-3495(15)00127-7 [pii]10.1016/j.bpj.2015.02.001.
90. Sivanandam, V.N.; Jayaraman, M.; Hoop, C.L.; Kodali, R.; Wetzel, R.; van der Wel, P.C. The aggregation-enhancing huntingtin N-terminus is helical in amyloid fibrils. *J Am Chem Soc* **2011**, *133*, 4558-4566, doi:10.1021/ja110715f.
91. Scherzinger, E.; Sittler, A.; Schweiger, K.; Heiser, V.; Lurz, R.; Hasenbank, R.; Bates, G.P.; Lehrach, H.; Wanker, E.E. Self-assembly of polyglutamine-containing huntingtin fragments into amyloid-like fibrils: implications for Huntington's disease pathology. *Proc Natl Acad Sci U S A* **1999**, *96*, 4604-4609.
92. Nucifora, L.G.; Burke, K.A.; Feng, X.; Arbez, N.; Zhu, S.; Miller, J.; Yang, G.; Ratovitski, T.; Delannoy, M.; Muchowski, P.J.; et al. Identification of novel potentially toxic oligomers formed in vitro from mammalian-derived expanded huntingtin exon-1 protein. *J Biol Chem* **2012**, *287*, 16017-16028, doi:10.1074/jbc.M111.252577.
93. Arndt, J.R.; Brown, R.J.; Burke, K.A.; Legleiter, J.; Valentine, S.J. Lysine residues in the N-terminal huntingtin amphipathic alpha-helix play a key role in peptide aggregation. *J Mass Spectrom* **2015**, *50*, 117-126, doi:10.1002/jms.3504.
94. Nagarajan, A.; Jawahery, S.; Matysiak, S. the effects of flanking sequences in the interaction of polyglutamine peptides with a membrane bilayer. *J. Phys. Chem B* **2013**, *118*, 6368-6379.
95. Terakawa, M.S.; Lin, Y.; Kinoshita, M.; Kanemura, S.; Itoh, D.; Sugiki, T.; Okumura, M.; Ramamoorthy, A.; Lee, Y.H. Impact of membrane curvature on amyloid aggregation. *Biochim Biophys Acta Biomembr* **2018**, *2736*, doi:10.1016/j.bbamem.2018.04.012.

-
96. Matsuzaki, K.; Kato, K.; Yanagisawa, K. Ganglioside-Mediated Assembly of Amyloid beta-Protein: Roles in Alzheimer's Disease. *Prog Mol Biol Transl Sci* **2018**, *156*, 413-434, doi:10.1016/bs.pmbts.2017.10.005.
97. van Meer, G.; Voelker, D.R.; Feigenson, G.W. Membrane lipids: where they are and how they behave. *Nat.Rev.Mol Cell Biol.* **2008**, *9*, 112-124.
98. Kobayashi, T.; Menon, A.K. Transbilayer lipid asymmetry. *Curr Biol* **2018**, *28*, R386-R391, doi:10.1016/j.cub.2018.01.007.