Biomass production and physiology of *Chlorella vulgaris* during the early stages of immobilized state are affected by light intensity and inoculum cell density

Su Fang Li, Andrea Fanesi, Thierry Martin, Filipa Lopes*

Laboratoire Génie des Procédés et Matériaux (LGPM), CentraleSupélec, Université Paris-Saclay, 91190 Gif-sur-Yvette, France

** Corresponding author.*

E-mail address: filipa.lopes@centralesupelec.fr Tel: [+33 0617607634](tel:+330617607634)

ABSTRACT

The interest for biofilm-based systems for microalgae and related compounds production has been increasing lately. Although extensive literature has been reported on productivity, the physiological characterization (photosynthetic activity and composition) of attached cells at early stages of biofilm development has seldom been investigated. In this work, the effect of light intensity and inoculum cell density on 3-days *Chlorella vulgaris* biofilms developed on membranes was studied. Biomass production was clearly impacted by mechanism of photo-limitation occurring in biofilms acclimated to low light intensity ($50 \mu\text{mol m}^{-2} \text{s}^{-1}$). A higher electron transport capacity and lower chlorophyll content in biofilms at high light intensity ($500 \mu\text{mol m}^{-2} \text{s}^{-1}$) were also measured which are in line with patterns observed for suspended microalgae cultures. In addition, optimal conditions in terms of light ($250 \mu\text{mol m}^{-2} \text{s}^{-1}$) combined with low ($4.8 \times 10^6 \text{ cells cm}^{-2}$) or high inoculum density ($28.8 \times 10^6 \text{ cells cm}^{-2}$) were identified to optimize biomass and lipids production, respectively. On the whole, measuring physiological profiles of immobilized cells at the initial stages of biofilm development provides information to efficiently operate and optimize biofilm-based systems.

Keywords: *Chlorella vulgaris*, FTIR-spectroscopy, Macromolecular composition, Microalgae biofilms, Photosynthetic activity

1.Introduction

Microalgae are a promising source of valuable compounds (e.g. proteins, pigments, carbohydrates, lipids) produced by the conversion of photons into chemical energy via photosynthesis [1–4]. Nowadays, biofilm-based systems for microalgae cultivation, where cells are growing attached to a substrate, have gained attention due to their higher productivity, lower water requirement and harvesting costs compared with conventional suspended culture technology [5,6]. These systems have been widely used for biomass/biofuel production, CO₂ fixation, and wastewater treatment [6–9]. Among those, the twin-layer system [10,11], in which cells grow on porous supports such as membranes, printing paper or synthetic nonwoven/textile combinations, is often used at the lab and pilot scales [12,13]. In particular, a direct exposure to gas and light are supposed to enhance gas mass transfer and light utilization due to the absence of the liquid phase [14,15] improving therefore productivity. On the other hand, nutrients are transported by diffusion through the porous material into the biofilm. Limitation in nutrients can therefore occur in thick and/or compact biofilms, affecting in turn their behavior and productivity [11,16].

Recent studies investigated environmental and operational parameters that typically affect the formation and development of microalgae biofilms [17,18]. Among those factors, the impact of light, nutrients availability, temperature, pH and shear stress on growth rate and productivity have been studied, but mostly over long-term period of development (i.e. from weeks to months) [6,19–21]. On the other hand, other factors such as substrate properties (e.g. surface energy, hydrophobicity, micro-pattern, etc.) have been studied on the short-term formation of biofilms, especially during the initial adhesion phase (from

several hours to 2-3 days) [18,22–26]. However, little information is available regarding the influence of process operational factors, such as light intensity and inoculum density, on biofilm growth, activity and composition at the early stages of biofilm development which in turn may affect biomass and compounds production. In our work, early stage of biofilm development refers to a period during which the cells attach firmly to the support, grow and fully acclimate to the new environmental conditions imposed by the sessile growth associated to the support.

Ji et al. [27] have reported that both biomass production and growth rate could be improved by increasing the inoculum density (from 0.05 to 3 g DW m⁻²). However, it must be kept in mind that over a certain density threshold, the cells would not grow further due to light attenuation and nutrients transfer limitation [14,16,18]. In this context, light attenuation could be buffered by tuning the photon flux density (PFD) used for cultivation. However care must be taken since like for planktonic cultures, the growth rate and biomass productivity of microalgae biofilms increase with light intensity within a favorable range, but decrease if the PFD exceeds the light saturation point because of photo-inhibition [28]. In order to avoid such operational problems, monitoring the physiological properties of cells (activity and composition) in the early stages of immobilized growth might be of great help to rapidly identify possible limiting operational factors (e.g. light, nutrients, CO₂, humidity, etc.) and thus adjust them to optimize the bioprocess performance [11,14,15,29]. Nevertheless, although extensive literature has been reported on the productivity of biofilm-based systems as summarized in [30] and reference therein, the characterization of cell physiological parameters at early stages has seldom been mentioned.

The goal of our work was to assess the impact of light and initial cell density on biofilm/compounds production and on cells physiology at early stages (3 days) of biofilm formation in a twin layer system. In particular, we aimed at better understanding photosynthetic mechanisms on biofilms that certainly allow a reasoned choice of process operational parameters to optimize productivity. In order to do that, we immobilized *C. vulgaris* on membranes with two initial cell densities and illuminated the biofilms with three PFDs. A complete characterization of physiological parameters of the immobilized cells, from photosynthetic performance to macromolecular composition was conducted through which biofilm growth patterns and composition under different light and inoculum density combinations were explained.

2. Materials and Methods

2.1. Planktonic culture maintenance

Chlorella vulgaris SAG 211–11b (Göttingen, Germany) was cultured at 25°C semi-continuously in 1 L transparent bottles filled with 800 mL 3N-Bristol medium [31]. The cultures were bubbled with filtered air under a continuous illumination of either 50, 250 or 500 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Viugreum 50W LED outdoor floodlights, the PFDs were measured by QSL-2100 quantum scalar irradiance sensor, Biospherical Instruments, San Diego, CA, USA). The cultures were kept in exponential phase with a max cell concentration of 6.9×10^6 cells mL^{-1} (Flow cytometer, Guava EasyCyte HT; Millipore, USA) by daily dilution in order to maintain a chlorophyll (Chl) *a* concentration of 0.2-1.5 mg Chl *a* L^{-1} to ensure

optimal light penetration. The planktonic cultures were pre-acclimated to each light condition for at least 8 days before starting any experiment (for *C. vulgaris*, typically 5 days were enough to have a stabilization of Chl *a* content and growth rate).

2.2. Immobilization and growth of *C. vulgaris* on membranes

Sterile Petri dishes (55-mm diameter) filled with 3 mL of 3N Bristol medium were used as bioreactors for *C. vulgaris* immobilized growth (Fig. 1). The support system consisted of two glass fiber filters (working as an absorbing material for the medium; 47-mm diameter, Whatman) and on top of that a nitrate cellulose filter (0.2- μ m pore size; NC membrane, 25-mm diameter, Whatman) on which the cells of *C. vulgaris* were immobilized.

Figure 1. Schematic representation of the cultivation system used for the immobilized culture of *C. vulgaris* (NC membrane represents cellulose nitrate membrane filter).

132

133 In order to test the effect of PFD and the initial cell density on the immobilized growth
134 of *C. vulgaris*, the pre-acclimated planktonic cells (grown at 50, 250 and 500 $\mu\text{mol m}^{-2} \text{s}^{-1}$)
135 were vacuum-filtered on the NC membranes with an effective colonization area of 2.01 cm^2 .
136 Two initial cell densities corresponding to $\sim 4.8 \times 10^6 \text{ cells cm}^{-2}$ (low initial cell density,
137 LC, $0.4 \pm 0.1 \text{ g m}^{-2}$) and $\sim 28.8 \times 10^6 \text{ cells cm}^{-2}$ (high initial cell density, HC, $2.6 \pm 0.8 \text{ g m}^{-2}$)
138 were obtained by filtrating specific volumes of planktonic cultures on the membranes.
139 Once the cells were immobilized, the membranes were placed on the glass fiber filters and
140 placed in Petri dishes illuminated at 50 (low light, LL), 250 (moderate light, ML) or 500
141 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (high light, HL; Hansatech Instruments Quantitherm light meter/thermometer,
142 Norfolk, England) depending on the experiments. The membranes were incubated for three
143 days and the medium was completely renewed every day. After three days of growth, the
144 cells were harvested from the membranes using Bristol medium and further measurements
145 were conducted to characterize a series of physiological parameters.

146

147 2.3. Microscopic observation of initially immobilized cultures

148

149 At day 0, the immobilized cultures at two cell density were scanned with a confocal
150 laser scanning microscope (CLSM) to acquire z-stacks over the whole culture depth (Fig.
151 S1). The confocal microscope set-up was as described in [19]. An inverted Zeiss LSM700
152 confocal microscope (Carl Zeiss microscopy GmbH, Jena, Germany) equipped with a LD

Plan-Neofluar 20×/0.4 Korr M27 was used to acquire images. Microalgae cells were detected on the base of chlorophyll *a* auto-fluorescence which was excited at 639 nm.

2.4. Relative biomass increase

Cells were harvested with 10 mL fresh Bristol medium. For the dry weight determination, the cell suspension was centrifuged and further washed in 10 mL of Bristol medium. After a second centrifugation step, the pellet was dried at 100 °C until a constant weight over time was reached. The relative biomass increase with respect to the initial level (R_b) was determined according to Eq. (1):

$$R_b = (X_t - X_0) / X_0 \quad (1)$$

Where X_t is biomass areal density (g m^{-2}) after three days ($t = 3$ days), and X_0 is biomass areal density (g m^{-2}) at the beginning of the immobilization.

2.5. Variable chlorophyll a fluorescence measurements and relative electron transport rate (rETR) estimation

Photosynthetic parameters were determined using a portable pulse amplitude modulation (PAM) fluorometer (AquaPen, AP 110-C, Photon Systems Instruments, Drasov, Czech Republic). Measurements were performed in a 4 mL cuvette (light path of 10 mm). Illumination was provided by a blue LED (455 nm), the measuring light was $0.02 \mu\text{mol m}^{-2}$

² s⁻¹ and saturation pulses had an intensity of 3000 μmol m⁻² s⁻¹. Prior to measurements, all samples were diluted to an appropriate concentration (1 × 10⁶ cells mL⁻¹). After 10 min of dark-adaptation, the samples were exposed to a stepwise increase of seven actinic lights (from 0 to 1000 μmol m⁻² s⁻¹) applied every 60s to construct the electron transport rate versus photon flux density (ETR/PFD) curves. The maximum quantum yield (F_v/F_m) and the effective quantum yield (ΔF/F'_m) were calculated according to Eq. (2) and Eq. (3):

$$F_v/F_m = (F_m - F_0) / F_m \quad (2)$$

$$\Delta F/F'_m = (F'_m - F) / F'_m \quad (3)$$

where F₀ and F_m are the minimum and max fluorescence determined after 10 min dark-adaptation, whereas F and F'_m are the minimum and max fluorescence during illumination.

The relative electron transport rate (rETR) was calculated using the Eq. (4):

$$rETR = \Delta F/F'_m \times PFD \times 0.5 \quad (4)$$

Where PFD is the incident light and 0.5 is a factor assuming that two photons are required for linear electron transfer [32]. Light curves were quantitatively compared using the parameters of maximum rate of relative ETR (rETR_{max}), α and E_k (E_k = rETR_{max}/α) obtained by fitting the rETR/PFD curves with the function rETR = ETR_{max} (1 - e^{-α I} /rETR_{max}) described by [33].

2.6. Determination of cellular compounds

Chl *a* was extracted from cells using dimethyl sulfoxide (DMSO) [34], and quantified by measuring the absorption at 649 nm and 665 nm with an Evolution 60S UV–visible spectrophotometer (Thermo Scientific, Madison, WI, USA). The Chl *a* concentration was calculated by Eq. (5) [34]:

$$\text{Chl } a \text{ (}\mu\text{g mL}^{-1}\text{)} = 12.19 \times \text{OD}_{665} - 3.45 \times \text{OD}_{649} \quad (5)$$

The Chl *a* content was normalized to the average cell bio-volume, which was estimated with an AxioSkop 2 plus microscope (Carl Zeiss, Oberkochen, Germany).

The macromolecular composition of the cells was analyzed by means of an ATR-FTIR PerkinElmer Spectrum-two spectrometer (PerkinElmer, Waltham, MA, USA). Biofilm cells were re-suspended in 1 mL Milli-Q water and washed twice. 1-2 μL of the concentrated sample was deposited on the crystal of the spectrometer, and dried at room temperature for 20 min. Infrared spectra were recorded in the range of 4000 to 400 cm^{-1} using an accumulation of 32 scans at a spectral resolution of 4 cm^{-1} . Before loading algal samples, the empty crystal was measured as background. The spectra were baselined and maximum absorption values in the spectral ranges corresponding to specific macromolecular pool: carbohydrates (C–O–C; 1200–950 cm^{-1}), lipids (C=O; 1750–1700 cm^{-1}), proteins (Amide I; 1700–1630 cm^{-1}) were used to calculate the relative carbohydrates and lipids contents with respect to the proteins signal (Fig. S2) [20,35].

Carbon and nitrogen contents of the biofilm samples were determined with an Elemental Analyzer (Organic Elemental Analyzer FLASH 2000 CHNS/O, Thermo Scientific) using

1~2 mg of dried biomass collected from biofilms that have been previously washed twice with miliQ water and dried at 100°C.

2.7. Statistics analysis

Statistical analysis was performed using IBM SPSS Statistics 25.0 (SPSS Inc., Chicago, IL). Two-way ANOVA was used to test the influence of PFD and initial cell density on the relative biomass increase, photosynthetic performance and cellular compounds of immobilized microalgae. Bonferroni significant difference test for pairwise comparisons testing was performed after the tests of normality and variance homogeneity. $P < 0.05$ indicates a statistically significant difference among tested values. Standard deviations were calculated from at least four independent biological replicates.

3. Results and discussion

3.1. Biomass production as a function of light intensity and initial inoculum density

In biofilms, microalgae cell behavior is strongly impacted by the local conditions (light, nutrients, biofilm properties such as density, thickness, gas exchange, ...) imposed by their new life style (sessile mode) [14,20,21,36]. This may impact biomass and compounds production and should be therefore studied in-depth.

Fig. 2 presents the relative biomass increase for the biofilms grown on membranes. Overall, our results show that when inoculated with lower cell density (LC), the immobilized cultures of *C. vulgaris* presented 3~5 times higher relative biomass increase than HC regardless of the light intensity ($P < 0.05$). On the other hand, light intensity stimulated biomass production, which is in agreement with other works [14,37].

Figure 2. Effect of light intensity and initial cell density on the relative biomass increase of *C. vulgaris* biofilms after 3-days cultivation. All the results were shown as mean value $\pm$ SD ($n \geq 4$). Bars with different letters represent the statistical differences among immobilized cultures under different light conditions at level of $P < 0.05$, while *** depicts the differences between biofilms with two inoculum cell densities at $P < 0.001$.

Indeed, regardless of the initial cell density the immobilized cells under LL showed lower relative biomass increase compared with biofilms exposed to ML and HL ($P < 0.05$). This suggests that the immobilized growth of *C. vulgaris* under $50 \mu\text{mol m}^{-2} \text{s}^{-1}$ may be photo-limited while that at higher light intensity were photo-saturated. This is not surprising as

similar results have been reported by Grenier et al. [28] for *Chlorella autotrophica* biofilms.

It also appears that biomass increase was negatively affected by the initial cell density (Fig. 2). These data can be explained by the fact that high cell density may affect light penetration and/or limit the diffusion of nutrients [14,21,38]. Indeed, studies reported that in immobilized cultures the biomass productivity may depend on the biofilm thickness due to an exponential decrease of light [21,39]. In our study, the two initial cell conditions did not produce biofilms with significantly different thickness (40 - 60 μm) but the HC culture presented a denser initial population as detected by CLSM (Fig. S1) which might have impacted biofilm growth. Finally, cell density dependent feedbacks (quorum-sensing) in cyanobacteria are also known to influence biofilm development as a function of the extracellular concentration of autoinducers (secreted by cells) which in turn affects the expression of biofilm-related gene [40,41]. Therefore, we cannot rule out that similar mechanisms may have occurred in our immobilized cultures. Other studies should be further carried out to test such hypothesis and verify possible synergistic interactions of light and nutrients availability and, quorum sensing mechanisms.

On the whole, these results show that with the proper combination of light intensity and initial inoculum size the biofilm growth performance could be improved. For example, by illuminating the biofilms with $250 \mu\text{mol m}^{-2} \text{s}^{-1}$ at low initial cell density we obtained a 28-fold enhancement of biomass after 3 days of cultivation, whereas in [29], the authors only obtained a 3-fold increase of *Chlorella kessleri* after 4-day cultivation.

281 *3.2. Monitoring the physiological state of cells in order to improve biomass and energy-*
282 *rich compounds production*

283
284 Although photosynthetic performance and macromolecular composition of microalgae
285 have been measured in suspended cultures [42,43], only few studies estimated these
286 parameters for microalgae biofilm-based systems [44,45]. Indeed, estimating the health-
287 state of early-stage biofilms by monitoring their physiological state would be of paramount
288 importance to help the operator identifying stressful parameters in advance and to do
289 reasoned operation choices. For instance, if limiting factors are identified, cultivating
290 parameters or production strategies can be rapidly adjusted (light tuning, medium supply,
291 etc.) to prevent adverse effects in the long-term biofilm cultivation and process productivity.
292 Moreover, monitoring cell physiology also allows a better understanding of acclimation
293 strategies in photosynthetic biofilms that still remains unclear (such as the interplay
294 between photosynthetic activity and pools of macromolecules).

295 In biofilms, light availability and nutrient supply vary as a function of the inoculum size
296 and/or with the biofilm thickness [14,21]. From studies on the phytoplankton, we know that
297 variations of these two factors induce a physiological reorganization spanning from light
298 absorption to electron transport and to the final synthesis of macromolecules to acclimate
299 themselves to the new conditions and maximize growth [46–48]. Interestingly, similar
300 responses occurred in our immobilized cultures. Like their planktonic counterparts,
301 immobilized cells appear to acclimate to a range of light intensities and likely nutrients
302 availability through a series of physiological adjustments in a short term (3 days). This is

typically done to balance the incoming flux of energy and their energetic demand for growth in order to increase their fitness under each specific set of conditions [46,47,49].

Figure 3. Effect of light intensity and initial cell density on Chl *a* content of *C. vulgaris* biofilms after 3-days cultivation. All the results were shown as mean value $\pm$ SD ($n \geq 4$). Bars with different letters represent the statistical differences among immobilized cultures under different light conditions at level of $P < 0.05$ while ** and * respectively depict the differences between biofilms with two inoculum cell densities at $P < 0.01$ and $P < 0.05$, and *ns* represents no difference.

The first level at which microalgae can regulate excitation pressure is the absorption of light [50]. This is typically reached by changing the amount of pigment content in the cells. Fig. 3 shows that Chl *a* content declined with increasing light intensity (from 7 to 2 $\text{fg } \mu\text{m}^{-3}$) ($P < 0.05$), suggesting a lower amount of photons absorbed by the cells as a protective strategy to diminish photons absorption and to avoid photo-damage [51,52]. This is well described for planktonic microalgae cultures but no report of such a mechanism exists for

microalgae biofilms. However, changes in accessory pigments content in natural biofilms have been reported as a photo-protective mechanism [53].

Figure 4. Effect of light intensity and initial cell density on photosynthetic parameters (a, F_v/F_m ; b, $rETR_{\text{max}}$; c, α ; d, E_k) of *C. vulgaris* biofilms after 3-days cultivation. All the results were shown as mean value $\pm$ SD ($n \geq 4$). Bars with different letters represent statistical differences among immobilized cultures under different light conditions at $P < 0.05$ while *ns* depicts no difference between biofilms with two inoculum cell densities.

Another level at which microalgae can regulate excitation pressure is by modulating their photosynthetic capacity in order to meet the incoming flux of photons [48]. In this context, photosynthetic parameters such as F_v/F_m , α , $rETR_{max}$ and E_k are often used to describe the photo-acclimation state of microalgae when environmental conditions change, especially with respect to irradiance and nutrient levels [42,49,54]. As depicted in Fig. 4, the cells exposed to the LL and ML presented a higher α (*c.a* 0.4), lower $rETR_{max}$ and E_k compared to those of HL acclimated cells, indicating that low-irradiance acclimated cells modified their photo-physiology to maximize light harvesting efficiency [55]. According to our expectations, algae photo-acclimated to LL and ML were more efficient in light utilization than those at HL, as shown by the initial slope (α) of the $rETR/PFD$ curves. Conversely, the higher $rETR_{max}$ (180 ± 10) and E_k ($575 \pm 5 \mu\text{mol m}^{-2} \text{s}^{-1}$) typically associating with HL demonstrated that the cells exposed to $500 \mu\text{mol m}^{-2} \text{s}^{-1}$ presented a higher electron transport capacity to face the high flux of incoming photons. A similar acclimation strategy with respect to light intensity has been described for fluvial biofilms [56], suggesting that the cells at $500 \mu\text{mol m}^{-2} \text{s}^{-1}$ were in a high-light acclimation state. The maximum quantum yield (F_v/F_m) is typically used as a stress indicator to evaluate the health state of suspended cultures and in natural photosynthetic biofilms [52,57,58]. After 3 days, the immobilized cells grown at $500 \mu\text{mol m}^{-2} \text{s}^{-1}$ presented only a 0.14 lower F_v/F_m value with respect to the cultures grown at LL ($P < 0.05$, Fig 4a), indicating that the changes in Chl *a* content and in the photosynthetic parameters allowed the cells to optimize their energy harvesting ability and using efficiency to protect from over-excitation and photo-damage [48,55,59].

Figure 5. Effect of light intensity and initial cell density on carbohydrates (a) or lipids (b) to proteins ratio of *C. vulgaris* biofilms after 3-days cultivation. All the results were shown as mean value $\pm$ SD ($n \geq 4$). Bars with different letters represent statistical differences among immobilized cultures under different light conditions at $P < 0.05$ while ***, ** and * respectively depict the differences between biofilms with two inoculum cell densities at $P < 0.001$, $P < 0.01$ and $P < 0.05$, and *ns* represents no difference.

In planktonic cultures of microalgae, the photo-acclimation state of the cells is not only reflected in different pigment contents and photosynthetic efficiency, but often also in changes of their macromolecular and elemental composition [47]. Storage pools such as carbohydrates and lipids serve typically as carbon and energy sinks during unbalanced growth as a results of high excitation pressure and/or nutrient limitation [60,61]. In this study, a significant increase of the carbohydrates to proteins ratio with light intensity was observed (46.4% and 24.4% increases at LC and HC biofilms exposed to $500 \mu\text{mol m}^{-2} \text{s}^{-1}$,

respectively; Fig. 5a, $P < 0.05$). This is consistent with the high-light acclimation state of the cells and in accordance with other works [62,63].

On the other hand, the relative lipids content (Fig. 5b) and the C/N ratio (Fig. S3), two parameters that in microalgae are highly sensitive to N concentration in the environment seemed to be affected by the initial cell density. The lipid to protein ratio increased by 47.3%, 76.3% and 30.4% when biofilms at HC were exposed to LL, ML and HL, respectively, compared to those at LC (Fig. 5b). These results suggest an uncoupling of carbon assimilation from nitrogen uptake which could be associated with nitrogen limitation, in agreement with data reported in suspended cultures [64–66].

On the whole, the results show that the Chl *a* content, photosynthetic activity and the carbohydrates to proteins ratio were highly related to light intensity, while the initial cell density mostly affected the C/N and lipids to proteins ratio. In addition, optimal conditions for biofilm-based cultivation of *C. vulgaris* were identified. Combined conditions of $250 \mu\text{mol m}^{-2} \text{s}^{-1}$ and low inoculum density ($4.8 \times 10^6 \text{ cells cm}^{-2}/0.4 \text{ g m}^{-2}$) promote biomass production by avoiding effects caused by photo-limitation at $50 \mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 2). On the other hand, if lipids are the target, combined conditions of $250 \mu\text{mol m}^{-2} \text{s}^{-1}$ and high inoculum density ($28.8 \times 10^6 \text{ cells cm}^{-2}/2.6 \text{ g m}^{-2}$) should be applied. In conclusion, monitoring photosynthetic activity and cellular composition help better understanding photosynthetic mechanisms in biofilms but also allow to identify optimal strategies for biomass and compounds production.

4. Conclusion

389

390 In this work, biofilm production and the physiological properties (photosynthetic
391 activity and composition) of sessile cells were assessed in response to two culture
392 operational factors, PFD and inoculum cell density. Results showed that light intensity
393 impacts biomass production. Mechanism of photo-limitation of biofilms exposed to LL was
394 highlighted. Acclimation of sessile cells to light and probably to nutrients were confirmed
395 by changes in the photosynthetic activity parameters and microalgae composition (Chl *a*
396 content and relative lipid/carbohydrates pools). Optimal conditions to produce biomass or
397 lipids were determined: 250 $\mu\text{mol m}^{-2} \text{s}^{-1}$ combined with LC or HC initial inoculum,
398 respectively. On the whole, monitoring physiological profiles at the early stage of biofilms
399 development provides information to better understand photosynthetic mechanisms in
400 biofilms and to operate efficiently biofilm-based systems in order to optimize biomass and
401 macromolecules productions.

402

403 **Author contributions**

404 S. F. Li, A. Fanesi, T. Martin and F. Lopes designed the experiments, S. F. Li conducted
405 the experiments, S. F. Li, A. Fanesi and F. Lopes analyzed the data, S. F. Li and A. Fanesi
406 wrote the manuscript, F. Lopes revised the article. All authors read and approved the final
407 manuscript.

408

409 **Acknowledgements**

S. F. Li was supported by the China Scholarship Council (CSC) and A. Fanesi was founded from the LabeX LaSIPS project Greenbelt managed by the French National Research Agency (ANR).

Competing interests

The authors declare no competing interests.

Reference:

- [1] M.J. Fernández-Reiriz, A. Perez-Camacho, M.J. Ferreiro, J. Blanco, M. Planas, M.J. Campos, U. Labarta, Biomass production and variation in the biochemical profile (total protein, carbohydrates, RNA, lipids and fatty acids) of seven species of marine microalgae, *Aquaculture*. 83 (1989) 17–37. [https://doi.org/10.1016/0044-8486\(89\)90057-4](https://doi.org/10.1016/0044-8486(89)90057-4).
- [2] A. Kumar, S. Ergas, X. Yuan, A. Sahu, Q. Zhang, J. Dewulf, F.X. Malcata, H. van Langenhove, Enhanced CO₂ fixation and biofuel production via microalgae: recent developments and future directions, *Trends Biotechnol.* 28 (2010) 371–380. <https://doi.org/10.1016/j.tibtech.2010.04.004>.
- [3] J.S. Tan, S.Y. Lee, K.W. Chew, M.K. Lam, J.W. Lim, S.-H. Ho, P.L. Show, A review on microalgae cultivation and harvesting, and their biomass extraction processing using ionic liquids, *Bioengineered*. 11 (2020) 116–129. <https://doi.org/10.1080/21655979.2020.1711626>.

- [4] T.D.P. Nguyen, D.H. Nguyen, J.W. Lim, C.-K. Chang, H.Y. Leong, T.N.T. Tran, T.B.H. Vu, T.T.C. Nguyen, P.L. Show, Investigation of the Relationship between Bacteria Growth and Lipid Production Cultivating of Microalgae *Chlorella Vulgaris* in Seafood Wastewater, *Energies*. 12 (2019) 2282.
<https://doi.org/10.3390/en12122282>.
- [5] A. Ozkan, K. Kinney, L. Katz, H. Berberoglu, Reduction of water and energy requirement of algae cultivation using an algae biofilm photobioreactor, *Bioresour. Technol.* 114 (2012) 542–548. <https://doi.org/10.1016/j.biortech.2012.03.055>.
- [6] J. Roostaei, Y. Zhang, K. Gopalakrishnan, A.J. Ochocki, Mixotrophic microalgae biofilm: a novel algae cultivation strategy for improved productivity and cost-efficiency of biofuel feedstock production, *Sci. Rep.* 8 (2018) 12528.
<https://doi.org/10.1038/s41598-018-31016-1>.
- [7] L. Christenson, R. Sims, Production and harvesting of microalgae for wastewater treatment, biofuels, and bioproducts, *Biotechnol. Adv.* 29 (2011) 686–702.
<https://doi.org/10.1016/j.biotechadv.2011.05.015>.
- [8] H. Yuan, Y. Wang, Y. Xi, Z. Jiang, X. Zhang, X. Wang, X. Zhang, Light-emitting diode power conversion capability and CO₂ fixation rate of microalgae biofilm cultured under different light spectra, *Energies*. 13 (2020) 1536.
<https://doi.org/10.3390/en13071536>.
- [9] A.-A. Mohd-Sahib, J.-W. Lim, M.-K. Lam, Y. Uemura, M.H. Isa, C.-D. Ho, S.R.M. Kutty, C.-Y. Wong, S.-S. Rosli, Lipid for biodiesel production from attached growth *Chlorella vulgaris* biomass cultivating in fluidized bed bioreactor packed with

polyurethane foam material, Bioresour. Technol. 239 (2017) 127–136.

<https://doi.org/10.1016/j.biortech.2017.04.118>.

- [10] D.A. Carbone, G. Olivieri, A. Pollio, Gabriele, M. Melkonian, Growth and biomass productivity of *Scenedesmus vacuolatus* on a twin layer system and a comparison with other types of cultivations, Appl. Microbiol. Biotechnol. 101 (2017) 8321–8329. <https://doi.org/10.1007/s00253-017-8515-y>.

- [11] T.E. Murphy, H. Berberoglu, Flux balancing of light and nutrients in a biofilm photobioreactor for maximizing photosynthetic productivity, Biotechnol. Prog. 30 (2014) 348–359. <https://doi.org/10.1002/btpr.1881>.

- [12] B. Podola, T. Li, M. Melkonian, Porous substrate bioreactors: a paradigm shift in microalgal biotechnology?, Trends Biotechnol. 35 (2017) 121–132. <https://doi.org/10.1016/j.tibtech.2016.06.004>.

- [13] M.E. Venable, M.R. Podbielski, Impact of substrate material on algal biofilm biomass growth, Environ. Sci. Pollut. Res. 26 (2019) 7256–7262. <https://doi.org/10.1007/s11356-019-04148-8>.

- [14] Y. Huang, W. Xiong, Q. Liao, Q. Fu, A. Xia, X. Zhu, Y. Sun, Comparison of *Chlorella vulgaris* biomass productivity cultivated in biofilm and suspension from the aspect of light transmission and microalgae affinity to carbon dioxide, Bioresour. Technol. 222 (2016) 367–373. <https://doi.org/10.1016/j.biortech.2016.09.099>.

- [15] J. Wang, J. Liu, T. Liu, The difference in effective light penetration may explain the superiority in photosynthetic efficiency of attached cultivation over the conventional open pond for microalgae, Biotechnol. Biofuels. 8 (2015) 49. <https://doi.org/10.1186/s13068-015-0240-0>.

- 477 [16] Y. Meng, A. Li, H. Li, Z. Shen, T. Ma, J. Liu, Z. Zhou, Q. Feng, Y. Sun, Effect of
478 membrane blocking on attached cultivation of microalgae, *J. Clean. Prod.* 284 (2021)
479 124695. <https://doi.org/10.1016/j.jclepro.2020.124695>.
- 480 [17] T.E. Irving, D.G. Allen, Species and material considerations in the formation and
481 development of microalgal biofilms, *Appl. Microbiol. Biotechnol.* 92 (2011) 283–294.
482 <https://doi.org/10.1007/s00253-011-3341-0>.
- 483 [18] P.J. Schnurr, D.G. Allen, Factors affecting algae biofilm growth and lipid production:
484 A review, *Renew. Sustain. Energy Rev.* 52 (2015) 418–429.
485 <https://doi.org/10.1016/j.rser.2015.07.090>.
- 486 [19] A. Fanesi, M. Lavayssière, C. Breton, O. Bernard, R. Briandet, F. Lopes, Shear stress
487 affects the architecture and cohesion of *Chlorella vulgaris* biofilms, *Sci. Rep.* 11
488 (2021) 4002. <https://doi.org/10.1038/s41598-021-83523-3>.
- 489 [20] A. Fanesi, A. Paule, O. Bernard, R. Briandet, F. Lopes, The architecture of
490 monospecific microalgae biofilms, *Microorganisms.* 7 (2019) 352.
491 <https://doi.org/10.3390/microorganisms7090352>.
- 492 [21] P.J. Schnurr, G.S. Espie, D.G. Allen, The effect of light direction and suspended cell
493 concentrations on algal biofilm growth rates, *Appl. Microbiol. Biotechnol.* 98 (2014)
494 8553–8562. <https://doi.org/10.1007/s00253-014-5964-4>.
- 495 [22] S.N. Genin, J. Stewart Aitchison, D. Grant Allen, Design of algal film
496 photobioreactors: Material surface energy effects on algal film productivity,
497 colonization and lipid content, *Bioresour. Technol.* 155 (2014) 136–143.
498 <https://doi.org/10.1016/j.biortech.2013.12.060>.

- [23] Y. Huang, Y. Zheng, J. Li, Q. Liao, Q. Fu, A. Xia, J. Fu, Y. Sun, Enhancing microalgae biofilm formation and growth by fabricating microgrooves onto the substrate surface, *Bioresour. Technol.* 261 (2018) 36–43.
<https://doi.org/10.1016/j.biortech.2018.03.139>.
- [24] A. Ozkan, H. Berberoglu, Adhesion of *Chlorella vulgaris* on hydrophilic and hydrophobic surfaces, in: Vol. 4 Energy Syst. Anal. Thermodyn. Sustain. Combust. Sci. Eng. Nanoeng. Energy Parts B, ASMEDC, Denver, Colorado, USA, 2011: pp. 169–178. <https://doi.org/10.1115/IMECE2011-64133>.
- [25] S.U. Gerbersdorf, S. Wieprecht, Biostabilization of cohesive sediments: revisiting the role of abiotic conditions, physiology and diversity of microbes, polymeric secretion, and biofilm architecture, *Geobiology*. 13 (2015) 68–97.
<https://doi.org/10.1111/gbi.12115>.
- [26] A.-A. Mohd-Sahib, J.-W. Lim, M.-K. Lam, Y. Uemura, C.-D. Ho, W.-D. Oh, W.-N. Tan, Mechanistic kinetic models describing impact of early attachment between *Chlorella vulgaris* and polyurethane foam material in fluidized bed bioreactor on lipid for biodiesel production, *Algal Res.* 33 (2018) 209–217.
<https://doi.org/10.1016/j.algal.2018.05.017>.
- [27] B. Ji, W. Zhang, N. Zhang, J. Wang, G.A. Lutz, T. Liu, Biofilm cultivation of the oleaginous microalgae *Pseudochlorococcum* sp., *Bioprocess Biosyst. Eng.* 37 (2014) 1369–1375. <https://doi.org/10.1007/s00449-013-1109-x>.
- [28] J. Grenier, H. Bonnefond, F. Lopes, O. Bernard, The impact of light supply to moving photosynthetic biofilms, *Algal Res.* 44 (2019) 101674.
<https://doi.org/10.1016/j.algal.2019.101674>.

- [29] T. Shiratake, A. Sato, A. Minoda, M. Tsuzuki, N. Sato, Air-drying of cells, the novel conditions for stimulated synthesis of triacylglycerol in a green alga, *Chlorella kessleri*, PLoS ONE. 8 (2013) e79630. <https://doi.org/10.1371/journal.pone.0079630>.
- [30] L.-L. Zhuang, The characteristics and influencing factors of the attached microalgae cultivation: A review, Renew. Sustain. Energy Rev. 94 (2018) 1110–1119. <https://doi.org/10.1016/j.rser.2018.06.006>.
- [31] H.W. Bischo, H.C. Bold, Phycological studies IV. some soil algae from enchanted rock and related algal species, University of Texas Publication: Austin, TX, USA, 1963.
- [32] A. Guzzon, F.D. Pippo, R. Congestri, Wastewater biofilm photosynthesis in photobioreactors, Microorganisms. 7 (2019) 252. <https://doi.org/10.3390/microorganisms7080252>.
- [33] W.L. Webb, M. Newton, D. Starr, Carbon dioxide exchange of *Alnus rubra*, Oecologia. 17 (1974) 281–291. <https://doi.org/10.1007/BF00345747>.
- [34] A.R. Wellburn, The spectral determination of chlorophylls *a* and *b*, as well as total carotenoids, using various solvents with spectrophotometers of different resolution, J. Plant Physiol. 144 (1994) 307–313. [https://doi.org/10.1016/S0176-1617\(11\)81192-2](https://doi.org/10.1016/S0176-1617(11)81192-2).
- [35] D.Y. Duygu, A.U. Udoh, T.B. Ozer, A. Akbulut, I.A. Erkaya, K. Yildiz, D. Guler, Fourier transform infrared (FTIR) spectroscopy for identification of *Chlorella vulgaris* Beijerinck 1890 and *Scenedesmus obliquus* (Turpin) Kützing 1833, Afr. J. Biotechnol. 11 (2012) 3817–3824.
- [36] H. Yuan, Y. Wang, Z. Lai, X. Zhang, Z. Jiang, X. Zhang, Analyzing microalgal biofilm structures formed under different light conditions by evaluating cell–cell

interactions, J. Colloid Interface Sci. 583 (2021) 563–570.

<https://doi.org/10.1016/j.jcis.2020.09.057>.

- [37] S. Kim, M. Moon, M. Kwak, B. Lee, Y.K. Chang, Statistical optimization of light intensity and CO₂ concentration for lipid production derived from attached cultivation of green microalga *Ettlia sp.*, Sci. Rep. 8 (2018) 15390.

<https://doi.org/10.1038/s41598-018-33793-1>.

- [38] C. Picioreanu, Effect of diffusive and convective substrate transport on biofilm structure formation: A two- dimensional modeling study, Biotechnol. Bioeng. 69 (2000) 504–515. [https://doi.org/10.1002/1097-0290\(20000905\)69:5<504::AID-BIT5>3.0.CO;2-S](https://doi.org/10.1002/1097-0290(20000905)69:5<504::AID-BIT5>3.0.CO;2-S).

- [39] C. Barranguet, S. van Beusekom, B. Veuger, T. Neu, E. Manders, J. Sinke, W. Admiraal, Studying undisturbed autotrophic biofilms: still a technical challenge, Aquat. Microb. Ecol. 34 (2004) 1–9. <https://doi.org/10.3354/ame034001>.

- [40] E. Nagar, R. Schwarz, To be or not to be planktonic? Self-inhibition of biofilm development: Self-inhibition of biofilm development, Environ. Microbiol. 17 (2015) 1477–1486. <https://doi.org/10.1111/1462-2920.12583>.

- [41] D. Schatz, E. Nagar, E. Sendersky, R. Parnasa, S. Zilberman, S. Carmeli, Y. Mastai, E. Shimoni, E. Klein, O. Yeager, Z. Reich, R. Schwarz, Self-suppression of biofilm formation in the cyanobacterium *Synechococcus elongatus*: Self-suppression of biofilm formation in a cyanobacterium, Environ. Microbiol. 15 (2013) 1786–1794. <https://doi.org/10.1111/1462-2920.12070>.

- [42] E. Cointet, G. Wielgosz-Collin, G. Bougaran, V. Rabesaotra, O. Gonçalves, V. Méléder, Effects of light and nitrogen availability on photosynthetic efficiency and

568 fatty acid content of three original benthic diatom strains, PLOS ONE. 14 (2019)
569 e0224701. <https://doi.org/10.1371/journal.pone.0224701>.

570 [43] Y. Jiang, T. Yoshida, A. Quigg, Photosynthetic performance, lipid production and
571 biomass composition in response to nitrogen limitation in marine microalgae, Plant
572 Physiol. Biochem. 54 (2012) 70–77. <https://doi.org/10.1016/j.plaphy.2012.02.012>.

573 [44] H. Yuan, X. Zhang, Z. Jiang, X. Wang, Y. Wang, L. Cao, X. Zhang, Effect of light
574 spectra on microalgal biofilm: Cell growth, photosynthetic property, and main organic
575 composition, Renew. Energy. 157 (2020) 83–89.
576 <https://doi.org/10.1016/j.renene.2020.04.109>.

577 [45] X. Zhang, H. Yuan, L. Guan, X. Wang, Y. Wang, Z. Jiang, L. Cao, X. Zhang,
578 Influence of photoperiods on microalgae biofilm: photosynthetic performance,
579 biomass yield, and cellular composition, Energies. 12 (2019) 3724.
580 <https://doi.org/10.3390/en12193724>.

581 [46] M.J. Behrenfeld, K.H. Halsey, A.J. Milligan, Evolved physiological responses of
582 phytoplankton to their integrated growth environment, Philos. Trans. R. Soc. B Biol.
583 Sci. 363 (2008) 2687–2703. <https://doi.org/10.1098/rstb.2008.0019>.

584 [47] K.H. Halsey, B.M. Jones, Phytoplankton strategies for photosynthetic energy
585 allocation, Annu. Rev. Mar. Sci. 7 (2015) 265–297. [https://doi.org/10.1146/annurev-](https://doi.org/10.1146/annurev-marine-010814-015813)
586 [marine-010814-015813](https://doi.org/10.1146/annurev-marine-010814-015813).

587 [48] H.L. MacIntyre, T.M. Kana, R.J. Geider, The effect of water motion on short-term
588 rates of photosynthesis by marine phytoplankton, Trends Plant Sci. 5 (2000) 12–17.
589 [https://doi.org/10.1016/S1360-1385\(99\)01504-6](https://doi.org/10.1016/S1360-1385(99)01504-6).

- [49] A. McMinn, E.N. Hegseth, Quantum yield and photosynthetic parameters of marine microalgae from the southern Arctic Ocean, Svalbard, J. Mar. Biol. Assoc. U. K. 84 (2004) 865. <https://doi.org/10.1017/S0025315404010112h>.
- [50] A. Fanesi, H. Wagner, A. Becker, C. Wilhelm, Temperature affects the partitioning of absorbed light energy in freshwater phytoplankton, Freshw. Biol. 61 (2016) 1365–1378. <https://doi.org/10.1111/fwb.12777>.
- [51] X. Chen, Q.Y. Goh, W. Tan, I. Hossain, W.N. Chen, R. Lau, Lumostatic strategy for microalgae cultivation utilizing image analysis and chlorophyll *a* content as design parameters, Bioresour. Technol. 102 (2011) 6005–6012. <https://doi.org/10.1016/j.biortech.2011.02.061>.
- [52] Q. He, H. Yang, L. Wu, C. Hu, Effect of light intensity on physiological changes, carbon allocation and neutral lipid accumulation in oleaginous microalgae, Bioresour. Technol. 191 (2015) 219–228. <https://doi.org/10.1016/j.biortech.2015.05.021>.
- [53] X. Timoner, T. Buchaca, A. Vicenc, S. Sergi, Photosynthetic pigment changes and adaptations in biofilms in response to flow intermittency, Aquat. Sci. 76 (2014) 565–578.
- [54] P.J. Ralph, R. Gademann, Rapid light curves: A powerful tool to assess photosynthetic activity, Aquat. Bot. 82 (2005) 222–237. <https://doi.org/10.1016/j.aquabot.2005.02.006>.
- [55] S.M. Rincon, N.F. Urrego, K.J. Avila, H.M. Romero, H. Beyenal, Photosynthetic activity assessment in mixotrophically cultured *Chlorella vulgaris* biofilms at various developmental stages, Algal Res. 38 (2019) 101408. <https://doi.org/10.1016/j.algal.2019.101408>.

- [56] N. Corcoll, B. Bonet, M. Leira, B. Montuelle, A. Tlili, H. Guasch, Light history influences the response of fluvial biofilms to Zn exposure, *J. Phycol.* 48 (2012) 1411–1423. <https://doi.org/10.1111/j.1529-8817.2012.01223.x>.
- [57] N. Häubner, R. Schumann, U. Karsten, Aeroterrestrial microalgae growing in biofilms on facades—response to temperature and water stress, *Microb. Ecol.* 51 (2006) 285–293. <https://doi.org/10.1007/s00248-006-9016-1>.
- [58] J. Malapascua, C. Jerez, M. Sergejevová, F. Figueroa, J. Masojídek, Photosynthesis monitoring to optimize growth of microalgal mass cultures: application of chlorophyll fluorescence techniques, *Aquat. Biol.* 22 (2014) 123–140. <https://doi.org/10.3354/ab00597>.
- [59] C.G. Jerez, J.R. Malapascua, M. Sergejevová, J. Masojídek, F.L. Figueroa, *Chlorella fusca* (Chlorophyta) grown in thin-layer cascades: Estimation of biomass productivity by in-vivo chlorophyll *a* fluorescence monitoring, *Algal Res.* 17 (2016) 21–30. <https://doi.org/10.1016/j.algal.2016.04.010>.
- [60] A. Fanesi, H. Wagner, C. Wilhelm, Phytoplankton growth rate modelling: can spectroscopic cell chemotyping be superior to physiological predictors?, *Proc. R. Soc. B Biol. Sci.* 284 (2017) 20161956. <https://doi.org/10.1098/rspb.2016.1956>.
- [61] H. Wagner, T. Jakob, A. Fanesi, C. Wilhelm, Towards an understanding of the molecular regulation of carbon allocation in diatoms: the interaction of energy and carbon allocation, *Philos. Trans. R. Soc. B Biol. Sci.* 372 (2017) 20160410. <https://doi.org/10.1098/rstb.2016.0410>.
- [62] S.-H. Ho, C.-Y. Chen, J.-S. Chang, Effect of light intensity and nitrogen starvation on CO₂ fixation and lipid/carbohydrate production of an indigenous microalga

Scenedesmus obliquus CNW-N, Bioresour. Technol. 113 (2012) 244–252.

<https://doi.org/10.1016/j.biortech.2011.11.133>.

[63] C. Jebsen, A. Norici, H. Wagner, M. Palmucci, M. Giordano, C. Wilhelm, FTIR spectra of algal species can be used as physiological fingerprints to assess their actual growth potential, *Physiol. Plant.* 146 (2012) 427–438. <https://doi.org/10.1111/j.1399-3054.2012.01636.x>.

[64] B. Clément-Larosière, F. Lopes, A. Gonçalves, B. Taidi, M. Benedetti, M. Minier, D. Pareau, Carbon dioxide biofixation by *Chlorella vulgaris* at different CO₂ concentrations and light intensities, *Eng. Life Sci.* 14 (2014) 509–519. <https://doi.org/10.1002/elsc.201200212>.

[65] M.J. Griffiths, R.P. van Hille, S.T.L. Harrison, The effect of nitrogen limitation on lipid productivity and cell composition in *Chlorella vulgaris*, *Appl. Microbiol. Biotechnol.* 98 (2014) 2345–2356. <https://doi.org/10.1007/s00253-013-5442-4>.

[66] H. Wagner, C. Jebsen, C. Wilhelm, Monitoring cellular C:N ratio in phytoplankton by means of FTIR - spectroscopy, *J. Phycol.* 55 (2019) 543–551. <https://doi.org/10.1111/jpy.12858>.