The role of sulfur in molybdenum transport in hydrothermal fluids: insight from *in situ* synchrotron XAS experiments and molecular dynamics simulations

¹*Weihua Liu, ²Barbara Etschmann, ¹Yuan Mei, ²Qiushi Guan, ³Denis Testemale, and ²Joël Brugger

1. CSIRO Mineral Resources, Clayton, Vic, 3168, Australia

2. School of Earth, Atmosphere and Environment, Monash University, Clayton, Vic, 3800, Australia

3. Université Grenoble Alpes, CNRS, Grenoble INP, Institut Néel, 38000 Grenoble, France

*Corresponding author

Abstract

Knowledge of the identity and thermodynamic properties of aqueous molybdenum species at elevated temperature is important for understanding the extraction of Mo from source rocks and the formation of hydrothermal deposits. Chlorine and sulfur are two major components in crustal fluids. Several studies have investigated Mo speciation and solubility in chloride-bearing hydrothermal fluids, but little is known about the behaviour of Mo in sulfur-bearing systems at elevated temperature. We used *in-situ* synchrotron X-ray absorption spectroscopy (XAS) measurements and molecular dynamics (MD) simulations to investigate the predominant Mo species in S-bearing (0.04-2 m NaHS) solutions at 30-394 °C (XAS) and 77-300 °C (MD) and 800bar. The XAS data show that, similar to previous room temperature studies, the S²⁻ ion progressively replaces O²⁻ in molybdate (MoO₄²⁻) to form thiomolybdates (MoO_{4-x}S_x²⁻, x = 1, 2, 3, 4) with increasing NaHS concentrations at temperature up to 394 °C. The MD simulations confirm the identity and structure of the thiomolybdate species

determined from the XAS experimental data. The speciation calculations based on the formation constants for the thiomolybdate species extrapolated from the room temperature data agree well with the experimental results. This study indicates that thiomolybdate species are likely to be important in transporting Mo in sulfur-bearing, weak-acidic to alkaline, hydrothermal fluids under reduced (sulfide stable) conditions in the Earth crust. Current models of Mo transport in hydrothermal ore fluids need to be re-evaluated to take into account the thiomolybdate species.

1. Introduction

Molybdenum (Mo) is mainly mined from magmatic hydrothermal deposits (e.g., porphyry Cu-Mo-Au deposits), where the metals including Mo are transported in high temperature-pressure geological fluids and deposited to form ores under favourable physiochemical conditions (e.g., [Seedorff et al., 2005](#)). Molybdenum is also enriched in other hydrothermal systems, including IOCG deposits (e.g., Olympic Dam, South Australia, [Ehrig et al. 2012](#); Uranium roll-front deposits, [Bonnetti et al. 2020](#); [Bullock and Parnell 2017](#)). Furthermore, elevated molybdenum concentrations are also found in sedimentary rocks such as shales, and Mo is commonly used as a paleoredox indicator (e.g., [Asael et al. 2018](#); [Tessin et al. 2018](#)). Therefore, the identity and thermodynamic properties of the aqueous Mo species prevalent in crustal fluids are important for understanding the formation of hydrothermal Mo deposits and constrain the use of molybdenum as a geological tracer.

A number of experimental and theoretical studies have investigated Mo speciation and solubility in hydrothermal fluids, and the partitioning of Mo between melt, vapour and aqueous phases (e.g., [Candela and Holland 1984](#), [Dadze et al. 2017, 2018](#); [Keppler and Wyllie, 1991](#); [Kudrin, 1989](#), [Rempel, et al., 2006, 2008, 2009](#); [Ulrich and Mavrogenes, 2008](#), [Zajacz et al., 2017](#)). These studies suggest that Mo(VI) species predominate under a wide range of redox conditions from oxidised to $\text{MoO}_3(\text{s})/\text{MoO}_2(\text{s})$ stable conditions. In pure water, the hydrogen molybdate (HMoO_4^-) and molybdate (MoO_4^{2-}) ions are thought to be dominant, and their ionization constants were determined up to 300 °C by [Minubayeva and Seward \(2010\)](#); molybdic acid, $\text{H}_2\text{MoO}_4(\text{aq})$, predominates only under highly acidic conditions, especially at $T > 100$ °C ([Borg et al. 2012](#)). Recently, [Dadze et al. \(2017\)](#) obtained the solubility constants for crystalline $\text{MoO}_3(\text{s})$ in acidic solutions up to 350 °C as $\text{H}_2\text{MoO}_4(\text{aq})$ and HMoO_4^- , and [Saha et al. \(2017\)](#) used a hydrothermal diamond anvil cell to

measure MoO₂(s) solubilities in pure water of 44 ppm and 658 ppm at 600 and 800 °C, respectively; however, they were not able to determine the oxidation state and predominant Mo species in the solutions. In pure, low density water vapour, a neutral MoO₃(H₂O)_n(aq) with hydration number n up to 3 at 360 °C was proposed based on the solubility of MoO₃(s) (Rempel et al. , 2006).

The structures of the molybdate and hydrogen molybdate ions show little change with increasing temperature, as evidenced by the synchrotron X-ray absorption study (up to 385 °C) of Borg et al. (2012); both species share a tetrahedral coordination, and confirmed by the *first principle* molecular simulations study of Liu et al. (2013). Liu et al. (2013) also predicts the change of the structure of molybdic acid from 6-coordinated MoO₂(OH)₂(H₂O)₂(aq) to 4-coordinated, unhydrated MoO₂OH₂(aq) with increasing temperatures to 300 °C. They calculated the acidity constants of molybdic acid up to 300 °C with reasonably good agreement with the experimental results by Minubayeva and Seward (2010), in particular for pKa1.

The possibility of Mo(VI) complexation with chloride, the major ligand in many hydrothermal fluids, has also been investigated. Wood et al. (1987) noted that Mo did not complex with chloride in their slightly acidic (pH>3) chloride-bearing experimental solutions (0-5 m NaCl, 25-350 °C), and Kurin (1989) proposed that K₂HMoO₄(aq) and NaHMoO₄(aq) are responsible for the Mo solubility in 0.05-1 m KCl and NaCl solutions (with 0.1 m HCl to 0.005 m KOH) at 300-394 °C. Under much more acidic conditions, Borg et al. (2012) identified oxo-chloro complexes MoO_mCl_n^{6-2m-n} (n up to 5) at the experimental P-T conditions (up to 385°C at 600 bar), consistent with the solubility study by Dadze et al. (2018), in which they found MoO₂(OH)₂Cl⁻ predominant in HCl-HClO₄-NaCl solutions at 300 °C. In HCl-bearing vapour phase, hydrated species MoO₃(H₂O)_y(g) predominate in less acidic vapour (log_f_{HCl(g)} < 0.1, Hurtig and Williams-Jones, 2014) and MoO₂Cl₂(g) is dominant in more acidic conditions (Rempel et al., 2008).

Sulfur is one of the major components in magmatic hydrothermal fluids and volcanic gases, and molybdenite, MoS₂, is the main form of Mo in ore deposits. Recent experimental and theoretical studies have shown that sulfur complexes with ore metals, and can be an important transport agent for Au, Cu, Zn and Pd (Pokrovski et al., 2009, Etschmann et al., 2010, 2019, Mei, 2013, 2015 2016; Liu et al., 2014; Brugger et al. 2016). Therefore, it is desirable to investigate the complexation between Mo and S and its contribution to Mo

transport in hydrothermal fluids. Most previous studies of the hydrothermal transport of Mo focused on Mo speciation in pure water and chloride-bearing systems, with much less work on the behaviour of Mo in hydrothermal sulfur-bearing fluids. Wood et al. (1987) didn't find evidence of Mo-S species in their multi-metal solubility experiments in 0-5 molal NaCl solution with pyrite-pyrrhotite-magnetite buffer for log fO_2 and log fS_2 at 200-350 °C and 0.69-172 bar, whereas Zhang et al. (2012) suggests $NaHMoO_2S_2$ predominant in their experimental solutions at 600-800 °C and 2000 bar, but find the solubility of MoS_2 increases with decreasing log fS_2 .

Under the ambient condition, Mo(VI)-S species or thiomolybdates have been suggested to be important in sediment and surface water environments (e.g., Helz et al., 1996, Erickson and Helz, 2000). Thiomolybdate anions ($MoO_{4-x}S_x^{2-}$, $x = 1, 2, 3, 4$) have been studied extensively with different methods (see reviews by Muller et al., 1981; Coucouvanis, 1998; Laurie, 2000; and Smedley and Kinniburgh, 2017). The kinetics and reaction constants of progressive replacement of oxygen by sulfur was established using mainly UV-Vis spectrophotometry (e.g., Harmer and Sykes, 1980; Erickson and Helz, 2000) and theoretical studies (e.g., Bernholc and Stiefel, 1985, El-Issa et al., 1989, Tossell, 2005). These studies generally agree that thiomolybdate species are important in surface waters under reduced environments. In contrast to detailed information available at room temperature, the identity and stability of the thiomolybdates in hydrothermal fluids at elevated temperature and pressure remain unknown. Furthermore, the thiomolybdate anion notably tends to form stable complexes with some transition metal ions such as Cu(I) (Helz et al., 2011) and Fe(II) (Helz et al., 1996, 2014) at room temperature, indicating the potential for co-transport of Fe and Cu with Mo in reduced, sulfur-rich fluids.

This work aimed to determine the speciation of Mo(VI) in hydrosulfide solutions at elevated temperature and pressure (up to 394 °C and 800 bar). We used *in-situ* Extended X-ray Absorption Fine Structure (EXAFS) spectroscopy and X-ray Absorption Near-edge Structure (XANES) data to obtain structural information for the predominant Mo species, combined with *ab initio* molecular dynamics simulations and *ab initio* XANES calculations to provide new insights for Mo transport and deposition in natural and engineered hydrothermal fluids relevant to the hydrothermal (re)mobilisation of Mo and formation of Mo deposits, as well as the hydrometallurgical processing of Mo-bearing ores.

2. Methods

2.1. XAS experiment

2.1.1. Sample solutions

All sample solutions were prepared from deionized water, $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}(\text{s})$ (analytical grade, Sigma AldrichTM) and a stock of 2 m NaHS solution. The NaHS stock solution was prepared by dissolving sodium hydroxide in deoxygenated water, and then bubbling $\text{H}_2\text{S}(\text{g})$ through the solution. This method was used by [Helz et al. \(1993\)](#), [Mosselmans et al. \(1999\)](#), [Etschmann et al. \(2010\)](#) and [Mei et al. \(2016\)](#) for studies on Cu(I) and Zn(II) bisulfide complexes, and it was demonstrated that the oxidation state of the sample solutions (loaded in glassy carbon cells) were within the stability field of the reduced sulfur over a large temperature range (up to 592 °C) ([Etschmann et al., 2010](#)). The solution compositions are compiled in [Table 1](#).

2.1.2. Sample cell and XAS measurement

Molybdenum K-edge (20,000 eV) X-ray Absorption Near Edge Structure (XANES) and Extended X-ray absorption Fine Structure (EXAFS) spectra were measured at beamline 30-BM (FAME) at the European Synchrotron Radiation Facility (ESRF) in Grenoble, France. The setup at the FAME beam line has been described by Proux et al. (2005). A focused beam size of FWHM $300 \times 150 \text{ } \mu\text{m}^2$ was used. The incident and transmitted beam intensity I_0 and I_t were measured with Si diodes; and fluorescence data were collected concurrently using a 30 element solid state fluorescence detector. The beam energy was calibrated with a Mo foil, such that the maximum of the first derivative of the XANES spectrum was at 20,000 eV.

The high temperature-high pressure cell developed by the “Laboratoire de Cristallographie” (CNRS) was used for XAS measurements of solutions up to supercritical conditions ([Testemale et al., 2005](#)). The cell consists of an external water-cooled high-pressure vessel equipped with three 1.5 mm thick beryllium windows enabling collection of fluorescence and transmission signals at a pressure of 800 bar. A sample solution (mass $\sim 0.1 \text{ g}$) was contained inside a glassy carbon tube with an internal diameter of 4 mm. The pressure was applied to the sample by two glassy carbon pistons, using helium as a pressure medium. The glassy carbon tube was placed inside a small cylindrical resistive heater; the heater and tube are installed inside in a water-cooled stainless-steel high-pressure vessel. The temperature of the fluid at the beam location was controlled by a Eurotherm PID controller to

a stability within 0.1 °C, and calibrated by correlating the density of a 2 molal NaCl solution at the beam position as a function of temperature, based on measurements of X-ray transmission (e.g., [Tooth et al., 2013](#)):

$$\mu_{20\text{ keV}}^{\text{AGS}} - \mu_{20\text{ keV}}^{\text{AGS}} = \left(\frac{\mu}{\rho} \right)_{20\text{ keV}}^S \mu^S x$$

(1),

where $\mu_{20\text{ keV}}^{\text{AGS}}$ is the measured transmission of the autoclave + glassy carbon cell + 800 bar He + solution at 20 keV, $\mu_{20\text{ keV}}^{\text{AGS}}$ is the measured baseline transmission without solution, $\left(\frac{\mu}{\rho} \right)_{20\text{ keV}}^S$ is the tabulated X-ray mass attenuation coefficient corresponding to the composition of solution S at 20 keV (Chantler, 1995), ρ^S is the density of the solution (program Sowat, [Driesner, 2007](#); [Driesner and Heinrich, 2007](#)), and x is the path length of the beam in the solution. The error from the calibration curve is $\sim \pm 3$ °C. The details of the autoclave can be found in several previous studies (e.g., [Testemale et al., 2005](#); [2009](#); [Liu et al., 2011](#); [Tooth et al., 2013](#)).

XANES and EXAFS data were analysed with the HORAE package ([Ravel and Newville, 2005](#)), using FEFF version 9 ([Rehr et al., 2009](#)). The reported errors are at the 1 ° level and taken from the HORAE package.

2.2. *Ab initio* molecular dynamics simulations

The simulations of molybdic acid, molybdates and thiomolybdates were performed with solutions containing 1 H₂MoO₄, 4 Na⁺, 4 HS⁻ and 111 H₂O to represent Na and S concentrations of 2 molal. The nominal Mo concentration is 0.5 molal, but because there is only one Mo atom in the simulation box, there is no Mo-Mo interaction, i.e. Mo behaves as if at infinite dilution in the simulated solutions. Periodic boundary conditions were applied in the *ab initio* MD simulations. We performed simulations at 77 °C, 200 °C and 300 °C at 800 bar to compare with the XAS experiments, and at near-saturation pressure at 200 °C (20 bar) and 300°C (100 bar). The simulations were run in the NVT ensemble with the temperature controlled by the Nosé thermostat for both the ions and electrons ([Nosé, 1984](#), [Hoover, 1985](#)). For these constant volumes calculations, the densities were taken from the equations of state of a 2 molal NaCl solution ([Driesner, 2007](#); [Driesner and Heinrich, 2007](#)),

an approximation that has been widely used in hydrothermal aqueous geochemical systems. Details of the physical conditions for each simulation are listed in Table 2. To test the stability of molybdic acid and thiomolybdates, simulations with six different initial species, i.e., H_2MoO_4 , MoO_4^{2-} , $\text{MoO}_3\text{S}^{2-}$, $\text{MoO}_2\text{S}_2^{2-}$, MoOS_3^{2-} , MoS_4^{2-} , were conducted at each T-P condition, for a total of 25 simulations

Ab initio molecular dynamics simulations were performed using the CP2K/QUICKSTEP package, which implements density functional theory (DFT) using a mixed Gaussian basis sets and plane-waves approach (Hutter et al., 2014). Goedecker-Teter-Hutter (GTH) pseudopotentials (Goedecker et al., 1996; Hartwigsen et al., 1998) were used for the core electrons with a cutoff of 280 Ry (3809.6 eV) for the electron density, and the double-zeta valence polarized (DZVP) basis sets were utilised for Mo, Na, S, O and H (VandeVondele and Hutter 2007) together with BLYP exchange-correlation functional (Lee et al., 1988; Becke, 1988). The BLYP functional generally provides a good description of the structural properties of water including O–O interactions, H-bond statistics, angular distributions, coordination numbers, and radial distribution functions when compared with experiments (Gillan et al., 2016; Lin et al., 2012). However, as discussed in our previous studies using the BLYP exchange correlation functional (Mei et al., 2018; Guan et al., 2020), we note that there is a well-known tendency for most functionals, including BLYP, to over-structure liquid water (Lee and Tuckerman, 2006; Lin et al., 2012). For example, Bankura et al. (2014) showed that BLYP, like most other functionals, best reproduces the hydrogen-bonded network of water at ambient conditions when using an elevated temperature (80 °C in their case), and it could be argued that there is a temperature offset of approximately 50-55 °C between the computed data as compared to the experimental structure of the solvent at room temperature. Therefore, here we performed simulations at 77 °C (350K) to reproduce the solution at room temperature. A simulation timestep of 0.5 femtosecond (fs, 10^{-15} second) was chosen for all simulations. To achieve good statistics, all the simulations were calculated for 9-15 ps including a 2 ps equilibration. The radial distribution functions (RDF) and the coordination number (integrals of RDF) were calculated using VMD (Humphrey et al., 1996) to extract the averaged structural information and to compare with the XAS data. As discussed previously (e.g., Mei et al., 2020), *ab initio* MD simulations are computationally intensive. *Ab initio* MD studies published by researchers from various groups (Jahn and Schmidt, 2010; Sulpizi et al., 2013; Liu et al., 2013a, b; Wagner et al., 2017; Stefanski and

212 [Jahn 2020](#)) indicate that it is viable to represent the geometrical and thermodynamic
213 properties of solute and solvent molecules within manageable simulation times of tens of
214 picoseconds.

215 2.3. *Ab initio* XANES calculations

216 *Ab initio* XANES spectra were calculated with the FDMNES program ([Joly, 2001](#)).
217 The final states were calculated by the Finite Difference Method (FDM) to solve the
218 Schrodinger equation in the cluster. The FDM can provide accurate calculation especially for
219 structure of low-symmetry aqueous complexes ([Joly, 2001](#); [Brugger et al., 2007](#)), as in this
220 study. To compare with the experimental spectra, the calculated spectra were convoluted with
221 a Lorentzian function of energy-dependent width to account for the core-hole lifetime
222 broadening (4.52 eV; [Krause and Oliver, 1979](#)) and the inelastic plasmon interactions with
223 the photoelectron. The energy resolution of the monochromator was taken into account by
224 convoluting the calculated XANES spectra with a Gaussian function (width of 1.024 eV). We
225 also calculated the XANES spectra of the five Mo-O-S complexes simulated with molecular
226 dynamics ($\text{MoO}_{4-x}\text{S}_x^{2-}$, $x = 0-4$) at 77 and 300 °C. 40 frames were chosen at every 0.2 ps from
227 each MD simulation box and the final spectra were obtained by taking the average of
228 FDMNES simulations of the 40 frames. The radius cutoff of all calculations was 5 Å.

3. Results

Molybdenum K-edge XANES and EXAFS spectra for the sample solutions with variable NaHS concentrations between 0.07 to 2 molal (Table 1) were collected between 30-394 °C at 800 bar, as well as the spectra of molybdenum reference compounds, including the molybdate solution for comparison. In general, the data collected from the solution are of good quality for EXAFS analysis at temperature up to 286-330 °C, and then only XANES data are used at higher temperatures up to the maximal temperature values for each solution listed in Table 1. At higher temperatures than the maximal temperatures, the intensity of the spectra dropped dramatically, and the signal become noisier due to a reduced metal solubility and precipitates on the wall of the cell, as a result of the decrease in solution density (e.g., Pokrovski et al. 2005; Liu et al. 2008; Mei et al. 2014). Most spectra of precipitates are too noisy to identify, but the spectrum of the highest NaHS concentrations (Sol6) has similarity with the spectrum of molybdenite (MoS_2) (see insert in Fig. 1b); therefore, the precipitates are likely to be molybdenite. For all the samples, only the good quality XANES and EXAFS of the solution phase are used in the data analysis.

In the following sections we describe the qualitative analysis of the XANES spectra, followed by the quantitative refinement of the EXAFS data for the determination of the coordination structure (bond distance, ligand types and coordination numbers) and predominating species. The results of the molecular dynamics simulations and XANES calculations of the major Mo-O and Mo-S species are then described and compared with the experimental results.

3.1. XANES

Effect of NaHS concentration: The XANES spectra of sample solutions as a function of NaHS concentration at 30 and 286 °C are plotted in Fig. 1a and 1b respectively, together with the spectra of two end-member references, Na_2MoO_4 solid (containing the $[\text{MoO}_4]^{2-}$ moiety – we use [] to refer to the moiety of a species throughout the text) and $(\text{NH}_4)_2\text{MoS}_4(\text{s})$ (containing the $[\text{MoS}_4]^{2-}$ moiety). The spectra of the sample solutions plot between the spectra of these two references, with the dominant absorbing species in the sample solutions evolving from $[\text{MoO}_4]^{2-}$ -like to $[\text{MoS}_4]^{2-}$ -like structures with increasing bisulfide concentration. The $[\text{MoO}_4]^{2-}$ structure has a distinctive pre-edge peak at ~20,005 eV (Feature A), which is associated with the 1s – 4d transition, and related to (i) the number of Mo=O bonds per Mo atom and (ii) the symmetry of the structure, the more symmetrical structures having a smaller

pre-edge (Borg et al., 2012 and references therein). The pre-edge peak of the spectrum of the $[\text{MoS}_4]^{2-}$ moiety at 20,003 eV is smaller than that of the $[\text{MoO}_4]^{2-}$ moiety at 20,005 eV (Fig. 1a). At both temperatures (30 and 286 °C), the pre-edge feature decreases with increasing NaHS concentration, indicating an increased contribution of the $[\text{MoS}_4]^{2-}$ -like species. The features B, C, and D also show systematic changes as a function of NaHS concentration, consistent with the features of the two end-member spectra.

Effect of increasing temperature. Temperature has a smaller effect on the XANES spectra than HS^- concentration (Fig. 2). The spectral changes with increasing temperatures are mainly reflected in the growing pre-edge peak (feature of the $[\text{MoO}_4]^{2-}$ moiety) with increasing temperature for most solutions (except for low NaHS solutions such as Sol14 and Sol12), indicating increased contribution from $[\text{MoO}_4]^{2-}$ with increasing temperature.

3.2. EXAFS

The changes of the EXAFS spectra as a function of NaHS concentration (Fig. 3) are similar to the trend of the XANES spectra. In particular, the (phase uncorrected) Fourier transform of the EXAFS shows a decreasing intensity of the first peak at ~ 1.3 Å and increasing intensity of the second peak at ~ 1.8 Å (Fig. 3) with increasing NaHS concentration. Quantitative EXAFS refinements (Table 3) reveal that the first peak is due to Mo-O interaction and the second peak is due to Mo-S.

EXAFS refinement of a molybdate standard ($\text{Na}_2\text{MoO}_{4(\text{s})}$) containing the $[\text{MoO}_4]^{2-}$ moiety) indicates 4 oxygen around Mo atom at average distance of 1.775(4) Å, similar to the Mo-O distance in molybdate solution (1.771-1.777 Å, Borg et al. 2012). The EXAFS refinement of the $(\text{NH}_4)_2\text{MoS}_4(\text{s})$ reference compound ($[\text{MoS}_4]^{2-}$ moiety) yields an average Mo-S bond distance of 2.192(5) Å. This value is similar to the published individual Mo(VI)-S bond distances in $[\text{MoS}_4]^{2-}$ moiety determined by EXAFS refinements (2.20 Å in $(\text{NH}_4)_2\text{MoS}_4(\text{s})$, Bostick et al., 2003; 2.18 Å in MoS_4^{2-} solution, Dahl et al, 2013) and by single crystal diffraction (2.1773(5) -2.2013(8) Å in $(\text{NH}_4)_2\text{MoS}_4(\text{s})$; average = 2.1844(6) Å; Hill et al., 2010). The fitted bond distance of the Mo-S bonds for solutions S5 and S6 with the highest NaHS concentrations, where the Mo-S peaks are most significant (Fig. 4), are also around 2.20 -2.21 Å at 30°C (Table 3); however, the refined Mo-S bond distances of solutions with a low concentration of NaHS and higher temperatures have longer distances of ~ 2.25 -2.29 Å (Table 3). This is due in part to the larger uncertainties in Mo-S distance because of the lower number of coordinated S, but also reflects the effect of the dominating

Mo-O bond. This is confirmed by the molecular dynamics simulations that indicate slightly longer (0.02-0.03 Å) Mo-S distances at elevated temperatures for mixed Mo-S and Mo-O species (See section 3.3).

The fitted number of S ligands increases with increasing NaHS concentrations at all temperatures, at the expense of the number of O atoms, reflecting an increased contribution from thio species (Table 3; the examples at 30 and 286 °C are plotted in Fig. 5). At the highest NaHS concentration (Sol6), the fitted number of S ligands is 3.1(2) at 30 °C; with increasing temperature to 200 °C, the S ligand number increases to 3.4, but at 286 °C and higher temperature, the S peak decreases notably and the O peak increases (Fig. 3b); this reflects an increase in O ligand numbers at those temperatures relative to lower temperatures (Table 3). The temperature dependence of the EXAFS spectra is consistent with, but more apparent than the change of the pre-edge feature in the XANES (Fig. 2). It is notable that while EXAFS analysis provides the average numbers of S and O ligands, it cannot distinguish explicitly the identity of the species present, i.e., for an equal ratio of S:O ligands obtained from the EXAFS refinement, it is not possible to tell whether it is 1:1 mixture of MoO_4^{2-} and MoS_4^{2-} , or predominantly a mixed ligand complex such as $\text{MoS}_2\text{O}_2^{2-}$.

3.3. *Ab initio* MD

The results of simulations of Mo(VI) speciation in solutions containing 2 m NaHS at experimental pressures (800 bar) and at near saturation pressure (200 °C, 20 bar and 300 °C, 100 bar) are listed in Table 4. Overall, the simulations show that fourfold tetrahedral complexes are stable in all simulations. Apart from some deprotonation for simulations that started with $\text{H}_2\text{MoO}_4(\text{aq})$, no ion exchange was observed during the simulation (up to 25 ps). Our previous studies show that in the MD simulations on the scale of picoseconds, it is difficult to observe simple ligand exchange between ions with charge of -1 such as Cl^- , HS^- and OH^- due to the slow exchange kinetics even at elevated temperatures (Mei et al. 2015, 2016, 2020).

3.3.1. Molybdate and hydrogen molybdate ions

As shown in Table 4, jobs 1A, 2A, 3A, 4A, 5A that started with $\text{H}_2\text{MoO}_4(\text{aq})$ established that the hydrogen molybdate ion, HMoO_4^- , was the predominant species over the entire temperature range, with a minor amount of $\text{H}_2\text{MoO}_4(\text{aq})$ at 77 °C (representing room T), 800 bar and a minor amount of molybdate ion, MoO_4^{2-} , at 300 °C, 100 bar. The molybdate

ion is stable in most of the simulations that started with Na_2MoO_4 (simulation 6-11, Table 4), except one solution (simulation 11) at 300 °C in which the stable species is HMoO_4^- . Snapshots of the structures for the observed molybdate and hydrogen molybdate ions are shown in Fig. 6. The Mo-O distance in MoO_4^{2-} is $\sim 1.80(3)$ Å at 77 °C, which is consistent with Liu (2013)'s values of 1.79 Å for Mo-O distance in MoO_4^{2-} . For HMoO_4^- , our MD results show that the Mo-OH bond distances (~ 1.92 - 1.94 Å) are notably longer than the Mo-O distances (~ 1.77 Å) (Table 4), also similar to the previous MD results ($d_{\text{Mo-O}} = 1.76$ Å and $d_{\text{Mo-OH}} = 1.79$ Å, Liu et al., 2013a; $d_{\text{Mo-O}} = 1.72$ Å and $d_{\text{Mo-OH}} = 1.8$ Å, Zhang et al., 2018). Fig. 7 shows the radial distribution functions (RDF) and the coordination numbers (CN; integral of RDF) of Mo-O/S at 300 °C, 800 bar. For simulation 5A, where HMoO_4^- is the predominant species, the RDF peak ranges from 1.7 – 2.0 Å (blue solid line) reflecting the Mo-O bond, which was split into one large peak at 1.77 Å and one small peak at 1.92 Å. The CN (dashed blue) curve shows the first peak results in a CN of 3 (corresponding to three Mo-O bonds) and the second peak adds one more ligand (corresponding to one Mo-OH bond), with a total CN of four. The simulated Mo-O bond distances for molybdate are also consistent with the experimental EXAFS results for molybdate ($d_{\text{Mo-O}} = 1.771$ - 1.777 Å, Borg et al., 2012; $1.775(4)$ Å, this study).

3.3.2. Thiomolybdates

All simulations of thiomolybdates show that fourfold tetrahedral complexes are stable. The geometrical properties of all MD runs are listed in Table 4. Fig. 8 shows the snapshots of thiomolybdates ($\text{MoO}_{4-x}\text{S}_x^{2-}$, $x = 1, 2, 3, 4$) calculated at 300 °C, 800 bar from MD simulations. As shown in Table 4, the Mo-O and Mo-S bonds are easily distinguishable by their lengths (~ 0.45 Å difference), with Mo-O distances of 1.78-1.79 Å and Mo-S distances of 2.23-2.28 Å. This is reflected in the RDF plot (Fig. 7), consistent with the systematic change of Mo-O and Mo-S peaks in the Fourier transforms of the EXAFS data (Figs. 3, 4), and the bond distances from the MD simulations agree well with the fitted EXAFS results of Mo-S distances of 2.20-2.3 Å (Table 3). In particular, both the EXAFS refinements and the MD results predict slightly longer (0.02-0.05 Å) Mo-S distances at elevated temperatures for mixed thiomolybdates ($\text{MoO}_{4-x}\text{S}_x^{2-}$, $x = 1, 2, 3$). This may indicate that temperature favours the loosening of the Mo-S bond in the mixed species.

Similar to the simulations for molybdates, no ligand exchange or proton exchange was observed during these simulations. As mentioned before, ligand exchange is unlikely to be observed in such short simulation times, and in this case, it is even more difficult to see the exchange between O^{2-} and S^{2-} given their strong affinity with Mo^{6+} due to their differences in electronegativity. The MD results shown here provide geometrical properties of thiomolybdates for *ab initio* XANES calculations and provide an independent check and comparison with EXAFS data.

3.4. *Ab initio* XANES calculations

The XANES spectra calculated using FDMNES and selected experimental spectra are plotted in Fig. 9. Although the peaks in the experimental spectra are broader than the calculated spectra, the major spectral features observed in the experimental data are confirmed by *ab initio* XANES simulations. As shown in Fig. 9, the pre-edge peak (Feature A) has a slight shift to higher energy (blue shift) when the structure changes from $[MoS_4]^{2-}$ to $[MoO_4]^{2-}$, with increased intensity for the $[MoO_4]^{2-}$ structure. The peak (Feature B) that is prominent in the spectrum of the $[MoS_4]^{2-}$ moiety decreases with decreasing the number of S ligand and is absent in the spectrum of the $[MoO_4]^{2-}$ structure. These features are consistent with the experimental XANES spectra of the two end member references and the sample solutions (Fig. 9), though the after-edge features C and D in the experimental spectra are not reproduced in the XANES simulations. The spectral changes from 77 °C to 300 °C are minimal, confirming the rigid structure of the thiomolybdate structures.

4. Discussions and conclusions

4.1. Comparison to previous studies at room temperature

It has long been documented that thiomolybdate species are important in surface environments (Muller et al., 1981; Coucouvanis, 1998; Laurie, 2000). Our results from EXAFS analysis, *ab initio* MD simulations and XANES calculations confirmed the formation of thiomolybdates in reduced sulfur-bearing solutions. Several UV-Vis spectroscopic studies (e.g., Harmer and Sykes, 1980; Erickson and Helz, 2000) suggest that thiomolybdate anions prevail in reduced sulfur-bearing fluids, and have established the kinetics of stepwise replacement of O^{2-} ions by S^{2-} . In particular, Erickson and Helz's (2000) UV-Vis study provided the formation constants of the thiomolybdates, and proposed that a sharp transition

occurred between MoO_4^{2-} to MoS_4^{2-} , where a 3-fold change in H_2S concentration results in a 100-fold change in the $\text{MoO}_4^{2-}/\text{MoS}_4^{2-}$ ratio, and MoS_4^{2-} becomes predominant in the solution. In our experiment we identified a mixture of MoO_4^{2-} and MoS_4^{2-} in the most solutions, with the MoS_4^{2-} predominant in the high NaHS concentration solutions. The fitted maximum S ligand number is 3.4 in the 2 m NaHS solution at 200 °C (Table 3). We did not observe 100% MoS_4^{2-} in any solutions, as we were not able to prepare a sufficiently concentrated solution, due to the precipitation of MoS_2 , for XAS measurement at the lower pH (6-8) where MoS_4^{2-} would be more predominant as shown in the previous studies at room temperatures (e.g., Erickson and Helz, 2000),

Using the formation constants for the $\text{MoO}_{4-x}\text{S}_x^{2-}$ ($x=1, 2, 3, 4$) species determined by Erickson and Helz (2000) and the thermodynamic properties of molybdate species (Minubayeva and Seward, 2010) and molybdenite by Robie and Hemingway (1995), we calculated the stability fields of predominant Mo species in a solution of 0.01 m Mo and 0.1 m total sulfur at 25 °C (Fig. 10a). Under reduced conditions, molybdenite predominates in solutions with a wide range of pH, and molybdate species under more oxidised conditions; which is contradictory to the thiomolybdate species observed in our experiments and previous UV-Vis studies (e.g., Harmer and Sykes, 1980; Erickson and Helz, 2000). For the sake of showing the only predominating aqueous species, we removed the molybdenite from the calculations, and the results show that MoS_4^{2-} predominates when $\text{pH}<11$, and mixed $\text{MoS}_4^{2-}/\text{MoO}_4^{2-}$ species in pH 11-12 (Fig. 10b), which is qualitatively consistent with previous studies and our experimental results, where no precipitation of molybdenite is observed in the experimental solutions. There are two possible explanations for the absence of molybdenite in the experimental solution: either slow precipitation of molybdenite precipitation is too slow to be observed, and/or uncertainties in the thermodynamic properties of molybdenite causing overestimation of its stability.

4.2. Stability of thiomolybdate at high P-T

Our experimental results show that the thiomolybdate species with various ratios of sulfur and oxygen ligands are stable in our experimental solutions up to 330 °C. In the absence of experimental determination of thermodynamic properties for thiomolybdate species at elevated P-T conditions, it is useful to extrapolate the available data at ambient temperatures to elevated temperatures. The validity of these extrapolations can be evaluated

by comparison with the experimental EXAFS results, and the models can be used to assess the role of the thiomolybdates in transporting molybdenum in ore fluids.

[Erickson and Helz \(2000\)](#) determined the stability constants for $(\text{MoO}_{4-x}\text{S}_x^{2-})$ in H_2S -bearing solutions at 25 °C. Based on their data, we used the isocoulombic approach ([Gu et al., 1994](#)) to extrapolate the formation constants of the $\text{MoO}_{4-x}\text{S}_x^{2-}$ species up to 300 °C and water saturated pressure ([Table 5](#)).

The basic assumption of the iso-coulombic extrapolation is that when a reaction has equal numbers of like-charged species on both sides of the reaction (iso-coulombic reaction), both heat capacity and entropy changes are small and tend to cancel one another, and the change of free energy is constant and independent of temperature ([Gu et al., 1994](#), [Liu et al., 2012](#)):

$$DG_T^0 = DG_{T_r}^0 = -R \cdot T \cdot \ln K_T \quad (2)$$

where $DG_{T_r}^0$ and DG_T^0 refer to the standard Gibbs free energy ($\text{J} \cdot \text{mol}^{-1}$) of reaction at reference temperature T_r (298.15 K) and the experimental temperature T (in Kelvin), R is the ideal gas constant ($\text{J} \cdot \text{K}^{-1} \cdot \text{mol}^{-1}$) and K_T is the equilibrium constant for the iso-coulombic reaction. Therefore, the $\log K$ for an iso-coulombic reaction at experimental temperature can be estimated from the $\log K$ values at the reference temperature where the thermodynamic properties are available (e.g., [Liu et al., 2012](#)):

$$\log K_T = \frac{T_r}{T} \log K_{T_r} \quad (3)$$

In the case of the thiomolybdate species, it is convenient that all the reactions of the replacement of O^{2-} by S^{2-} meet the requirement of iso-coulombic reactions, i.e., species on both sides of the reactions are equally charged:

Therefore, the stepwise formation constants for these species at elevated temperatures can be estimated with the room temperature data of [Erickson and Helz \(2000\)](#) using Eq. (3), and the results are compiled in [Table 5](#).

Based on the extrapolated data, the stability of the thiomolybdate and molybdate species were estimated as a function of pH and Eh at 300 °C in a solution with 0.1 m total sulfur (Fig. 10c). The thiomolybdate species, in particular MoS_4^{2-} , is predicted to be stable in the solution with slightly acidic to neutral pH (pH 6-8) under reduced conditions (sulfide stable), and molybdate species (HMoO_4^- and MoO_4^{2-}) under more oxidised (sulfate stable), and acidic conditions (sulfide stable) (pH<6 at 300 °C),. Similar to the room temperature situation, our experimental results are qualitatively consistent with the extrapolation based on the room temperature data, i.e., mixed ligand species ($\text{MoO}_3\text{S}^{2-}$ to MoOS_3^{2-}) are present in the solution with pH values of 8.5-9, and MoS_4^{2-} would predominate in wide pH ranges (pH 5 - 8.5 at 300 °C). Such pH range covers the major pH buffers in geological fluids, such as K-feldspar-Muscovite-Quartz (KMq), $\text{H}_2\text{S}/\text{HS}^-$ and $\text{CO}_{2(\text{aq})}/\text{HCO}_3^-$, as shown in Fig. 10.

Therefore, both our experimental data and the extrapolations from room temperature thermodynamic properties indicate that the thiomolybdate species may contribute to Mo transport in hydrothermal sulfur-rich fluids with neutral to alkaline pH range under reduced conditions, and current models of Mo transport in hydrothermal ore fluids need to be re-evaluated to take into account the thiomolybdate species. It is notable that in very acidic solutions the molybdenite would precipitate in sulfide-stable solutions (Fig. 10c). Previous solubility study of Zhang et al. (2012) suggested NaHMo_2S_2 as predominant Mo – S species but they found negative trend of molybdenite solubility with increasing $\log f\text{S}_2$ at 600-800 °C and 2000 bar. Apart from the difference in the experimental temperatures and pressures between the two studies, their experimental solutions have pH range of 1-2 (measurement of quenched solutions, Zhang et al., 2012), which is too acidic to see the impact of thiomolybdate on the solubility of molybdenite, as indicate in Fig 10c. To this end, further experimental studies are needed to improve the thermodynamic properties for the thiomolybdate species at wide range of P-T conditions in order to accurately assess the role of thiomolybdate in transporting Mo (and possibility of transporting other metal ions such as Cu and Fe) in hydrothermal fluids. Finally, Our experiments show that Mo(VI) species predominate at high P-T in solutions with reduced sulfur, consistent with the previous studies in chloride solutions and vapours cited; therefore, Mo(VI) is likely to be the main oxidation state of molybdenum in typical hydrothermal ore fluids. Under extremely reduced conditions Mo(V) and/or Mo(IV) species may occur, but they are not considered due to a lack of experimental evidence and thermodynamic data for these species.

477 *Acknowledgement:* Research funding is from CSIRO and ARC. The EXAS measurements
478 were conducted at the BM30B FAME beamline of the European Synchrotron Radiation
479 Facility (ESRF), with travel funding from The International Synchrotron Access Program
480 (ISAP) administered by the Australian Synchrotron. We thank Dr Peter Kappen and Dr
481 Jeremy Wykes for helping in collect spectra of some standard compound at the XAS
482 beamline of the Australian Synchrotron. The MD simulations were supported by the Pawsey
483 Supercomputing Centre with funding from the Australian and Western Australian, and the
484 National Computational Infrastructure (NCI) supported by the Australian Government.

485 5. References

- 486 Asael, D., Rouxel, O., Poulton, S.W., Lyons, T.W. and Bekker, A. (2018) Molybdenum
487 record from black shales indicates oscillating atmospheric oxygen levels in the early
488 Paleoproterozoic. *American Journal of Science* 318, 275-299.
- 489 Bankura, A., Karmakar, A., Carnevale, V., Chandra, A. and Klein, M.L. (2014) Structure,
490 Dynamics, and Spectral Diffusion of Water from First-Principles Molecular
491 Dynamics. *The Journal of Physical Chemistry C* 118, 29401-29411.
- 492 Becke, A.D. (1988) Density-functional exchange-energy approximation with correct
493 asymptotic behavior. *Physical review. A, General physics* 38, 3098-3100.
- 494 Bernholc, J. and Stiefel, E.I. (1985) Electronic structure of thiomolybdates (MoS_4^{2-} and
495 $\text{Mo}_3\text{S}_9^{2-}$). *Inorganic Chemistry* 24, 1323-1330.
- 496 Bethke, C.M. (2007) *Geochemical and Biogeochemical Reaction Modeling*, 2 ed. Cambridge
497 University Press, Cambridge.
- 498 Bonnetti, C., Zhou, L., Riegler, T., Brugger, J. and Fairclough, M. (2020) Large S isotope and
499 trace element fractionations in pyrite of uranium roll front systems result from
500 internally-driven biogeochemical cycle. *Geochimica Cosmochimica Acta* 282, 113–
501 132.
- 502 Borg, S., Liu, W.H., Etschmann, B., Tian, Y. and Brugger, J. (2012) An XAS study of
503 molybdenum speciation in hydrothermal chloride solutions from 25-385 °C and
504 600 bar. *Geochimica Et Cosmochimica Acta* 92, 292-307.

505 Bostick, B.C., Fendorf, S. and Helz, G.R. (2003) Differential Adsorption of Molybdate and
 506 Tetrathiomolybdate on Pyrite (FeS₂). Environmental Science & Technology 37, 285-
 507 291.

508 Brugger, J., Etschmann, B., Liu, W., Testemale, D., Hazemann, J.L., Emerich, H., van Beek,
 509 W. and Proux, O. (2007) An XAS study of the structure and thermodynamics of Cu(I)
 510 chloride complexes in brines up to high temperature (400 °C, 600 bar). Geochimica Et
 511 Cosmochimica Acta 71, 4920-4941.

512 Brugger, J., Liu, W., Etschmann, B., Mei, Y., Sherman, D.M. and Testemale, D. (2016) A
 513 review of the coordination chemistry of hydrothermal systems: do coordination
 514 changes make ore deposits ? Chemical Geology 447, 219–253.

515 Bullock, L.A. and Parnell, J. (2017) Selenium and molybdenum enrichment in uranium roll-
 516 front deposits of Wyoming and Colorado, USA. Journal of Geochemical Exploration
 517 180, 101-112.

518 Candela, P.A. and Holland, H.D. (1984) The partitioning of copper and molybdenum
 519 between silicate melts and aqueous fluids. Geochimica et Cosmochimica Acta 48,
 520 373-3

521 Chantler, C.T. (1995) Theoretical Form Factor, Attenuation, and Scattering Tabulation for
 522 Z=1–92 from E=1–10 eV to E=0.4–1.0 MeV. Journal of Physical and Chemical
 523 Reference Data 24, 71-643.80.

524 Combariza, J.E., Enemark, J.H., Barfield, M. and Facelli, J.C. (1989) *Ab initio* molecular
 525 orbital studies of chemical shielding in transition-metal compounds: molybdenum-95
 526 shielding in molybdate and thiomolybdate [MoO_nS_(4-n)]²⁻ anions. J Am Chem Soc 111,
 527 7619-7621.

528 Coucouvanis, D. (1998) Syntheses, Structures, and Reactions of Binary and Tertiary
 529 Thiomolybdate Complexes Containing the (O)Mo(S_x) and (S)Mo(S_x) Functional
 530 Groups (x = 1, 2, 4), in: Sykes, A.G. (Ed.), Advances in Inorganic Chemistry.
 531 Academic Press, pp. 1-73.

532 Dadze, T.P., Kashirtseva, G.A., Novikov, M.P. and Plyasunov, A.V. (2017) Solubility of
 533 MoO₃ in acid solutions and vapor-liquid distribution of molybdic acid. Fluid Phase
 534 Equilib 440, 64-76.

535 Dadze, T.P., Kashirtseva, G.A., Novikov, M.P. and Plyasunov, A.V. (2018) Solubility of
 536 MoO_3 in Aqueous Acid Chloride-Bearing Solutions at 573 K. *Journal of Chemical &*
 537 *Engineering Data* 63, 1827-1832.

538 Dahl, T.W., Chappaz, A., Fitts, J.P. and Lyons, T.W. (2013) Molybdenum reduction in a
 539 sulfidic lake: Evidence from X-ray absorption fine-structure spectroscopy and
 540 implications for the Mo paleoproxy. *Geochimica et Cosmochimica Acta* 103, 213-
 541 231.

542 Driesner, T. and Heinrich, C.A. (2007) The system H_2O – NaCl . Part I: Correlation formulae
 543 for phase relations in temperature–pressure–composition space from 0 to 1000 °C, 0
 544 to 5000 bar, and 0 to 1 X_{NaCl} . *Geochimica et Cosmochimica Acta* 71, 4880-4901.

545 Driesner, T. (2007) The system H_2O – NaCl . Part II: Correlations for molar volume, enthalpy,
 546 and isobaric heat capacity from 0 to 1000 °C, 1 to 5000 bar, and 0 to 1 X_{NaCl} .
 547 *Geochimica et Cosmochimica Acta* 71, 4902-4919.

548 Ehrig, K., Mcphie, J. and Kamenetsky, V.S. (2012) Geology and mineralogical zonation of
 549 the Olympic Dam iron oxide Cu-U-Au-Ag deposit, South Australia. *Society of*
 550 *Economic Geologists, Special Publications* 16, 237–267.

551 El-Issa, B.D., Ali, A.A.M. and Zanati, H. (1989) Theoretical study of the bonding in sulfur-
 552 and oxygen-containing molybdenum species of the general formula MoSnO_{4-n}^{2-} ($n =$
 553 $0-4$). *Inorganic Chemistry* 28, 3297-3305.

554 Erickson, B.E. and Helz, G.R. (2000) Molybdenum(VI) speciation in sulfidic waters::
 555 Stability and lability of thiomolybdates. *Geochimica et Cosmochimica Acta* 64, 1149-
 556 1158.

557 Etschmann, B.E., Liu, W., Testemale, D., Muller, H., Rae, N.A., Proux, O., Hazemann, J.L.
 558 and Brugger, J. (2010) An in situ XAS study of copper(I) transport as hydrosulfide
 559 complexes in hydrothermal solutions (25-592 °C, 180-600 bar): Speciation and
 560 solubility in vapor and liquid phases. *Geochimica Et Cosmochimica Acta* 74, 4723-
 561 4739.

562 Gillan, M.J., Alfè, D. and Michaelides, A. (2016) Perspective: How good is DFT for water?
 563 *The Journal of Chemical Physics* 144, 130901.

564 Goedecker, S., Teter, M. and Hutter, J. (1996) Separable dual-space Gaussian
 565 pseudopotentials. *Physical Review B* 54, 1703-1710.

566 Gu, Y., Gammons, C.H. and Bloom, M.S. (1994) A one-term extrapolation method for
 567 estimating equilibrium constants of aqueous reactions at elevated temperatures.
 568 *Geochimica et Cosmochimica Acta* 58, 3545-3560.

569 Guan, Q. Mei, Y., Etschmann, B., Testemale, D., Louvel, M. and Brugger, J. (2020) Yttrium
 570 complexation and hydration in chloride-rich hydrothermal fluids: a combined *ab initio*
 571 molecular dynamics and in situ X-ray absorption spectroscopy study. *Geochimica et*
 572 *Cosmochimica Acta* 281, 168-189.

573 Harmer, M.A. and Sykes, A.G. (1980) Kinetics of the interconversion of sulfido- and
 574 oxomolybdate(VI) species $\text{MoO}_x\text{S}_{4-x}^{2-}$ in aqueous solutions. *Inorganic Chemistry* 19,
 575 2881-2885.

576 Hartwigsen, C., Goedecker, S. and Hutter, J. (1998) Relativistic separable dual-space
 577 Gaussian pseudopotentials from H to Rn. *Physical Review B* 58, 3641-3662.

578 Helz, G.R., Miller, C.V., Charnock, J.M., Mosselmans, J.F.W., Patrick, R.A.D., Garner, C.D.
 579 and Vaughan, D.J. (1996) Mechanism of molybdenum removal from the sea and its
 580 concentration in black shales: EXAFS evidence. *Geochimica et Cosmochimica Acta*
 581 60, 3631-3642.

582 Helz, G.R. and Erickson, B.E. (2011) Extraordinary stability of copper(I)–tetrathiomolybdate
 583 complexes: Possible implications for aquatic ecosystems. *Environmental Toxicology*
 584 *and Chemistry* 30, 97-102.

585 Helz, G.R., Erickson, B.E. and Vorlicek, T.P. (2014) Stabilities of thiomolybdate complexes
 586 of iron; implications for retention of essential trace elements (Fe, Cu, Mo) in sulfidic
 587 waters. *Metallomics* 6, 1131-1140.

588 Hoover, W.G. (1985) Canonical dynamics: Equilibrium phase-space distributions. *Physical*
 589 *Review A* 31, 1695-1697.

590 Humphrey, W., Dalke, A. and Schulten, K. (1996) VMD: visual molecular dynamics. *Journal*
 591 *of molecular graphics* 14, 33-38, 27-38.

592 Hurtig, N.C. and Williams-Jones, A.E. (2014) An experimental study of the transport of gold
 593 through hydration of AuCl in aqueous vapour and vapour-like fluids. *Geochimica et*
 594 *Cosmochimica Acta* 127, 305-325.

595 Hutter, J., Iannuzzi, M., Schiffmann, F. and VandeVondele, J. (2014) cp2k: atomistic
 596 simulations of condensed matter systems. WIREs Computational Molecular Science
 597 4, 15-25.

598 Jahn, S. and Schmidt, C. (2010) Speciation in Aqueous MgSO₄ Fluids at High Pressures and
 599 High Temperatures from *Ab Initio* Molecular Dynamics and Raman Spectroscopy.
 600 The Journal of Physical Chemistry B 114, 15565-15572.

601 Joly, Y. (2001) X-ray absorption near-edge structure calculations beyond the muffin-tin
 602 approximation. Phys. Rev. B 63, art. no.-125120.

603 Krause, M.O. and Oliver, J.H. (1979) Natural widths of atomic K and L levels, K_α X-ray
 604 lines and several KLL Auger lines. Journal of Physical and Chemical Reference Data
 605 8, 329-338.

606 Keppler, H. and Wyllie, P.J. (1991) Partitioning of Cu, Sn, Mo, W, U, and Th between Melt
 607 and Aqueous Fluid in the Systems Haplogranite-H₂O-HCl and Haplogranite-H₂O-Hf.
 608 Contributions to Mineralogy and Petrology 109, 139-150.

609 Kudrin, A.V. (1989) Behavior of Mo in aqueous NaCl and KCl solutions at 300-450 °C.
 610 Geochem. Int. 26, 87-99.

611 Laurie, Stuart H. (2000) Thiomolybdates — Simple but Very Versatile Reagents. European
 612 Journal of Inorganic Chemistry 2000, 2443-2450.

613 Lee, C., Yang, W. and Parr, R.G. (1988) Development of the Colle-Salvetti correlation-
 614 energy formula into a functional of the electron density. Physical Review B 37, 785-
 615 789.

616 Lee, H.S. and Tuckerman, M.E. (2006) Structure of liquid water at ambient temperature from
 617 *ab initio* molecular dynamics performed in the complete basis set limit. The Journal of
 618 Chemical Physics 125, 154507.

619 Lin, I.C., Seitsonen, A.P., Tavernelli, I. and Rothlisberger, U. (2012) Structure and Dynamics
 620 of Liquid Water from *Ab Initio* Molecular Dynamics-Comparison of BLYP, PBE, and
 621 revPBE Density Functionals with and without van der Waals Corrections. Journal of
 622 chemical theory and computation 8, 3902-3910.

623 Liu, W., Brugger, J., Etschmann, B., Testemale, D. and Hazemann, J.-L. (2008) The
 624 solubility of nantokite (CuCl(s)) and Cu speciation in low density fluids near the

critical isochore: an in-situ XAS study. *Geochimica Et Cosmochimica Acta* 72, 4094–
4106.

Liu, W., Borg, S., Testemale, D., Etschmann, B., Hazemann, J. and Brugger, J. (2011)
Speciation and thermodynamic properties for cobalt chloride complexes in
hydrothermal fluids at 35-440 °C and 600 bar: An in-situ XAS study. *Geochimica Et
Cosmochimica Acta* 75, 1227-1248.

Liu, W., Borg, S., Etschmann, B., Mei, Y. and Brugger, J. (2012) An XAS study of
speciation and thermodynamic properties of aqueous zinc bromide complexes at 25-
150 °C. *Chemical Geology* 298, 57-69.

Liu, W., Etschmann, B., Testemale, D., Hazemann, J.-L., Rempel, K., Mueller, H. and
Brugger, J. (2014) Gold transport in hydrothermal fluids: Competition among the Cl⁻,
Br⁻, HS⁻ and NH₃(aq) ligands. *Chemical Geology* 376, 11-19.

Liu, X., Cheng, J., Sprik, M. and Lu, X. (2013a) Solution structures and acidity constants of
molybdic acid. *Journal of Physical Chemistry Letters* 4, 2926-2930.

Liu, X., Sprik, M. and Cheng, J. (2013b) Hydration, acidity and metal complexing of
polysulfide species: A first principles molecular dynamics study. *Chemical Physics
Letters* 563, 9-14.

Mei, Y., Sherman, D.M., Liu, W. and Brugger, J. (2013) *Ab initio* molecular dynamics
simulation and free energy exploration of copper(I) complexation by chloride and
bisulfide in hydrothermal fluids. *Geochimica Et Cosmochimica Acta* 102, 45-64.

Mei, Y., Sherman, D., Liu, W. and Brugger, J. (2014) Metal complexation and ion hydration
in low density hydrothermal fluids: *ab initio* molecular dynamics simulation of Cu(I)
and Au(I) in chloride solutions (25-1000 °C, 1-5000 bar). *Geochimica et
Cosmochimica Acta*, 131, 196–212.

Mei, Y., Etschmann, B., Liu, W., Sherman, D.M., Barnes, S.J., Fiorentini, M.L., Seward,
T.M., Testemale, D. and Brugger, J. (2015) Palladium complexation in chloride- and
bisulfide-rich fluids: Insights from *ab initio* molecular dynamics simulations and X-
ray absorption spectroscopy. *Geochimica Et Cosmochimica Acta* 161, 128-145.

Mei, Y., Etschmann, B., Liu, W., Sherman, D.M., Testemale, D. and Brugger, J. (2016)
Speciation and thermodynamic properties of zinc in sulfur-rich hydrothermal fluids:

655 Insights from *ab initio* molecular dynamics simulations and X-ray absorption
 656 spectroscopy. *Geochimica Et Cosmochimica Acta* 179, 32-52.

657 Mei, Y., Liu, W., Brugger, J., Sherman, D.M. (2018) The dissociation mechanism and
 658 thermodynamic properties of HCl(aq) in hydrothermal fluids (to 700 °C, 60 kbar) by
 659 *ab initio* molecular dynamics simulations. *Geochimica Cosmochimica Acta* 226, 84-
 660 106.

661 Mei, Y., Liu, W., Brugger, J., Guan, Q. (2020) Gold solubility in alkaline and ammonia-rich
 662 hydrothermal fluids: insights from *ab initio* molecular dynamics simulations.
 663 *Geochimica Cosmochimica Acta*. In press, <https://doi.org/10.1016/j.gca.2019.12.031>.

664 Minubayeva, Z. and Seward, T.M. (2010) Molybdic acid ionisation under hydrothermal
 665 conditions to 300 °C. *Geochimica Et Cosmochimica Acta* 74, 4365-4374.

666 Mosselmans, J.F.W., Pattrick, R.A.D., Charnock, J.M. and Sole, V.A. (1999) EXAFS of
 667 copper in hydrosulfide solutions at very low concentrations: implications for the
 668 speciation of copper in natural waters. *Mineralogical Magazine* 63, 769-772.

669 Müller, A., Diemann, E., Jostes, R. and Bögge, H. (1981) Transition Metal Thiometalates:
 670 Properties and Significance in Complex and Bioinorganic Chemistry. *Angewandte*
 671 *Chemie International Edition in English* 20, 934-955.

672 Nosé, S. (1984) A unified formulation of the constant temperature molecular dynamics
 673 methods. *The Journal of Chemical Physics* 81, 511-519.

674 Pokrovski, G.S., Roux, J. and Harrichoury, J.C. (2005) Fluid density control on vapor-liquid
 675 partitioning of metals in hydrothermal systems. *Geology*, 33, 657-660.

676 Pokrovski, G.S., Tagirov, B.R., Schott, J., Hazemann, J.L. and Proux, O. (2009) A new view
 677 on gold speciation in sulfur-bearing hydrothermal fluids from in situ X-ray absorption
 678 spectroscopy and quantum-chemical modeling. *Geochimica et Cosmochimica Acta*
 679 73, 5406-5427.

680 Proux, O., Biquard, X., Lahera, E., Menthonnex, J.J., Prat, A., Ulrich, O., Soldo, Y.,
 681 Trevisson, P., Kapoujyan, G., Perroux, G., Taunier, P., Grand, D., Jeantet, P.,
 682 Deleglise, M., Roux, J.P. and Hazemann, J.L. (2005) FAME: A new beamline for X-
 683 ray absorption investigations of very-diluted systems of environmental, material and
 684 biological interests. *Physica Scripta* T115, 970-973.

685 Ravel, B. and Newville, M. (2005) ATHENA, ARTEMIS, HEPHAESTUS: data analysis for
686 X-ray absorption spectroscopy using IFEFFIT. *J. Synchrotr. Radiat.* 12, 537-541.

687 Rehr, J.J., Kas, J.J., Prange, M.P., Sorini, A.P., Takimoto, Y. and Vila, F. (2009) *Ab initio*
688 theory and calculations of X-ray spectra. *Comptes Rendus Physique* 10, 548-559.

689 Rempel, K.U., Migdisov, A.A. and Williams-Jones, A.E. (2006) The solubility and speciation
690 of molybdenum in water vapour at elevated temperatures and pressures: Implications
691 for ore genesis. *Geochimica Et Cosmochimica Acta* 70, 687-696.

692 Rempel, K.U., Williams-Jones, A.E. and Migdisov, A.A. (2008) The solubility of
693 molybdenum dioxide and trioxide in HCl-bearing water vapour at 350 °C and
694 pressures up to 160 bars. *Geochimica et Cosmochimica Acta* 72, 3074-3083.

695 Rempel, K.U., Williams-Jones, A.E. and Migdisov, A.A. (2009) The partitioning of
696 molybdenum(VI) between aqueous liquid and vapour at temperatures up to 370 °C.
697 *Geochimica et Cosmochimica Acta* 73, 3381-3392.

698 Robie, R.A. and Hemingway, B.S. (1995) Thermodynamic properties of minerals and related
699 substances at 298.15 K and 1 bar (10^5 Pascals) pressure and at higher temperatures.
700 US Geological Survey Bulletin 2131, 461.

701 Smedley, P.L. and Kinniburgh, D.G. (2017) Molybdenum in natural waters: A review of
702 occurrence, distributions and controls. *Appl Geochem* 84, 387-432.

703 Pokrovski, G.S., Tagirov, B.R., Schott, J., Hazemann, J.L. and Proux, O. (2009) A new view
704 on gold speciation in sulfur-bearing hydrothermal fluids from in situ X-ray absorption
705 spectroscopy and quantum-chemical modeling. *Geochimica Et Cosmochimica Acta*
706 73, 5406-5427.

707 Rempel, K.U., Migdisov, A.A. and Williams-Jones, A.E. (2006) The solubility and speciation
708 of molybdenum in water vapour at elevated temperatures and pressures: Implications
709 for ore genesis. *Geochimica Et Cosmochimica Acta* 70, 687-696.

710 Rempel, K.U., Williams-Jones, A.E. and Migdisov, A.A. (2008) The solubility of
711 molybdenum dioxide and trioxide in HCl-bearing water vapour at 350 °C and
712 pressures up to 160 bars. *Geochimica et Cosmochimica Acta* 72, 3074-3083.

713 Rempel, K.U., Williams-Jones, A.E. and Migdisov, A.A. (2009) The partitioning of
714 molybdenum (VI) between aqueous liquid and vapour at temperatures up to 370 °C.
715 *Geochimica et Cosmochimica Acta* 73, 3381-3392.

716 Saha, P., Anderson, A.J., Lee, T. and Klemm, M. (2017) The Solubility of Tugarinovite
717 (MoO₂) in H₂O at Elevated Temperatures and Pressures. *Geofluids* 2017, 12.

718 Seedorff, E., Dilles, J.H., Proffett, J.M., Jr., Einaudi, M.T., Zurcher, L., Stavast, W.J.A.,
719 Johnson, D.A., Barton, M.D., Hedenquist, J.W., Thompson, J.F.H., Goldfarb, R.J. and
720 Richards, J.P. (2005) Porphyry Deposits: Characteristics and Origin of Hypogene
721 Features, One Hundredth Anniversary Volume. Society of Economic Geologists, p. 0.

722 Stefanski, J. and Jahn, S. (2020) Yttrium speciation in subduction zone fluids from *ab initio*
723 molecular dynamics simulations. *Solid Earth Discuss.* 2020, 1-36.

724 Sulpizi, M., Salanne, M., Sprik, M. and Gaigeot, M.-P. (2013) Vibrational Sum Frequency
725 Generation Spectroscopy of the Water Liquid–Vapor Interface from Density
726 Functional Theory-Based Molecular Dynamics Simulations. *The Journal of Physical*
727 *Chemistry Letters* 4, 83-87.

728 Tessin, A., Chappaz, A., Hendy, I. and Sheldon, N. (2018) Molybdenum speciation as a
729 paleo-redox proxy: A case study from late cretaceous western interior seaway black
730 shales. *Geology* 47, 59-62

731 Testemale, D., Argoud, R., Geaymond, O. and Hazemann, J.L. (2005) High pressure high
732 temperature cell for x-ray absorption and scattering techniques. *Rev. Sci. Instrum.* 76,
733 043905.

734 Testemale, D., Brugger, J., Liu, W., Etschmann, B. and Hazemann, J.-L. (2009) In-situ X-ray
735 absorption study of Iron(II) speciation in brines up to supercritical conditions.
736 *Chemical Geology* 264, 295-310.

737 Tooth, B., Etschmann, B., Pokrovski, G.S., Testemale, D., Hazemann, J.L., Grundler, P.V.
738 and Brugger, J. (2013) Bismuth speciation in hydrothermal fluids: An x-ray
739 absorption spectroscopy and solubility study. *Geochimica et Cosmochimica Acta*, 101
740 156-172.

741 Tossell, J.A. (2005) Calculation of the UV-visible spectra and the stability of Mo and Re
742 oxysulfides in aqueous solution. *Geochimica et Cosmochimica Acta* 69, 2497-2503.

743 Ulrich, T. and Mavrogenes, J. (2008) An experimental study of the solubility of molybdenum
744 in H₂O and KCl-H₂O solutions from 500 °C to 800 °C, and 150 to 300 MPa.
745 *Geochimica et Cosmochimica Acta* 72, 2316-2330.

746 VandeVondele, J. and Hutter, J. (2007) Gaussian basis sets for accurate calculations on
 747 molecular systems in gas and condensed phases. *The Journal of Chemical Physics*
 748 127, 114105.

749 Wagner, J., Haigis, V., Künzel, D. and Jahn, S. (2017) Trace element partitioning between
 750 silicate melts – A molecular dynamics approach. *Geochimica et Cosmochimica Acta*
 751 205, 245-255.

752 Wood, S.A., Crerar, D.A. and Borcsik, M.P. (1987) Solubility of the assemblage pyrite-
 753 pyrrhotite-magnetite-sphalerite-galena-gold-stibnite-bismuthinite-argen- tite-
 754 molybdenite in H₂O-NaCl-CO₂ solutions from 200 to 350 °C degrees. *Economic*
 755 *Geology* 82, 1864-1887.

756 Zajacz, Z., Candela, P.A. and Piccoli, P.M. (2017) The partitioning of Cu, Au and Mo
 757 between liquid and vapor at magmatic temperatures and its implications for the
 758 genesis of magmatic-hydrothermal ore deposits. *Geochimica et Cosmochimica Acta*
 759 207, 81-101.

760 Zhang, N., Yi, H., Zeng, D., Zhao, Z., Wang, W. and Costanzo, F. (2018) Structure evolution
 761 of mononuclear tungsten and molybdenum species in the protonation process: Insight
 762 from FPMD and DFT calculations. *Chemical Physics* 502, 77-86.

763

764 Table 1 Sample compositions. pH values are calculated from the compositions of NaHS,
765 Na₂MoO₄ and NaCl using HCh modelling package and the default database (Shvarov, 2008).

Solution	Composition (m: mole/kg)	T (°C)	pH (30 °C, 800 bar)	pH (286°C, 800 bar)
Sol 5	0.004 m Na ₂ MoO ₄ , 1.13 m NaHS	30, 103, 194, 286	11.5	8.5
Sol 6	0.006 m Na ₂ MoO ₄ , 2.00 m NaHS	30, 103, 194, 286	11.4	8.5
Sol 8	0.008 m Na ₂ MoO ₄ , 0.28 m NaHS	30, 103, 194, 286, 370, 394	11.9	8.8
Sol 10	0.009 m Na ₂ MoO ₄ , 0.28 m NaHS, 1.0 m NaCl	30, 103, 194, 286, 330	11.8	8.7
Sol 12	0.009 m Na ₂ MoO ₄ , 0.070 m NaHS	30, 103, 194, 286, 330, 370	11.8	8.6
Sol 14	0.009 m Na ₂ MoO ₄ , 0.04 m NaHS	30, 103, 194, 286, 330, 370	12.0	8.8
Sol 15	0.007 m Na ₂ MoO ₄ , 0.43 m NaHS	30, 103, 194, 286, 330, 370, 394	11.8	8.7

766

767 Table 2. Details of ab initio MD simulation using CP2K.

solution compositio n	Jo b ID	T (°C)	P (bar)	Density of box (g/cm ³)	Box size (Å)
1 H ₂ MoO ₄ , 4 Na ⁺ , 4 HS ⁻ , 111 H ₂ O	1	77	800	1.1114 6	15.277
	2	200	20	0.9911	15.872
	3	200	800	1.0291 5	15.674
	4	300	100	0.8844	16.486
	5	300	800	0.9421 6	16.142
1 MoO ₄ ²⁻ , 2 Na ⁺ , 111 H ₂ O	6	77	800	1.0740	15.053
	7	200	800	0.9859	15.489
	8	300	800	0.8887	16.034
1 MoO ₄ ²⁻ , 4Na ⁺ , 2HS ⁻ , 111 H ₂ O	9	77	800	1.1039	15.165
	10	200	800	1.0183	15.578
	11	300	800	0.9260	16.079

768

769

770 Table 3 EXAFS refinement (nO and nS; number of oxygen and sulfur ligands; rO and rS: Mo-O and Mo-S bond distances, n_{tot} : number
771 of total ligands; σ^2 : Debye-waller factor; E0: shift of energy; χ^2_{red} : reduced χ^2 , R-factor: goodness of fit. Fitting range in R space: 1-3
772 (Å)).

.T	nO	rO (Å)	nS	rS (Å)	σ^2	E0	χ^2_{red}	R-factor	k-range
Sol 14	0.04 m HS-								
30	3.8(1)	1.766(5)	0.2(1)	2.276 (fix)	0.002(1)	6.6(6)	128	0.016	3-11
103	3.8(1)	1.769(5)	0.2(1)	2.276 (fix)	0.002(1)				3-11
194	3.3(2)	1.760(6)	0.7(2)	2.29(2)	0.002(1)				3-11
286	3.4(1)	1.758(5)	0.6(1)	2.28(2)	0.002(1)				3-11
330	3.8(1)	1.764(5)	0.2(1)	2.31(8)	0.002(1)				
330	4.0(2)	1.763 (fix)	0.0(2)		0.0029(8)	2.2(9)			3-11
Sol 12	0.07 m HS-								
30	3.3(2)	1.763(9)	0.7(2)	2.276 (fix)	0.002(1)	6.0(7)	221	0.031	3-11
103	3.4(2)	1.765(6)	0.6(2)	2.266 (fix)	0.002(1)				3-11
194	3.2(1)	1.763(6)	0.8(1)	2.286 (fix)	0.004(1)				3-11
286	3.3(2)	1.768(6)	0.7(2)	2.286 (fix)	0.003(1)				3-11
330	3.5(2)	1.767(8)	0.5(2)	2.286 (6)	0.003(1)				3-11
Sol 8	0.28 m HS-								
30	3.0(1)	1.76(2)	1.0(1)	2.24(3)	0.002(1)	1(2)	431	0.038	3-8.5
103	2.7(3)	1.75(1)	1.3(3)	2.24(2)	0.002(1)	5(2)			3-12
194	2.8(3)	1.75(1)	1.2(3)	2.25(2)	0.002(1)	5(2)			3-12
286	3.1(2)	1.75(1)	0.9(2)	2.27(3)	0.003(1)	5(2)			3-12
Sol 10	0.28 m HS- + 1 m NaCl								
30	2.6(2)	1.760(9)	1.4(2)	2.26(1)	0.0023(8)	4(1)	281	0.042	3-11

103	2.5(2)	1.760(9)	1.5(2)	2.27(1)	0.0026(8)				3-11
194	2.4(3)	1.75(2)	1.6(3)	2.27(2)	0.003(2)				3-10
286	2.8(2)	1.750(9)	1.2(2)	2.28(2)	0.0025(9)				3-11
330	3.0(2)	1.75(1)	1.0(2)	2.28(3)	0.003(1)	6(2)			3-11

Sol 15	0.43 m HS-								
30	2.4(2)	1.75(1)	1.6(2)	2.25(1)	0.002(1)	5.2(7)	153	0.024	3-11
103	2.3(1)	1.746(6)	1.7(1)	2.239(7)	0.0014(5)				3-11
194	2.6(2)	1.741(8)	1.4(2)	2.25(1)	0.002(1)				3-11
286	2.9(2)	1.747(7)	1.1(2)	2.26(1)	0.002(2)				3-11
330	3.1(2)	1.750(7)	0.9(2)	2.26(2)	0.002(1)				3-10

Sol 5	1.1 m HS-								
30	1.0(1)	1.688 (fix)	3.0(1)	2.211(5)	0.0025(5)	6.4(5)	171	0.018	3-10
103	0.4(1)	1.668 (fix)	3.6(1)	2.206(4)	0.0017(4)				3-11
194	1.0(2)	1.738 (fix)	3.0(2)	2.212(7)	0.0025(9)				3-11
286	2.4(2)	1.748(9)	1.6(2)	2.26(1)	0.0035(9)				3-11

Sol 6	2 m HS-								
30	0.9(1)	1.708 (fix)	3.1(1)	2.210(5)	0.0015(6)	6.1(4)	157	0.016	3-12
103	0.7(2)	1.718 (fix)	3.3(2)	2.210(5)	0.0018(6)				3-12
194	0.6(1)	1.738 (fix)	3.4(1)	2.212(4)	0.0021(5)				3-12
286	1.7(1)	1.738 (fix)	2.3(1)	2.229(6)	0.0032(6)				3-11

773

774

775

776 Table 4. Simulation results of molybdate and thiomolybdate species.

Initial configuration of Mo	Job ID	T (°C)	P bar	MD time (ps)	species	Species time (ps)	Mo-S			Mo-O		
							CN	R (Å)	σ_s^2 (Å ²)	CN	R (Å)	σ_o^2 (Å ²)
H ₂ MoO ₄	1A	77	800	13.64	H ₂ MoO ₄ & HMoO ₄ ⁻	12.64	0	-	-	4	1.81(4)	0.0057
	2A	200	20	15.55	HMoO ₄ ⁻	14.05	0	-	-	3(O) 1(OH)	1.77(4) 1.93(4)	0.0014 0.0019
	3A	200	800	10.30	HMoO ₄ ⁻	9.30	0	-	-	3(O) 1(OH)	1.77(4) 1.92(5)	0.0013 0.0021
	4A	300	100	13.49	HMoO ₄ ⁻ & MoO ₄ ²⁻	11.99	0	-	-	4	1.81(5)	0.0058
	5A	300	800	10.17	HMoO ₄ ⁻	9.17	0	-	-	3(O) 1(OH)	1.77(3) 1.92(5)	0.0012 0.0027
MoO ₃ S ²⁻	1B	77	800	10.16	MoO ₃ S ²⁻	9.16	1	2.25(5)	0.0026	3	1.79(3)	0.0011
	2B	200	20	11.13	MoO ₃ S ²⁻	10.13	1	2.27(6)	0.0031	3	1.79(4)	0.0013
	3B	200	800	10.39	MoO ₃ S ²⁻	9.39	1	2.27(5)	0.0025	3	1.79(4)	0.0017
	4B	300	100	10.02	MoO ₃ S ²⁻	9.02	1	2.27(6)	0.0031	3	1.79(4)	0.0020
	5B	300	800	10.45	MoO ₃ S ²⁻	9.45	1	2.28(6)	0.0036	3	1.79(4)	0.0018
MoO ₂ S ₂ ²⁻	1C	77	800	10.07	MoO ₂ S ₂ ²⁻	9.07	2	2.24(4)	0.0017	2	1.79(3)	0.0011
	2C	200	20	9.51	MoO ₂ S ₂ ²⁻	8.51	2	2.27(5)	0.0030	2	1.78(3)	0.0012
	3C	200	800	10.20	MoO ₂ S ₂ ²⁻	9.20	2	2.25(6)	0.0030	2	1.79(4)	0.0014
	4C	300	100	10.02	MoO ₂ S ₂ ²⁻	9.02	2	2.26(6)	0.0036	2	1.79(4)	0.0020
	5C	300	800	10.38	MoO ₂ S ₂ ²⁻	9.38	2	2.26(6)	0.0038	2	1.78(4)	0.0017
MoOS ₃ ²⁻	1D	77	800	10.55	MoOS ₃ ²⁻	9.55	3	2.23(4)	0.0016	1	1.78(3)	0.0009
	2D	200	20	9.23	MoOS ₃ ²⁻	8.23	3	2.24(6)	0.0033	1	1.78(4)	0.0014
	3D	200	800	10.02	MoOS ₃ ²⁻	9.02	3	2.24(5)	0.0023	1	1.78(4)	0.0013
	4D	300	100	9.89	MoOS ₃ ²⁻	11.02	3	2.25(6)	0.0034	1	1.77(4)	0.0017
	5D	300	800	10.54	MoOS ₃ ²⁻	9.54	3	2.25(6)	0.0037	1	1.78(4)	0.0014
MoS ₄ ²⁻	1E	77	800	10.51	MoS ₄ ²⁻	9.51	4	2.23(5)	0.0024	0	-	-
	2E	200	20	10.21	MoS ₄ ²⁻	9.21	4	2.23(5)	0.0024	0	-	-
	3E	200	800	10.26	MoS ₄ ²⁻	9.26	4	2.23(5)	0.0027	0	-	-
	4E	300	100	9.90	MoS ₄ ²⁻	8.90	4	2.23(6)	0.0031	0	-	-
	5E	300	800	10.63	MoS ₄ ²⁻	9.63	4	2.23(6)	0.0033	0	-	-
MoO ₄ ²⁻	6	77	800	17.73	MoO ₄ ²⁻	16.73	-	-	-	4	1.80(3)	0.0011
	7	200	800	24.89	MoO ₄ ²⁻	23.89	-	-	-	4	1.80(4)	0.0016
	8	300	800	15.48	MoO ₄ ²⁻	14.48	-	-	-	4	1.81(4)	0.0018
MoO ₄ ²⁻	9	77	800	14.78	MoO ₄ ²⁻	13.78	-	-	-	4	1.80(3)	0.0011
	10	200	800	13.04	MoO ₄ ²⁻	12.04	-	-	-	4	1.80(4)	0.0017
	11	300	800	10.37	HMoO ₄ ⁻	9.37	-	-	-	3(O) 1(OH)	1.77(3) 1.94(5)	0.0012 0.0025

778 Table 5 The stepwise formation constants for thiomolybdate species extrapolated from
779 [Erickson and Helz \(2000\)](#).

Temperature (°C)	0	25	60	100	150	200	250	300
$\text{MoO}_4^{2-} + \text{H}_2\text{S}(\text{aq}) = \text{MoO}_3\text{S}^{2-} + \text{H}_2\text{O}(\text{l})$	5.67	5.19	4.64	4.15	3.66	3.27	2.96	2.70
$\text{MoO}_3\text{S}^{2-} + \text{H}_2\text{S}(\text{aq}) = \text{MoO}_2\text{S}_2^{2-} + \text{H}_2\text{O}(\text{l})$	5.24	4.80	4.30	3.84	3.38	3.02	2.74	2.50
$\text{MoO}_2\text{S}_2^{2-} + \text{H}_2\text{S}(\text{aq}) = \text{MoOS}_3^{2-} + \text{H}_2\text{O}(\text{l})$	5.46	5.00	4.47	4.00	3.52	3.15	2.85	2.60
$\text{MoOS}_3^{2-} + \text{H}_2\text{S}(\text{aq}) = \text{MoS}_4^{2-} + \text{H}_2\text{O}(\text{l})$	5.33	4.88	4.37	3.90	3.44	3.08	2.78	2.54

780

781

782 Figure captions

783 Fig. 1. Molybdenum K-edge XANES spectra as a function of NaHS concentration at 30 °C
784 (a) and 286 °C (b). Dotted lines in (a) represent standard structures: Also shown in (a)
785 are the measured spectra of $[\text{MoO}_4]^{2-}$ moiety in $\text{Na}_2\text{MoO}_{4(\text{solid})}$, and $[\text{MoS}_4]^{2-}$ in
786 $(\text{NH}_4)_2\text{MoS}_{4(\text{solid})}$.

787 Fig. 2. Evolution of the XANES spectra for all sample solutions as a function of temperature,
788 from 30 °C (blue line) to the maximum experimental temperature as labelled (red
789 line). The temperature increments for each sample are listed in Table 1.

790 Fig. 3. Fourier transforms of the EXAFS spectra at 30 °C (a) and 286 °C (b). The solid lines
791 are measured data and dashed lines are the fits using the parameters listed in the
792 Table 3. Also shown in (a) are the spectra of $[\text{MoO}_4]^{2-}$ moiety in $\text{Na}_2\text{MoO}_{4(\text{solid})}$, and
793 $[\text{MoS}_4]^{2-}$ in $(\text{NH}_4)_2\text{MoS}_{4(\text{solid})}$.

794 Fig. 4. Molybdenum K-edge EXAFS spectra in K-space (a) and Fourier transforms of
795 EXAFS (b) collected at 30 °C to the maximum temperature as labelled for all sample
796 solutions. The solid lines are measured data, and dashed lines are the fits using the
797 parameters listed in Table 3. The composition and temperature increment for each
798 sample are listed in Table 1.

799 Fig. 5. The change of ligand number derived from the EXAFS refinement as a function of
800 NaHS concentration at 30 °C (a) and 286 °C (b) and 800 bar.

801 Fig. 6. Snapshots of hydrogen molybdate ions and molybdate species, determined from the
802 MD simulations.

803 Fig. 7. Radial distribution function (RDF; solid lines) and coordination number (CN; dashed
804 lines) for simulation 5A, 5B, 5C, 5D, 5E and 8.

805 Fig. 8. Snapshots of thiomolybdates determined from the MD simulations (bond distances see
806 Table 3).

Fig. 9. The calculated XANES spectra using FDMNES and select experimental spectra. The XANES calculations are based on the molecular dynamic simulations of the structures for the species (see legend) at 77 and 300 °C. The Molybdenum K-edge (20,000 eV) was subtracted from the experimental energy for the x-axis of experimental spectra. The $[\text{MoO}_4]^{2-}$ standard is Na_2MoO_4 and $[\text{MoS}_4]^{2-}$ is $(\text{NH}_4)_2\text{MoS}_4$ solids.

Fig. 10. Predominant Mo species in sulfur-bearing solutions (total Mo = 1 ppm, total sulfur = 0.1 m) as a function of pH and oxygen fugacity at 25 (a, b) and 300 °C (c, d). The calculations are done using The Geochemist's Workbench (Bethke, 2007) and the default thermodynamic database. The presence of molybdenite is suppressed in (b) and (d) to show only the fields of aqueous Mo(VI) species. The solubility contour of molybdenite of 10 and 100 ppb are also shown in (c). Blue dashed lines show the boundaries between stability fields of sulfur species, and the greyed areas show the estimated pH ranges of the experimental solutions (Table 1). Also shown in Fig c and d are the positions of the pH buffers of $\text{CO}_2(\text{aq})/\text{HCO}_3^-$ and K-feldspar-Quartz-Muscovite (KMq, $a_{\text{K}^+} = 0.1$).