Oxetanes and Oxetenes: Fused-ring Derivatives

Nicolas Blanchard, Vincent Bizet, Nicolas Brach, Lénaïc Rummler, Krishna P Kaliappan

► To cite this version:

Nicolas Blanchard, Vincent Bizet, Nicolas Brach, Lénaïc Rummler, Krishna P Kaliappan. Oxetanes and Oxetenes: Fused-ring Derivatives. Reference Module in Chemistry, Molecular Sciences and Chemical Engineering, Elsevier, 2021, 10.1016/B978-0-12-818655-8.00150-5 . hal-03358269

HAL Id: hal-03358269

<https://hal.science/hal-03358269>

Submitted on 29 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

2.06

Oxetanes and Oxetenes: Fused-ring Derivatives

Authors:

Corresponding Author:

Nicolas Blanchard

Université de Haute-Alsace, Université de Strasbourg, CNRS, LIMA UMR 7042, Mulhouse, France

n.blanchard@unistra.fr

Co-Authors:

Vincent Bizet

Université de Haute-Alsace, Université de Strasbourg, CNRS, LIMA UMR 7042, Mulhouse, France

vbizet@unistra.fr

Nicolas Brach

Université de Haute-Alsace, Université de Strasbourg, CNRS, LIMA UMR 7042, Mulhouse, France

Nicolas.brach@etu.unistra.fr

Lénaïc Rummler

Université de Haute-Alsace, Université de Strasbourg, CNRS, LIMA UMR 7042, Mulhouse, France

Lenaic.rummler@etu.unistra.fr

Krishna P. Kaliappan

Department of Chemistry, Indian Institute of Technology Bombay, Powai, Mumbai 400076, India

kpk@chem.iitb.ac.in

Abstract

Fused and spiro-oxetanes, oxetenes and β -lactones are important classes of compounds that have found numerous applications in various areas of chemistry including medicinal chemistry and total synthesis of natural products. In this chapter, the main achievements from the theoretical and synthetic point of views since its last edition are discussed.

Keywords

Fused oxetane; Fused oxetene; Spiro-oxetane; Spiro-oxetene; β -Lactone; Spiro- β -Lactone; Heterocycles; Four-Membered rings.

2.06.1 Introduction

This chapter reviews developments related to fused and spiro-oxetanes, oxetenes and β -lactones from 2008 to mid-2020. Their physical properties, the methods for their synthesis, their reactivity trends and their applications to the preparation of biologically relevant compounds are discussed.

2.06.2 Theoretical Methods

A physical organic chemistry study of the photochemical generation of *ortho*-naphthoquinone methides **5** from **1** and their reactivity in aqueous solution was reported by Arumugam and Popik.¹ Naphthoxete **3** was proposed as intermediate in the formation of *ortho*-naphthoquinone methide by irradiation of 3-hydroxy-2-naphthalenemethanol **1**. The ring opening of naphthoxete **3** is a thermal reaction, slower than the decay of the excited state of **2a**. All structures were investigated at the (B3LYP/6-311++G(d,p)//B3LYP/6-31++G(d,p) and MP2/6-31++G(d,p)//FC-MP2/6-31++G(d,p) levels of theory.

<Scheme 1 near here>

1,8-Peribridged naphthalenes are highly strained structures that are often unstable although a few molecules have succumbed to X-ray diffraction. Novak reported ring strain energies (RSE) of series of single atom peribridged naphthalenes (SAPN) using high level quantum chemical methods. SAPN **6** has the highest RSE of 289.7 kJ/mol. A comparison was also proposed with compounds **3**, **7** and **8** possessing a fused-oxetene at different positions of a naphthalene core.²

<Scheme 2 near here>

Ground state and excited state potential energy surfaces of the photoexcitation of *ortho*-hydroxybenzaldehyde have been studied in silico by Migani.³ A potential product is an oxetene adduct arising from a decay at a twisted-pyramidalized intersection seam, although the barrier for its formation was higher than for other decay products. Oxetenes undergo conrotatory electrocyclic ring opening as demonstrated as early as 1971.⁴ In 2010, Venuvanalingam and Chattaraj reported a quantum chemical mechanical study at the B3LYP level with 6-31G(d) and 6-311+G(d,p) basis sets on this kind of perturbed pericyclic isomerizations.⁵ This study proposed hardness indices based on closely lying frontier molecular orbitals that reflect much better the relative reactivity of oxetene and its analogues (phosphetene, thiete and azete). Jayaprakash has reported an *ab initio* molecular orbital and DFT study of the electrocyclic ring-opening of oxetene in the gas phase.⁶ It was shown

that this pseudo-pericyclic process was characterized by a more planar transition state and an inferior barrier of activation compared to the electrocyclic ring-opening of cyclobutene.

Ruthenium-catalyzed decarbonylative couplings of aromatic aldehydes and terminal alkynes lead to alkenes, as demonstrated by Li in 2009.⁷ An intramolecular variant was also reported by Saá (using protic acid additives)⁸ and the mechanism of the latter transformation was explored through quantum mechanical calculations by Li.⁹ Recently, Li and Moitessier studied in depth the mechanism of this reaction through computational chemistry and isotope-labeling experiments. Zhang's fused oxetene intermediate was ruled out in favor of an aldol-type reaction of a ruthenium–vinylidene complex.¹⁰

<Scheme 3 near here>

A detailed study of the directing role of hydroxy functions in the Paternò-Büchi [2+2] photocycloaddition was reported by Abe in 2011.¹¹ Through a combination of experimental and theoretical studies, it was shown that cooperative hydrogen-bonding stabilization and steric interactions were crucial.

Fused oxetenes are intermediates in the Fe(III)-catalyzed cyclization of yne-acetals.¹² Yu and Li have reported a DFT investigation of the reaction mechanism using the B3LYP method and the 6-311+G** basis set, excluding [2+2] cycloaddition of yne-oxonium **13** in favor of cyclization of the vinylic carbocation **14** to the fused oxetenium **15** with an activation barrier of 17.2 kcal/mol. These alkyne-oxonium reactions are discussed in Section 2.06.6.6.

<Scheme 4 near here>

A parametric/DFT hybrid computational method DU8+ was recently applied to the analysis of published NMR data of naturally occurring oxetanes.¹³ A high rate of structure misassignment was found with more than 30 proposed structures requiring revisions, selected examples being highlighted in Section 2.06.11.

2.06.3 Experimental Structural Methods

2.06.3.1 Electron momentum spectroscopy

Electron momentum spectroscopy of oxetane was reported by Chen, leading to the binding energy spectra and the electron momentum distribution for the outer-valence molecular orbitals.^{14,15} It was shown that the ring-puckering motion of oxetane has a dramatic influence on the electron density distribution of the highest occupied molecular orbital.

2.06.3.2 Electronic and vibrational absorption spectroscopies

Ashfold reported the use of broadband transient electronic and vibrational absorption spectroscopies for an insightful study of the Paternò-Büchi reaction between benzaldehyde and cyclohexene.¹⁶ Quantum yields and rates constants for competing pathways were determined.

2.06.3.3 IR spectroscopy

Benzoxetone **19** was formed quite easily from 6-oxocyclohexa-2,4-dienylideneketene **18** by irradiation at 340 nm, **18** being prepared by photolysis of **17** at 315 nm. **19** can reverse to **18** upon photolysis at 315 nm. No differences were observed in the IR spectra of **18** and **19** for at least 6 photolysis cycles.¹⁷

<Scheme 5 near here>

Benzoxete chemistry has been reviewed by Meier and the reader is redirected to this excellent article for details related to their isolation and reactivity.¹⁸ Many erroneous or questionable benzoxetes are also discussed. Other reviews were also recently published.¹⁹ 2*H*-Benz[*b*]oxetes were first isolated by Tomioka and Matsushita in 1997, by irradiation of *ortho*-hydroxyphenyldiazomethane (at $\lambda > 300$ nm) or benzofuranone (at $\lambda > 254$ nm) in an argon matrix at 10 K.²⁰ These highly strained compounds (38 kcal/mol)²¹ are in valence equilibrium with their *ortho*-quinoid structures, the latter being much more favored from a thermodynamic point of view. Theoretical and experimental IR spectrum of benzoxetes were compared and found in very good agreement. Benzoxete was also detected by irradiation of a vapor deposition prepared sample of 2-

hydroxymethylphenol at 254 nm and 14 K. The characteristic signal at $\nu = 1668\text{ cm}^{-1}$ (scissoring vibration of the CH_2 motif) was clearly observed.²²

2.06.3.4 X-ray crystallography

X-ray crystallographic studies on a range of fused or spiro-oxetanes and β -lactones have been reported over the period of interest to this chapter, as in the cases of the natural products hawaiiinolide E,²³ hawaiiinolide A and C,²⁴ aphanalide A and C,²⁵ and rubesanolide A.²⁶ X-ray structures of various synthetic compounds have also been reported²⁷ including fused oxetanes resulting from the photocycloaddition of electronically excited carbonyl to isoxazoles (e.g. **20**),²⁸ a planar oxetane arising from the Paternò–Büchi reaction of triplet benzophenone with 4-aryl-4*H*-pyrans,²⁹ an unprecedented tricyclic 3*H*-oxeto[2,3,4-*ij*]isochromene featuring a rare oxetane acetal reported in 2017 during the course of synthetic investigations of sigillin A, an insect repellent natural product isolated from springtails.³⁰ Spirocyclohexene-oxetanes were prepared through catalytic enantioselective cycloaddition to α,β,β -trisubstituted nitroalkenes, and a single structure **21** was reported, allowing the assignment of absolute configuration.³¹ Spiro-oxetane embedded in heterocyclic structures have been disclosed recently.³² Structures of spiro- β -lactones prepared via the [2+2] cycloaddition of disubstituted ketenes with isatins, such as **22**, were also characterized by X-ray crystallography.³³

<Scheme 6 near here>

2.06.4 Thermodynamic Aspects

Oxetanes are prone to ring-opening³⁴ under acidic conditions at similar rates than epoxides,³⁵ an experimental observation that can be rationalized by their similar ring-strain (106 kJ/mol for oxetanes vs. 112 kJ/mol for epoxides).^{36,37} On the other hand, oxetanes are hydrolyzed at slower rates than epoxides under basic conditions. Theoretical studies have addressed this reduction in hydrolysis rates.³⁸⁻⁴² Strain energies and equilibrium geometries of 2,4-oxytwistane **23** were computed *ab initio* using the MP2(fc)/6-31G(d) theoretical model and compared with oxetane.⁴³ The ring-strain energies (RSEs) of spiroepoxy- β -lactone **24** and spiroepoxy-oxetane **25** have been calculated from the energies of fully-optimized structures at the MP2 + ZP/cc-pVDZ level. The stability/reactivity of spiroepoxy- β -lactone **24** has been discussed and found to not correlate with RSE.⁴⁴

Gas-phase enthalpies of formation of oxetane **26** (-80 kJ/mol) and oxetene **10** (-52.8 kJ/mol) were determined by high-level quantum chemical calculations at the G3(MP2)//B3LYP level by Taskinen in 2009.⁴⁵ Closely related results were reported by Green in 2012 using an application of a QCISD(T) quantum chemistry strategy.⁴⁶

<Scheme 7 near here>

2.06.5 Reactivity of Fully Conjugated Rings

No examples of this class of compounds have been found in the literature in the period covered.

2.06.6 Reactivity of Nonconjugated Rings

2.06.6.1 Reaction with Nucleophiles

As shown by Carreira and Roger-Evans,⁴⁷⁻⁵⁰ oxetanes are relevant bioisosters for *gem*-dimethyl substituents or carbonyl groups. α -Functionalization of oxetan-3-ones can be conveniently performed using Enders' SAMP/RAMP synthetic methodology⁵¹ as shown by Shipman.⁵² Hydrazone **28** was obtained in 68% yield and could be further hydrolyzed to the fused oxetane **29** in 60% yield and 84% enantiomeric excess.

<Scheme 8 near here>

Bode introduced the SnAP (stannyl (Sn) amine protocol) concept for the rapid synthesis of diversely substituted heterocycles from stannyl methylether and aldehydes or ketones.⁵³⁻⁵⁵ SnAP reagent **30** serves as a cross-coupling partner with 3-quinolinecarboxaldehyde **31**, leading to **32** in 56% yield.⁵⁶

<Scheme 9 near here>

Spiro-oxetanes can be obtained using classical Pictet-Spengler cyclization of L-tryptophan ethyl ester **33** and oxetan-3-one **34** using iodine in hot acetonitrile. Spiro-oxetane **35** was obtained in 76% as a mixture of two diastereomers (dr = 88:12).⁵²

<Scheme 10 near here>

Fused N,O-acetals of oxetan-3-one, such as **36**, can be transformed into a myriad of interesting heterocyclic scaffolds via the addition of carbon-based nucleophiles. As an example, Carreira showed that potassium trifluoroorganoborates were competent partners under Lewis acid catalysis leading to oxetanes such as **37** containing a quaternary center.⁵⁷ The pendant alcohol function can then cyclize onto the oxetane, leading to morpholines (e.g. **38**) by ring-expansion. Introduction of cyanide as a nucleophile is also possible, using trimethylsilyl cyanide under indium triflate catalysis in acetonitrile.⁵⁸

<Scheme 11 near here>

Spirocyclohexene-oxetanes were obtained by Jørgensen via an impressive catalytic enantioselective [4+2] cycloaddition of α,β,β -trisubstituted nitroolefin **40** with $\alpha,\beta,\gamma,\delta$ -unsaturated aldehyde **39**.³¹ The reaction proceeds through trienamine organocatalysis⁵⁹ and delivers a broad range of spirocyclic oxetanes, such as **41**, containing two contiguous stereogenic centers including a quaternary one. It should be pointed out that the exact mechanism is still a matter of debate, whether being a pericyclic or a stepwise process.^{59,60}

<Scheme 12 near here>

Intramolecular [4+2] cycloaddition reactions of sydnone **44** have been reported by Harrity as an efficient tool to prepare spiro-oxetane containing fused pyrazoles such as compound **45** obtained in 93% yield. Cycloaddition precursor **44** itself is prepared in a straightforward sequence from sydnone **42** and 3-oxetanone **43**.⁶¹

<Scheme 13 near here>

4- π -Photocyclization of 2-pyrone, a process reported for the first time in 1964, leads to a valuable bicyclic β -lactone⁶² whose chemistry has been reviewed early 2020 by Coote.⁶³ Selected examples over the 2008-2020 period demonstrate the versatility of this building block for the straightforward synthesis of relevant cyclobutenes or cyclobutanes.⁶⁴⁻⁷²

2.06.6.2 Elimination

Elimination of spiro-oxetanes is a facile process that has been central to the synthesis of homoallylic alcohols. Catalytic processes have also been reported, such as the tris(pentafluorophenyl)borane-catalyzed isomerization of 2,2-disubstituted oxetanes **46** or **48**. Under very mild conditions (0.5 mol% of Lewis acid, room temperature, short reaction times), good yields of the desired homoallylic products **47** and **49** were obtained.⁷³

<Scheme 14 near here>

2.06.6.3 Ring Expansion

Ghosh reported the synthesis of high-affinity P2 ligand for HIV-1 protease inhibitors such as (—)-**52** using a short synthetic sequence relying on the Paternò-Büchi photoadduct (±)-**50**. Hydrogenation of the latter followed by an acid-catalyzed transacetalization using *para*-toluenesulfonic acid in warm methanol delivered (±)-**51** in good yield.⁷⁴

<Scheme 15 near here>

3-Oxetanone **43** has been introduced as a versatile building block for medicinal chemistry. The corresponding amins, hemiaminals and hemithioacetals **53** were found to react with trimethylsilyl cyanide and a catalytic amount of indium(III) triflate in acetonitrile at room temperature in a smooth ring-expansion process, going through postulated intermediate **54**. The functional group tolerance of this cascade reaction is quite large and morpholines, thiomorpholines and piperazines **55** were obtained in moderate to excellent yields and diastereoselective ratios.⁵⁸

<Scheme 16 near here>

2.06.6.4 Reduction

Photoadduct **56**, arising from a Paternò-Büchi reaction between benzaldehyde and trimethylsilyloxycyclopentene, was found to undergo clean reduction to the corresponding tertiary alcohol **57** or a regioselective rearrangement to cyclohexanone **58** via an aluminum(III)-mediated pinacol type process.⁷⁵

<Scheme 17 near here>

One step reduction of β -lactones **59** to oxetanes **60** has not been reported in the literature, which prompted the development of a 2-step sequence implying reduction to the diol followed by cyclization under basic conditions.⁷⁶ The first step of the process turned out to be quite difficult, leading to decomposition via retro-aldol reactions. Finally, diisobutylaluminium hydride in dichloromethane proved to be the best reducing agent after an extensive screening. Cyclization

reaction of the diol could be efficiently promoted by tosyl chloride and potassium *tert*-butoxide in refluxing tetrahydrofuran.

<Scheme 18 near here>

2.06.6.5 Cycloreversion

Cycloreversion of oxetanes has been comprehensively reviewed in 2019.⁷⁷ Decarboxylation of β -lactone⁷⁸ is a fundamental process in organic synthesis that has been exploited by Chi in 2017. Fused β -lactone **61** could be transformed into the corresponding epoxide **62** without any loss in enantiopurity, via a decarboxylation in DMF at $110\text{ }^\circ\text{C}$ followed by epoxidation with *m*-CPBA.⁷⁹

<Scheme 19 near here>

2.06.6.6 Alkyne-Carbonyl Metathesis

Carbonyl-alkyne metathesis reactions are powerful tools to build in a straightforward manner α,β -unsaturated carbonyls through the cycloreversion of oxetenes intermediates.⁸⁰⁻⁸⁵ As evocated before, metal-catalyzed cyclization reactions of ω -alkynyl carbonyls constitute efficient transformations for the synthesis of α,β -unsaturated ketones or aldehydes. As an example, Yamamoto⁸⁶ reported the gold-catalyzed carbocyclization⁸⁷ of a series of internal ketones **63** that delivered enones **66** in good to excellent yields. The fused carbocationic oxetane **65**, arising from the π -activation of the alkyne followed by cyclization of the carbonyl, is postulated as an intermediate.

<Scheme 20 near here>

Yu and Li reported a Fe(III) -catalyzed cyclisation of yne-acetal **67** that proceeds via fused oxetanium **70**.¹² Ring opening of the latter followed by hydrolysis of the oxocarbenium leads to α,β -unsaturated ketones **68**. A dramatic solvent effect was found, and nitromethane or acetone delivered the highest yields. The alkynyl motif could be substituted by aromatic, alkyl, vinyl or alkynyl groups with the same efficiency. Intermolecular versions are also possible as exemplified by the Fe(III) -catalyzed transformation of alkyne **72** into α,β -unsaturated ketone **74**.

<Scheme 21 near here>

Saá reported the easy preparation of hydroazulenones **76** and **77** from the corresponding acyclic enyne acetals **75** using tetrafluoroboric acid in dichloromethane at 0°C .⁸⁸ The scope of the transformation is quite large and the reaction is postulated to go through formation of oxonium **78**, followed by alkyne-oxonium metathesis. The fused O-substituted oxetanium **79** would then undergo an electrocyclic ring-opening to the divinyl ketone **80**. Subsequent Nazarov electrocyclization and proton elimination would then deliver the observed products. A dramatic influence of the starting material alkene substitution on the regioselective course of the process was found.

<Scheme 22 near here>

Silylated fused oxeteniums **85** were postulated as precursors to α,β -unsaturated ketones en route to 3,4-disubstituted pyrroles **83**.⁸⁹ Initial semi-pinacol rearrangement leads to a ring-contracted intermediate **84** which undergoes an alkyne-ketone metathesis yielding the substituted pyrrole after spontaneous oxidation.

<Scheme 23 near here>

Fused oxetenium ions are intermediates in the synthesis of medium and large ring lactones from acetals and silylated ynol ethers. Sun reported that treating acetal **86** with a Lewis acid in the presence of 2,4,6-collidine led to the formation of oxocarbenium such as **91** that underwent a spontaneous electrocyclic ring-opening to **88**.⁹⁰ This unusual and elegant ring expansion strategy overcomes the current limitations encountered for the preparation of medium to large ring lactones, whether entropic or enthalpic by nature.

<Scheme 24 near here>

Spiro-oxetenes have been proposed as intermediates in the synthesis of 3-substituted oxindoles **97**.⁹¹ Indeed, copper(I)-catalyzed reaction of phenylacetylene **95**, tosylazide **96** and isatin **94** led to a metal ynamide **98** which could undergo a stepwise cyclization to spiro-oxetene intermediate **99**. Ring opening of the latter finally afforded product **97** in excellent yield.

<Scheme 25 near here>

The inherent strain of benzoxetenes has been judiciously exploited by Miyabe who reported the synthesis of *ortho*-disubstituted aromatics such as **101** from an aryne precursor **100**, a formamide and diphenylzinc.⁹² Transient formation of benzoxetene **103**, arising from reaction of benzyne **102** and DMF, undergoes a ring opening reaction to **104** which is immediately trapped *in situ* by diphenylzinc.

<Scheme 26 near here>

Diels-Alder cycloadditions of the intermediate *ortho*-quinone methide similar to **104** are also possible as shown by Yoshida.⁹³ When the aryne precursor **105** is reacted with active methylene derivatives such as **106** in the presence of potassium fluoride, coumarins **107** are obtained in moderate to good yields.

<Scheme 27 near here>

Ortho-quinomethide **108** could also act as a nucleophile by delocalization of the lone pair of electrons of the nitrogen atom, and then react with an electrophile. Lu and Wang have demonstrated that benzoyl cyanide **110** was a relevant electrophile, leading to the formation of α -amino carbonitriles **112** via an elegant three-component reaction.⁹⁴

<Scheme 28 near here>

2.06.6.7 Cycloisomerisation of Oxetane-Containing Enynes

1 α ,25-Dihydroxyvitamin D₃, a *seco*-steroid having a key role in the regulation of calcium homeostasis, has been structurally edited to include a spiro-oxetane at C3 position.⁹⁵⁻⁹⁷ The latter compound **115** is conveniently obtained via a Trost palladium-catalyzed cycloisomerisation⁹⁸ of enyne **113** in the presence of vinyl bromide **114** in 66% yield. It was shown that the introduction of a spiro-oxetane modified the conformation of **115** compared to the parent 1 α ,25-dihydroxyvitamin D₃, and thus favored the formation of ligand-vitamin D receptor complex.

<Scheme 29 near here>

2.06.7 Reactivity of Substituents Attached to Ring Carbon Atoms

2.06.7.1 Exocyclic Alkenes

1,3-Dipolar cycloaddition of azomethine ylide derived from **116** with *exo*-methylene oxetanes **40b,c** leads to spiro-oxetanyl pyrrolidines **117a,b** in excellent yields. This copper(I)-catalyzed process relies on a planar-chiral ferrocene P,N-ligand that allows for a perfect control of *exo*-selectivity and high enantiomeric excesses (94-99%).⁹⁹ Silver-catalyzed processes were also reported.¹⁰⁰

<Scheme 30 near here>

2.06.7.2 Exocyclic Epoxides and Cyclopropanes

2-Methyleneoxetane **118** could be converted into the corresponding epoxide **119** with good diastereoselectivity (8:1) and quantitative yield using anhydrous dimethyldioxirane (DMDO). Epoxide-opening was best conducted with diisobutylaluminum hydride in dichloromethane, leading to **120** in 56% yield as a 3:1 mixture of diastereomers.¹⁰¹

<Scheme 31 near here>

Spirocyclopropyl oxetanes **121** were reported to rearrange to 3-methylenetetrahydrofurans **122** upon treatment with a catalytic amount of Zeise's dimer and 20 mol% of tricyclohexylphosphine in dichloromethane at 45 °C.¹⁰² The mechanism of this new platinum-catalyzed rearrangement was extensively studied (including with ¹³C NMR-labeling) and it was found that oxidative addition of the substrate to platinum(II) was taking place at the least substituted carbon-carbon bond of the spirocyclopropyl unit.

<Scheme 32 near here>

Spirocyclic oxetanes **123**, obtained by photocycloaddition of *N*-acylisatins and bicyclopropylidene, could be ring-opened by an excess of acid, such as hydrochloric or hydrobromic acid in acetonitrile at room temperature, yielding spirocyclopropyl butenolide **124** in good yield.¹⁰³ On the other hand, γ -butyrolactones **125** possessing the two spirocyclopropane units were obtained upon action of hydroiodic acid and iodine.

<Scheme 33 near here>

2.06.7.3 Reactivity of Exocyclic Radicals

Mono-debromination of spirooxetane **126** was performed with tributyltin hydride in toluene at room temperature. A single diastereomer **127** was obtained, although the yield was only 21%.¹⁰⁴

<Scheme 34 near here>

2.06.8 Reactivity of Substituents Attached to Ring Heteroatoms

No examples of this class of compounds have been found in the literature in the period covered.

2.06.9 Ring Synthesis Classified by the Number of Ring Atoms

2.06.9.1 Intermolecular [2+2] Cycloaddition Reactions

Paternò-Büchi [2+2] photocycloaddition^{77,105} is one of the most direct route to oxetanes and this well-established transformation can be transposed to flow conditions, as demonstrated in the synthesis of the cytotoxic lactone (+)-goniofufurone **132**, isolated from *Goniothalamus* trees.¹⁰⁶ Thanks to fluorinated ethylene polymer (FEP) continuous flow reactors in conjunction with a mercury medium-pressure lamp (400 W), the [2+2] reaction of **128** with benzaldehyde could be upscaled to deliver 42 g of photocycloadducts **129** and **130** (97% combined yield).

<Scheme 35 near here>

The Paternò-Büchi [2+2] photocycloaddition of oxindoles has also been reported using a visible-light microfluidic photoreactor.¹⁰⁷ Optimization of photocycloadditions under flow conditions can also be conducted in an automatic fashion using self-optimizing algorithms. Rueping demonstrated that the [2+2] photocyclization of benzophenone and furan in a capillary-type continuous photo-flow reactor could indeed be optimized without human intervention.¹⁰⁸

Paternò-Büchi [2+2] photocycloaddition proceeds via direct carbonyl excitation or photosensitization followed by alkene interception as in Scheme 35. In this regard, alkyl substituted ketones are not the most reactive partners and require more energetic irradiation and specialized glassware. However, it was demonstrated in 2019 by Schmidt that a copper(I)-norbornene complex was able to catalyze the [2+2] carbonyl-olefin photocycloaddition with a broad range of alkyl ketones.¹⁰⁹ A relevant example is described in Scheme 36 with the reaction of ketone **133** and norbornene **134**. In the presence of 10 mol% of copper(I) tri(1*H*-pyrazol-1-yl)hydroborate (TpCu) and under irradiation with a 100 W mercury lamp, a good yield of adduct **135** is obtained. The reaction pathway is mechanistically distinct from classical Paternò-Büchi reactions and relies on the excitation of an *in situ* formed TpCu(Norb) complex.

<Scheme 36 near here>

Unprecedented angular tricyclic 4:4:4 oxetane-containing compounds were obtained by Aitken through a triple photochemical process, starting from 4-substituted cyclopentenones **136** and alkenes.¹¹⁰ Under irradiation at 300 nm for 40 h in acetonitrile, 12 different angular tricyclic oxetanes were obtained with moderated to good diastereoselectivities. A plausible mechanism calls for an initial [2+2] photocycloaddition followed by a Norrish-I/ γ -H transfer. A final intramolecular Paternò-Büchi reaction would then deliver **137**.

<Scheme 37 near here>

Photocycloadditions of isoquinoline-1,3,4-trione with 5-methoxyoxazoles were reported by Zhang. Fused oxetanes were obtained in very good yields and could be rearranged to spiroisoquinoline-oxazolines under acidic conditions.¹¹¹

2.06.9.2 Intramolecular Paternò-Büchi Cycloadditions

Transposed Paternò-Büchi [2+2] photocycloadditions of atropoisomeric enamides **141** have been reported by Sunoj and Sivaguru, during which a switch in excited states from $n-\pi^*$ of the carbonyl group to $\pi-\pi^*$ of the alkene occurs.¹¹² After cycloaddition under sensitized conditions, chiral oxetanes **142** were obtained.

<Scheme 38 near here>

2.06.9.3 Norrish II (Biradical) Cyclization

Photoreaction of 2-isopropoxy naphthoquinone **143** in the presence of bicyclopropylidene leads to the expected cyclobutane **144** after 3 hours. However, upon extended irradiation time, a γ -hydrogen abstraction occurs from the triplet state of alkoxyketone **144** leading to the fused-oxetane **145** in 54% yield.¹¹³

<Scheme 39 near here>

2.06.9.4 Cycloadditions of Ketene Equivalents

Bode reported in 2009 that chiral triazolium-derived NHCs were able to catalyze the formation of fused β -lactones with an excellent enantioselectivity via the intermolecular reaction of an enal **147** and an α -hydroxyketone **148**.¹¹⁴ Quite surprisingly, the use of chiral imidazolium-derived NHC led to a different fused lactone, also with excellent enantioselectivity. A detailed mechanistic discussion proposes that the different reactivities of the intermediate acylazolium species involved in the lactone-forming step, such as **153** derived from catalyst **150**, might be responsible for this outcome.

<Scheme 40 near here>

Related NHC-catalyzed synthesis of fused β -lactones from enals and 2-enolpyridines¹¹⁵ or oxindolyl β,γ -unsaturated α -ketoesters¹¹⁶ were reported. Scheidt reported the first NHC-catalyzed dynamic kinetic resolution (DKR) of β -ketoesters **154** substituted in the α -position, leading to fused β -lactones.^{117,118} When applied to electron-rich ketones (such as **156**), this DKR strategy is followed by a spontaneous decarboxylation process that afforded cyclopentene **158**.¹¹⁹ This concerted asynchronous *retro*-[2+2] reaction was studied using quantum mechanical computations and it was found that the *anti*-selective process was favored based on steric interactions between the carboxylate and the α -substituent in the *syn*-transition state.

<Scheme 41 near here>

Chiral NHCs developed by Lupton were shown to catalyze the formation of β -lactones fused to cyclopentanes **161**, starting from α,β -unsaturated acyl fluorides **159** and donor-acceptor cyclopropanes **160**.^{120,121} The process is general and R^1 could be an aryl, alkyl, alkenyl group whereas R^2 could be alkyl or cycloalkyl. Good yields and enantioselectivities up to 97% were obtained for compounds **161**. Mechanistically, two pathways were proposed based on the different modes of addition of enolate **162** onto intermediate **163**.

<Scheme 42 near here>

Besides donor-acceptor cyclopropanes, β -diketones, β -ketoesters and malonates (such as compound **165**) proved to be competent partners in oxidative carbene catalysis, leading to fused β -lactones **166**. An interesting cooperative Lewis acid effect was observed, yields and enantioselectivities being optimal in the presence of 50 mol% of lithium chloride.¹²²

<Scheme 43 near here>

Chi reported that bicyclic pyrrolidine-fused β -lactones **169** could be obtained with excellent yields and stereoselectivities from aldehydes **147** and amino esters **167** via a chiral NHC-catalyzed reaction that created the desired β -lactone as well as a new carbon–nitrogen bond in β -position relative to the carbonyl.⁷⁹ The proposed mechanism relies on the generation of an acyl-azolium intermediate **170** that undergoes an intermolecular Michael addition of the sulfonamide **167** followed by aldol/ β -lactonization to **169**.

<Scheme 44 near here>

α,β-Unsaturated acylammonium intermediates are also competent partners as shown by Romo in 2013¹²³ who proposed a multicomponent organocascade reaction coined Nucleophile-Catalyzed Michael-Aldol-β-Lactonization (NCMAL), which led to the synthesis of β-lactone **174** in excellent yields and enantioselectivities.

<Scheme 45 near here>

2.06.9.5 Intramolecular Nucleophilic Displacements

Brown reported a novel palladium-catalyzed deallylative cyclization of **175** to synthesize β-lactone rings **176**, an elegant strategy that was applied to the total synthesis of (±)-vibrallactone, a natural product isolated from the culture of Basidiomycete *Borestereum vibrans*.¹²⁴ Palladium-catalyzed deprotection of β-mesyloxy ester **175** led to the desired β-lactone **176** in only 2 minutes in 95% yield. Although this reaction does not strictly fall into the area covered by this chapter focused on fused β-lactones, it would be interesting to study the formation of bicyclic β-lactones via this novel type of intramolecular cyclization.

<Scheme 46 near here>

Bull reported the two-step preparation of diversely substituted oxetanes **180** from functionalized diazo-compounds **177** by rhodium-catalyzed O–H insertion followed by a base-mediated cyclization of intermediate **179**. Several fused [3.2.0] ring systems **180** were obtained, in addition to an oxabicyclo[4.2.0]octane, in excellent yields.^{125,126}

<Scheme 47 near here>

Benzoxetes **182** were efficiently prepared by Qin through a sulfonyl fluoride-mediated conversion of 2-hydroxy acetophenones **181** in the presence of potassium carbonate in DMSO at 90 °C.¹²⁷

<Scheme 48 near here>

Researchers at Merck reported an optimized synthetic route towards spiro-oxetane containing piperidines and pyrrolidines via a simple intramolecular $\text{S}_{\text{N}}2$ -type displacement of a primary tosylate.^{128,129} The latter is prepared *in situ* from diol **183** using *n*-butyllithium and tosyl chloride; the addition of a second equivalent of *n*-butyllithium then delivered the desired spiro-oxetane piperidine **185** in 66% yield on a 37 g scale. Similar strategy led to compounds **186** and **187** from ethyl nipecotate and methyl 1-benzylpyrrolidine-3-carboxylate (respectively).

<Scheme 49 near here>

2.06.9.6 Dehydration of Hydroxy Acids

Bis(2-oxo-3-oxazolidinyl)phosphinic chloride (BOPCI) proved to be a reliable and efficient reagent for the synthesis of fused β -lactones within densely functionalized scaffolds. It has been applied to the

ring closure of numerous β -hydroxyacids in the context of total synthesis of natural products, such as (+)-lactacystin,¹³⁰ a rocaglate-derived β -lactone,¹³¹ (-)-7-methylomuralide,¹³² (-)-omuralide,¹³³ or salinosporamide A¹³⁴⁻¹³⁶ to cite only a few. As an example, Burton reported a high-yielding sequence from β -hydroxyester **188** to salinosporamide A in 3 steps (61% overall), using BOPCl as the key coupling agent.¹³⁷ As noted in previous versions of this chapter, this reaction leads to the fused β -lactone only and no traces of spiro- β -lactone (arising from the cyclization of the exocyclic secondary alcohol) were detected.

<Scheme 50 near here>

2.06.9.7 Electrophilic Cyclizations onto Alkenes

Bromolactonization of β,γ -unsaturated carboxylic acid is a reliable and simple method to synthesize β -lactones, especially on large scale. Millenium Pharmaceuticals has reported a practical synthesis of Pevonedistat, a NEDD8-activating enzyme (NAE) inhibitor using the bromolactonization of **190** to bicyclic intermediate **191** as one of the key steps.¹³⁸

<Scheme 51 near here>

Acetal-substituted spiro-oxetanes **193** having a 1,5-dioxaspiro[3.4]octane scaffold were prepared through a bromocation-induced cyclization of bis-homopropargylalcohols **192** with pyridinium tribromide.¹⁰⁴

<Scheme 52 near here>

2.06.9.8 Cationic Cyclizations

Oxidative dearomatization of phenols is an interesting class of reactions that is relevant to the formation of fused oxetane, as shown for the first time by Li in 2014.¹³⁹ Treatment of substituted phenol **194** by $\text{PhI}(\text{OAc})_2$ in aqueous acetonitrile led to the strained oxetane **195** in moderate yield.

<Scheme 53 near here>

2.06.9.9 Intramolecular C-H Insertion

Moody reported a straightforward preparation of spiro- β -lactones **197** based on the selective 1,4-C-H insertion reactions of rhodium carbenes derived from α -diaoesters **196**.¹⁴⁰ The latter could be generated in flow or in classical batch chemistry. The obtained β -lactones could then be reduced to oxetanes using a two-steps protocol discussed in Section 2.06.6.3.

<Scheme 54 near here>

Metalloporphyrin catalysts are also efficient for C-H functionalization reactions, as shown by Che. For example, the methylated iridium(III) porphyrin complex $[\text{Ir}(\text{TTP})\text{Me}(\text{L})]$ (H_2TTP = *meso*-tetrakis(*p*-tolyl)porphyrin; L = H_2O or solvent) was able to catalyze the formation of spiro- β -lactone **197f** from α -diaoester **196f** at room temperature in only 1 h.¹⁴¹

<Scheme 55 near here>

2.06.10 Ring Synthesis by Transformation of Another Ring

2.06.10.1 Rearrangement of Epoxides

Highly strained fused oxetane, such as the one embedded in merrilactone A **199**, a natural product isolated from the pericarps of *Illicium merrillianum* in 2000, has been prepared by the Fukuyama procedure¹⁴² involving a stereoselective epoxidation of the trisubstituted alkene of **198** followed by a homo-Payne rearrangement.^{143,144} This classical reaction has been used in several syntheses of the neurotrophic active natural product.¹⁴⁵

<Scheme 56 near here>

2.06.10.1 Carbonylation of Epoxides

Enantioenriched fused β -lactones **201** can be prepared via an asymmetric ring-expansion/carbonylation reaction of *meso*-epoxides **200**. Combination of catalytic amounts of a chiral chromium(III) complex and dicobalt octacarbonyl proved to be optimal to promote this important class of reactions.¹⁴⁶

<Scheme 57 near here>

Spiro-epoxides such as **202a** and **202b** were shown to react with high regioselectivity under carbon monoxide atmosphere (900 psi) in diisopropylether at room temperature in presence of an octaethylporphyrin catalyst. Mechanistic investigations demonstrated that the rate-limiting step of the reaction was the epoxide ring-opening, leading to **203a,b** as the final products.¹⁴⁷

<Scheme 58 near here>

2.06.10.3 Rearrangement of Diazoketones

A rare example of Wolff rearrangement of a rhodium carbene was reported by Chiu in 2017, leading to the fused β -lactone **205** in 39% yield alongside two other products. It was hypothesized that the conformation of the [4.4.0] bicyclic diazoketone **204** could be responsible for this specific outcome.¹⁴⁸

<Scheme 59 near here>

2.06.11 Important Compounds and Applications

Oxetanes are found in more than 600 naturally occurring structures produced by microorganisms or plants.¹⁴⁹

2.06.11.1 (-)-Stemoamine

β -Lactones can be transient intermediates en route to cyclopentenones via a tandem sequence of ynolate addition onto a ketone followed by intramolecular condensation onto an ester and decarboxylation. Such a sequence was used by Shindo in 2018 in the total synthesis of (-)-stemoamine, an alkaloid isolated from the root of *Stemona japonica* Miq. that has been used in traditional Chinese medicine.¹⁵⁰ Addition of ynolate **208** (generated from **207** and ^tBuLi) to ketone **206** led to the formation of **211**, that can undergo silica gel-assisted decarboxylation in refluxing benzene.¹⁵¹ The very efficient sequence forms the A-ring of (-)-stemoamine.

<Scheme 60 near here>

2.06.11.2 *ent*-Dichrocephone B

Fully substituted oxetanes are found in natural products such as dichrocephone B **217**. In 2018, the total synthesis and structure revision of this compound was reported by Christmann and Tantillo, featuring a Lewis acid-mediated cyclization of *ent*-dichrocephone A **216** into *ent*-dichrocephone B **217** in 88% yield. A low yield (12%) was obtained for the acid-mediated cyclization of **215**.¹⁵²

<Scheme 61 near here>

2.06.11.3 (±)-Spongiolactone

Fused β -lactones are obtained through an elegant procedure reported by Romo starting from aldehydic acids.^{153,154} Intramolecular Nucleophile-Catalyzed Aldol Lactonization (NCAL) reactions delivers β -lactones with excellent functional group tolerance; chiral amines were also shown to induce high enantioselectivity to the process¹⁵³ and applications to the total synthesis of natural products were disclosed, as in the (±)-spongiolactone case (Scheme 60). Extension to the dynamic kinetic resolution of keto-acids was also reported.¹⁵⁵

<Scheme 62 near here>

2.06.11.4 (-)-Curcumanolide A and (-)-Curcumalactone

Desymmetrization of diones was shown to deliver fused β-lactone **222** with excellent enantiomeric excesses via an asymmetric nucleophile (Lewis base)-catalyzed aldol lactonization of **221**. Baeyer-Villiger oxidation of the tricyclic lactone followed by dyotropic 1,2-acyl migration led to bislactone **224**, a common intermediate in the synthesis of (-)-curcumanolide A and (-)-curcumalactone.¹⁵⁶⁻¹⁵⁸

<Scheme 63 near here>

A similar strategy was used by Romo for several other natural products, such as (±)-salinosporamide A,¹⁵⁹ (±)-cinnabaramide A¹⁶⁰ and (+)-omphadiol.¹⁶¹

<Scheme 64 near here>

2.06.11.5 (+)-Dictyoxetane

(+)-Dictyoxetane was isolated from a brown alga *Dictyota dichotoma* (Indian ocean) in 1985 and was found to possess a unique oxetane motif embedded in a 2,7-dioxatricyclo[4.2.1.0^{3,8}]nonane ring system.¹⁶² Magauer reported the very first synthesis of (+)-dictyoxetane in 2016.^{163,164} The challenging oxetane in **226** was constructed through the intramolecular displacement of an allylic mesylate, generated from **225**, under basic conditions in 88% yield.

<Scheme 65 near here>

2.06.11.6 (+)-Pleuromutilin

Carbonyl-alkyne metathesis reactions⁸⁰⁻⁸⁵ have found numerous applications in the context of total synthesis of natural products and only a few examples will be discussed here. During the course of investigations aiming at the total synthesis of (+)-pleuromutilin, Herzon reported the unexpected formation of the transient intermediate fused oxetene **230** from the oxidative cyclization/C-O reductive elimination of yne-aldehyde **227** using Ni(COD)₂ (40 mol%), 4,5-dichloro-1,3-bis(2,6-diisopropylphenyl)-1,3-dihydro-2*H*-imidazol-2-ylidene (40 mol%) and triisopropylsilane as a reductant.¹⁶⁵

<Scheme 66 near here>

2.06.11.7 Phomopsidone A, Artoheterophyllin E, Morusalbin B, Fusariumin D

Shen reported the isolation of phomopsidone A in 2014 from the mangrove endophytic fungus *Phomopsis* sp. A123.¹⁶⁶ The same year, artoheterophyllin E was isolated from the wood of *Artocarpus heterophyllus*, an evergreen fruit tree collected in Hainan province (PR China).¹⁶⁷ Although extensive NMR and mass spectrometry experiments were performed and discussed, the structure of these compounds are questionable in light of the well-known instability of benzoxetes that are in equilibrium with their thermodynamically favored *ortho*-quinoïds valence isomers. Indeed, phomopsidone A was recently suggested to possess excelsione's structure.¹³

<Scheme 67 near here>

Morusalbin B was extracted from the root bark of *M. alba* L. (Moraceae) in the Republic of Korea. Its quite complex structure is proposed to possess a fused benzoxete.¹⁶⁸ Recently, Zhang and Wang extracted from the fermentation broth of *Fusarium oxysporum* ZYP-R1 two new antimicrobial agents, among which Fusariumin D.¹⁶⁹ The structure of the latter was recently proposed to actually be a 4-pyranone scaffold.¹³

<Scheme 68 near here>

2.06.11.8 (-)-Mitrephorone A

Mitrephorone A is a member of the family of *ent*-trachylobanes which was isolated in 2005 from *Mitrephora glabra*, a Bornean shrub. Its structure features a highly congested hexacycle with an embedded oxetane. The central oxetane of mitrephorone A was prepared thanks to an impressive oxidative 4-*exo-trig* cyclization of a hydroxy enol ketone, derived from **231**, activated with Koser's reagent.¹⁷⁰ An alternative synthetic blueprint was reported a year later,¹⁷¹ featuring a direct conversion of mitrephorone B to mitrephorone A via a bioinspired C-H oxidation with White-Chen catalyst. Electrochemical oxidation or photocatalyzed formation of the oxetane lead to lower yields.

<Scheme 69 near here>

Acknowledgments

The authors thank the Université de Strasbourg, the Université de Haute-Alsace, the CNRS and the Indo-French Centre for the Promotion of Advanced Research (IFCPAR, project 6105-1) for their financial support.

References

- ¹ Arumugam, S.; Popik, V. V. *J. Am. Chem. Soc.* **2009**, *131*, 11892-11899.
- ² Novak, I.; Harrison, L. J.; Li, W.; Kovač, B. *J. Phys. Chem. A* **2007**, *111*, 2619-2624.
- ³ Migani, A.; Blancafort, L.; Robb, M. A.; DeBellis, A. D. *J. Am. Chem. Soc.* **2008**, *130*, 6932-6933
- ⁴ Friedrich, L. E.; Schuster, G. B. *J. Am. Chem. Soc.* **1971**, *93*, 4602-4604.
- ⁵ Jaccob, M.; Sheeba Jem, I.; Giri, S.; Venuvanalingam, P.; Chattaraj, P. K. *J. Phys. Org. Chem.* **2011**, *24*, 460-465.
- ⁶ Jayaprakash, S.; Jeevanandam, J.; Subramani, K. *J. Mol. Model.* **2014**, *20*, 2494.

-
- ⁷ Guo, X.; Wang, J.; Li, C.-J. *J. Am. Chem. Soc.* **2009**, *131*, 15092-15093.
- ⁸ Varela, J. A.; González-Rodríguez, C.; Rubín, S. G.; Castedo, L.; Saá, C. *J. Am. Chem. Soc.* **2006**, *128*, 9576-9577.
- ⁹ Zhang, J.; Shen, W.; Liu, R.; Yu, Y.; Wu, H.; Li, M. *Int. J. Quantum Chem.* **2009**, *109*, 679-687.
- ¹⁰ Tomberg, A.; Kundu, S.; Zhou, F.; Li, C.-J.; Moitessier, N. *ACS Omega* **2018**, *3*, 3218-3227.
- ¹¹ Yabuno, Y.; Hiraga, Y.; Takagi, R.; Abe, M. *J. Am. Chem. Soc.* **2011**, *133*, 2592-2604.
- ¹² Xu, T.; Yang, Q.; Li, D.; Dong, J.; Yu, Z.; Li, Y. *Chem. Eur. J.* **2010**, *16*, 9264-9272.
- ¹³ Kutateladze, A. G.; Holt, T.; Reddy, D. S. *J. Org. Chem.* **2019**, *84*, 7575-7586.
- ¹⁴ Yang, J.; Shan, X.; Zhang, Z.; Tang, Y.; Zhao, M.; Chen, X. *J. Phys. Chem. A* **2014**, *118*, 11780-11786.
- ¹⁵ Tang, Y.; Shan, X.; Yang, J.; Niu, S.; Zhang, Z.; Watanabe, N.; Yamazaki, M.; Takahashi, M.; Chen, X. *J. Phys. Chem. A* **2016**, *120*, 6855-6863.
- ¹⁶ Harris, S. J.; Murdock, D.; Grubb, M. P.; Clark, I. P.; Greetham, G. M.; Towrie, M.; Ashfold, M. N. R. *J. Phys. Chem. A* **2014**, *118*, 10240-10245.
- ¹⁷ Koch, R.; Blanch, R. J.; Wentrup, C. *J. Org. Chem.* **2014**, *79*, 6978-6986.
- ¹⁸ Meier, H. *Molecules* **2012**, *17*, 1548-1570.
- ¹⁹ Balaban, A. T.; Greer, A.; Liebman, J. F. In *Advances in Heterocyclic Chemistry*; Katritzky, A. R., Ed.; Academic Press, 2014; Vol. 113, 111-142.
- ²⁰ Hideo, T.; Takeshi, M. *Chem. Lett.* **1997**, *26*, 399-400.
- ²¹ Shanks, D.; Frisell, H.; Ottosson, H.; Engman, L. *Org. Biomol. Chem.* **2006**, *4*, 846-852.
- ²² Škalamera, Đ.; Antol, I.; Mlinarić-Majerski, K.; Vančik, H.; Phillips, D. L.; Ma, J.; Basarić, N. *Chem. Eur. J.* **2018**, *24*, 9426-9435.
- ²³ Chen, S.; Zhang, Y.; Zhao, C.; Ren, F.; Liu, X.; Che, Y. *Fitoterapia* **2014**, *99*, 236-242.
- ²⁴ Chen, S.; Zhang, Y.; Niu, S.; Liu, X.; Che, Y. *J. Nat. Prod.* **2014**, *77*, 1513-1518.
- ²⁵ Wang, J.-S.; Zhang, Y.; Wang, X.-B.; Kong, L.-Y. *Tetrahedron* **2012**, *68*, 3963-3971.
- ²⁶ Zou, J.; Pan, L.; Li, Q.; Zhao, J.; Pu, J.; Yao, P.; Gong, N.; Lu, Y.; Kondratyuk, T. P.; Pezzuto, J. M. et al. *Org. Lett.* **2011**, *13*, 1406-1409.
- ²⁷ Sautier, B.; Lyons, S. E.; Webb, M. R.; Procter, D. *J. Org. Lett.* **2012**, *14*, 146-149.
- ²⁸ Griesbeck, A. G.; Franke, M.; Neudörfl, J.; Kotaka, H. *Beilstein J. Org. Chem.* **2011**, *7*, 127-134.
- ²⁹ Xin, H.; Zhu, X.; Yan, H.; Song, X. *Tetrahedron Lett.* **2013**, *54*, 3325-3328.
- ³⁰ Schmidt, W.; Jones, P. G.; Herrmann, J.; Müller, R.; Schulz, S. *Synlett* **2017**, *13*, 467-470.
- ³¹ Monleón, A.; Glaus, F.; Vergura, S.; Jørgensen, K. A. *Angew. Chem. Int. Ed.* **2016**, *55*, 2478-2482.
- ³² Conboy, D.; Mirallai, S. I.; Craig, A.; McArdle, P.; Al-Kinani, A. A.; Barton, S.; Aldabbagh, F. *J. Org. Chem.* **2019**, *84*, 9811-9818.
- ³³ Hao, X.; Liu, X.; Li, W.; Tan, F.; Chu, Y.; Zhao, X.; Lin, L.; Feng, X. *Org. Lett.* **2014**, *16*, 134-137.
- ³⁴ Ahmad, S.; Yousaf, M.; Mansha, A.; Rasool, N.; Zahoor, A. F.; Hafeez, F.; Rizvi, S. M. A. *Synth. Commun.* **2016**, *46*, 1397-1416.
- ³⁵ Pritchard, J. G.; Long, F. A. *J. Am. Chem. Soc.* **1958**, *80*, 4162-4165.
- ³⁶ Ringnér, B.; Sunner, S.; Watanabe, H. *Acta. Chem. Scand.* **1971**, *25*, 141-146.
- ³⁷ Eigenmann, H. K.; Golden, D. M.; Benson, S. W. *J. Phys. Chem.* **1973**, *77*, 1687-1691.
- ³⁸ Sella, A.; Basch, H.; Hoz, S. *J. Am. Chem. Soc.* **1996**, *118*, 416-420.
- ³⁹ Wolk, J. L.; Sprecher, M.; Basch, H.; Hoz, S. *Org. Biomol. Chem.* **2004**, *2*, 1065-1069.
- ⁴⁰ Wolk, J. L.; Hoz, T.; Basch, H.; Hoz, S. *J. Org. Chem.* **2001**, *66*, 915-918.
- ⁴¹ Banks, H. D. *J. Org. Chem.* **2003**, *68*, 2639-2644.
- ⁴² Sawicka, D.; Houk, K. N. *Mol. Model.* **2000**, *6*, 158-165.
- ⁴³ Rosenberg, M. G.; Billing, P.; Brecker, L.; Brinker, U. H. *J. Org. Chem.* **2014**, *79*, 8786-8799.
- ⁴⁴ Stedjan, M. K.; Augspurger, J. D. *J. Phys. Org. Chem.* **2015**, *28*, 298-303.

-
- ⁴⁵ Taskinen, E. J. *Phys. Org. Chem.* **2009**, *22*, 42-51.
- ⁴⁶ Goldsmith, C. F.; Magoon, G. R.; Green, W. H. *J. Phys. Chem. A* **2012**, *116*, 9033-9057.
- ⁴⁷ Wuitschik, G.; Carreira, E. M.; Wagner, B.; Fischer, H.; Parrilla, I.; Schuler, F.; Rogers-Evans, M.; Müller, K. *J. Med. Chem.* **2010**, *53*, 3227-3246.
- ⁴⁸ Burkhard, J. A.; Wuitschik, G.; Rogers-Evans, M.; Müller, K.; Carreira, E. M. *Angew. Chem. Int. Ed.* **2010**, *49*, 9052-9067.
- ⁴⁹ Wuitschik, G.; Rogers-Evans, M.; Buckl, A.; Bernasconi, M.; Märki, M.; Godel, T.; Fischer, H.; Wagner, B.; Parrilla, I.; Schuler, F. et al. *Angew. Chem. Int. Ed.* **2008**, *47*, 4512-4515.
- ⁵⁰ Wuitschik, G.; Rogers-Evans, M.; Müller, K.; Fischer, H.; Wagner, B.; Schuler, F.; Polonchuk, L.; Carreira, E. M. *Angew. Chem. Int. Ed.* **2006**, *45*, 7736-7739.
- ⁵¹ Job, A.; Janeck, C. F.; Bettray, W.; Peters, R.; Enders, D. *Tetrahedron* **2002**, *58*, 2253-2329.
- ⁵² Geden, J. V.; Beasley, B. O.; Clarkson, G. J.; Shipman, M. *J. Org. Chem.* **2013**, *78*, 12243-12250.
- ⁵³ Vo, C.-V. T.; Mikutis, G.; Bode, J. W. *Angew. Chem. Int. Ed.* **2013**, *52*, 1705-1708.
- ⁵⁴ Siau, W.-Y.; Bode, J. W. *J. Am. Chem. Soc.* **2014**, *136*, 17726-17729.
- ⁵⁵ Vo, C.-V. T.; Luescher, M. U.; Bode, J. W. *Nat. Chem.* **2014**, *6*, 310-314.
- ⁵⁶ Geoghegan, K.; Bode, J. W. *Org. Lett.* **2015**, *17*, 1934-1937.
- ⁵⁷ Brady, P. B.; Carreira, E. M. *Org. Lett.* **2015**, *17*, 3350-3353.
- ⁵⁸ Ruider, S. A.; Müller, S.; Carreira, E. M. *Angew. Chem. Int. Ed.* **2013**, *52*, 11908-11911.
- ⁵⁹ Curti, C.; Battistini, L.; Sartori, A.; Zanardi, F. *Chem. Rev.* **2020**, *120*, 2448-2612.
- ⁶⁰ Dieckmann, A.; Breugst, M.; Houk, K. N. *J. Am. Chem. Soc.* **2013**, *135*, 3237-3242.
- ⁶¹ Nassoy, A.-C.; Raubo, P.; Harrity, J. P. A. *Tetrahedron Lett.* **2013**, *54*, 3094-3096.
- ⁶² Corey, E. J.; Streith, J. *J. Am. Chem. Soc.* **1964**, *86*, 950-951.
- ⁶³ Coote, S. C. *Eur. J. Org. Chem.* **2020**, 1405-1423.
- ⁶⁴ Frébault, F.; Luparia, M.; Oliveira, M. T.; Goddard, R.; Maulide, N. *Angew. Chem. Int. Ed.* **2010**, *49*, 5672-5676.
- ⁶⁵ Gutekunst, W. R.; Baran, P. S. *J. Am. Chem. Soc.* **2011**, *133*, 19076-19079.
- ⁶⁶ Audisio, D.; Luparia, M.; Oliveira, M. T.; Klütt, D.; Maulide, N. *Angew. Chem. Int. Ed.* **2012**, *51*, 7314-7317.
- ⁶⁷ Souris, C.; Frébault, F.; Patel, A.; Audisio, D.; Houk, K. N.; Maulide, N. *Org. Lett.* **2013**, *15*, 3242-3245.
- ⁶⁸ Souris, C.; Luparia, M.; Frébault, F.; Audisio, D.; Farès, C.; Goddard, R.; Maulide, N. *Chem. Eur. J.* **2013**, *19*, 6566-6570.
- ⁶⁹ Souris, C.; Misale, A.; Chen, Y.; Luparia, M.; Maulide, N. *Org. Lett.* **2015**, *17*, 4486-4489.
- ⁷⁰ Misale, A.; Niyomchon, S.; Maulide, N. *Acc. Chem. Res.* **2016**, *49*, 2444-2458.
- ⁷¹ Bauer, A.; Nam, J.-H.; Maulide, N. *Synlett* **2019**, *30*, 413-416.
- ⁷² Yamashita, T.; Nishikawa, H.; Kawamoto, T. *Tetrahedron* **2019**, *75*, 617-623.
- ⁷³ Cabré, A.; Rafael, S.; Sciortino, G.; Ujaque, G.; Verdaguer, X.; Lledós, A.; Riera, A. *Angew. Chem. Int. Ed.* **2020**, *59*, 7521-7527.
- ⁷⁴ Ghosh, A. K.; Robinson, W. L. *J. Org. Chem.* **2019**, *84*, 9801-9805.
- ⁷⁵ Melis, N.; Luridiana, A.; Guillot, R.; Secci, F.; Frongia, A.; Boddaert, T.; Aitken, D. *J. Eur. J. Org. Chem.* **2017**, *2017*, 5896-5902.
- ⁷⁶ Nicolle, S. M.; Nortcliffe, A.; Bartrum, H. E.; Lewis, W.; Hayes, C. J.; Moody, C. J. *Chem. Eur. J.* **2017**, *23*, 13623-13627.
- ⁷⁷ D'Auria, M. *Photochem. Photobiol. Sci.* **2019**, *18*, 2297-2362.
- ⁷⁸ Mulzer, J.; Zippel, M.; Brüntrup, G. *Angew. Chem. Int. Ed.* **1980**, *19*, 465-466.
- ⁷⁹ Wu, X.; Hao, L.; Zhang, Y.; Rakesh, M.; Reddi, R. N.; Yang, S.; Song, B.-A.; Chi, Y. R. *Angew. Chem. Int. Ed.* **2017**, *56*, 4201-4205.
- ⁸⁰ Becker, M. R.; Reid, J. P.; Rykaczewski, K. A.; Schindler, C. S. *ACS Catal.* **2020**, *10*, 4387-4397.

- ⁸¹ Hanson, C. S.; Psaltakis, M. C.; Cortes, J. J.; Siddiqi, S. S.; Devery, J. J. *J. Org. Chem.* **2020**, *85*, 820-832.
- ⁸² McFarlin, A. T.; Watson, R. B.; Zehnder, T. E.; Schindler, C. S. *Adv. Synth. Catal.* **2020**, *362*, 365-369.
- ⁸³ Riehl, P. S.; Nasrallah, D. J.; Schindler, C. S. *Chem. Sci.* **2019**, *10*, 10267-10274.
- ⁸⁴ Becker, M. R.; Watson, R. B.; Schindler, C. S. *Chem. Soc. Rev.* **2018**, *47*, 7867-7881.
- ⁸⁵ Saito, A.; Tateishi, K. *Heterocycles* **2016**, *92*, 607.
- ⁸⁶ Jin, T.; Yamamoto, Y. *Org. Lett.* **2007**, *9*, 5259-5262.
- ⁸⁷ Liu, L.; Xu, B.; Hammond, G. B. *Beilstein J. Org. Chem.* **2011**, *7*, 606-614.
- ⁸⁸ Escalante, L.; González-Rodríguez, C.; Varela, J. A.; Saá, C. *Angew. Chem. Int. Ed.* **2012**, *51*, 12316-12320.
- ⁸⁹ Yeh, M.-C. P.; Lin, M.-N.; Hsu, C.-H.; Liang, C.-J. *J. Org. Chem.* **2013**, *78*, 12381-12396.
- ⁹⁰ Zhao, W.; Li, Z.; Sun, J. *J. Am. Chem. Soc.* **2013**, *135*, 4680-4683.
- ⁹¹ Cheng, D.; Ling, F.; Zheng, C.; Ma, C. *Org. Lett.* **2016**, *18*, 2435-2438.
- ⁹² Yoshioka, E.; Kohtani, S.; Miyabe, H. *Org. Lett.* **2010**, *12*, 1956-1959.
- ⁹³ Yoshida, H.; Ito, Y.; Ohshita, J. *Chem. Commun.* **2011**, *47*, 8512-8514.
- ⁹⁴ Zhou, C.; Wang, J.; Jin, J.; Lu, P.; Wang, Y. *Eur. J. Org. Chem.* **2014**, *2014*, 1832-1835.
- ⁹⁵ Fujishima, T.; Suenaga, T.; Nozaki, T. *Tetrahedron Lett.* **2014**, *55*, 3805-3808.
- ⁹⁶ Fujishima, T.; Nozaki, T.; Suenaga, T. *Bioorg. Med. Chem.* **2013**, *21*, 5209-5217.
- ⁹⁷ Suenaga, T.; Fujishima, T. *Tetrahedron* **2018**, *74*, 1461-1467.
- ⁹⁸ Trost, B. M.; Masters, J. T. *Chem. Soc. Rev.* **2016**, *45*, 2212-2238.
- ⁹⁹ Deng, H.; Jia, R.; Yang, W.-L.; Yu, X.; Deng, W.-P. *Chem. Commun.* **2019**, *55*, 7346-7349.
- ¹⁰⁰ Jones, B.; Proud, M.; Sridharan, V. *Tetrahedron Lett.* **2016**, *57*, 2811-2813.
- ¹⁰¹ Keshipeddy, S.; Martínez, I.; Castillo, B. F.; Morton, M. D.; Howell, A. R. *J. Org. Chem.* **2012**, *77*, 7883-7890.
- ¹⁰² Malapit, C. A.; Chitale, S. M.; Thakur, M. S.; Taboada, R.; Howell, A. R. *J. Org. Chem.* **2015**, *80*, 5196-5209.
- ¹⁰³ Wu, D.-D.; Huang, C.-M.; Wu, Y.-H.; Fun, H.-K.; Xu, J.-H.; Zhang, Y. *RSC Advances* **2013**, *3*, 7529-7536.
- ¹⁰⁴ Nakazaki, A.; Nakane, Y.; Ishikawa, Y.; Nishikawa, T. *Heterocycles* **2015**, *91*, 1157-1163.
- ¹⁰⁵ Fréneau, M.; Hoffmann, N. *J. Photochem. Photobiol. C: Photochem. Rev.* **2017**, *33*, 83-108.
- ¹⁰⁶ Ralph, M.; Ng, S.; Booker-Milburn, K. I. *Org. Lett.* **2016**, *18*, 968-971.
- ¹⁰⁷ Franceschi, P.; Mateos, J.; Vega Peñaloza, A.; Dell'Amico, L. *Eur. J. Org. Chem.* **2020**, 6718-6722.
- ¹⁰⁸ Poscharny, K.; Fabry, D. C.; Heddrich, S.; Sugiono, E.; Liauw, M. A.; Rueping, M. *Tetrahedron* **2018**, *74*, 3171-3175.
- ¹⁰⁹ Flores, D. M.; Schmidt, V. A. *J. Am. Chem. Soc.* **2019**, *141*, 8741-8745.
- ¹¹⁰ Buendia, J.; Chang, Z.; Eijsberg, H.; Guillot, R.; Frongia, A.; Secci, F.; Xie, J.; Robin, S.; Boddaert, T.; Aitken, D. *J. Angew. Chem. Int. Ed.* **2018**, *57*, 6592-6596.
- ¹¹¹ Huang, C.; Yu, H.; Miao, Z.; Zhou, J.; Wang, S.; Fun, H.-K.; Xu, J.; Zhang, Y. *Org. Biomol. Chem.* **2011**, *9*, 3629-3631.
- ¹¹² Kumarasamy, E.; Raghunathan, R.; Kandappa, S. K.; Sreenithya, A.; Jockusch, S.; Sunoj, R. B.; Sivaguru, J. *J. Am. Chem. Soc.* **2017**, *139*, 655-662.
- ¹¹³ Wang, W.; Zhang, W.-J.; Wang, L.; Quah, C. K.; Fun, H.-K.; Xu, J.-H.; Zhang, Y. *J. Org. Chem.* **2013**, *78*, 6211-6222.
- ¹¹⁴ Kaeobamrung, J.; Bode, J. W. *Org. Lett.* **2009**, *11*, 677-680.
- ¹¹⁵ Mukherjee, S.; Mondal, S.; Patra, A.; Gonnade, R. G.; Biju, A. T. *Chem. Commun.* **2015**, *51*, 9559-9562.
- ¹¹⁶ Zhang, J.-Q.; Li, N.-K.; Yin, S.-J.; Sun, B.-B.; Fan, W.-T.; Wang, X.-W. *Adv. Synth. Catal.* **2017**, *359*, 1541-1551.
- ¹¹⁷ Cohen, D. T.; Eichman, C. C.; Phillips, E. M.; Zarefsky, E. R.; Scheidt, K. A. *Angew. Chem. Int. Ed.* **2012**, *51*, 7309-7313.
- ¹¹⁸ Johnston, R. C.; Cohen, D. T.; Eichman, C. C.; Scheidt, K. A.; Ha-Yeon Cheong, P. *Chem. Sci.* **2014**, *5*, 1974-1982.

- ¹¹⁹ Cohen, D. T.; Johnston, R. C.; Rosson, N. T.; Cheong, P. H.-Y.; Scheidt, K. A. *Chem. Commun.* **2015**, 51, 2690-2693.
- ¹²⁰ Candish, L.; Forsyth, C. M.; Lupton, D. W. *Angew. Chem. Int. Ed.* **2013**, 52, 9149-9152.
- ¹²¹ Candish, L.; Lupton, D. W. *J. Am. Chem. Soc.* **2013**, 135, 58-61.
- ¹²² Bera, S.; Samanta, R. C.; Daniliuc, C. G.; Studer, A. *Angew. Chem. Int. Ed.* **2014**, 53, 9622-9626.
- ¹²³ Liu, G.; Shirley, M. E.; Van, K. N.; McFarlin, R. L.; Romo, D. *Nat. Chem.* **2013**, 5, 1049-1057.
- ¹²⁴ Leeder, A. J.; Heap, R. J.; Brown, L. J.; Franck, X.; Brown, R. C. D. *Org. Lett.* **2016**, 18, 5971-5973.
- ¹²⁵ Davis, O. A.; Croft, R. A.; Bull, J. A. *Chem. Commun.* **2015**, 51, 15446-15449.
- ¹²⁶ Davis, O. A.; Bull, J. A. *Angew. Chem. Int. Ed.* **2014**, 53, 14230-14234.
- ¹²⁷ Lekkala, R.; Lekkala, R.; Moku, B.; Rakesh, K. P.; Qin, H.-L. *Beilstein J. Org. Chem.* **2019**, 15, 976-980.
- ¹²⁸ Xu, R.; Czarniecki, M.; Man, J. d.; Pan, J.; Qiang, L.; Root, Y.; Ying, S.; Su, J.; Sun, X.; Zhang, Y.; Hu, T.; Chen, S.-H. *Tetrahedron Lett.* **2011**, 52, 3266-3270.
- ¹²⁹ For a review on spiro-piperidines, see: Griggs, S. D.; Tape, D. T.; Clarke, P. A. *Org. Biomol. Chem.* **2018**, 16, 6620-6633.
- ¹³⁰ Yoshioka, S.; Nagatomo, M.; Inoue, M. *Org. Lett.* **2015**, 17, 90-93.
- ¹³¹ Lajkiewicz, N. J.; Cognetta, A. B.; Niphakis, M. J.; Cravatt, B. F.; Porco, J. A. *J. Am. Chem. Soc.* **2014**, 136, 2659-2664.
- ¹³² Shenvi, R. A.; Corey, E. J. *J. Am. Chem. Soc.* **2009**, 131, 5746-5747.
- ¹³³ Rullière, P.; Cannillo, A.; Grisel, J.; Cividino, P.; Sébastien, C.; Poisson, J.-F. *Org. Lett.* **2018**, 20, 4558-4561.
- ¹³⁴ Fukuda, T.; Sugiyama, K.; Arima, S.; Harigaya, Y.; Nagamitsu, T.; Ōmura, S. *Org. Lett.* **2008**, 10, 4239-4242.
- ¹³⁵ Satoh, N.; Yokoshima, S.; Fukuyama, T. *Org. Lett.* **2011**, 13, 3028-3031.
- ¹³⁶ Gholami, H.; Kulshrestha, A.; Favor, O. K.; Staples, R. J.; Borhan, B. *Angew. Chem. Int. Ed.* **2019**, 58, 10110-10113.
- ¹³⁷ Marx, L. B.; Burton, J. W. *Chem. Eur. J.* **2018**, 24, 6747-6754.
- ¹³⁸ Armitage, I.; Elliott, E. L.; Hicks, F.; Langston, M.; McCarron, A.; McCubbin, Q. J.; O'Brien, E.; Stirling, M.; Zhu, L. *Org. Proc. Res. Dev.* **2015**, 19, 1299-1307.
- ¹³⁹ Tong, G.; Liu, Z.; Li, P. *Org. Lett.* **2014**, 16, 2288-2291.
- ¹⁴⁰ Nicolle, S. M.; Nortcliffe, A.; Bartrum, H. E.; Lewis, W.; Hayes, C. J.; Moody, C. J. *Chem. Eur. J.* **2017**, 23, 13623-13627.
- ¹⁴¹ Wang, J.-C.; Zhang, Y.; Xu, Z.-J.; Lo, V. K.-Y.; Che, C.-M. *ACS Catal.* **2013**, 3, 1144-1148.
- ¹⁴² Huang, J.-M.; Yang, C.-S.; Tanaka, M.; Fukuyama, Y. *Tetrahedron* **2001**, 57, 4691-4698.
- ¹⁴³ Birman, V. B.; Danishefsky, S. J. *J. Am. Chem. Soc.* **2002**, 124, 2080-2081.
- ¹⁴⁴ Chen, J.; Gao, P.; Yu, F.; Yang, Y.; Zhu, S.; Zhai, H. *Angew. Chem. Int. Ed.* **2012**, 51, 5897-5899.
- ¹⁴⁵ He, W.; Huang, J.; Sun, X.; Frontier, A. J. *J. Am. Chem. Soc.* **2008**, 130, 300-308.
- ¹⁴⁶ Ganji, P.; Ibrahim, H. *Chem. Commun.* **2012**, 48, 10138-10140.
- ¹⁴⁷ Hubbell, A. K.; LaPointe, A. M.; Lamb, J. R.; Coates, G. W. *J. Am. Chem. Soc.* **2019**, 141, 2474-2480.
- ¹⁴⁸ Lam, S. M.; Wong, W.-t.; Chiu, P. *Org. Lett.* **2017**, 19, 4468-4471.
- ¹⁴⁹ Vil, V.; Terent'ev, A. O.; Al Quntar, A. A. A.; Glorizova, T. A.; Savidov, N.; Dembitsky, V. M. *Appl. Microbiol. Biotechnol.* **2019**, 103, 2449-2467.
- ¹⁵⁰ Fujita, S.; Nishikawa, K.; Iwata, T.; Tomiyama, T.; Ikenaga, H.; Matsumoto, K.; Shindo, M. *Chem. Eur. J.* **2018**, 24, 1539-1543.
- ¹⁵¹ Shindo, M.; Sato, Y.; Shishido, K. *Tetrahedron* **1998**, 54, 2411-2422.
- ¹⁵² Schmiedel, V. M.; Hong, Y. J.; Lentz, D.; Tantillo, D. J.; Christmann, M. *Angew. Chem. Int. Ed.* **2018**, 57, 2419-2422.
- ¹⁵³ Harvey, N. L.; Krysiak, J.; Chamni, S.; Cho, S. W.; Sieber, S. A.; Romo, D. *Chem. Eur. J.* **2015**, 21, 1425-1428.

-
- ¹⁵⁴ Cortez, G. S.; Tennyson, R. L.; Romo, D. *J. Am. Chem. Soc.* **2001**, *123*, 7945-7946.
- ¹⁵⁵ Mondal, S.; Mukherjee, S.; Das, T. K.; Gonnade, R.; Biju, A. T. *ACS Catal.* **2017**, *7*, 3995-3999.
- ¹⁵⁶ Leverett, C. A.; Purohit, V. C.; Johnson, A. G.; Davis, R. L.; Tantillo, D. J.; Romo, D. *J. Am. Chem. Soc.* **2012**, *134*, 13348-13356.
- ¹⁵⁷ Liu, G.; Shirley, M. E.; Romo, D. *J. Org. Chem.* **2012**, *77*, 2496-2500.
- ¹⁵⁸ Kong, W.; Romo, D. *J. Org. Chem.* **2017**, *82*, 13161-13170.
- ¹⁵⁹ Nguyen, H.; Ma, G.; Romo, D. *Chem. Commun.* **2010**, *46*, 4803-4805.
- ¹⁶⁰ Ma, G.; Nguyen, H.; Romo, D. *Org. Lett.* **2007**, *9*, 2143-2146.
- ¹⁶¹ Liu, G.; Romo, D. *Angew. Chem. Int. Ed.* **2011**, *50*, 7537-7540.
- ¹⁶² Pullaiah, K. C.; Surapaneni, R. K.; Rao, C. B.; Albizati, K. F.; Sullivan, B. W.; Faulkner, D. J.; He, C. H.; Clardy, J. *J. Org. Chem.* **1985**, *50*, 3665-3666.
- ¹⁶³ Hugelshofer, C. L.; Magauer, T. *J. Am. Chem. Soc.* **2016**, *138*, 6420-6423.
- ¹⁶⁴ Hugelshofer, C. L.; Magauer, T. *Chem. Eur. J.* **2016**, *22*, 15125-15136.
- ¹⁶⁵ Zeng, M.; Murphy, S. K.; Herzon, S. B. *J. Am. Chem. Soc.* **2017**, *139*, 16377-16388.
- ¹⁶⁶ Zhang, W.; Xu, L.; Yang, L.; Huang, Y.; Li, S.; Shen, Y. *Fitoterapia* **2014**, *96*, 146-151.
- ¹⁶⁷ Zheng, Z.-P.; Xu, Y.; Qin, C.; Zhang, S.; Gu, X.; Lin, Y.; Xie, G.; Wang, M.; Chen, J. *J. Agr. Food Chem.* **2014**, *62*, 5519-5527.
- ¹⁶⁸ Ha, M. T.; Seong, S. H.; Nguyen, T. D.; Cho, W.-K.; Ah, K. J.; Ma, J. Y.; Woo, M. H.; Choi, J. S.; Min, B. S. *Phytochem.* **2018**, *155*, 114-125.
- ¹⁶⁹ Chen, J.; Bai, X.; Hua, Y.; Zhang, H.; Wang, H. *Fitoterapia* **2019**, *134*, 1-4.
- ¹⁷⁰ Richter, M. J. R.; Schneider, M.; Brandstätter, M.; Krautwald, S.; Carreira, E. M. *J. Am. Chem. Soc.* **2018**, *140*, 16704-16710.
- ¹⁷¹ Wein, L. A.; Wurst, K.; Angyal, P.; Weisheit, L.; Magauer, T. *J. Am. Chem. Soc.* **2019**, *141*, 19589-19593.