

The proline cycle as a eukaryotic redox valve

Yao Zheng, Cécile Cabassa-Hourton, Séverine Planchais, Sandrine Lebreton, Arnould Savouré

▶ To cite this version:

Yao Zheng, Cécile Cabassa-Hourton, Séverine Planchais, Sandrine Lebreton, Arnould Savouré. The proline cycle as a eukaryotic redox valve. Journal of Experimental Botany, 2021, 72 (20), pp.6856-6866. 10.1093/jxb/erab361 . hal-03360252

HAL Id: hal-03360252 https://hal.sorbonne-universite.fr/hal-03360252

Submitted on 30 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Fig. 1. Proline metabolism in plants and its links to the ornithine and pentose phosphate pathways. The proline cycle is highlighted by blue arrows and the putative proline-P5C cycle by a square in light blue. Asp, aspartate; BAC, basic amino acid carrier; BOU, a bout de souffle; ETC, electron transfer chain; Glu, glutamate; KG, α -ketoglutarate; OAT, ornithine δ - aminotransferase; Orn, ornithine; PPP shunt, pentose phosphate pathway; P5C, pyrroline-5-carboxyate; P5CDH, P5C dehydrogenase; P5CS, P5C synthetase; ProDH, proline dehydrogenase; Pro, proline; UCP, mitochondrial uncoupling protein; ? indicates unknown transporter(s).

The proline cycle as a eukaryotic redox valve

Yao Zheng, Cécile Cabassa-Hourton, Séverine Planchais, Sandrine Lebreton, Arnould Savouré¹

Sorbonne Université, UPEC, CNRS, IRD, INRAE Institute of Ecology and Environmental Sciences of Paris (iEES), F-75005 Paris, France

1 Corresponding author arnould.savoure@sorbonne-universite.fr

Running title

The proline cycle

Highlight

The proline cycle regulates energy and redox power in a shuttle mechanism with a valve function during plant development and in stress conditions

1 Abstract

2 The amino acid proline has been known for many years to be a component of proteins as well as an osmolyte. Many recent studies have demonstrated that proline has other roles such as regulating redox 3 balance and energy status. In animals and plants, the well-described proline cycle is concomitantly 4 5 responsible for the preferential accumulation of proline and shuttling of redox equivalents from the cytosol to mitochondria. The impact of the proline cycle goes beyond regulating proline levels. In 6 7 this review, we focus on recent evidence of how the proline cycle regulates redox status in relation to 8 other redox shuttles. We discuss how the interconversion of proline to glutamate shuttles reducing 9 power between cellular compartments. Spatial aspects of the proline cycle in the entire plant are 10 considered in terms of proline transport between organs with different metabolic regimes 11 (photosynthesis versus respiration). Furthermore, we highlight the importance of this shuttle in the 12 regulation of energy and redox power in plants, through a particularly intricate coordination notably 13 between mitochondria and cytosol.

- 14
- 15

16 Keywords

Mitochondria; Proline metabolism; Proline cycle; Proline transport; Redox status; Redox shuttle;
Redox valve

- 19
- 20

21 Introduction

22 Redox homeostasis during development is vital for all branches of the tree of life, particularly in 23 response to environmental changes (Scheibe et al., 2005). Compartmentalization of metabolism and 24 other cellular functions in eukaryotic cells implies that pools of metabolites, ions and reducing 25 equivalents are concentrated locally in a regulated manner (Lunn, 2006). Maintenance of metabolism 26 as environmental conditions vary, whether regularly, rapidly or to extremes, is a challenge for all 27 eukaryotic organisms, and particularly plants for several reasons. As sessile organisms, plants cannot 28 move away to more clement conditions, so they are subject to dynamic environmental alterations. In 29 eukaryotic photosynthetic organisms, the regulation of energy metabolism is particularly complex 30 because it involves two organelles, the mitochondrion and the chloroplast, as major sources of ATP 31 and reducing power in the form of pyridine nucleotides NADH and NADPH (Foyer and Noctor, 32 2020). Additional complexity of redox and energy metabolism arises from the daily cycles of light 33 and dark periods, and the presence of green and non-green tissues within the same individual. In daylight, photon energy is transformed in green tissues by photosystems into ATP and NADPH, 34 35 which in turn are used to fix CO₂ by the Calvin–Benson cycle. In mitochondria, oxidative

36 phosphorylation breaks down carbohydrates to generate ATP in all tissues at any time, but remains 37 the only powerhouse system operative during the night and in non-green tissues. Tight spatiotemporal 38 coordination of supply and demand to the two powerhouse systems of plant cells is required to meet 39 the energy needs of anabolic processes such as primary assimilation of carbon and nitrogen, as well 40 as transport of substrates, intermediates and products.

41 Specific translocators enable the direct or indirect exchange of reducing power between cell 42 compartments (Palmieri et al., 2009). The cofactor NAD⁺, the oxidized form of NADH, is synthesized 43 in the cytosol and must be imported into organelles to ensure timely supply to many enzymatic 44 reactions. In Arabidopsis, the two NAD⁺ carrier proteins NDT1 and NDT2 have been shown to 45 respectively localize in plastid and mitochondrial membranes, allowing direct transport of NAD⁺ into 46 these organelles (Palmieri et al., 2009). Recent data locate NDT1 exclusively in the mitochondrial membrane (de Souza Chaves et al., 2019). On the contrary, no transporter has been identified as a 47 48 carrier of reduced form NADH from one subcellular compartment to another. Membranes are also 49 impermeable to both NADPH cofactor and its oxidized form NADP⁺. An NADP⁺ carrier has not been identified yet, suggesting that this compound is likely to be formed by an NAD⁺ kinase. Indeed NAD⁺ 50 51 kinases have been found to be localized in the cytosol and chloroplasts (Gakière et al., 2018; 52 Dell'Aglio et al., 2019). Furthermore, NAD⁺ kinase has been also detected in the mitochondrial 53 matrix of yeasts and mammals (Outten and Culotta, 2003; Ohashi et al., 2012), but not of plants 54 (Gakière et al., 2018).

55 As organellar membranes are not permeable to reduced forms of pyridine nucleotides, the 56 compartmentalized redox couples (NAD+/NADH and NADP+/NADPH) rely on shuttles for their 57 translocation between subcellular compartments. A shuttle system is a series of biochemical 58 interconversions that temporarily bind the redox or other molecular entity in a permeable form that 59 can be ferried across the membrane barrier, releasing or reconstituting the entity on the other side of the barrier. The number of steps, enzymes, substrates, and transporters involved depends on the type, 60 61 site and function of the shuttle. Shuttles are thus dynamic control systems that redirect metabolic flux 62 so certain compounds can be overproduced. Organellar and cytosolic NADH and NADPH pools have 63 been shown to be regulated by multiple shuttles or metabolic valves, such as the glycerol-3-phosphate 64 (G3P) shuttle and the malate-aspartate shuttle for the NADH pools, and the malate-oxaloacetate 65 (OAA) shuttle for NADPH pools. The concept of a metabolic valve is that build-up of an entity, such 66 as a reducing equivalent, on one side of a membrane barrier can be released or purged in a regulated 67 manner, redistributing the entity in the cell while resetting the conditions on either side of the barrier. In this sense, a valve might not be expected to operate stoichiometrically. Such shuttle and valve 68 mechanisms enable redox and energy homeostasis in both normal and stressed states. Proline 69 70 metabolism involves the interconversion of glutamate and proline in a process linked to cellular

compartments and energetics. Therefore, enzymes involved in proline metabolism could participate
 in NAD(P)⁺/NAD(P)H homeostasis in a shuttle mechanism with a valve function during plant
 development and in stress conditions.

74

75 **Proline metabolism**

76 Abiotic stresses perturb cellular redox homeostasis and proline metabolism is thought to act as an 77 important redox buffer in both plants and animals (Krishman et al., 2008; Sharma et al., 2011; Ben 78 Rejeb et al., 2014, 2015; Phang, 2019). Proline plays important roles in various developmental stages 79 of plants as well as in stress tolerance (Kavi Khishor and Sreenivasulu, 2014; Bhaskara et al., 2015; 80 Trovato *et al.*, 2019). Indeed most plant species have been shown to accumulate proline in response 81 to abiotic and biotic stresses. In plants, proline biosynthesis occurs in the cytosol starting from 82 glutamate (Fig. 1). Glutamate is phosphorylated and reduced by the Δ^1 -pyrroline-5-carboxylate 83 synthetase (P5CS) using NADPH as a cofactor to form glutamate semialdehyde (GSA) in equilibrium 84 with pyrroline-5-carboxylate (P5C). P5C is further reduced into proline by P5C reductase (P5CR) 85 also using NAD(P)H as a cofactor. Székely et al. (2008) presented evidence that P5CS1 may be 86 localized in chloroplasts during salt stress, indicating that proline may also be synthesized in this 87 organelle. Recent work using spectrally resolved fluorescence imaging showed exclusive cytosolic 88 localization of P5CS1 and P5CS2 in Arabidopsis protoplasts, suggesting that plastids do not 89 contribute to P5CS-mediated proline biosynthesis (Funck et al., 2020). The subcellular localization 90 of P5CS1 therefore remains ambiguous and may depend on growth and/or stress conditions. Further 91 research is required to unravel these discrepancies.

In animals, the conversion of P5C to proline is also mediated by P5CR, commonly named PYCR.
The organization of the proline biosynthesis pathway is more complex at least in humans where
PYCR is encoded by three different genes, *PYCR1*, 2, and *L*, with both PYCR1 and 2 localized in
mitochondria and PYCRL in the cytosol (De Ingeniis *et al.*, 2012). P5CR was shown to be exclusively
cytosolic in Arabidopsis cells (Funck *et al.*, 2012).

97 When stress is relieved, proline is rapidly transported from the cytosol into mitochondria by either a 98 mitochondrial proline symporter or a proline/glutamate antiporter (Di Martino et al., 2006); however, 99 genes encoding these transporters have yet to be identified. Mitochondrial proline is then oxidized by 100 the sequential action of proline dehydrogenase (ProDH) and P5C dehydrogenase (P5CDH) to release 101 glutamate. Mitochondrial glutamate transporters "a bout de souffle" (BOU) and mitochondrial 102 uncoupling proteins UCP1/2 have now been identified and it is conceivable that they could participate 103 in the shuttling of glutamate during the proline cycle (Monné et al., 2018; Porcelli et al., 2018). 104 ProDH, also known as proline oxidase (POX), catalyzes the first and rate-limiting step of proline 105 catabolism using FAD as a cofactor (Servet et al., 2012). In both animals and plants, ProDH is

localized on the matrix side of the mitochondrial inner membrane (Elthon and Stewart, 1981;
Cabassa-Hourton *et al.*, 2016) where it transfers electrons released from proline oxidation to the
mitochondrial electron chain. While P5CDH activity was detected in the mitochondrial inner
membrane in *Zea mays* (maize) (Elthon and Stewart, 1981), this enzyme is localized in the
mitochondrial matrix in animals (Brunner and Neupert, 1969) and other plants (Deuschle *et al.*, 2001).
Recently, Ren *et al.* (2018) have shown that the two enzymes may physically interact with the
inhibitory protein DFR1 in Arabidopsis to regulate proline catabolism.

- 113 In mitochondria, ornithine δ -aminotransaminase (OAT) contributes to the formation of GSA and glutamate by transferring the ornithine δ -amino group to α -ketoglutarate (KG). There have been some 114 115 reports of proline biosynthesis from ornithine in plants (Delauney et al., 1993; Roosens et al., 1998) and in animals (De Ingeniis et al., 2012). An increase in proline content was also shown to be 116 117 correlated with higher OAT activity in salt-stressed cashew plants (da Rocha et al., 2012) and in rice overexpressing OAT (You et al., 2012). In comparison, Funck et al. (2008) demonstrated that, at least 118 119 in Arabidopsis, P5C is involved in glutamate formation through the action of P5CDH rather than 120 proline biosynthesis.
- 121 In bacteria, invertebrates and plants such as halophytes, proline accumulation has been shown to 122 contribute to osmotic adjustment of cells to counterbalance water loss (Slama et al., 2015; Forlani et 123 al., 2019). Proline can act as a metabolic signaling molecule to modulate mitochondrial functions and 124 influence cell death (Szabados and Savouré, 2010; Zhang and Becker, 2015; Phang, 2019; Senthil-125 Kumar and Mysore, 2012; Rizzi et al., 2016). Proline has also been shown to participate in the redox 126 balance and energy status of the cell. Proline metabolism may provide stress protection by 127 maintaining the NADPH/NADP⁺ balance and the levels of antioxidants (Ben Rejeb et al., 2014). On a cellular scale, proline metabolism may bridge compartments to allow redox changes across 128 membranes. Given this reach and multifunctionality, we now focus on the characteristics of three 129 redox shuttle systems to provide a framework in which to consider whether proline metabolism could 130 131 fulfil a similar purpose in eukaryotes.
- 132

133 Eukaryotic redox shuttles

134

135 *The glycerol-3-phosphate (G3P) shuttle*

In mitochondria, electrons released from NADH oxidation are transferred to the electron transport chain, which in turn feeds the energy necessary to pump protons across the inner mitochondrial membrane. This creates an electrochemical proton gradient that drives ATP synthesis. In addition to ATP generation, mitochondrial membrane potential drives the transport of metabolites between mitochondrial and cytosolic compartments. In yeasts, plants and animals, a G3P shuttle involved in

redox homeostasis has been described (Ansell et al., 1997; Shen et al., 2003, 2006; Mráček et al., 141 142 2013). A key component of the shuttle is a G3P dehydrogenase on the outer surface of the inner mitochondrial membrane which donates electrons directly to the ubiquinone pool. The G3P shuttle 143 144 irreversibly consumes cytosolic NADH generating mitochondrial FADH₂ (Fig. 2A; Shen et al., 2006; 145 McKenna et al., 2006). Briefly, electrons from NADH are transferred to dihydroxyacetone phosphate (DHAP) via the cytosolic G3P dehydrogenase (cGPDH) to form G3P. Then, G3P is converted to 146 147 DHAP on the outer surface of the inner mitochondrial membrane by mitochondrial G3P dehydrogenase (mGPDH). Next, electrons are transferred to FAD to form FADH₂ within the 148 149 mitochondria, coupled with the reduction of ubiquinone (McKenna et al., 2006). This G3P shuttle is 150 especially prominent in muscle cells enabling an extremely high rate of oxidative phosphorylation to 151 be sustained. Some insects that lack lactate dehydrogenase are completely dependent on the G3P shuttle for the regeneration of cytoplasmic NAD⁺ (Mráček et al., 2013). Generally, the G3P shuttle 152 provides a less important mechanism of redox state regulation in mammalian cells than the malate-153 154 aspartate shuttle (described in detail below) because most tissues have only low levels of mGPDH 155 (Mráček et al., 2013). In plants, the FAD-GPDH gene has been reported to be highly expressed during 156 seed germination and repressed upon water stress (Shen et al., 2006; Quettier et al., 2008). However, 157 the role of G3P shuttle remains elusive in plants.

158

159 The malate-aspartate shuttle

160 Like the G3P shuttle, the more complex malate-aspartate shuttle acts to irreversibly transfer reducing 161 equivalents from cytosolic NADH to mitochondria (Borst, 2020), but in contrast it transfers NADH 162 from the cytosol into mitochondrial matrix forming NADH again. This system involves two cytosolic 163 and two matrix enzymes as well as two inner membrane transporters. In the cytosol, electrons from 164 NADH are transferred to OAA by cytosolic malate dehydrogenase (cMDH) to generate malate (Fig. 165 2B). Malate is multifunctional because it is an intermediate in energy metabolism that is also involved 166 in the transfer of reducing equivalents, and possibly metabolic trafficking between different cell types 167 in eukaryotes. Malate enters the mitochondrial matrix via the malate-KG carrier (OGC) in exchange 168 for KG. The mitochondrial malate dehydrogenase (mMDH) converts malate to OAA, transferring the 169 electrons to NAD⁺ to form NADH in the matrix. Subsequently, OAA is transaminated to aspartate 170 via mitochondrial glutamate OAA aminotransferase (GOT) with the simultaneous conversion of 171 glutamate to KG. Finally, aspartate is exported from the mitochondria via the aspartate-glutamate 172 carrier (AGC) in exchange for cytosolic glutamate, which brings one proton into the mitochondria. 173 The efflux of aspartate from mitochondria is stoichiometric with entry of glutamate plus a proton in

an electrogenic exchange that provides directionality while controlling the rate (LaNoue *et al.*, 1974a,
b). Consequently, the malate-aspartate shuttle is powered by the electrochemical proton gradient to

- 176 export aspartate and so aids the import of reducing equivalents into mitochondria. It is not clear what
- 177 regulates the balance between malate-aspartate shuttle and G3P shuttle.

The malate-aspartate shuttle is found in yeast and animals (Cavero *et al.*, 2003; McKenna *et al.*, 2006; Satrústegui and Bak, 2015). However, it is still unclear whether this shuttle also occurs in plant mitochondria. There is evidence for this type of shuttle in plant-bacteria interactions in root nodules of *Alnus glutinosa* (Akkermans *et al.*, 1981) and *Pisum sativum* with an associated role in nitrogen fixation (Appels and Haaker, 1991). This shuttle has also been described as operating between glyoxysomes and mitochondria from the endosperm of germinating castor bean (Mettler and Beevers, 1980).

185

186 The malate-OAA shuttle

187 In plant cells, the regulation of energy metabolism is particularly complex because both mitochondria and chloroplasts are involved. The malate-OAA shuttle, sometimes referred to as the malate valve, is 188 189 a powerful system for balancing metabolic fluxes through indirect transport of reducing 190 equivalents. The valve is composed of cytosolic, mitochondrial, plastid, and peroxisomal malate 191 dehydrogenases (MDH) in cooperation with malate-OAA translocators (MOT) (Fig. 3). For example, 192 higher levels of NAD(P)H drive plastid MDH to convert OAA to malate. Then, malate is transferred 193 to cytosol or mitochondria by MOTs. In the mitochondrial matrix, malate is oxidized by 194 mitochondrial MDH which reduces NAD⁺ to NADH (Selinski and Scheibe, 2019). Theoretically, the 195 malate-OAA shuttle should be able to transfer reducing equivalents in either direction depending on the prevailing redox conditions on either side of the membrane. Oxidation of malate to OAA by 196 197 mitochondrial MDH generates NADH, which is then oxidized by Complex I producing ROS. 198 Generation of ROS from malate oxidation can trigger cell death in both animals and plants suggesting 199 that such mechanisms are conserved (Zhao et al., 2020). However, in plant cells this shuttle may only 200 serve to export reducing equivalents from mitochondria and plastids in normal growth conditions 201 (Selinski and Scheibe, 2019). It should be noted that the exchange of malate and OAA mediated by 202 MOTs across different cellular compartments is not driven by the electrochemical proton gradient, 203 so this exchange can only move reducing equivalents from a relatively reduced compartment to a 204 relatively oxidized compartment. Since the mitochondrial matrix is much more reduced than the 205 cytosol with respect to the NADH/NAD $^+$ ratio, the OAA transporter is likely to work primarily to 206 export reducing equivalents from mitochondria to the cytosol (Selinski and Scheibe, 2019).

207

208 The proline cycle as a reducing-equivalent shuttle

209 Considering that metabolism of proline involves the oxidation of NAD(P)H in the cytosol and the 210 reduction of NAD⁺ in mitochondria, we postulate that proline metabolism may also play an important role as a redox regulator in eukaryotic cells. Indeed, P5CS and P5CR oxidize NAD(P)H in the cytosol
while in mitochondria ProDH and P5CDH reduce FAD and NAD⁺ respectively.

213 In the cytosol, induction of P5CS and/or P5CR expression may help to maintain a low 214 NADPH/NADP⁺ ratio (Liang *et al.*, 2013). Coordination between proline and redox metabolism can 215 be deduced from the phenotype of the Arabidopsis *p5cs1* mutant because it displays hypersensitivity 216 to salt stress, accumulation of ROS and strongly enriched expression of genes involved in redox 217 metabolism of the mitochondria and chloroplasts (Székely et al., 2008; Shinde et al., 2016). Using a 218 forward genetic screen based on a ProDH1-promoter luciferase reporter, Shinde et al. (2016) found 219 that mutants affected in the synthesis of very-long-chain fatty acids (VLCFA) or in cuticle deposition 220 also accumulate more proline. This implies a strong coordination between proline and lipid 221 metabolism in relation to redox status. Proline and lipid metabolism both help buffer cellular redox 222 status under stress.

223 Regulation of P5CR activity in Arabidopsis seems to depend on a complex pattern of regulation 224 including the ratios of reduced to oxidized pyridine nucleotides, the preference for phosphorylated 225 over non-phosphorylated pyridine nucleotides, and the presence of ions (Giberti et al., 2014; Ruszkowski et al., 2015). Equimolar concentrations of NADP⁺ completely suppress the NADH-226 227 dependent activity of P5CR, whereas the NADPH-dependent reaction is mildly affected (Giberti et 228 al., 2014). Moreover, NADH-dependent but not NADPH-dependent activity of P5CR is inhibited by 229 an excess of proline. Excess of salt inhibits the NADH-dependent and activates the NADPH-230 dependent activity of P5CR. Similarly, in the protozoan Trypanosoma cruzi, TcP5CR is an NADPH-231 dependent cytosolic enzyme whose activity is fine-tuned by NADPH cytosolic pools (Marchese et 232 al., 2020). Thus, the cytosolic reduced status dependent on proline cycle functioning would be mostly 233 driven by NADPH homeostasis regulation.

234 In mitochondria, regulation of the reduced state by the proline cycle occurs through the production 235 of both NADH and FADH₂, which impact the electron transfer chain functioning as well as ROS 236 production. Several studies in both animal and plant cells have shown that increasing the reduced 237 state of mitochondria by metabolic or mutational manipulation leads to proline biosynthesis, while decreasing mitochondrial reduced state and/or ATP synthesis impairs proline accumulation (Huang 238 239 et al., 2013; Schwörer et al., 2020). A decrease in the activity of the mitochondrial electron transfer 240 chain caused by a defect in one of its super protein complexes leaves cells prone to the damage of 241 excess reducing power. Enhancing proline biosynthesis and down-regulating its degradation would 242 be an alternative way to trap reducing power and limit saturation of the electron transfer chain and 243 formation of ROS. Recently, SSR1, a mitochondrial protein with a tetratricopeptide repeat domain, was shown to be involved in maintaining the function of the mitochondrial electron transfer chain 244 245 (Han *et al.*, 2021). When the *ssr1-1* mutant was treated with proline, the ROS content was higher, the

246 ATP level was lower and AOX function was enhanced. ProDH was hypothesized to directly generate 247 ROS because this enzyme was shown to reduce oxygen in vitro (White et al., 2007) and its 248 overexpression increased ROS production in cancers and apoptosis (Pandhare et al., 2009; Moxley 249 et al., 2011). However, Goncalves et al. (2014) demonstrated that superoxides are not produced by 250 ProDH itself but by specific components of the electron transfer chain. ProDH/POX was found to 251 bind directly to Coenzyme Q suggesting that proline-derived electrons can directly reduce oxygen at 252 Complex III to generate superoxides (Hancock et al., 2016). On the other hand, p5cs1-4, an Arabidopsis mutant impaired in stress-induced proline accumulation, was shown to be associated 253 254 with strongly upregulated expression of a number of genes encoding NAD(P)H dehydrogenases 255 and/or related to mitochondrial respiration (Shinde et al., 2016). Similarly, Lovell et al. (2015) 256 revealed that proline concentration was strongly affected by cytoplasmic genome variation in 257 Arabidopsis mapping population in response to drought. Mitochondrial DNA polymorphisms were 258 linked to two genes coding NADH dehydrogenase subunits indicating that proline accumulation is 259 tightly regulated by cellular redox status and that proline catabolism is important in drought tolerance. 260 These results combined support the idea of a tight regulation between proline metabolism and cellular 261 redox status with mitochondria having a key role, possibly dissipating energy for drought stress tolerance (Atkin and Macherel, 2009). This would be consistent with the view that the dynamics of 262 263 proline metabolism, probably through a proline cycle between biosynthesis and catabolism, is 264 important for cell homeostasis.

265

266 The importance of the proline cycle in development and stress

267 Various environmental stresses such as drought or salt stress have been shown to trigger proline 268 accumulation by upregulating proline biosynthesis and downregulating proline catabolism (Peng et 269 al., 1996; Parre et al., 2007; Leprince et al., 2015). If ProDH is absent, the proline cycle may not be 270 active in such stress conditions, although a low level of ProDH may be sufficient to allow cycling 271 between proline and glutamate to support the maintenance of a proper $NAD(P)^+$ to NAD(P)H ratio. 272 For example, expression of ProDH2 has been shown to be triggered by salt stress whereas ProDH1 273 expression is repressed (Funck et al., 2010). Other studies have shown the importance of 274 mitochondrial proline catabolism in the regulation of cell redox status in response to drought stress 275 (Sharma *et al.*, 2011). In such stress conditions, the presence of ProDH1, including some with putative 276 post-translational modifications, was observed in western blots of protein extracts (Bhaskara et al., 2015). This discrepancy could be due to the different growth conditions and/or sensitivities of the 277 278 antibodies used, and it would be important to determine the activity of ProDH with an appropriate 279 assay to clarify this issue (Lebreton et al., 2020). In addition, transcriptome analyses have revealed 280 that *ProDH1* transcript levels oscillate during light-dark cycles, being downregulated during the day

and upregulated during the night. Drought downregulates ProDH1 transcript levels but only during the day. Interestingly, VLCA biosynthesis genes had a similar expression profile to the *ProDH1* gene expression profile (Dubois *et al.*, 2017), which corroborates the idea that proline metabolism is coordinated with VLCA biosynthesis when cellular redox status is altered.

285 Transgenic plants overexpressing P5CS1 under the control of a heat shock promoter accumulated proline in a heat-dependent manner and, interestingly, were less tolerant to heat shock. Indeed, higher 286 287 *ProDH1* expression and ROS contents were measured in these transgenic plants suggesting that the proline cycle had been stimulated (Lv et al., 2011). High levels of P5C were also detected in the yeast 288 289 put2 mutant (mutated in the yeast gene for P5CDH) (Nomura and Takagi, 2004) and the Arabidopsis 290 *p5cdh* mutant in addition to high levels of ROS upon proline treatment (Deuschle *et al.*, 2004; Miller 291 et al., 2009). These works revealed the importance of the proline cycle in maintaining cellular ROS 292 homeostasis.

The proline cycle is also involved in NADP⁺/NADPH homeostasis during plant development through a spatial and temporal compartmentalization of proline synthesis and proline degradation. Such a distribution of metabolism between different organs implies that proline as well as glutamate must be transported somehow. Proline is known to move between tissues through vascular vessels (Girousse *et al.*, 1996) and from cell to cell by plasma membrane transporters such as the amino acid/auxin permease (AAP) and proline transporters (ProT) (Schwacke *et al.*, 1999; Hirner *et al.*, 2006; Lehmann *et al.*, 2011).

300 Interestingly proline is mainly synthesized in photosynthetically active tissues, although its oxidation 301 is localized in sink non-green tissues (Sharma et al., 2011). Proline is transported from source to sink 302 tissues such as growing regions of root and shoot where the oxidation of proline will provide energy 303 to support growth. Source-sink interactions are also key determinants of leaf senescence (Dellero et 304 al., 2020). During plant senescence, specific degradation of Calvin cycle enzymes is triggered, 305 lowering the NADPH levels in chloroplasts or senescent organs. Expression of ProDH1 and to a 306 lesser extent *ProDH2* has been shown to be triggered when leaf senescence is induced by darkness, 307 leading to higher ProDH activity (Launay et al., 2019). In addition, proline was demonstrated to be 308 an alternative metabolic substrate whose oxidation fuels the electron transfer chain to generate energy 309 (Launay et al., 2019). Proline oxidation is therefore important for restoring the NADP⁺ to NADPH 310 ratio of the cell.

In another example, transport of proline was shown to be important during the formation of plant reproductive organs. Proline content was 56 times more concentrated in flowers (sink organs) than in source leaves (Schwacke et al., 1999). Proline accumulated in pollen grains accounted for 70% of the total free amino acids and was essential for pollen fertility (Mattioli *et al.*, 2018). The *LeProT1* gene, which encodes a proline transporter, is expressed in the germinating pollen tube. However, the proline 316 transport through this route is insufficient to fulfill the demand of developing microspore cells 317 (Mattioli *et al.*, 2018). The proline that accumulates in pollen derives from local synthesis inside 318 developing microspores and mature pollen grains. In this case, it was also suggested that proline 319 would have a role in supplying energy during the rapid growth of the pollen tube.

320 Proline accumulation may also be a very important mechanism to balance redox potential oscillating 321 during the day. Indeed, light is known to affect transcript levels of both proline biosynthesis and 322 degradation genes in plants (Hanson and Tully, 1979; Sanada et al., 1995). P5CS gene expression is promoted during the light period and repressed during dark periods, whereas *ProDH* transcript levels 323 324 show the opposite behavior (Hayashi et al., 2000; Dubois et al., 2017). It is conceivable that light 325 regulation also occurs at the protein level. By using a proteomic approach, Marchand et al. (2010) 326 have shown that P5CDH could be a target of thioredoxin. Thioredoxins (Trx) are ubiquitous proteins 327 catalyzing reversible disulfide-bond formation involved in enzyme redox and activity regulation. 328 Different Trx isoforms are present in mitochondria (Trx o, Trx h) and cytosol (Trx h) and the reducing 329 power for Trx reduction is provided by NADPH through NAPDH-thioredoxin reductase (NTR) 330 (Geigenberger et al., 2017). Interestingly, less proline and more glutamate and malate accumulated in trxol and ntra ntrb double mutants compared to the wild type (Daloso et al., 2015). A 30% 331 332 reduction in proline accumulation was also observed in a *trx1* mutant under low water potential 333 conditions by Verslues et al. (2014). All these data reflect a probable regulation of proline metabolic 334 enzymes by Trx, that are able to transmit the light signal from the chloroplast to other organelles like 335 mitochondria. Moreover, as the proline cycle may shuttle reducing power from cytosol to mitochondria, we can hypothesize that it also contributes to the NTR/Trx system control. Thus, the 336 337 proline cycle may be important during the night when photosynthesis is not active to generate redox 338 power and energy.

339

340 The proline-P5C cycle and its functions

341 Interestingly, when it is in equilibrium with GSA in a reversible non-enzymatic reaction, P5C is an 342 intermediate for both proline biosynthesis from either glutamate or ornithine and proline oxidation. 343 It is a product of ProDH activity in mitochondria and a substrate of P5CR in the cytosol. If P5C could 344 move back and forth between the cytosol and the mitochondrial matrix, P5CR and ProDH would then 345 be said to form a metabolic ring, the proline-P5C cycle. Notionally, the prerequisite for a functional 346 proline-P5C cycle is for both P5CR and ProDH to be expressed at the same time and for either GSA or P5C to be transported through mitochondrial membranes. The existence of this cycle in plants is 347 348 far from proven, not least because no transporter of P5C has been identified yet.

349

350 The proline-P5C cycle was first proposed by Hagedorn and Phang (1986) when they demonstrated 351 the catalytic functioning of the shuttle to generate cytosolic NADP⁺ without stoichiometric 352 consumption of proline. Proline is exported into mitochondria by a transporter whose gene has not 353 been identified yet. Proline is then oxidized to P5C, which may be transported back into the cytosol, with the help of a putative unknown transporter, to be further reduced to proline by P5CR. By this 354 route, mitochondrial P5C would replenish cytosolic proline biosynthesis and avoid at the same time 355 356 the consumption of cytosolic ATP and production of mitochondrial NADH by bypassing P5CS and P5CDH, respectively. This proline-P5C cycle would then allow NADPH to be consumed in the 357 358 cytosol, while FAD⁺ would be reduced in mitochondria. Thus, reducing power would be shuttled in 359 a single direction from the cytosol to the mitochondrial matrix in this cycle (Figure 1). The proline-P5C cycle has been proposed to maintain the redox balance between mitochondria and the cytosol. 360

In animals, the proline-P5C cycle has been shown to play important roles in the regulation of cell growth and cell death. This cycle acts by transferring the reducing power formed by the pentose phosphate pathway into mitochondria for the production of either ROS or ATP. ROS signaling mediated by ProDH was shown to trigger apoptosis, suggesting that ProDH is a mitochondrial tumor suppressor (Liu and Phang, 2012).

Miller *et al.* (2009) proposed that a proline-P5C cycle also operates in plants with P5C transport. This plant biochemical cycle may support the preferential accumulation of proline as well as the shuttling of redox equivalents from the cytosol to mitochondria. This cycle would maintain a high cellular proline to P5C ratio and direct electrons to the electron transfer chain.

Increases in ProDH expression and activity have been observed when the oxidative burst occurs during the hypersensitive response (Cecchini *et al.*, 2011; Monteoliva *et al.*, 2014). In this instance, ProDH was proposed to participate in the hypersensitive response through the proline-P5C cycle by generating ROS to trigger cell death in order to prevent pathogen spreading in plants. It is possible that proline toxicity may be due to an overflow of electrons in the mitochondrial electron transfer chain.

376 P5C may also originate from the arginine pathway. In most organisms, arginine catabolism 377 contributes to the formation of ornithine and urea through the action of arginase. In general, OAT 378 catalyzes the transfer of the δ -amino group from ornithine to KG yielding GSA/P5C and glutamate, 379 contributing to P5C input into mitochondria to serve as the substrate for P5CDH (Figure 1). In 380 intestine cells, OAT also participates in the degradation of ornithine to form citrulline, which results 381 in a consumption of P5C (Ginguay et al., 2017). In plants, a direct contribution of OAT to stress-382 induced proline accumulation is still debated. A study of salt-stressed Arabidopsis oat knockout 383 mutants provided strong evidence that OAT does not contribute to stress-induced proline 384 accumulation (Funck et al., 2008). Therefore, it is unlikely that the OAT pathway contributes to the proline-P5C cycle, at least in salt stress conditions. It has nevertheless been proposed that high concentrations of free arginine may increase the affinity of P5CDH for P5C, possibly influencing the balance between P5C export into the cytosol and its mitochondrial oxidation to glutamate (Forlani *et al.*, 2015).

389

390 **Conclusions and perspectives**

As mitochondrial membranes are not permeable to $NAD(P)^+$ or NAD(P)H, eukaryotes have evolved redox shuttles, like the G3P, malate-aspartate and malate-OAA shuttles, to allow transfer of reducing power from one compartment to another. It is not clear yet whether a malate-aspartate shuttle operates in plant cells.

395 The availability of a redox shuttle that allows redox exchange between cytosol and mitochondria is 396 of particular interest for plant cells. Proline metabolism is a unique pathway because it involves the 397 interconversion of proline and glutamate, a process linked to consumption of reducing power for 398 proline biosynthesis in the cytosol. The existence of the proline cycle allows the biosynthesis of 399 proline and concomitant oxidation of NADPH molecules in the cytosol. Furthermore, glucose 400 oxidation from the pentose phosphate pathway in the cytosol can be linked to ATP production via the 401 sequential transfer of the generated NADPH to proline, which would be then oxidized by ProDH in the mitochondrion. 402

403 When proline is transported into mitochondria, its oxidation supports oxidative phosphorylation by a 404 mechanism independent of NADH oxidation because electrons enter the electron transport system at 405 the level of the flavoenzyme ProDH, which directly feeds the mitochondrial electron transfer chain 406 and drives the synthesis of ATP. This proline cycle enables electrons from cytosolic NADPH to be 407 transported into mitochondria against the NADH concentration gradient. It is unlikely that the proline 408 cycle acts as a continuous major source of energy because firstly ProDH has to be expressed and 409 active, and secondly this cycle produces only a limited amount of the ATP produced by succinate 410 entering the tricarboxylic acid cycle. Nevertheless, there may be special conditions when this shuttle 411 acts as a transient yet significant source of energy for development or survival. The proline cycle may 412 be required when the tricarboxylic acid cycle cannot operate maximally, as postulated to occur during 413 the initiation of leaf senescence. The proposed proline cycle appears to play an important role in 414 regulating both cytosolic and mitochondrial redox states. Regulation at the level of ProDH and P5CR 415 enzymes and/or transport of proline, P5C and glutamate across mitochondrial membranes requires 416 more study, but given the influence and reach of the proline shuttle, any new findings may be relevant 417 in the study and development of resilient crops.

- 418
- 419

420

421 Acknowledgements

- 422 This work received financial support from Sorbonne Université and a PhD fellowship to Y.Z. from
- 423 the China Scholarship Council. We would like to thank the reviewers for their constructive comments
- 424 and efforts for improving our review. Members of the APCE, iEES team are gratefully acknowledged
- 425 for their stimulating discussions.
- 426

427 Author contribution

- 428 AS conceived and planned the review topic. AS wrote the manuscript with support from YZ, CC,
- 429 SP and SL. All authors approved the final version of the manuscript.
- 430

431 **Conflict of interest**

- 432 The authors declare that this research was conducted in the absence of any commercial or financial
- 433 relationships that would cause any potential conflicts of interest.
- 434
- 435

References

Akkermans ADL, Huss-Danell K, Roelofsen W. 1981. Enzymes of the tricarboxylic acid cycle and the malate-aspartate shuttle in the N2-fixing endophyte of *Alnus glutinosa*. Physiologia Plantarum **53**, 289–294.

Ansell R, Granath K, Hohmann S, Thevelein JM, Adler L. 1997. The two isoenzymes for yeast NAD⁺-dependent glycerol 3-phosphate dehydrogenase encoded by GPD1 and GPD2 have distinct roles in osmoadaptation and redox regulation. The EMBO journal **16**, 2179–2187.

Appels MA, Haaker H. 1991. Glutamate Oxaloacetate Transaminase in Pea Root Nodules : Participation in a Malate/Aspartate Shuttle between Plant and Bacteroid. Plant Physiology **95**, 740–747.

Atkin OK, Macherel D. 2009. The crucial role of plant mitochondria in orchestrating drought tolerance. Annals of Botany **103**, 581–97.

Ben Rejeb K, Abdelly C, Savouré A. 2014. How reactive oxygen species and proline face stress together. Plant Physiology and Biochemistry **80**, 278–284.

Ben Rejeb K, Lefebvre-De Vos D, Le Disquet I, Leprince A-S, Bordenave M, Maldiney R, Jdey A, Abdelly C, Savouré A. 2015. Hydrogen peroxide produced by NADPH oxidases increases proline accumulation during salt or mannitol stress in *Arabidopsis thaliana*. New Phytologist **208**, 1138–1148.

Bhaskara GB, Yang T-H, Verslues PE. 2015. Dynamic proline metabolism: importance and regulation in water limited environments. Frontiers in Plant Science **6**, 484.

Borst P. 2020. The malate-aspartate shuttle (Borst cycle): How it started and developed into a major metabolic pathway. IUBMB life **72**, 2241–2259.

Brunner G, Neupert W. 1969. Localisation of proline oxidase and Delta-pyrroline-5-carboxylic acid dehydrogenase in rat liver. FEBS letters **3**, 283–286.

Cabassa-Hourton C, Schertl P, Bordenave-Jacquemin M, *et al.* 2016. Proteomic and functional analysis of proline dehydrogenase 1 link proline catabolism to mitochondrial electron transport in Arabidopsis thaliana. The Biochemical Journal **473**, 2623–2634.

Cavero S, Vozza A, Arco AD, *et al.* 2003. Identification and metabolic role of the mitochondrial aspartate-glutamate transporter in Saccharomyces cerevisiae. Molecular Microbiology **50**, 1257–1269.

Cecchini NM, Monteoliva MI, Alvarez ME. 2011. Proline dehydrogenase contributes to pathogen defense in Arabidopsis. Plant Physiology **155**, 1947–59.

Daloso DM, Müller K, Obata T, et al. 2015. Thioredoxin, a master regulator of the tricarboxylic acid cycle in plant mitochondria. Proceedings of the National Academy of Sciences of the United States of America **112**, E1392-1400.

De Ingeniis J, Ratnikov B, Richardson AD, *et al.* 2012. Functional Specialization in Proline Biosynthesis of Melanoma. PLoS ONE **7**, e45190.

Delauney AJ, Hu CA, Kishor PB, Verma DP. 1993. Cloning of ornithine delta-aminotransferase cDNA from Vigna aconitifolia by trans-complementation in *Escherichia coli* and regulation of proline biosynthesis. The Journal of Biological Chemistry **268**, 18673–8.

Delauney AJ, Verma DPS. 1993. Proline biosynthesis and osmoregulation in plants. The Plant Journal **4**, 215–223.

Dell'Aglio E, Giustini C, Kraut A, *et al.* 2019. Identification of the Arabidopsis Calmodulin-Dependent NAD⁺ Kinase That Sustains the Elicitor-Induced Oxidative Burst. Plant Physiology **181**, 1449–1458.

Dellero Y, Clouet V, Marnet N, Pellizzaro A, Dechaumet S, Niogret M-F, Bouchereau A. 2020. Leaf status and environmental signals jointly regulate proline metabolism in winter oilseed rape. Journal of Experimental Botany **71**, 2098–2111.

Deuschle K, Funck D, Forlani G, Stransky H, Biehl A, Leister D, van der Graaff E, Kunze R, Frommer WB. 2004. The role of [Delta]1-pyrroline-5-carboxylate dehydrogenase in proline degradation. The Plant Cell 16, 3413–25.

Deuschle K, Funck D, Hellmann H, Daschner K, Binder S, Frommer WB. 2001. A nuclear gene encoding mitochondrial Delta-pyrroline-5-carboxylate dehydrogenase and its potential role in protection from proline toxicity. The Plant Journal **27**, 345–56.

Di Martino C, Pizzuto R, Pallotta ML, De Santis A, Passarella S. 2006. Mitochondrial transport in proline catabolism in plants: the existence of two separate translocators in mitochondria isolated from durum wheat seedlings. Planta, **223**, 1123–1133.

Dubois M, Claeys H, Van den Broeck L, Inzé D. 2017. Time of day determines Arabidopsis transcriptome and growth dynamics under mild drought. Plant, Cell & Environment **40**, 180–189.

El Moukhtari A, Cabassa-Hourton C, Farissi M, Savouré A. 2020. How Does Proline Treatment Promote Salt Stress Tolerance During Crop Plant Development? Frontiers in Plant Science **11**, 1127.

Elthon TE, Stewart CR. 1981. Submitochondrial Location and Electron Transport Characteristics of Enzymes Involved in Proline Oxidation. Plant Physiology 67, 780–784.

Forlani G, Bertazzini M, Zarattini M, Funck D. 2015. Functional characterization and expression analysis of rice $\delta(1)$ -pyrroline-5-carboxylate dehydrogenase provide new insight into the regulation of proline and arginine catabolism. Frontiers in Plant Science **6**, 591.

Forlani G, Trovato M, Funck D, Signorelli S. 2019. Regulation of Proline Accumulation and Its Molecular and Physiological Functions in Stress Defence. In: Hossain MA, Kumar V, Burritt DJ, Fujita M, Mäkelä PSA, eds. Osmoprotectant-Mediated Abiotic Stress Tolerance in Plants: Recent Advances and Future Perspectives. Cham: Springer International Publishing, 73–97.

Foyer CH, Noctor G. 2020. Redox Homeostasis and Signaling in a Higher-CO₂ World. Annual Review of Plant Biology **71**, 157–182.

Funck D, Baumgarten L, Stift M, von Wirén N, Schönemann L. 2020. Differential Contribution of P5CS Isoforms to Stress Tolerance in Arabidopsis. Frontiers in Plant Science **11**.

Funck D, Eckard S, Muller G. 2010. Non-redundant functions of two proline dehydrogenase isoforms in Arabidopsis. BMC Plant Biol **10**, 70.

Funck D, Stadelhofer B, Koch W. 2008. Ornithine-delta-aminotransferase is essential for arginine catabolism but not for proline biosynthesis. BMC plant biology **8**, 40.

Funck D, Winter G, Baumgarten L, Forlani G. 2012. Requirement of proline synthesis during Arabidopsis reproductive development. BMC Plant Biology **12**, 191.

Gakière B, Hao J, Bont L de, Pétriacq P, Nunes-Nesi A, Fernie AR. 2018. NAD⁺ Biosynthesis and signaling in plants. Critical Reviews in Plant Sciences **37**, 259–307.

Geigenberger P, Thormählen I, Daloso DM, Fernie AR. 2017. The Unprecedented Versatility of the Plant Thioredoxin System. Trends in Plant Science 22, 249–262.

Giberti S, Funck D, Forlani G. 2014. Δ 1-Pyrroline-5-carboxylate reductase from *Arabidopsis thaliana*: stimulation or inhibition by chloride ions and feedback regulation by proline depend on whether NADPH or NADH acts as co-substrate. The New Phytologist **202**, 911–919.

Ginguay A, Cynober L, Curis E, Nicolis I. 2017. Ornithine Aminotransferase, an Important Glutamate-Metabolizing Enzyme at the Crossroads of Multiple Metabolic Pathways. Biology **6**, 18.

Goncalves RLS, Rothschild DE, Quinlan CL, Scott GK, Benz CC, Brand MD. 2014. Sources of superoxide/H₂O₂ during mitochondrial proline oxidation. Redox Biology **2**, 901–909.

Hagedorn CH, Phang JM. 1986. Catalytic transfer of hydride ions from NADPH to oxygen by the interconversions of proline and Δ 1-pyrroline-5-carboxylate. Archives of Biochemistry and Biophysics **248**, 166–174.

Han HL, Liu J, Feng XJ, Zhang M, Lin QF, Wang T, Qi SL, Xu T, Hua XJ. 2021. SSR1 is involved in maintaining the function of mitochondria electron transport chain and iron homeostasis upon proline treatment in Arabidopsis. Journal of Plant Physiology **256**, 153325.

Hancock CN, Liu W, Alvord WG, Phang JM. 2016. Co-regulation of mitochondrial respiration by proline dehydrogenase/oxidase and succinate. Amino Acids **48**, 859–872.

Hanson A, Tully R, 1979. Light stimulation of proline synthesis in water-stressed barley leaves. Planta 145, 45-51.

Hayashi F, Ichino T, Osanai M, Wada K. 2000. Oscillation and Regulation of Proline Content by P5CS and ProDH Gene Expressions in the Light/Dark Cycles in *Arabidopsis thaliana* L. Plant and Cell Physiology **41**, 1096–1101.

Hirner A, Ladwig F, Stransky H, Okumoto S, Keinath M, Harms A, Frommer WB, Koch W. 2006. Arabidopsis LHT1 Is a High-Affinity Transporter for Cellular Amino Acid Uptake in Both Root Epidermis and Leaf Mesophyll. The Plant Cell **18**, 1931–1946.

Huang S, Taylor NL, Ströher E, Fenske R, Millar AH. 2013. Succinate dehydrogenase assembly factor 2 is needed for assembly and activity of mitochondrial complex II and for normal root elongation in Arabidopsis. The Plant Journal **73**, 429–441.

Kishor PBK, Sreenivasulu N. 2014. Is proline accumulation per se correlated with stress tolerance or is proline homeostasis a more critical issue? Plant, Cell & Environment **37**, 300–311.

Krieger-Liszkay A, Krupinska K, Shimakawa G. 2019. The impact of photosynthesis on initiation of leaf senescence. Physiologia Plantarum 166, 148–164.

Krishnan N, Dickman MB, Becker DF. 2008. Proline modulates the intracellular redox environment and protects mammalian cells against oxidative stress. Free Radical Biology & Medicine 44, 671–681.

LaNoue KF, Bryla J, Bassett DJ. 1974*a*. Energy-driven aspartate efflux from heart and liver mitochondria. The Journal of Biological Chemistry **249**, 7514–7521.

LaNoue KF, Meijer AJ, Brouwer A. 1974*b*. Evidence for electrogenic aspartate transport in rat liver mitochondria. Archives of Biochemistry and Biophysics **161**, 544–550.

Launay A, Cabassa-Hourton C, Eubel H, *et al.* 2019. Proline oxidation fuels mitochondrial respiration during dark-induced leaf senescence in *Arabidopsis thaliana*. Journal of Experimental Botany. **70**, 6203–6214.

Lebreton S, Cabassa-Hourton C, Savouré A, Funck D, Forlani G. 2020. Appropriate Activity Assays Are Crucial for the Specific Determination of Proline Dehydrogenase and Pyrroline-5-Carboxylate Reductase Activities. Frontiers in Plant Science **11**, 602939.

Lehmann S, Gumy C, Blatter E, Boeffel S, Fricke W, Rentsch D. 2011. In planta function of compatible solute transporters of the AtProT family. Journal of Experimental Botany **62**, 787–796.

Leprince A-S, Magalhaes N, De Vos D, Bordenave M, Crilat E, Clément G, Meyer C, Munnik T, Savouré A. 2015. Involvement of Phosphatidylinositol 3-kinase in the regulation of proline catabolism in Arabidopsis thaliana. Frontiers in Plant Science 5.

Liang X, Zhang L, Natarajan SK, Becker DF. 2013. Proline mechanisms of stress survival. Antioxidants & Redox Signaling 19, 998–1011.

Liu W, Phang JM. 2012. Proline dehydrogenase (oxidase), a mitochondrial tumor suppressor, and autophagy under the hypoxia microenvironment. Autophagy **8**, 1407–1409.

Lovell JT, Mullen JL, Lowry DB, Awole K, Richards JH, Sen S, Verslues PE, Juenger TE, McKay JK. 2015. Exploiting Differential Gene Expression and Epistasis to Discover Candidate Genes for Drought-Associated QTLs in Arabidopsis thaliana. The Plant Cell **27**, 969–983.

Lunn JE. 2006. Compartmentation in plant metabolism. Journal of Experimental Botany 58, 35-47.

Lv W-T, Lin B, Zhang M, Hua X-J. 2011. Proline Accumulation Is Inhibitory to Arabidopsis Seedlings during Heat Stress. Plant Physiology **156**, 1921–1933.

Mani S, Van De Cotte B, Van Montagu M, Verbruggen N. 2002. Altered levels of proline dehydrogenase cause hypersensitivity to proline and its analogs in Arabidopsis. Plant Physiology **128**, 73–83.

Marchand CH, Vanacker H, Collin V, Issakidis-Bourguet E, Maréchal PL, Decottignies P. 2010. Thioredoxin targets in Arabidopsis roots. Proteomics **10**, 2418–2428.

Mattioli R, Biancucci M, El Shall A, Mosca L, Costantino P, Funck D, Trovato M. 2018. Proline synthesis in developing microspores is required for pollen development and fertility. BMC Plant Biology 18, 356.

Marchese L, Olavarria K, Mantilla BS, Avila CC, Souza ROO, Damasceno FS, Elias MC, Silber AM. 2020. Trypanosoma cruzi synthesizes proline via a Δ 1-pyrroline-5-carboxylate reductase whose activity is fine-tuned by NADPH cytosolic pools. The Biochemical Journal **477**, 1827–1845.

Maxwell SA, Davis GE. 2000. Differential gene expression in p53-mediated apoptosis-resistant vs. apoptosis-sensitive tumor cell lines. Proceedings of the National Academy of Sciences of the United States of America **97**, 13009–13014.

McKenna MC, Waagepetersen HS, Schousboe A, Sonnewald U. 2006. Neuronal and astrocytic shuttle mechanisms for cytosolic-mitochondrial transfer of reducing equivalents: Current evidence and pharmacological tools. Biochemical Pharmacology **71**, 399–407.

Mettler IJ, Beevers H. 1980. Oxidation of NADH in Glyoxysomes by a Malate-Aspartate Shuttle. Plant Physiology **66**, 555–560.

Miller G, Honig A, Stein H, Suzuki N, Mittler R, Zilberstein A. 2009. Unraveling delta1pyrroline-5-carboxylate-proline cycle in plants by uncoupled expression of proline oxidation enzymes. The Journal of Biological Chemistry **284**, 26482–26492.

Monné M, Daddabbo L, Gagneul D, Obata T, Hielscher B, Palmieri L, Miniero DV, Fernie AR, Weber APM, Palmieri F. 2018. Uncoupling proteins 1 and 2 (UCP1 and UCP2) from *Arabidopsis thaliana* are mitochondrial transporters of aspartate, glutamate, and dicarboxylates. The Journal of Biological Chemistry **293**, 4213–4227.

Monteoliva MI, Rizzi YS, Cecchini NM, Hajirezaei M-R, Alvarez ME. 2014. Context of action of proline dehydrogenase (ProDH) in the Hypersensitive Response of Arabidopsis. BMC Plant Biology 14, 21.

Moxley MA, Tanner JJ, Becker DF. 2011. Steady-state kinetic mechanism of the proline:ubiquinone oxidoreductase activity of proline utilization A (PutA) from *Escherichia coli*. Archives of Biochemistry and Biophysics **516**, 113–120.

Mráček T, Drahota Z, Houštěk J. 2013. The function and the role of the mitochondrial glycerol-3phosphate dehydrogenase in mammalian tissues. Biochimica et Biophysica Acta (BBA) -Bioenergetics 1827, 401–410.

Nanjo T, Fujita M, Seki M, Kato T, Tabata S, Shinozaki K. 2003. Toxicity of free proline revealed in an Arabidopsis T-DNA-tagged mutant deficient in proline dehydrogenase. Plant & Cell Physiology 44, 541–548.

Nomura M, Takagi H. 2004. Role of the yeast acetyltransferase Mpr1 in oxidative stress: regulation of oxygen reactive species caused by a toxic proline catabolism intermediate. Proceedings of the National Academy of Sciences U S A **101**, 12616–21.

Ohashi K, Kawai S, Murata K. 2012. Identification and characterization of a human mitochondrial NAD kinase. Nature Communications **3**, 1248.

Outten CE, Culotta VC. 2003. A novel NADH kinase is the mitochondrial source of NADPH in Saccharomyces cerevisiae. The EMBO journal **22**, 2015–2024.

Palmieri F, Rieder B, Ventrella A, *et al.* 2009. Molecular Identification and Functional Characterization of Arabidopsis thaliana Mitochondrial and Chloroplastic NAD+ Carrier Proteins. The Journal of Biological Chemistry **284**, 31249–31259.

Pandhare J, Donald SP, Cooper SK, Phang JM. 2009. Regulation and function of proline oxidase under nutrient stress. Journal of Cellular Biochemistry **107**, 759–768.

Parre E, Ghars MA, Leprince AS, Thiery L, Lefebvre D, Bordenave M, Richard L, Mazars C, Abdelly C, Savouré A. 2007. Calcium signaling via phospholipase C is essential for proline accumulation upon ionic but not nonionic hyperosmotic stresses in Arabidopsis. Plant Physioly **144**, 503–12.

Peng Z, Lu Q, Verma DP. 1996. Reciprocal regulation of delta 1-pyrroline-5-carboxylate synthetase and proline dehydrogenase genes controls proline levels during and after osmotic stress in plants. Molecular Genetics and Genomic **253**, 334–41.

Phang JM. 2019. Proline Metabolism in Cell Regulation and Cancer Biology: Recent Advances and Hypotheses. Antioxidants & Redox Signaling **30**, 635–649.

Porcelli V, Vozza A, Calcagnile V, Gorgoglione R, Arrigoni R, Fontanesi F, Marobbio CMT, Castegna A, Palmieri F, Palmieri L. 2018. Molecular identification and functional characterization of a novel glutamate transporter in yeast and plant mitochondria. Biochimica et Biophysica Acta (BBA) - Bioenergetics 1859, 1249–1258.

Quettier A-L, Shaw E, Eastmond PJ. 2008. SUGAR-DEPENDENT6 Encodes a Mitochondrial Flavin Adenine Dinucleotide-Dependent Glycerol-3-P Dehydrogenase, Which Is Required for Glycerol Catabolism and Postgerminative Seedling Growth in Arabidopsis. Plant Physiology 148, 519–528.

Ren Y, Miao M, Meng Y, Cao J, Fan T, Yue J, Xiao F, Liu Y, Cao S. 2018. DFR1-Mediated Inhibition of Proline Degradation Pathway Regulates Drought and Freezing Tolerance in Arabidopsis. Cell Reports 23, 3960–3974.

Rizzi YS, Cecchini NM, Fabro G, Alvarez ME. 2016. Differential control and function of Arabidopsis ProDH1 and ProDH2 genes on infection with biotrophic and necrotrophic pathogens. Molecular Plant Pathology **18**, 1164–1174

da Rocha IMA, Vitorello VA, Silva JS, Ferreira-Silva SL, Viégas RA, Silva EN, Silveira JAG. 2012. Exogenous ornithine is an effective precursor and the δ -ornithine amino transferase pathway contributes to proline accumulation under high N recycling in salt-stressed cashew leaves. Journal of Plant Physiology **169**, 41–49.

Roosens NH, Thu TT, Iskandar HM, Jacobs M. 1998. Isolation of the ornithine-deltaaminotransferase cDNA and effect of salt stress on its expression in *Arabidopsis thaliana*. Plant Physiology **117**, 263–71.

Ruszkowski M, Nocek B, Forlani G, Dauter Z. 2015. The structure of *Medicago truncatula* δ 1-pyrroline-5-carboxylate reductase provides new insights into regulation of proline biosynthesis in plants. Frontiers in Plant Science **6**, 869.

Sanada Y, Ueda H, Kuribayashi K, Andoh T, Hayashi F, Tamai N, Wada K. 1995. Novel lightdark change of proline levels in halophyte (*Mesembryanthemum crystallinum* L.) and Glycophytes (*Hordeum vulgare* L. and *Triticum aestivum* L.) leaves and roots under salt stress. Plant Cell Physiology 36, 965-970

Satrústegui J, Bak LK. 2015. Fluctuations in Cytosolic Calcium Regulate the Neuronal Malate-Aspartate NADH Shuttle: Implications for Neuronal Energy Metabolism. Neurochemical Research 40, 2425–2430.

Scheibe R, Backhausen JE, Emmerlich V, Holtgrefe S. 2005. Strategies to maintain redox homeostasis during photosynthesis under changing conditions. Journal of Experimental Botany 56, 1481–1489.

Schwacke R, Grallath S, Breitkreuz KE, Stransky E, Stransky H, Frommer WB, Rentsch D. 1999. LeProT1, a Transporter for Proline, Glycine Betaine, and γ-Amino Butyric Acid in Tomato Pollen. The Plant Cell **11**, 377–391. **Schwörer S, Berisa M, Violante S, Qin W, Zhu J, Hendrickson RC, Cross JR, Thompson CB**. 2020. Proline biosynthesis is a vent for TGFβ-induced mitochondrial redox stress. The EMBO Journal **39**, e103334.

Selinski J, Scheibe R. 2019. Malate valves: old shuttles with new perspectives. Plant Biology 21 Suppl 1, 21–30.

Senthil-Kumar M, Mysore KS. 2012. Ornithine-delta-aminotransferase and proline dehydrogenase genes play a role in non-host disease resistance by regulating pyrroline-5-carboxylate metabolism-induced hypersensitive response. Plant, Cell & Environment **35**, 1329–1343.

Servet C, Ghelis T, Richard L, Zilberstein A, Savouré A. 2012. Proline dehydrogenase: a key enzyme in controlling cellular homeostasis. Frontiers In Bioscience **17**, 607–620.

Sharma S, Villamor JG, Verslues PE. 2011. Essential role of tissue-specific proline synthesis and catabolism in growth and redox balance at low water potential. Plant Physiology **157**, 292–304.

Shen W, Wei Y, Dauk M, Tan Y, Taylor DC, Selvaraj G, Zou J. 2006. Involvement of a Glycerol-3-Phosphate Dehydrogenase in Modulating the NADH/NAD+ Ratio Provides Evidence of a Mitochondrial Glycerol-3-Phosphate Shuttle in Arabidopsis. The Plant Cell **18**, 422–441.

Shen W, Wei Y, Dauk M, Zheng Z, Zou J. 2003. Identification of a mitochondrial glycerol-3-phosphate dehydrogenase from *Arabidopsis thaliana*: evidence for a mitochondrial glycerol-3-phosphate shuttle in plants. FEBS Letters **536**, 92–96.

Shinde S, Villamor JG, Lin W, Sharma S, Verslues PE. 2016. Proline Coordination with Fatty Acid Synthesis and Redox Metabolism of Chloroplast and Mitochondria. Plant Physiology **172**, 1074–1088.

Slama I, Abdelly C, Bouchereau A, Flowers T, Savouré A. 2015. Diversity, distribution and roles of osmoprotective compounds accumulated in halophytes under abiotic stress. Annals of Botany **115**, 433–447.

de Souza Chaves I, Feitosa-Araújo E, Florian A, *et al.* 2019. The mitochondrial NAD+ transporter (NDT1) plays important roles in cellular NAD⁺ homeostasis in *Arabidopsis thaliana*. The Plant Journal **100**, 487–504.

Szabados L, Savouré A. 2010. Proline: a multifunctional amino acid. Trends in Plant Science 15, 89–97.

Székely G, Abraham E, Cseplo A, *et al.* 2008. Duplicated P5CS genes of Arabidopsis play distinct roles in stress regulation and developmental control of proline biosynthesis. The Plant Journal **53**, 11–28.

Trovato M, Forlani G, Signorelli S, Funck D. 2019. Proline Metabolism and Its Functions in Development and Stress Tolerance. In: Hossain MA, Kumar V, Burritt DJ, Fujita M, Mäkelä PSA,

eds. Osmoprotectant-Mediated Abiotic Stress Tolerance in Plants: Recent Advances and Future Perspectives. Cham: Springer International Publishing, 41-72.

Verslues PE, Lasky JR, Juenger TE, Liu T-W, Kumar MN. 2014. Genome-wide association mapping combined with reverse genetics identifies new effectors of low water potential-induced proline accumulation in Arabidopsis. Plant Physiology **164**, 144-159.

White TA, Krishnan N, Becker DF, Tanner JJ. 2007. Structure and kinetics of monofunctional proline dehydrogenase from *Thermus thermophilus*. The Journal of Biological Chemistry **282**, 14316–14327.

You J, Hu H, Xiong L. 2012. An ornithine δ -aminotransferase gene OsOAT confers drought and oxidative stress tolerance in rice. Plant Science **197**, 59–69.

Zhang L, Becker DF. 2015. Connecting proline metabolism and signaling pathways in plant senescence. Frontiers in Plant Science 6, 552.

Zhao Y, Yu H, Zhou J-M, Smith SM, Li J. 2020. Malate Circulation: Linking Chloroplast Metabolism to Mitochondrial ROS. Trends in Plant Science **25**, 446–454.

Figure legends

Fig. 1. Proline metabolism in plants and its links to the ornithine and pentose phosphate pathways. The proline cycle is highlighted by blue arrows and the putative proline-P5C cycle by a square in light blue. Asp, aspartate; BAC, basic amino acid carrier; BOU, a bout de souffle; ETC, electron transfer chain; Glu, glutamate; KG, α -ketoglutarate; OAT, ornithine δ - aminotransferase; Orn, ornithine; PPP shunt, pentose phosphate pathway; P5C, pyrroline-5-carboxyate; P5CDH, P5C dehydrogenase; P5CS, P5C synthetase; ProDH, proline dehydrogenase; Pro, proline; UCP, mitochondrial uncoupling protein; ? indicates unknown transporter(s).

Fig. 2. The glycerol-3-phosphate (G3P) (A) and the malate–aspartate (B) shuttles for transporting reducing equivalents from cytosolic NADH into the mitochondrial matrix or mitochondrial oxidative phosphorylation pathway respectively. AGC, aspartate-glutamate carrier; Asp, aspartate; DHAP, dihydroxyacetone phosphate; GPDH, glycerol-3-phosphate dehydrogenase; G3P, glycerol-3-phosphate; Glu, glutamate; GOT, glutamate oxaloacetate aminotransferase; IMS, intermembrane space; KG, α -ketoglutarate; Mal, malate; MDH, malate dehydrogenase; OAA, oxaloacetate; OGC, KG-malate carrier.

Fig. 3. Malate-oxaloacetate shuttle, or the malate valve, involves the interconversion of malate and oxaloacetate by malate dehydrogenase (MDH). Isoenzymes of MDH are present in each cellular compartment, the cytosol, chloroplasts and mitochondria. MDH, malate dehydrogenase; OAA,



Fig. 1. Proline metabolism in plants and its links to the ornithine and pentose phosphate pathways. The proline cycle is highlighted by blue arrows and the putative proline-P5C cycle by a square in light blue. Asp, aspartate; BAC, basic amino acid carrier; BOU, a bout de souffle; ETC, electron transfer chain; Glu, glutamate; KG, α -ketoglutarate; OAT, ornithine δ - aminotransferase; Orn, ornithine; PPP shunt, pentose phosphate pathway; P5C, pyrroline-5-carboxyate; P5CDH, P5C dehydrogenase; P5CS, P5C synthetase; ProDH, proline dehydrogenase; Pro, proline; UCP, mitochondrial uncoupling protein; ? indicates unknown transporter(s).

oxaloacetate; MOT, malate- α -ketoglutarate transporter.



Fig. 2. The glycerol-3-phosphate (G3P) (A) and the malate–aspartate (B) shuttles for transporting reducing equivalents from cytosolic NADH into the mitochondrial matrix or mitochondrial oxidative phosphorylation pathway respectively. AGC, aspartate-glutamate carrier; Asp, aspartate; DHAP, dihydroxyacetone phosphate; GPDH, glycerol-3-phosphate dehydrogenase; G3P, glycerol-3-phosphate; Glu, glutamate; GOT, glutamate oxaloacetate aminotransferase; IMS, intermembrane space; KG, α -ketoglutarate; Mal, malate; MDH, malate dehydrogenase; OAA, oxaloacetate; OGC, KG-malate carrier.



Fig. 3. Malate-oxaloacetate shuttle, or the malate valve, involves the interconversion of malate and oxaloacetate by malate dehydrogenase (MDH). Isoenzymes of MDH are present in each cellular compartment, the cytosol, chloroplasts and mitochondria. MDH, malate dehydrogenase; OAA, oxaloacetate; MOT, malate- α -ketoglutarate transporter.