Miniaturization of an extraction protocol for the monitoring of pesticides and polar transformation products in biotic matrices

Vincent Dufour, Laure Wiest, Sylvain Slaby, François Le Cor, Lucile Auger, Olivier Cardoso, Laurence Curtet, Laure Pasquini, Xavier Dauchy, Emmanuelle Vullet, et al.

► To cite this version:

Vincent Dufour, Laure Wiest, Sylvain Slaby, François Le Cor, Lucile Auger, et al.. Miniaturization of an extraction protocol for the monitoring of pesticides and polar transformation products in biotic matrices. Chemosphere, 2021, 284, pp.131292. 10.1016/j.chemosphere.2021.131292 . hal-03364505

HAL Id: hal-03364505

<https://hal.science/hal-03364505>

Submitted on 13 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Miniaturization of an extraction protocol for the monitoring of pesticides and polar transformation products in biotic matrices

Vincent Dufour ^{ab}, Laure Wiest ^b, Sylvain Slaby ^a, François Le Cor ^{ac}, Lucile Auger ^{ad}, Olivier Cardoso ^e, Laurence Curtet ^d, Laure Pasquini ^c, Xavier Dauchy ^c, Emmanuelle Vulliet ^b, Damien Banas ^a

^a Université de Lorraine, INRAE, URAFPA, F-54000, Nancy, France

^b Univ Lyon, CNRS, Université Claude Bernard Lyon 1, Institut des Sciences Analytiques, UMR 5280, 5 rue de la Doua, F-69100, Villeurbanne, France

^c LHN, Laboratoire d'Hydrologie de Nancy, ANSES, 40 rue Lionnois, F-54000, Nancy, France

^d Office Français de la Biodiversité, Montfort, F-01330, Birieux, France

^e Office Français de la Biodiversité, 9 Avenue Buffon, F-45071, Orléans, France

1. Introduction

Pesticides, or plant protection products (PPPs), are widely used across the world to increase agricultural yields. As only some of the applied dose reaches target organisms, a significant amount of active substances ends up in the environment, depending on the PPPs, environmental conditions, and agricultural practices (Gaillard et al. 2016b). Aquatic ecosystems play the role of final receptacles for organic micropollutants, with concentration levels ranging from ultra-traces to $\mu\text{g.L}^{-1}$ (Gaillard et al. 2016a; Nowell et al. 2018). Gaillard et al. (2016a) estimated PPP concentrations of up to $20 \mu\text{g.L}^{-1}$ in streams located in headwater agricultural catchments. Moreover, PPPs can affect the health of wild organisms such as macroinvertebrates (MIs) (Macchi et al. 2018), amphibians (Slaby et al. 2019), fish (Clasen et al. 2018), and birds (Stanton et al. 2018; Brain and Anderson 2019), and biodiversity in general (McLaughlin and Mineau 1995; Geiger et al. 2010). Bioaccumulation and bioamplification along the trophic chain are of particular concern as they may result in increased contamination levels in top predators (Konwick et al. 2006; Rostron 2010). To overcome these phenomena, more polar PPPs have been synthesized. However, they can reach aquatic ecosystems and contaminate wild organisms even more easily because of their intrinsic mobility. In addition,

16 PPPs can undergo biotic and abiotic processes in the environment that lead to the generation of
17 transformation products (TPs) being usually more stable, polar, and mobile (Reemtsma et al. 2016),
18 and sometimes having more effects on organisms (Sinclair and Boxall 2003; Tousova et al. 2017)
19 than their parent compounds. There is a current gap in knowledge on these polar mobile organic
20 chemicals (PMOCs) since, compared to historic PPPs, their nature and dynamics in the environment
21 are poorly known (Reemtsma et al. 2016) despite their potential toxicity at low concentrations.
22 Filling this gap would require the use of specific analytical tools and method such as ion
23 chromatography, that are usually not compatible with multi-residues analysis of medium to non-
24 polar compounds.

25 The QuEChERS (quick, easy, cheap, effective, rugged, and safe) method (Anastassiades et al. 2003)
26 is one of the most universal extraction approach for multi-residue analysis of pesticides, including
27 polar pesticides. It usually ensures good extraction rates (ERs) for a large number of substances
28 (Knoll et al. 2020). This method was successfully applied to solid matrices with different levels of
29 complexity such as fruit and vegetables (Lehotay et al. 2010; Koesukwiwat et al. 2010), meat and
30 fish (Lazartigues et al. 2011; Lichtmannegger et al. 2015; Oliveira et al. 2018), or food from animal
31 origin (Giroud et al. 2013; Golge et al. 2018; Song et al. 2019). However, owing to their nature, more
32 polar substances such as TPs present other analytical challenges. Indeed, they are more soluble in the
33 aqueous phase than in acetonitrile, even if the latter is saturated with buffer, resulting in low or no
34 recoveries. (Dufour et al. 2020).

35 Another disadvantage of most current extraction methods is the use of high sample masses, usually
36 greater than 1 g, that can be very difficult to obtain for many environmental matrices. This hinders
37 the study of PPP transfers in food webs for two reasons: (1) it may not be possible to collect enough
38 biomass to analyze pesticides in smaller organisms such as insects or small fish (Knoll et al. 2020);
39 or (2) individuals must be pooled, which may lead to less accurate results (Roche et al. 2009). This is
40 typically the case for MIs (e.g., insect or crustacean larvae), which have small individual masses but

41 play a key role in trophic chains by ingesting plant matter and then being consumed by predators
42 (Four et al. 2019). Developing analytical methods to quantify the contamination of this trophic level
43 is a major challenge for modeling PPP transfers in ecosystems. Moreover, it may also represent an
44 interesting tool for the monitoring of marine mammals, reptiles and apex predators from a biopsy,
45 avoiding euthanasia or invasive surgery to obtain sufficient material for analysis, which is an actual
46 need concerning the assessment of micropollutants in marine wildlife (Sanganyado et al., 2020).
47 Providing a robust, reliable, and sensitive analytical method based on low matrix masses could
48 increase the number of studies in that research field and benefit the understanding of the global
49 dynamics of organic micropollutants in the environment. In this context, the aim of this work was to
50 study the effect of the miniaturization of a previously developed extraction protocol (Dufour et al.,
51 2020) on analytical performance and to test its applicability on two matrices: fish and MIs. As
52 masses of MI samples are very low, the effect of miniaturization was first investigated on fish, and
53 then applied and validated on MIs. Special attention was paid to the validation phase in order to
54 guarantee the reliability and accuracy of the method as it is well known that decreasing the reduction
55 of the sample mass increases the variability of the measurement (Han et al., 2018; Lehotay et al.,
56 2018).

57

58 **2. Materials and methods**

59 *2.1. Standards and reagents*

60 Ultrapure water (Fisher Chemical, Geel, BE), acetonitrile (ACN) (Honeywell, Seelze, DE), and
61 heptane (Merck, Darmstadt, DE) were of LC-MS quality. Magnesium sulfate (MgSO_4) was
62 purchased from Sigma-Aldrich (Saint Louis, USA). Formic acid (FA) and ammonium sulfate were
63 purchased from Biosolve-chemicals (Dieuze, FR). Fifty-milliliter polypropylene metal-free
64 centrifuge tubes were purchased from VWR (Fontenay-sous-Bois, FR) and 2 mL polypropylene
65 safe-lock centrifuge tubes were obtained from Eppendorf (Montesson, FR). Forty-one analytical

standards and 15 isotopically labeled standards (Table S1) were purchased from A2S (Martignas-sur-Jalle, FR), Honeywell (Seelze, DE), HPC standards (Cunnersdorf, DE), LGC group (Teddington, UK), Sigma-Aldrich (Saint-Louis, USA), and TCI Europe (Anvers, BE). All were of high purity (> 96%) with most > 99%. Individual solutions made in acetonitrile (100 mg.L⁻¹) were stored at -18°C. They were mixed to obtain a 5 µg.L⁻¹ solution used to develop and validate the extraction protocol. Mix solutions were regularly characterized to check their stability over time and data were compiled in a control chart. Stored at -18°C, they were stable for at least 6 months.

2.2. Sampling strategy and pre-treatment

Young tench (*Tinca tinca*) (n = 13) were provided by the experimental platform of aquaculture (URAFPA, University of Lorraine) and were used as a blank reference matrix for fish. The MI reference matrix consisted in chironomid larvae reared under laboratory conditions, kindly provided by INRAE (ECOTOX team, Centre Lyon-Villeurbanne, FR).

For this purpose, fish and MIs were both sampled in November 2018 in two fishponds of the Dombes region (FR), over 2 distinct days. Those fishponds have watershed strongly influenced by conventional farming with the presence of either grassland or maize, wheat or rape crops. Fish and MI are expected to be exposed to diverse pesticides at important levels, and more especially herbicides like metazachlor and metolachlor, which are often applied to this type of crop, as well as to their TPs. Three fish species were selected for their representativeness of the local production: carp (*Cyprinus carpio*, n = 6), roach (*Rutilus rutilus*, n = 6), and rudd (*Scardinius erythrophthalmus*, n = 5). On average (± SEM), they weighed 247.5 ± 11.2, 58.3 ± 3.5, and 37.6 ± 12.2 g wet weight (ww), respectively. Concerning MIs, organisms were pooled to constitute 19 samples according to taxonomic classification. They were sorted into Anisoptera (n = 2), Chironomidae (n = 5), Corixidae (n = 5), Daphniidae (n = 2), Dystiscidae (n = 1), and Physidae (n = 4) groups. All blank reference matrices and samples (whole organisms) were frozen at -20°C in aluminum tray. Then, they were

freeze-dried (Reacti-Vap, PIERCE, Bellafonte, USA), ground, and stored at room temperature until extraction.

2.3. Initial protocol

The initial protocol (IP), based on a previous work by Dufour et al. (2020) and designed for the analysis of pesticides and their degradation products in high-mass environmental samples (e.g., eggs), is illustrated in Fig. 1. Briefly, 500 mg of sample was placed in a 50 mL centrifuge tube and spiked with 50 μL of a 500 $\mu\text{g.L}^{-1}$ mix solution containing all standards to constitute artificial samples. The solvent of the solution was evaporated under gentle nitrogen flux for 10 min and the spiked matrix was mixed with 5 mL of LC-MS water at 16 Hz for 10 min (SamplePrep 2010 Geno/Grinder from SPEX, Costa Mesa, USA). Then, 5 mL of heptane and 10 mL of ACN (+ 0.2% FA) were added and the whole sample was mixed again at 16 Hz for 10 min before centrifugation at 9,500 RCF (Relative Centrifugal Force) at 20°C for 5 min. Six milliliters of the lower layer (ACN/water mix) was dehydrated over 2.5 g of MgSO_4 in a 50 mL centrifuge tube and manually mixed before being centrifuged again for 2 min (20°C, 9,500 RCF). The final extract was diluted 10 times in ultrapure water + 0.1% FA before injection in LC-MS/MS.

2.4. Miniaturized protocol

In order to make the analysis of small organisms (e.g., insect larvae, small fish) possible, a miniaturized protocol (MP) was developed (Fig. 1). For this MP, a 2 mL polypropylene centrifuge tube was filled with 30 mg of matrix spiked with 50 μL of a mix solution containing all analytes (30 $\mu\text{g.L}^{-1}$) to constitute artificial samples (concentrations and volumes used specifically for validation are given in section 2.6). Before extraction, the added solvent was evaporated under gentle nitrogen flux for 10 min. Next, 0.5 mL of LC-MS water was added, and the whole sample was shaken for 10 min at 16 Hz using Geno/Grinder. The tube was then completed with 1 mL of ACN +

0.2% FA and 0.5 mL of heptane before being shaken again for 10 min at 16 Hz. It was centrifuged for 5 min at 20°C and at 9,500 RCF. Then, 0.6 mL of the lower layer (ACN/water mix) was sampled and dehydrated in a new centrifuge tube containing 0.3 g of MgSO₄ and 3 zirconium oxide 3 mm grinding balls (Fisher Scientific, Illkirch, FR). Grinding balls were used to break up crystals formed in these small tubes, which slow down the dehydration process and lead to poor recoveries for the most polar analytes (e.g. -ESA or -OXA TPs). The tube was shaken for 1 min at 16 Hz to favor the reaction and was centrifuged again at 9,500 RCF and 20°C for 2 min. Finally, the extract was diluted 10 times in ultrapure water + 0.1% FA before injection in LC-MS/MS.

124

125 2.5. LC-MS/MS

The development of the LC-MS/MS method was described in Dufour et al. (2020). Analysis of the 41 PPP residues was performed using a 1290 Infinity UPLC system (Agilent Technologies, Massy, FR) coupled with a 5500 QTRAP mass spectrometer (Sciex, Villebon-sur-Yvette, FR) in both positive and negative modes using electrospray ionization. Forty microliters of sample was injected, and separation was performed at 0.4 mL.min⁻¹ and at 50°C on an XSelect® HSS T3 column (2.1 × 150 mm, 2.5 µm, Waters, Saint Quentin-en-Yvelines, FR). The column was protected with a KrudKatcher Ultra filter (Phenomenex, Le Pecq, FR) and an XSelect® HSS T3 precolumn (2.1 × 5 mm, 2.5 µm, Waters, Saint Quentin-en-Yvelines, FR). Ultrapure water + 0.1% FA (A) and ACN + 0.1% FA (B) were used as mobile phases in positive mode, while 1 mM ammonium acetate in ultrapure water (A) and ACN (B) were used in negative mode. The gradient was the same for both methods: it started with 10% B for 1 min, increased to 100% B over 7 min and held for 6 min, then decreased back to 10% B over 1 min and held for 4 min. The temperatures of the mass spectrometer source, the nebulizer gas pressure, and heating gas pressure were set at 550°C, 40 psi, and 50 psi, respectively. Ion spray voltages were 5500 V for positive ionization and -4500 V for negative ionization. Analytes and analytical parameters are available in Table S1.

2.6. Method validation

The MP was validated for the 41 molecules (21 PPPs and 20 TPs) for the fish matrix, using matrix-matched calibration and following European guidance recommendations (European Commission DG-SANTE 2019). The reference matrix spiked at 6 different concentration levels, namely 1 (C1), 2 (C2), 5 (C3), 10 (C4), 15 (C5), and 20 (C6) times the initial concentration (C1) (available in Table 1), and a protocol blank (C0), made up with reference matrix only, were extracted 5 times following the whole MP to validate response linearity. The matrix was spiked at C2, C4, and C6 before extraction and the corresponding areas were compared to the matrix spiked after extraction to calculate ERs. The areas in the matrix spiked after extraction were also compared to those of standard solutions of equivalent concentrations to calculate matrix effects (MEs). ER and ME calculations are available in Supplementary information (SI). Inter-day precision was determined over 5 days with the daily extraction in triplicate of one of the following concentration levels: C2 (n = 3), C4 (n = 2), and C6 (n = 3).

Because only low masses of the reference MI matrix were available, the validation of the method for MIs could not follow European guidelines (European Commission DG-SANTE 2019). Two triplicates of matrix spiked at C4 and 2 protocol blanks were extracted on 2 different days to assess protocol performance on this specific matrix. The limits of quantification (LOQs) were estimated to be 10 times the signal-to-noise ratio on C4 samples. No ER estimation could be performed but recoveries (RECs, Eq. 3 in SI), which reflect the trueness of the method, were calculated instead for each molecule on the C4 samples (n = 6), using matrix-matched calibration (C1 to C6). Chromatograms of both reference matrices spiked with monitored molecules at concentration C6 are available in SI (Fig. S1).

2.7. Method application

166 The developed miniaturized method was applied to 17 fish from 3 different species and 19 MI
167 samples from 6 different taxa (see Section 2.2 for details) to characterize contamination in field
168 samples. Following European guidelines (European Commission DG-SANTE 2019), quantification
169 was performed with a 6-point matrix-matched calibration curve (described in Section 2.6). A
170 protocol blank and a reference matrix spiked at C2 for fish or C4 for MIs were extracted with each
171 sample series to avoid false positives and control RECs of each extraction series. When protocol
172 blanks showed contamination, the LOQs were increased to 3 times their quantified concentrations to
173 avoid false positives.

174

175 *2.8. Statistical analysis*

176 Isotope-labeled standards (D11-acetochlor, D5-acetochlor-ESA, D5-atrazine, D5-atrazine-2-
177 hydroxy, D6-bentazone, D4-carbendazim, ¹³C-fipronil, D4-imidacloprid, D3-isoproturon, D5-
178 metazachlor, D6-metazachlor-ESA, D10-simazine, D9-tebuconazole, D5-terbutryn) were introduced
179 into the MP in parallel with the analytes in the artificial samples, or instead of them for the analysis
180 of the field samples (50 µL of a 30 µg.L⁻¹ mix solution). They are too few and do not cover all the
181 chemical families researched to be used as internal standards, or they would probably have led to a
182 decrease of the accuracy and an increase of uncertainties (Han et al., 2018; Lehotay et al., 2018).
183 However, they can be used for evaluating the complexity of the matrices analyzed and tracing matrix
184 effects.

185 The area ratio between the reference matrix and the field samples was calculated for each
186 isotopically labeled standard. This parameter allowed to evaluate matrix complexity because
187 quantities introduced in all samples were the same. Principal component analyses (PCAs) were
188 performed on this ratio, for both fish and MIs, to evaluate if clusters of matrix complexity could be
189 determined based on species (fish) and taxonomic groups (MIs). PCAs were based on Pearson
190 regression ($\alpha = 0.05$). All statistical analyses and graphical PCA illustrations were performed using

the software R (v. 4.0.0, © The R Foundation, 2020) with the following packages: “ggplot2” (Wickham 2016), “factoextra” (Kassambara and Mundt 2020), “FactoMineR” (Lê et al. 2008), and “RColorBrewer” (Neuwirth 2014).

3. Results and discussion

3.1. Performance comparison between the initial protocol and the miniaturized protocol

The IP and the MP were applied to a fish matrix ($n = 5$) with a C4 concentration level to assess protocol performance (ERs and LOQs). The results are shown in Fig. 2. ERs ranged from 53% (terbutryn) to 103% (benzamide) with the IP and from 50% (prosulfocarb) to 120% (methsulfuron-methyl) with the MP. According to European guidance recommendations (European Commission DG-SANTE 2019), the ERs were considered good ($70\% < \text{ERs} < 120\%$) for 39 and 38 molecules, respectively. They were similar for both protocols, with a slight advantage for the MP. The most noticeable increases in ER were observed for desethylterbuthylazine-hydroxy (61% to 97% for IP and MP, respectively), terbuthylazine (73% to 87%), and terbutryn (53% to 69%). The only outlier that did not benefit from the change in protocol was prosulfocarb with respective ERs of 54% and 50%. LOQs ranged from 0.1 (fipronil, fipronil sulfone, metolachlor) to 221.8 ng.g^{-1} dw (metazachlor-OXA) with the IP and from 0.1 (metolachlor) to 221.2 ng.g^{-1} dw (metazachlor-OXA) for the MP. Fourteen molecules showed LOQs of up to 10 ng.g^{-1} dw for the IP, as against 18 molecules for the MP, with 4 and 5 molecules having LOQs of up to 50 ng.g^{-1} dw, respectively. The LOQs were very similar for both protocols, following a near-perfect linear regression ($R^2 = 0.958$). However, they were on average 1.4 times higher with the MP than with the IP. As proportions of reactants were conserved between the two protocols, obtaining similar performance results in terms of LOQ and ER did not come as a surprise. Miniaturizing the IP yielded equivalent ERs and LOQs, and allowed trace-level quantification of pesticide residues using 30 mg instead of 500 mg of matrix.

3.2. Performance of the miniaturized protocol in fish

The performance of the MP was fully evaluated for fish, as described in Section 2.6. The results are shown in Table 1. Linear regression was determined for each molecule and R^2 values were always greater than 0.99. Only 6 chemicals out of the 41 tested had R^2 values below 0.99: acetochlor-OXA (0.987), chlorotoluron (0.988), fenthion (0.950), flufenacet-ESA (0.987), metolachlor-ESA (0.989), and terbutryn (0.989). However, second-order polynomial regressions were more adapted for those molecules and their R^2 values were found to be 0.994, 0.996, 0.991, 0.992, 0.999, 0.993, and 0.996, respectively.

Among the 41 targeted substances, 38 had LOQs lower than or equal to 60 ng.g⁻¹ dw, including 19 below 10 ng.g⁻¹ dw. Taking into account the dehydration of the matrix (82% ± 11%), these limits corresponded to about 2 and 10 ng.g⁻¹ ww, respectively. The overall performance was consistent with that reported in the literature (Lazartigues et al. 2011; Kaczyński et al. 2017; Inostroza et al. 2017; Barbieri et al. 2019) although some LOQs were higher probably because of the low matrix masses used to perform the extraction (0.9 to 10 g in the literature, as against 0.03 g in the present study).

MEs ranged from -77% (fenthion) to 131% (imidacloprid-desnitro). Signal suppression was observed for 11 molecules and signal enhancement concerned 8 substances (out of a -20/+20 range). This was consistent with other studies reporting that signal suppression is the most commonly observed ME in complex matrices (Lazartigues et al. 2011; Barbieri et al. 2019). The monitoring of this parameter was important for validating the method even though matrix-matched calibration is supposed to compensate for those effects.

ERs at C2 ranged from 57% (prosulfocarb) to 108% (TCP). According to the European guidelines (European Commission DG-SANTE 2019) 35 of the 42 molecules presented good ERs (between 70% and 120%), including 12 of up to 80%. At C4, acceptable ERs for fish and MI matrices were obtained for 31 and 25 molecules, respectively. The inter-day precision was considered good for 32

chemicals at C2 according to the European guidelines previously mentioned (< 20%). This variability tended to decrease for higher concentration levels as the inter-day precision was lower than or equal to 20% for 31 molecules at C4, which is in agreement with European guidelines (European Commission DG-SANTE 2019). Moreover, if we leave aside the nature of the tested matrix, the inter-day precision is better with IP (Dufour et al., 2020) than with MP. This was most probably due to the decrease of the sample mass which led to an increase of the measurement uncertainties, as shown by Han et al. (2018) and Lehotay et al. (2018) on food samples towards pesticide contaminations. The developed analytical method was suitable for accurate quantification, except for fenthion for which inter-day precision was of up to 50% for the 3 tested concentration levels.

A signal-to-noise ratio of more than 3 was observed in protocol blanks for acetochlor-alachlor-ESA, bentazon, benzamide, boscalid, CGA 50267, fipronil, fipronil sulfone, isoproturon, metolachlor, prosulfocarb, and terbutryn. Contamination ranged from 0.1 to 8.8 ng depending on the molecule considered. The quantification frequency in blanks was usually below 50%, but reached 60% for fipronil and prosulfocarb, while metolachlor and terbutryn were systematically quantified. The LOQs were increased to 3 times the contamination levels for these molecules to ensure the reliability of the method.

258

3.3. Performance of the miniaturized protocol in macroinvertebrates

Performance of the MP was good for fish and compatible with trace analysis, so the protocol was tested on MI samples. As very low mass was available for the reference matrix (chironomid larvae), only linearity, RECs, and LOQs were assessed (Fig. 3). The tested concentration domains, ERs, LOQs, and blank contamination of each molecule are shown in Table S2. The quantification of most molecules followed a linear regression with R^2 values greater than 0.99; only fenthion (0.986), flufenacet (0.943), and TCP (0.961) were slightly below, reaching 0.994, 0.961, and 0.992,

266 respectively, when a second-order polynomial regression was applied. Protocol blanks were clean (n
267 = 4) but traces (≤ 0.8 ng) of dimethenamid-ESA, MCPA, and terbutryn were quantified
268 (quantification frequency below 0.5). LOQs ranged from 0.1 (fipronil, fipronil sulfone, metolachlor)
269 to 356 ng.g⁻¹ dw (metazachlor-OXA); taking into account the dehydration (85% on average for this
270 matrix), the LOQs ranged from 0.015 to 53 ng.g⁻¹ ww. Among the 41 tested molecules, 29 had LOQs
271 below 10 ng.g⁻¹ dw, 14 of which being below 1 ng.g⁻¹ dw. These values were of the same order of
272 magnitude as those reported in the literature (Haroune et al. 2015; Althakafy et al. 2018; Miller et al.
273 2019). Most tested molecules had good RECs (80%-120%) and values above 120% were obtained
274 with four of them (i.e., bentazone 122%, boscalid 122%, desethylterbutylazine-hydroxy 121%, and
275 metazachlor-OXA 131%). The latter values were close to the acceptable limits, except for
276 metazachlor-OXA for which a correction was necessary for the quantification of MI field samples. It
277 confirmed that most pesticide residues can be monitored in complex biotic matrices with low matrix
278 masses using a single method that provides equivalent levels of performance.

279

280 3.4. Field samples

281 In order to validate the performance of the method, 17 fish (3 species) and 19 MI (6 orders) samples
282 were collected and analyzed with the MP. All samples were spiked before extraction with 14
283 isotopically labeled standards (see section 2.8). The areas of these molecules were compared
284 between field samples and reference matrices. The average ratio ranged from 0.9 (D5-acetochlor-
285 ESA, D6-bentazone, D6-metazachlor-ESA, D5-atrazine-2-hydroxy, D9-tebuconazole) to 1.4 (D5-
286 terbutryn) for fish and from 0.8 (D5-acetochlor-ESA, D6-metazachlor-ESA) to 1.5 (D11-acetochlor)
287 for MIs. The relative standard deviation (%RSD) was lower than 30% for all molecules, except for
288 D5-atrazine (34%) and 13C-fipronil (32%) in fish, and D5-acetochlor-ESA (50%) in MIs. Taking
289 into account that a single matrix was used to quantify 3 different fish species, and another single
290 matrix for 6 different MI groups, these ratios close to 1 confirmed the potential for the protocol to be

transposed to other biotic matrices with similar relative levels of fat and protein. PCAs performed on area ratios illustrated 77.6% and 66.5% of total variability for fish and MI matrices, respectively (Fig. 4). Clusters could be identified for fish, showing that differences in matrix complexity exist between the 3 species, probably based on their physiological constitution. However, there was considerable overlap between the clusters, indicating relatively similar responses for internal standards. The same overlap was observed for MIs, confirming that the chosen reference matrix is suitable for matrix-matched calibration. However, hindsight is relatively poor for Anisoptera, Daphniidae, and Dytiscidae as their number of samples is ≤ 2 .

Concentrations monitored in field samples are shown in Table 2. Among the 40 targeted molecules (fenthion being excluded because of its unreliability, see Section 3.3), 7 were detected: benzamide, imidacloprid-desnitro, fipronil, imidacloprid, prosulfocarb, tebuconazole, and terbutryn. Benzamide, imidacloprid-desnitro, and prosulfocarb were almost systematically quantified in fish (from 42 to 237 ng.g⁻¹ dw, 3 ng.g⁻¹ dw, and from 30 to 165 ng.g⁻¹ dw, respectively) and in MIs (from 62 to 438 ng.g⁻¹ dw, from 2 to 6 ng.g⁻¹ dw, and from 15 to 29 ng.g⁻¹ dw, respectively). Considering the dehydration ratio, the average quantified concentrations of these three molecules were of 21, <LOQ, and 12 ng.g⁻¹ ww in fish, respectively, and of 26, 0.4, and 3 ng.g⁻¹ ww in MIs, respectively.

Prosulfocarb is a herbicide known to drift away, leading to the contamination of non-target crops and waterbodies (Devault et al. 2019). As it possesses a log P of 4.5, it is likely to accumulate in organisms, which seemed to be the case in this study for MIs and fish. It has already been reported in ponds at concentrations of up to 0.5 µg.L⁻¹ (Gaillard et al. 2016a). Benzamide is a TP of folpet that accumulates in both fish and MIs. Despite having a lower log P value than prosulfocarb (log P = 0.6), it was found in higher concentrations. Imidacloprid-desnitro, a TP of imidacloprid, was quantified in 5% of the fish samples and in 42% of the MI samples, while the quantification frequency of the active substance was 0% and 16%, respectively. There is a lack of data in the literature investigating the toxicity of imidacloprid-desnitro in aquatic organisms. Nevertheless,

research revealed that it has lower toxicity than imidacloprid in bees (Suchail et al. 2001). It should be noted that, in the present study, field sampling was carried out in May 2018. A few months later, in September 2018, the outdoor use of the herbicide was banned by the European Commission, so biotic concentrations are expected to decrease over time. However, the monitoring of TPs remain relevant as they sometimes have higher environmental concentrations than their parent molecules, and there is little evidence for their safety (Kiefer et al., 2019).

4. Conclusion

A miniaturized extraction protocol was successfully developed and applied to two complex matrices: fish and MIs. It allowed a sensitive and consistent analysis of PPPs even when only low sample mass was available. A decrease in the required mass of matrix slightly increased the LOQs of the MP in fish compared to the IP, but final LOQs were still suitable for trace-level analysis with most below 10 ng.g⁻¹ dw. Moreover, ERs were in the range 70%-120% for 35 of the 41 molecules. The transposition of IP to MP did not altered its performances. The application of this protocol in the field has revealed low contamination of aquatic organisms, but it has also shown that they are contaminated by a cocktail of molecules and TPs. The harmlessness of such a combination of contaminants is not guaranteed, and it is important to take it into consideration in order to ensure the good status of aquatic ecosystems. The herbicide prosulfocarb and the transformation product benzamide were almost systematically quantified in both matrices. The MP, developed in this study, represents a potential tool for the characterization of polar pesticides in small complex biotic samples.

Acknowledgments

This project was funded by the “*Office Français pour la Biodiversité*”. The authors sincerely thank the fish farmers and owners for granting access to their ponds. They are also grateful to Alain Iurétig

(sampling and pre-treatment, Univ. of Lorraine), Pamela Hartmeyer (pre-treatment, Univ. of Lorraine), Aisha Nunoo (extractions, Univ. of Lorraine/ISA), and Maud Dessein-Lepasteur (extractions and analysis, Univ. of Lorraine/ISA) for their work, as well as to Arnaud Chaumot and Laura Garnero from the Ecotoxicology Team of the UR RIVERLY (INRAE, Centre de Lyon-Villeurbanne) for providing chironomid larvae. The authors sincerely thank ABC Translation for the proofreading work done.

Funding Information:

This project was funded by the “Office Français pour la Biodiversité”.

References

- Althakafy JT, Kulsing C, Grace MR, Marriott PJ (2018) Determination of selected emerging contaminants in freshwater invertebrates using a universal extraction technique and liquid chromatography accurate mass spectrometry. *J Sep Sci* 41:3706–3715. <https://doi.org/10.1002/jssc.201800507>
- Anastassiades M, Lehotay SJ, Štajnbaher D, Schenck FJ (2003) Fast and Easy Multiresidue Method Employing Acetonitrile Extraction/Partitioning and “Dispersive Solid-Phase Extraction” for the Determination of Pesticide Residues in Produce. *J AOAC Int* 86:412–431. <https://doi.org/10.1093/jaoac/86.2.412>
- Barbieri MV, Postigo C, Guillem-Argiles N, et al (2019) Analysis of 52 pesticides in fresh fish muscle by QuEChERS extraction followed by LC-MS/MS determination. *Sci Total Environ* 653:958–967. <https://doi.org/10.1016/j.scitotenv.2018.10.289>
- Brain RA, Anderson JC (2019) The agro-enabled urban revolution, pesticides, politics, and popular culture: a case study of land use, birds, and insecticides in the USA. *Environ Sci Pollut Res* 26:21717–21735. <https://doi.org/10.1007/s11356-019-05305-9>

366 Clasen B, Loro VL, Murussi CR, et al (2018) Bioaccumulation and oxidative stress caused by
 367 pesticides in *Cyprinus carpio* reared in a rice-fish system. *Sci Total Environ* 626:737–743.
 368 <https://doi.org/10.1016/j.scitotenv.2018.01.154>
 369 Devault DA, Guillemain J-P, Millet M, et al (2019) Prosulfocarb at center stage! *Environ Sci Pollut*
 370 *Res* 1–15. <https://doi.org/10.1007/s11356-019-06928-8>
 371 Dufour V, Wiest L, Slaby S, et al (2020) Development of a simple multiresidue extraction method
 372 for the quantification of a wide polarity range list of pesticides and transformation products in
 373 eggs by liquid chromatography and tandem mass spectrometry. *J Chromatogr A* 1628:461447.
 374 <https://doi.org/10.1016/j.chroma.2020.461447>
 375 European Commission DG-SANTE (2019) Analytical Quality Control and Method Validation
 376 Procedures for Pesticide Residues Analysis in Food and Feed, N° SANTE/12682/2019. 49
 377 Four B, Thomas M, Danger M, et al (2019) Using stable isotope approach to quantify pond dam
 378 impacts on isotopic niches and assimilation of resources by invertebrates in temporary streams:
 379 a case study. *Hydrobiologia* 834:163–181. <https://doi.org/10.1007/s10750-019-3920-0>
 380 Gaillard J, Thomas M, Iuretig A, et al (2016a) Barrage fishponds: Reduction of pesticide
 381 concentration peaks and associated risk of adverse ecological effects in headwater streams. *J*
 382 *Environ Manage* 169:261–271. <https://doi.org/10.1016/j.jenvman.2015.12.035>
 383 Gaillard J, Thomas M, Lazartigues A, et al (2016b) Potential of barrage fish ponds for the mitigation
 384 of pesticide pollution in streams. *Environ Sci Pollut Res* 23:23–35.
 385 <https://doi.org/10.1007/s11356-015-5378-6>
 386 Geiger F, Bengtsson J, Berendse F, et al (2010) Persistent negative effects of pesticides on
 387 biodiversity and biological control potential on European farmland. *Basic Appl Ecol* 11:97–105.
 388 <https://doi.org/10.1016/j.baae.2009.12.001>
 389 Giroud B, Vauchez A, Vulliet E, et al (2013) Trace level determination of pyrethroid and
 390 neonicotinoid insecticides in beebread using acetonitrile-based extraction followed by analysis

391 with ultra-high-performance liquid chromatography–tandem mass spectrometry. *J Chromatogr*
 392 *A* 1316:53–61. <https://doi.org/10.1016/j.chroma.2013.09.088>

393 Golge O, Koluman A, Kabak B (2018) Validation of a Modified QuEChERS Method for the
 394 Determination of 167 Pesticides in Milk and Milk Products by LC-MS/MS. *Food Anal Methods*
 395 11:1122–1148. <https://doi.org/10.1007/s12161-017-1066-0>

396 Han, L., Lehotay, S.J., Sapozhnikova, Y., 2018. Use of an Efficient Measurement Uncertainty
 397 Approach To Compare Room Temperature and Cryogenic Sample Processing in the Analysis of
 398 Chemical Contaminants in Foods. *J. Agric. Food Chem.* 66, 4986–4996.
 399 <https://doi.org/10.1021/acs.jafc.7b04359>

400 Haroune L, Cassoulet R, Lafontaine M-P, et al (2015) Liquid chromatography-tandem mass
 401 spectrometry determination for multiclass pesticides from insect samples by microwave-assisted
 402 solvent extraction followed by a salt-out effect and micro-dispersion purification. *Anal Chim*
 403 *Acta* 891:160–170. <https://doi.org/10.1016/j.aca.2015.07.031>

404 Inostroza PA, Massei R, Wild R, et al (2017) Chemical activity and distribution of emerging
 405 pollutants: Insights from a multi-compartment analysis of a freshwater system. *Environ Pollut*
 406 231:339–347. <https://doi.org/10.1016/j.envpol.2017.08.015>

407 Kaczyński P, Łozowicka B, Perkowski M, Szabuńko J (2017) Multiclass pesticide residue analysis in
 408 fish muscle and liver on one-step extraction-cleanup strategy coupled with liquid
 409 chromatography tandem mass spectrometry. *Ecotoxicol Environ Saf* 138:179–189.
 410 <https://doi.org/10.1016/j.ecoenv.2016.12.040>

411 Kassambara A, Mundt F (2020) factoextra: Extract and Visualize the Results of Multivariate Data
 412 Analyses

413 Kiefer, K., Müller, A., Singer, H., Hollender, J., 2019. New relevant pesticide transformation
 414 products in groundwater detected using target and suspect screening for agricultural and urban
 415 micropollutants with LC-HRMS. *Water Res.* 165, 114972.

416 <https://doi.org/10.1016/j.watres.2019.114972>

417 Knoll S, Rösch T, Huhn C (2020) Trends in sample preparation and separation methods for the
 418 analysis of very polar and ionic compounds in environmental water and biota samples. *Anal*
 419 *Bioanal Chem* 412:6149–6165. <https://doi.org/10.1007/s00216-020-02811-5>

420 Koesukwiwat U, Lehotay SJ, Miao S, Leepipatpiboon N (2010) High throughput analysis of 150
 421 pesticides in fruits and vegetables using QuEChERS and low-pressure gas chromatography–
 422 time-of-flight mass spectrometry. *J Chromatogr A* 1217:6692–6703.
 423 <https://doi.org/10.1016/j.chroma.2010.05.012>

424 Konwick BJ, Garrison AW, Black MC, et al (2006) Bioaccumulation, biotransformation, and
 425 metabolite formation of fipronil and chiral legacy pesticides in rainbow trout. *Environ Sci*
 426 *Technol* 40:2930–2936. <https://doi.org/10.1021/es0600678>

427 Lazartigues A, Wiest L, Baudot R, et al (2011) Multiresidue method to quantify pesticides in fish
 428 muscle by QuEChERS-based extraction and LC-MS/MS. *Anal Bioanal Chem* 400:2185–2193.
 429 <https://doi.org/10.1007/s00216-011-4945-z>

430 Lê S, Josse J, Husson F (2008) FactoMineR : An R Package for Multivariate Analysis. *J Stat Softw*
 431 25:. <https://doi.org/10.18637/jss.v025.i01>

432 Lehotay, S.J., Han, L., Sapozhnikova, Y., 2018. Use of a quality control approach to assess
 433 measurement uncertainty in the comparison of sample processing techniques in the analysis of
 434 pesticide residues in fruits and vegetables. *Anal. Bioanal. Chem.* 410, 5465–5479.
 435 <https://doi.org/10.1007/s00216-018-0905-1>

436 Lehotay SJ, Son KA, Kwon H, et al (2010) Comparison of QuEChERS sample preparation methods
 437 for the analysis of pesticide residues in fruits and vegetables. *J Chromatogr A* 1217:2548–2560.
 438 <https://doi.org/10.1016/j.chroma.2010.01.044>

439 Lichtmannegger K, Fischer R, Steemann FX ave., et al (2015) Alternative QuEChERS-based
 440 modular approach for pesticide residue analysis in food of animal origin. *Anal Bioanal Chem*

441 407:3727–3742. <https://doi.org/10.1007/s00216-015-8594-5>

442 Macchi P, Loewy RM, Lares B, et al (2018) The impact of pesticides on the macroinvertebrate
 443 community in the water channels of the Río Negro and Neuquén Valley, North Patagonia
 444 (Argentina). *Environ Sci Pollut Res* 25:10668–10678. [https://doi.org/10.1007/s11356-018-](https://doi.org/10.1007/s11356-018-1330-x)
 445 1330-x

446 McLaughlin A, Mineau P (1995) The impact of agricultural practices on biodiversity. *Agric Ecosyst*
 447 *Environ* 55:201–212. [https://doi.org/10.1016/0167-8809\(95\)00609-V](https://doi.org/10.1016/0167-8809(95)00609-V)

448 Miller TH, Ng KT, Bury ST, et al (2019) Biomonitoring of pesticides, pharmaceuticals and illicit
 449 drugs in a freshwater invertebrate to estimate toxic or effect pressure. *Environ Int* 129:595–606.
 450 <https://doi.org/10.1016/j.envint.2019.04.038>

451 Neuwirth E (2014) RColorBrewer: ColorBrewer Palettes

452 Nowell LH, Moran PW, Schmidt TS, et al (2018) Complex mixtures of dissolved pesticides show
 453 potential aquatic toxicity in a synoptic study of Midwestern U.S. streams. *Sci Total Environ*
 454 613–614:1469–1488. <https://doi.org/10.1016/j.scitotenv.2017.06.156>

455 Oliveira FA da S, Pereira ENC, Gobbi JM, et al (2018) Multiresidue method for detection of
 456 pesticides in beef meat using liquid chromatography coupled to mass spectrometry detection
 457 (LC-MS) after QuEChERS extraction. *Food Addit Contam Part A* 35:94–109.
 458 <https://doi.org/10.1080/19440049.2017.1395519>

459 Reemtsma T, Berger U, Arp HPH, et al (2016) Mind the Gap: Persistent and Mobile Organic
 460 Compounds—Water Contaminants That Slip Through. *Environ Sci Technol* 50:10308–10315.
 461 <https://doi.org/10.1021/acs.est.6b03338>

462 Roche H, Vollaire Y, Martin E, et al (2009) Rice fields regulate organochlorine pesticides and PCBs
 463 in lagoons of the Nature Reserve of Camargue. *Chemosphere* 75:526–533.
 464 <https://doi.org/10.1016/j.chemosphere.2008.12.009>

465 Rostron C (2010) Review of Environmental Contamination and Toxicology Volume 204. Springer

466 New York, New York, NY

467 Sanganyado, E., Bi, R., Teta, C., Buruaem Moreira, L., Yu, X., Yajing, S., Dalu, T., Rajput, I.R.,
 468 Liu, W., 2020. Toward an integrated framework for assessing micropollutants in marine
 469 mammals: Challenges, progress, and opportunities. *Crit. Rev. Environ. Sci. Technol.* 0, 1–48.
 470 <https://doi.org/10.1080/10643389.2020.1806663>

471 Sinclair CJ, Boxall ABA (2003) Assessing the Ecotoxicity of Pesticide Transformation Products.
 472 *Environ Sci Technol* 37:4617–4625. <https://doi.org/10.1021/es030038m>

473 Slaby S, Marin M, Marchand G, Lemiere S (2019) Exposures to chemical contaminants: What can
 474 we learn from reproduction and development endpoints in the amphibian toxicology literature?
 475 *Environ Pollut* 248:478–495. <https://doi.org/10.1016/j.envpol.2019.02.014>

476 Song N-E, Lee JY, Mansur AR, et al (2019) Determination of 60 pesticides in hen eggs using the
 477 QuEChERS procedure followed by LC-MS/MS and GC-MS/MS. *Food Chem* 298:125050.
 478 <https://doi.org/10.1016/j.foodchem.2019.125050>

479 Stanton RL, Morrissey CA, Clark RG (2018) Analysis of trends and agricultural drivers of farmland
 480 bird declines in North America: A review. *Agric Ecosyst Environ* 254:244–254.
 481 <https://doi.org/10.1016/j.agee.2017.11.028>

482 Suchail S, Guez D, Belzunces LP (2001) Discrepancy between acute and chronic toxicity induced by
 483 imidacloprid and its metabolites in *Apis mellifera*. *Environ Toxicol Chem* 20:2482–2486.
 484 <https://doi.org/10.1002/etc.5620201113>

485 Tousova Z, Oswald P, Slobodnik J, et al (2017) European demonstration program on the effect-based
 486 and chemical identification and monitoring of organic pollutants in European surface waters.
 487 *Sci Total Environ* 601–602:1849–1868. <https://doi.org/10.1016/j.scitotenv.2017.06.032>

488 Wickham H (2016) *GGPLOT2: Elegant Graphics for Data Analysis*. Springer-Verlag New York

489

490

491

492 Fig. 1. Step comparison between the initial and miniaturized protocol (ACN: acetonitrile, FA: Formic acid)

493

494

495

496 Fig. 2. Comparison between the initial (IP) and miniaturized protocol (MP) in terms of extraction rates (ERs)
497 and limits of quantification (LOQs) obtained for fish blank matrix spiked at C4 (n=5); see Table 1 for the
498 molecules associated with the numbers

499

500

501

502 Fig. 3. Quantification recoveries (%) and limits of quantification (LOQs) (ng.g⁻¹ dw) obtained for
503 macroinvertebrates spiked at C4 (n=6); see Table 1 for the molecules associated with the numbers

504

505

506

507 Fig. 4. Representation of isotope-labeled standard area ratios between reference and field samples for fish and
508 macroinvertebrates; PCA based on Pearson regression ($\alpha = 0.05$)

509 (USE COLORS FOR THIS FIGURE)

510

511 Table 1. Analytical performance of the validated method in fish

n°	Chemicals	Domain (ng.g ⁻¹ dw)	r ² ± %RSD (n = 5)	LOQ (ng.g ⁻¹ dw)	Matrix effect ± SD (%)	Recovery ± %RSD (%)			Inter-day precision (%RSD)			Blanks (n = 5)	
						C2 (n = 6)	C4 (n = 3)	C6 (n = 6)	C2 (n = 5)	C4 (n = 5)	C6 (n = 5)	Min - Max (ng)	QF
Fungicides													
1	Boscalid	60 - 1200	0.996 (4)	60	-6 ± 12	80 ± 15	87 ± 4	93 ± 8	27	18	23	5.1 - 5.1	0.2
2	Tebuconazole	3 - 60	0.993 (2)	3	-18 ± 15	69 ± 20	98 ± 17	89 ± 8	26	15	24	-	-
Herbicides													
3	Acetochlore	60 - 1200	0.995 (4)	60	-2 ± 14	76 ± 12	77 ± 2	80 ± 4	27	20	22	-	-
4	Alachlor	30 - 600	0.995 (3)	30	-10 ± 12	77 ± 9	79 ± 6	79 ± 7	24	19	22	-	-
5	Atrazine	3 - 60	0.996 (2)	3	-31 ± 12	74 ± 13	82 ± 5	86 ± 7	22	18	20	-	-
6	Bentazon	3 - 60	0.992 (8)	3	-8 ± 16	69 ± 33	88 ± 6	88 ± 3	22	17	18	0.1 - 0.1	0.2
7	Chlorotoluron	6 - 120	0.988 (14)	6	-20 ± 16	83 ± 19	86 ± 10	92 ± 7	20	21	62	-	-
8	Dimethachlor	6 - 120	0.996 (2)	6	-14 ± 17	78 ± 22	89 ± 5	89 ± 3	27	21	21	-	-
9	Dimethenamide	6 - 120	0.995 (1)	6	-20 ± 14	76 ± 25	86 ± 5	88 ± 10	22	20	22	-	-
10	Flufenacet	6 - 120	0.996 (4)	6	-6 ± 17	74 ± 18	85 ± 6	91 ± 5	27	18	21	-	-
11	Isoproturon	3 - 60	0.993 (4)	3	-23 ± 15	77 ± 18	92 ± 9	93 ± 7	22	18	19	0.1	0.2
12	MCPA	30 - 600	0.993 (4)	30	16 ± 9	72 ± 29	89 ± 1	86 ± 7	26	21	21	-	-
13	Metazachlor	3 - 60	0.992 (5)	3	-10 ± 16	83 ± 19	90 ± 10	92 ± 9	25	18	20	-	-
14	Metolachlor	3 - 60	0.997 (3)	3	-6 ± 12	88 ± 8	82 ± 12	83 ± 4	26	18	19	0.1 - 0.2	1.0
15	Metsulfuron-methyl	30 - 600	0.994 (5)	30	48 ± 29	71 ± 31	101 ± 7	106 ± 7	27	21	21	-	-
16	Prosulfocarb	30 - 600	0.99 (6)	30	-57 ± 16	102 ± 63	36 ± 11	45 ± 13	23	24	23	0 - 8.8	0.6
17	Terbuthylazine	3 - 60	0.994 (4)	3	-41 ± 23	84 ± 24	67 ± 16	74 ± 7	19	17	21	-	-
18	Terbutryn	3 - 60	0.989 (7)	3	-43 ± 7	86 ± 26	63 ± 10	66 ± 6	26	16	20	0.1 - 0.2	1.0
Insecticides													
19	Fenthion	20 - 400	0.95 (76)	80	-77 ± 21	87 ± 33	82 ± 8	121 ± 88	62	59	60	-	-
20	Fipronil	3 - 60	0.997 (2)	3	-13 ± 8	74 ± 22	85 ± 5	90 ± 6	23	16	19	0.1	0.6
21	Imidacloprid	30 - 600	0.993 (8)	30	19 ± 11	101 ± 70	94 ± 6	94 ± 4	27	19	23	-	-

Transformation products												
22	Acetochlor OXA	30 - 600	0.987 (10)	30	27 ± 12	70 ± 24	76 ± 10	78 ± 12	27	19	24	-
23	Alachlor-Acetochlor ESA	30 - 600	0.993 (5)	30	33 ± 11	77 ± 17	69 ± 5	76 ± 7	34	18	22	0.5 - 0.5
24	Atrazine-2-hydroxy	3 - 60	0.996 (2)	3	-3 ± 16	68 ± 17	83 ± 8	82 ± 6	26	18	19	-
25	Benzamide	30 - 600	0.991 (5)	30	61 ± 93	96 ± 16	105 ± 10	98 ± 10	16	18	22	0.2 - 0.3
26	CGA 50267	3 - 60	0.995 (4)	3	-10 ± 13	72 ± 25	92 ± 3	91 ± 10	25	19	19	0.2 - 0.2
27	Desethylterbutylazine	3 - 60	0.995 (3)	3	-18 ± 14	75 ± 18	86 ± 6	89 ± 5	23	14	20	-
28	Desethylterbutylazine -OH	50 - 1000	0.994 (4)	50	-4 ± 8	71 ± 20	86 ± 10	77 ± 10	24	23	62	-
29	Dimethachlor ESA	30 - 600	0.994 (3)	60	10 ± 12	79 ± 7	67 ± 14	70 ± 15	29	18	22	-
30	Dimethenamide ESA	30 - 600	0.99 (9)	30	15 ± 12	87 ± 25	60 ± 8	69 ± 11	33	17	22	-
31	Fipronil sulfone	3 - 60	0.99 (10)	3	-43 ± 21	78 ± 14	82 ± 7	87 ± 11	27	15	22	0.1
32	Flufenacet ESA	6 - 120	0.987 (10)	12	11 ± 15	66 ± 34	64 ± 7	71 ± 12	42	13	24	-
33	Flufenacet OXA	30 - 600	0.991 (8)	30	20 ± 16	74 ± 28	76 ± 10	73 ± 8	35	18	23	-
34	Imidacloprid-desnitro	3 - 60	0.994 (3)	3	131 ± 108	71 ± 28	89 ± 12	90 ± 6	20	14	20	-
35	Isoproturon-desmethyl	6 - 120	0.992 (4)	6	-22 ± 14	77 ± 27	87 ± 9	91 ± 8	31	20	23	-
36	Me-Desphenyl-Chloridazon	30 - 600	0.995 (3)	30	-20 ± 12	77 ± 21	83 ± 6	91 ± 5	18	19	21	-
37	Metazachlor ESA	30 - 600	0.99 (7)	30	11 ± 10	73 ± 23	68 ± 8	71 ± 15	62	21	22	-
38	Metazachlor OXA	30 - 600	0.993 (2)	120	35 ± 35	77 ± 35	62 ± 12	77 ± 10	33	22	28	-
39	Metolachlor ESA	30 - 600	0.989 (8)	30	14 ± 9	85 ± 36	88 ± 17	70 ± 12	20	13	19	-
40	Metolachlor OXA	30 - 600	0.993 (5)	60	30 ± 15	59 ± 15	97 ± 11	76 ± 7	29	13	20	-
41	Trichloropyridinol	60 - 1200	0.991 (4)	480	-7 ± 10	108 ± 30	80 ± 11	82 ± 12	31	22	21	-

n°: identification of molecules in Fig. 2 and Fig. 3, **Domain**: range of tested concentrations (ng.g⁻¹ dw), **dw**: dry weight, **LOQ**: estimated limit of quantification (ng.g⁻¹ dw), **ME**: matrix effect (%), **ER**: extraction rate (%), **QF**: quantification frequency, **C2 to C6**: tested concentration levels.

512

513

514 Table 2. Pesticide residues quantified in fish and macroinvertebrates in ng.g⁻¹ dw

Molecule	Fish (n = 17)			Macroinvertebrates (n = 19)		
	QF (%)	Min - Max	Average ± SEM	QF (%)	Min - Max	Average ± SEM
Benzamide	100	42 - 237	116 ± 12	100	62 - 438	174 ± 23
Imidacloprid-desnitro	5	3	-	42	2 - 6	3 ± 1
Fipronil	0	-	-	5	4	-
Imidacloprid	0	-	-	16	12 - 37	21 ± 8
Prosulfocarb	86	30 - 165	67 ± 8	100	15 - 29	21 ± 1
Tebuconazole	0	-	-	11	3 - 8	-
Terbutryn	5	61	-	0	-	-

QF: quantification frequency, **dw**: dry weight, **Min - Max**: minimal and maximal values quantified, **SEM**: standard error of the mean.

515