Pentafluorophenyl Esters as Exchangeable Stoppers for the Construction of Photoactive [2]rotaxanes

Marine Rémy,^[a] Iwona Nierengarten,^[a] Boram Park,^[a] Michel Holler,^[a] Uwe Hahn,^[a] and Jean-François Nierengarten^{*[a]}

Dedicated to Prof. Didier Astruc on the occasion of his 75th birthday

[a] M. Rémy, Dr. I. Nierengarten, B. Park, Dr. M. Holler, Dr. U. Hahn, Dr. J.-F. Nierengarten
Laboratoire de Chimie des Matériaux Moléculaires
Université de Strasbourg et CNRS (UMR 7402 LIMA), Ecole Européenne de Chimie, Polymères et Matériaux
25 rue Becquerel, 67087 Strasbourg Cedex 2, France
E-mail: nierengarten@unistra.fr

Supporting information for this article is given via a link at the end of the document.

Abstract: Stable pillar[5]arene-containing [2]rotaxane building blocks with pentafluorophenyl ester stoppers have been efficiently prepared on a multi-gram scale. Reaction of these building blocks with various nucleophiles gave access to a wide range of [2]rotaxanes with amide, ester or thioester stoppers in good to excellent yields. The rotaxane structure is fully preserved during these chemical transformations. Actually, the addition-elimination mechanism at work during these transformations totally prevents the unthreading of the axle moiety of the mechanically interlocked system. The stopper exchange reactions were optimized both in solution and under mechanochemical solvent-free conditions. While amide formation is more efficient in solution, the solvent-free conditions are more powerful for the transesterification reactions. Starting from a fullerene-functionalized pillar[5]arene derivative, this new strategy gave easy access to a photoactive [2]rotaxane incorporating a C₆₀ moiety and two Bodipy stoppers. Despite the absence of covalent connectivity between the Bodipy and the fullerene moieties in this photoactive molecular device, efficient through space excited state interactions have been evidenced in this rotaxane.

Introduction

Mechanically interlocked molecules (MIMs) played a key role in the development of molecular machines.^[1-2] On the other hand, such structures are also appealing scaffolds for the construction of photoactive molecular devices in which complementary chromophores are associated into a single molecular ensemble but without being connected through covalent chemical bonds.^[3-9] As a result, intercomponent photochemical events observed in such systems occur exclusively through space.^[3-9] Sauvage was a pioneer in this particular field and his team has reported spectacular photoactive MIMs mimicking the ultra-fast charge separation observed in the natural photosynthetic system through the peculiar orientation of complementary porphyrinic moieties in rotaxanes and catenanes constructed by a metal-templated approach.^[3-4] The same construction principle has been also intensively used to prepare related photoactive fullerene-containing systems.^[5-6] Further examples of photoactive MIMs in which through-space photoinduced electron or energy transfer events have been evidenced were also

prepared by using other synthetic approaches.^[7] As part of this research, we have recently shown that pillar[5]arene-containing rotaxanes are attractive scaffolds for the construction of photoactive MIMs.^[8-9] Their synthesis was however particularly difficult and new efficient methods for the preparation of pillar[5]arene-containing rotaxanes are needed to further develop this particular field. This prompted us to develop synthetic methodologies based on stopper exchange reactions from rotaxane building blocks incorporating 2,4-dinitrophenyl ester or 3,5-bis(trifluoromethyl) benzenesulfonate stoppers.^[10-12] While very efficient for the preparation of rotaxanes incorporating an ethoxy functionalized pillar[5]arene moiety, major limitations of these building blocks are related to their moderate stability in particular on silica gel. As a result, their purification is often challenging. In order to solve this stability issue, we decided to further explore this strategy with new exchangeable stoppers. In this paper, we now report the preparation of pillar[5]arene-containing rotaxanes with pentafluorophenyl ester stoppers. These building blocks have been obtained in almost quantitative yields and are perfectly stable on silica gel thus allowing easy purifications. When compared to the previous building block with 2,4-dinitrophenol leaving groups,^[10] the stopper exchange is not only limited to the reaction with amines to afford rotaxanes with amide stoppers. Effectively, rotaxanes with ester or thioester stoppers have been also efficiently prepared by treatment of the pentafluorophenyl ester derivatives with alcohol, phenol or thiol reagents. Moreover, all the reported chemical transformations could be achieved in solution but also under mechanochemical solvent-free conditions. Finally, this new methodology has been successfully applied to the construction of a photoactive MIM incorporating a fullerene-functionalized pillar[5]arene derivative.

Results and Discussion

Preparation of the [2]rotaxane building block. Pillar[5]arene-containing rotaxane building blocks incorporating activated ester stoppers are attractive building blocks for the preparation of a large variety of rotaxanes. We have recently shown that 2,4-dinitrophenyl esters are well suited for such a purpose.^[10] However, partial hydrolysis of the 2,4-dinitrophenol ester functions was observed on silica gel and purification of the

rotaxane building blocks was only possible by recrystallisation. This strategy was efficient when ethoxypillar[5]arene was used as starting material but the preparation of related rotaxanes from more sophisticated pillar[5]arene derivatives was found difficult. It was thus decided to explore alternative activated stoppers for the construction of pillar[5]arene-containing rotaxanes. Pentafluorophenyl ester derivatives attracted our attention. Owing to their easy preparation and their good stability, such derivatives are widely used to activate carboxylic acids for the efficient preparation of amides and esters.^[13-14] Based on molecular modeling, the pentafluorophenyl unit appears to be bulky enough to prevent the pillar[5]arene moiety from escaping the thread of the rotaxane. The preparation of a [2]rotaxane with pentafluorophenyl ester stoppers was first tested starting from ethoxypillar[5]arene **1** and diacyl chloride **2** (Scheme 1).

Scheme 1. Preparation of rotaxane building block **4** and dumbbell **5**. *Reagents and conditions:* (i) **2**, Et₃N, CHCl₃, -15°C (98%); (ii) **2**, CHCl₃, -15°C then evaporation; (iii) **3**, Et₃N, mixing in a Retsch MM400 mill at 30 Hz for 3 h (94%); (iv) **3**, Et₃N, CH₂Cl₂, 0°C (87%).

Successful rotaxane preparation relies on the ability of **1** and **2** to self-assemble into the appropriate pseudo-rotaxane precursor. The association constant (K_a) for the 1:1 supramolecular complex obtained from **1** and **2** is rather low ($\log K_a = 1.85$ in CDCl₃ at 25°C).^[15] It is therefore essential to properly adjust the experimental conditions used for the introduction of the stoppers in order to favor the assembly of **1** with **2**. For this reason, all the stoppering reactions were performed at the highest possible concentration and at the lowest possible temperature (-15°C). The reaction of diacyl chloride **2** with pentafluorophenol (**3**) in the presence of an excess of pillar[5]arene **1** (2 equiv.) and Et₃N in CHCl₃ afforded the desired rotaxane together with small amounts of by-product **5** resulting from the reaction of uncomplexed **2** with **3**. Compounds **4** and **5** were conveniently

separated by column chromatography on silica gel. Rotaxane **4** was thus obtained pure in 75% yield and by-product **5** in 19% yield. Compound **5** was also prepared independently in 87% yield by treatment of diacyl chloride **2** with pentafluorophenol (**3**). Importantly rotaxane **4** and dumbbell **5** were perfectly stable on silica gel and partial hydrolysis of their ester functions was not observed as in the case of the corresponding 2,4-dinitrophenyl derivatives. The preparation of **4** was further improved by starting from a larger excess of **1** (3 equiv.). Under these conditions, the formation of the inclusion complex between **1** and **2** was more favorable and compound **4** was isolated in a nearly quantitative yield (98%). The preparation of **4** was also achieved under mechanochemical solvent-free conditions (Scheme 1).^[16-18] In this particular case, the host-guest complex [**1**⊂**2**] was prepared first and used in the solid state as received for the stoppering reaction. Specifically, a 2:1 mixture of **1** and **2** was dissolved in a minimum of CHCl₃.^[16] The resulting solution was cooled at -15°C to favor the assembly of the desired inclusion complex and then evaporated to dryness without heating. The colorless solid thus obtained was transferred into a stainless-steel grinding vial containing pentafluorophenol (2.5 equiv.), Et₃N (2.5 equiv.) and four stainless steel balls. The resulting mixture was mixed in a Retsch MM400 mill at 30 Hz for 1 h. Rotaxane **4** was thus obtained in 78% yield. The corresponding dumbbell (**5**) was also isolated as a by-product in 20% yield. To further improve the formation of inclusion complex [**1**⊂**2**], the same procedure was applied starting from a 3:1 mixture of **1** and **2**. Under these conditions, 3 hours of mixing were necessary to complete the reaction. Rotaxane **4** was isolated in 94% yield and the corresponding unthreaded axle by-product (**5**) could not be detected anymore. The outcome of the solvent-free stoppering reactions is therefore mainly dependent on the initial conditions used for the preparation of the neat samples of [**1**⊂**2**] but dissociation of the host-guest complex initially formed does not occur in the solid state during the mechanochemical treatment.

Rotaxane **4** was found stable for several months under normal laboratory conditions and could be stored without particular precaution. As anticipated, the pentafluorophenyl ester groups are large enough to prevent the pillar[5]arene moiety of **4** from escaping its axle. Compound **4** was effectively stable in solution and no dissociation could be evidenced even upon heating THF or CHCl₃ solutions of **4** under reflux for several hours. The NMR and mass spectra recorded for **4** were in full agreement with the proposed structure (see ESI). The formation of rotaxane **4** was further evidenced by X-ray crystallography. Crystals suitable for X-ray crystal-structure analysis were obtained by slow diffusion of diethyl ether into a CH₂Cl₂ solution of compound **4**. As shown in Figure 1, the -(CH₂)₁₀- chain of the axle component of rotaxane **4** does not adopt a fully extended form with all the successive methylene groups in an anti-conformation. Effectively, gauche conformations are almost exclusively observed for the decyl chain of **4** in the solid state. This peculiar conformation is most probably not related to the maximization of the intramolecular C-H...π interactions between the two components of the rotaxane. Effectively, in most of the X-ray crystal structures reported to date for analogous rotaxanes,^[10-11] the methylene units of the alkyl chains are typically in a fully extended all-anti conformation. In the particular case of **4**, packing forces are likely responsible for the observed gauche conformations adopted by the decyl chain. The resulting

contraction of the axle moiety allows most probably for optimized intermolecular contacts in the crystal lattice and the strain energy resulting from the adopted gauche conformations is thus compensated. This view is supported by the observation of intermolecular π - π interactions of the electron-deficient pentafluorophenyl aromatic rings with electron-rich hydroquinone subunits of neighboring pillar[5]arenes leading to the formation of infinite supramolecular chains in the crystal lattice (Figure 1B).

Figure 1. (A) ORTEP plot of the structure of **4** (H: white, F: light green, O: red, C: grey for the dumbbell and pale blue for the pillar[5]arene moiety; the disorder of some methylene groups of the decyl chain has been omitted for clarity; thermal ellipsoids are shown at 30% probability level). (B) Stacking within the **4** lattice highlighting the intermolecular π - π interactions of the pentafluorophenyl stoppers with the pillar[5]arene subunits of neighbouring rotaxanes.

Stopper exchange reactions. The preparation of rotaxanes **7a-g** with amide stoppers was attempted by reaction of building block **4** with the corresponding amine reagents (**6a-g**, Scheme 2). All the reactions were performed in anhydrous THF. When primary amine reagents (**6a-c**) were used, the stopper exchange reactions with **4** were complete after 5 hours at room temperature. The desired amide rotaxanes **7a-c** were thus obtained in almost quantitative yields. In contrast, the stopper exchange reactions were very slow when secondary amine and aniline reagents (**6d-g**) were used. Rotaxane **7d** was only obtained in a moderate yield (26%) upon heating the reaction mixture under reflux for 3 days. By further increasing steric effects, the reactivity of the amine reagent was totally vanished thus preventing the formation of rotaxane **7e** from **4** and **6e**. On the other hand, the reaction of **4** with aniline **6f** was complete after heating the THF solution under reflux for 24 h and afforded rotaxane **7f** in 62% yield. Aniline **6g** is a weak nucleophile due to the presence of the electron-withdrawing CF_3 substituents and the reaction with **4** was particularly slow. After 7 days, the desired amide rotaxane (**7g**) was obtained in only 5% yield. Overall, the results obtained for the stopper exchange reactions of **4** are very similar to those reported for the corresponding rotaxane building block with 2,4-dinitrophenyl ester stoppers and similar limitations are observed.^[10] Almost any primary

alkylamine reagents can be efficiently grafted onto the rotaxane scaffold by reaction with both pentafluorophenyl and 2,4-dinitrophenyl ester building blocks. This strategy is however less efficient as soon as the nucleophilicity of the starting material is reduced by either steric hindrance in the case of secondary amines or electronic effects in the case of anilines. It can be noted that the reaction of **4** with amines **6a** and **6c-g** were also performed under mechanochemical solvent-free conditions. Whereas no stopper exchange reactions took place in the case of secondary amine **6e** and anilines **6f-g**, mixing **4** in the presence of an excess of the appropriate amine reagents (**6a**, **6c** or **6d**, 8 equiv.) in a Retsch MM400 mill at 30 Hz for 1.5 hours gave the corresponding rotaxanes (**7a**, **7c** and **7d**). Under these optimized conditions, the isolated yields were similar to the ones obtained for the reactions performed in THF solutions. The use of a large excess of amine reagents (8 equiv.) was mandatory to achieve the stopper exchanges in good yields under mechanochemical conditions. This was not the case in solution where stoichiometric amounts of the amine reagent are sufficient to complete the amide formation. For this reason, wet conditions are more convenient to achieve stopper exchange reactions from building block **4** and amine reagents.

Scheme 2. Stopper exchange reactions with amine reagents. *Reagents and conditions:* (i) **4**, THF (**7a**: quantitative, **7b**: 96%, **7c**: 99%, **7d**: 26%, **7e**: 0%, **7f**: 62%, **7g**: 5%); (ii) **4**, mixing in a Retsch MM400 mill at 30 Hz for 1.5 h (**7a**: 98%, **7c**: 92%, **7d**: 24%, **7e-g**: 0%).

The stopper exchange reactions of **4** with a series of phenol, alcohol and thiophenol reagents were also successfully achieved (Scheme 3). In our hands, related reactions starting from the rotaxane building block with 2,4-dinitrophenyl ester stoppers were at best low-yielding and decomposition of the starting material was typically faster than the formation of the desired products. The increased stability of the pentafluorophenyl ester subunits of **4** is therefore not only an advantage for purification purposes but it allows also to prepare

rotaxanes with a larger structural diversity. Under optimized conditions, treatment of compound **4** with an excess of phenol **8a** (4 equiv.) in the presence of 1,8-diazabicyclo(5.4.0)undec-7-ene (DBU, 2.05 equiv.) in THF as a base provided rotaxane **9a** in 57% yield. Applying these conditions to the other phenol or alcohol reagents gave however unsatisfactory results and the desired rotaxanes were generally thus obtained in low to moderate yields. Side products resulting from the partial hydrolysis of the rotaxanes were also typically formed under these conditions. This prompted us to test other organocatalysts for the transesterification reactions. The best results were obtained with 1,4,7-triazabicyclo[4.4.0]dec-5-ene (TBD).^[14] Reaction of **4** with the phenol or alcohol starting materials (**8a-h**, 4 equiv.) and TBD (2.05 equiv.) in THF at room temperature gave the corresponding rotaxanes **9a-h** without formation of by-products. The use of a large excess of **8a-h** was mandatory for a complete conversion. This observation suggests that an equilibrium is reached in solution and the excess of **8a-h** is necessary to fully drive the equilibrium towards the formation of the desired products. Rotaxanes **9a-h** were thus prepared in good yields. The only exception was **9b**. In this particular case the bis(trifluoromethyl)phenyl ester is also an activated ester^[12f] and the use of 4 equiv. of **8b** was certainly not sufficient to fully displace the equilibrium towards the complete formation of **9b**. The conditions used for the preparation of rotaxane diesters **9a-h** were also successful for the preparation of **9i** starting from **4** and thiophenol **8i**. Rotaxane **9i** with two thiophenyl ester stoppers was thus obtained in 71% yield.

Scheme 3. Stopper exchange reactions with phenol, alcohol and thiophenol reagents. *Reagents and conditions:* (i) **4**, TBD, THF (**9a**: 86%, **9b**: 19%, **9c**: 79%, **9d**: 97%, **9e**: 86%, **9f**: 63%, **9g**: 80%, **9h**: 81%, **9i**: 71%); (ii) **4**, TBD, mixing in a Retsch MM400 mill at 30 Hz for 2 h (**9a**: 82%, **9b**: 88%, **9c**: 92%, **9d**: 78%, **9e**: 86%, **9f**: 88%, **9g**: 87%, **9h**: 87%, **9i**: 97%).

Compounds **9a-i** were also prepared under mechanochemical solvent-free conditions (Scheme 3). A mixture of rotaxane **4**, the appropriate nucleophile (**8a-i**, 2.3 equiv.) and TBD (2.3 equiv.)

was mixed in a Retsch MM400 mill at 30 Hz for 2 h. Rotaxanes **9a-i** were thus obtained in 78-97% yields. The transesterification reactions were clearly more efficient under the solvent-free conditions. Moreover, only a slight excess of **8a-i** was necessary to complete the reactions thus making the purification of the resulting rotaxanes easier.

All the rotaxanes were fully characterized by NMR and IR spectroscopies and mass spectrometry. Their purity was confirmed by elemental analysis data. Crystals suitable for X-ray structure analysis were obtained for rotaxanes **9c** and **9d**. The structures are depicted in Figure 2.

Figure 2. (A) ORTEP plot of the structure of **9c** (the two co-crystallized Et₂O molecules are omitted for clarity, thermal ellipsoids are shown at 30% probability level). (B) Stacking within the **9c** lattice highlighting the intermolecular π - π interactions between the stoppers of neighbouring rotaxanes. (C) ORTEP plot of the two conformers of **9d** present in the crystal lattice (thermal ellipsoids are shown at 30% probability level). Colour code: H: white, O: red, C: grey for the dumbbell and pale blue for the pillar[5]arene moiety.

Both compounds are chiral owing to their *D*₅-symmetry imposed by the pillar[5]arene subunits. Both **9c** and **9d** crystallized as racemates. Close inspection of the packing in the crystal lattice of **9c** along the crystallographic *a* axis revealed alternated layers of both enantiomers. Within these layers, the rotaxanes are organized in infinite supramolecular chains through the establishment of π - π interactions between the stoppers of neighboring molecules (Figure 2B). As a result, this supramolecular organization does not generate specific packing constraints as observed in the case of **4**. Therefore, the decyl chain of the axle component of **9c** is not compressed by the packing forces and adopts a fully extended all-anti conformation in the solid state. In the case of rotaxane **9d**, the supramolecular organization is more complex due to the existence of two

conformers within the crystal lattice (Figure 2C). For the first one, the decyl chains adopts a linear all anti conformation. In contrast, the second one is twisted due to the gauche conformation adopted by two successive methylene subunits located outside the cavity of the pillar[5]arene moiety. In the case of **9d**, close inspections of the crystal lattice revealed no particular π - π interactions between neighboring molecules and the packing is mainly dominated by Van der Waals forces.

Preparation of fullerene-containing photoactive rotaxanes.

The preparation of a wide range of rotaxanes was successively achieved by stopper exchange reactions from key precursor **4**. To further evaluate the potential of this new methodology, it was decided to prepare building block **11** starting from fulleropillar[5]arene **10** in order to prepare a photoactive MIM by the grafting of complementary photoactive stoppers to the fullerorotaxane scaffold (Scheme 4). Compound **10** was prepared as previously described.^[9] Owing to the moderate solubility of **10** in CHCl_3 , the reaction conditions optimized for the preparation of compound **4** had to be adapted. In order to favor the incorporation of **10** in host-guest complexes, an excess of diacyl chloride **2** (3 equiv.) was used in this case. Treatment of the 3:1 mixture of **2** and **10** dissolved in a minimum of CHCl_3 with pentafluorophenol (**3**) in the presence of Et_3N at -15°C afforded **11**. After work-up and purification, rotaxane **11** was obtained in 82% yield. The ^1H NMR spectrum of **11** showing the diagnostic signals of both subunits was fully consistent with the proposed structure. Moreover, the dramatic shielding observed for the signals arising from the methylene groups of the axle moiety provides definitive evidence for the interlocked structure of **11**. The up-field shifted chemical shifts of these signals results actually from their location within the cavity of the macrocycle and is due to the ring current effect of the fulleropillar[5]arene hydroxyquinone subunits. The MALDI-TOF mass spectrum of **11** showing the expected molecular ion peak at $m/z = 2170.65$ ($[\text{M}]^+$, calcd for $\text{C}_{139}\text{H}_{88}\text{F}_{10}\text{O}_{14}$: 2170.60) as the base peak fully confirmed the structure of this compound. The reaction conditions developed for the preparation of diamide [2]rotaxanes from **4** were then applied to **11**. Treatment of **11** with primary amine **6a** in THF provided rotaxane **12a** in 67% yield. Finally,

incorporation of photoactive boron-dipyrromethene (Bodipy) stoppers was achieved by reaction of **11** with **6b**. For solubility reasons, the stopper exchange reaction was carried out in a 2:1 THF/ CHCl_3 mixture. Photoactive MIM **12b** was thus obtained in 71% yield. Rotaxanes **12a-b** were characterized by a combination of several analytical techniques. In both cases, the expected molecular ion peak was observed in their MALDI-TOF mass spectra. The NMR data were also fully consistent with the proposed structures. Compound **12b** combines complementary chromophore subunits, namely a fullerene moiety grafted onto the macrocyclic unit and two Bodipy subunits acting as stoppers.^[19] The ground state absorption spectra of rotaxane **12b** and model compounds **6b**, **7b** and **12a** recorded in CH_2Cl_2 are depicted in Figure 3. The absorption spectrum of model Bodipy compounds **6b** and **7b** are dominated by an intense band centered at 524 and 526 nm, respectively. This band is attributed to the lowest energy spin-allowed π - π^* transition of the Bodipy chromophore.^[20] The typical higher energy π - π^* transitions of the bodipy moieties^[20] are also observed in the 280-380 nm region for both compounds. Comparison of the molar extinction coefficients (ϵ) of **6b** and **7b** are fully consistent with the presence of two Bodipy subunits in the rotaxane and the differences observed for **6b** and **7b** between 270 and 340 nm are due to transitions centered on the pillar[5]arene moiety of **7b** which are also contributing to the overall absorption in the UV region. On the other hand, the absorption spectrum of model rotaxane **12a** displays the characteristic features of methanofullerene derivatives^[21] with a very intense band in the UV (259 nm, $\epsilon = 113800 \text{ M}^{-1}\text{cm}^{-1}$) and a much weaker and broad absorption in the all visible region. The very weak band observed at 702 nm ($\epsilon = 320 \text{ M}^{-1}\text{cm}^{-1}$) is attributable to the lowest singlet transition.^[21] The absorption spectrum of multi-component fullerene-Bodipy system **12b** shows the characteristic features of both chromophores. Comparison with the absorption spectra of the corresponding model compounds **7b** and **12a** revealed only very minor differences and ground-state electronic interactions between the Bodipy stoppers and the central C_{60} subunit are negligible.

Scheme 4. Preparation of fullerene-containing rotaxanes. *Reagents and conditions:* (i) **2**, **3**, Et₃N, CHCl₃, -15°C (82%); (ii) **6a**, THF (**12a**: 67%); (iii) **6b**, THF/CHCl₃ (**12b**: 71%).

Figure 3. (A) Absorption spectra of **6b** (blue), **7b** (black), **12a** (green) and **12b** (red) recorded in CH₂Cl₂ at 25°C. (B) Emission spectra of isoabsorbing solutions of compounds **7b** (black) and **12b** (red) in CH₂Cl₂ at 25°C ($\lambda_{\text{exc}} = 490$ nm) and schematic representation of the photoexcitation of the Bodipy moiety followed by a strong emission in the case of **7b** and by energy transfer (ET) in the case of **12b**.

Emission spectra were also recorded for compounds **6b**, **7b** and **12a-b** in CH₂Cl₂ solutions at room temperature. Under our experimental conditions, the extremely weak fullerene emission ($\phi_{\text{em}} < 0.0001$)^[21] could not be detected for compound **12a**. In contrast, model compounds **6b** and **7b** exhibit the typical strong emission of Bodipy dyes at 535 nm ($\lambda_{\text{exc}} = 490$ nm). As shown in Figure 3B, the bodipy fluorescence is still detected for compound **12b** upon excitation at 490 nm where most of the light is absorbed by the Bodipy. However, comparison of the emission intensity of isoabsorbing solutions of **7b** and **12b** revealed a dramatic quenching (ca. 96%) of the Bodipy emission by the fullerene moiety in **12b** despite the absence of covalent linkage between these chromophores (Figure 3B). The bodipy dyes used as stoppers in **12b** are not particularly strong electron donors neither in their ground nor excited states.^[22] The Bodipy singlet excited state in **12b** is most likely quenched by an efficient energy transfer to populate the first fullerene singlet

excited state. Whereas it was not possible to fully confirm the energy transfer mechanism by monitoring the weak fullerene emission with our experimental setup, the low electron donating ability of the Bodipy stoppers in **12b** was supported by DFT calculations (see ESI for the details). The highest occupied molecular orbital (HOMO) is indeed located on hydroquinone subunits of the pillar[5]arene subunit of **12b** and not on the Bodipy stoppers. Only the HOMO-1 and HOMO-2 were found on the Bodipy subunits (Figure 4). As expected, the lowest occupied molecular orbital (LUMO) is located on the fullerene moiety of **12b**.

Figure 4. Frontier molecular orbitals calculated at the B3LYP/6-31G* level for compound **12b**. In the calculated structure, the boron atoms of the two stoppers are separated by a distance of 32.18 Å.

Owing to possible shuttling motions of the pillar[5]arene moiety along the axle of rotaxane **12b** and to the overall conformational flexibility of the system, the Bodipy-fullerene distance is fluctuating over a large range. Based on the theoretical calculations, when the fullerenopillar[5]arene is close to one Bodipy stopper in **12b** (Bodipy-fullerene distance of ca. 4 Å), the distance to the second Bodipy unit is quite large (ca. 22 Å). The though space quenching of the Bodipy emission is very efficient and apparently insensitive to dynamic fluctuations of the Bodipy-fullerene distance in **12b**. This is also in favor for a Bodipy to fullerene singlet-singlet energy transfer mechanism upon photoexcitation of the Bodipy subunits.

Conclusion

Rotaxane **4** incorporating pentafluorophenol ester stoppers has been prepared in a nearly quantitative yield. This building block is easily affordable on a multi-gram scale. It is also perfectly stable under standard laboratory conditions thus allowing storage for several months before further transformation. Stopper exchange reactions have been carried out from **4** and various nucleophiles to generate a wide range of pillar[5]arene-containing rotaxanes in high yields. The pentafluorophenyl ester activation method has been also applied to the construction of more sophisticated [2]rotaxanes incorporating a fullerene-functionalized pillar[5]arene moiety. This allowed us to easily prepare a photoactive MIM in which through space intercomponent photoinduced events have been evidenced. The versatility of this new methodology and its application for the easy construction of sophisticated rotaxanes are clear advantages for future developments in the field as the preparation of new molecular machines, advanced materials or bioactive compounds based on MIMs will not be limited by their synthesis anymore.

Experimental Section

Synthesis

The preparation and characterization of all the new compounds are described in the Supporting Information. Compounds **1**^[23], **6b**^[24] and **10**^[9] were prepared according to previously reported procedures.

X-ray crystal structures

The crystallographic data and the refinement parameters are reported in the Supporting Information for all the compounds. The X-ray crystal structures have been deposited at the Cambridge Structural Database (CCDC deposition numbers: 2058646 for **4**, 2058647 for **9c**, and 2059867 for **9d**). These data are provided free of charge by the joint Cambridge Crystallographic Data Centre and Fachinformationszentrum Karlsruhe Access Structures service (<https://www.ccdc.cam.ac.uk/structures>).

Acknowledgements

Financial support by the ANR (projects FastGiant ANR-17-CE07-0012-01 and Pillar ANR-19-CE06-0032), the International Center for Frontier Research in Chemistry and the LabEx "Chimie des Systèmes Complexes" is gratefully acknowledged. We thank L. Karmazin and C. Bailly for the X-ray crystal structure resolution, E. Wasielewski for high-field NMR measurements and J.M. Strub for the mass spectra. We further thank A. N. Kursunlu for the initial preparation of starting material **6b**.

Keywords: Stopper exchange • Pillar[5]arene • Rotaxanes • Supramolecular chemistry • Photoactive systems

- [1] a) J.-P. Sauvage, *Angew. Chem. Int. Ed.* **2017**, *56*, 11080-11093; *Angew. Chem.* **2017**, *129*, 11228-11242; b) J. F. Stoddart, *Angew. Chem. Int. Ed.* **2017**, *56*, 11228-11242; *Angew. Chem.* **2017**, *129*, 11244-11277; c) B. L. Feringa, *Angew. Chem. Int. Ed.* **2017**, *56*, 11060-11078; *Angew. Chem.* **2017**, *129*, 11228-11242.
- [2] a) V. Balzani, A. Credi, M. Venturi, *Molecular Devices and Machines – A Journey into the Nano World*, Wiley-VCH, **2003**; b) J.-P. Sauvage, C. Dietrich-Buchecker (Eds.), *Molecular Catenanes, Rotaxanes and Knots – A Journey Through the World of Molecular Topology*, J. Wiley and Sons, **2008**; c) E. R. Kay, D. A. Leigh, *Angew. Chem. Int. Ed.* **2015**, *54*, 10080-10088; *Angew. Chem.* **2015**, *127*, 10218-10226; d) C. J. Brunts, J. F. Stoddart, *The Nature of the Mechanical Bond – From Molecules to Machines*, J. Wiley & Sons, **2016**; e) S. Erbas-Cakmak, D. A. Leigh, C. T. McTernan, A. L. Nussbaumer, *Chem. Rev.* **2015**, *115*, 10081-10206; f) D. A. Leigh, *Angew. Chem. Int. Ed.* **2016**, *55*, 14506-14508; *Angew. Chem.* **2016**, *128*, 14722-14724; g) S.-J. Rao, Q. Zhang, J. Mei, X.-H. Ye, C. Gao, Q.-C. Wang, D.-H. Qu, H. Tian, *Chem. Sci.* **2017**, *8*, 6777-6783; h) S.-J. Rao, K. Nakazono, X. Liang, K. Nakajima, T. Takata, *Chem. Commun.* **2019**, *55*, 5231-5234; i) Q. Zhang, S.-R. Rao, T. Xie, X. Li, T.-Y. Xu, D.-W. Li, D.-H. Qu, Y.-T. Long, H. Tian, *Chem* **2018**, *4*, 2670-2684.
- [3] a) J.-C. Chambron, V. Heitz, J.-P. Sauvage, *J. Chem. Soc., Chem. Commun.* **1992**, 1131-1133; b) J.-C. Chambron, A. Harriman, V. Heitz, J.-P. Sauvage, *J. Am. Chem. Soc.* **1993**, *115*, 6109-6114; c) J.-C. Chambron, A. Harriman, V. Heitz, J.-P. Sauvage, *J. Am. Chem. Soc.* **1993**, *115*, 7419-7425; d) J.-C. Chambron, V. Heitz, J.-P. Sauvage, *J. Am. Chem. Soc.* **1993**, *115*, 12378-12384; e) M. Linke, J.-C. Chambron, V. Heitz, J.-P. Sauvage, *J. Am. Chem. Soc.* **1997**, *119*, 11329-11330; f) M.-J. Blanco, J.-C. Chambron, V. Heitz, J.-P. Sauvage, *Org. Lett.* **2000**, *2*, 3051-3054; g) M. Linke, J.-C. Chambron, V. Heitz, J.-P. Sauvage, S. Encinas, F. Bargelletti, L. Flamigni, *J. Am. Chem. Soc.* **2000**, *122*, 11834-11844; h) M. Andersson, M. Linke, J.-C. Chambron, J. Davidsson, V. Heitz, L. Hammarström, J.-P. Sauvage, *J. Am. Chem. Soc.* **2002**, *124*, 4347-4362.
- [4] For reviews, see: a) A. Harriman, J.-P. Sauvage, *Chem. Soc. Rev.* **1996**, *25*, 41-48; b) M.-J. Blanco, M. C. Jiménez, J.-C. Chambron, V. Heitz, M. Linke, J.-P. Sauvage, *Chem. Soc. Rev.* **1999**, *28*, 293-305; c) J. A. Faiz, V. Heitz, J.-P. Sauvage, *Chem. Soc. Rev.* **2009**, *38*, 422-442.
- [5] a) F. Diederich, C. Dietrich-Buchecker, J.-F. Nierengarten, J.-P. Sauvage, *J. Chem. Soc., Chem. Commun.* **1995**, 781-782; b) N. Armaroli, F. Diederich, C. O. Dietrich-Buchecker, L. Flamigni, G. Marconi, J.-F. Nierengarten, J.-P. Sauvage, *Chem. Eur. J.* **1998**, *4*, 406-416; c) J. D. Megiatto Jr., R. Spencer, D. I. Schuster, *Org. Lett.* **2009**, *11*, 4152-4155; d) J. D. Megiatto Jr., K. Li, D. I. Schuster, A. Palkar, M. A. Herranz, L. Echegoyen, S. Abwandner, G. de Miguel, D. M. Guldi, *J. Phys. Chem. B* **2010**, *114*, 14408-14419; e) J. D. Megiatto Jr., D. I. Schuster, S. Abwandner, G. de Miguel, D. M. Guldi, *J. Am. Chem. Soc.* **2010**, *132*, 3847-3861.
- [6] For a review, see: J. D. Megiatto Jr., D. I. Schuster, D. M. Guldi, *Chem. Soc. Rev.* **2020**, *49*, 8-20.
- [7] For selected examples, see: a) S. Saha, A. H. Flood, J. F. Stoddart, S. Impellizzeri, S. Silvi, M. Ventura, A. Credi, *J. Am. Chem. Soc.* **2007**, *129*, 12159-12171; b) T. Ogoshi, D. Yamafuji, T.-a. Yamagishi, A. M. Brouwer, *Chem. Commun.* **2013**, *49*, 5468-5470; c) T. A. Barendt, I. Rasovic, M. A. Lebedeva, G. A. Farrow, A. Auty, D. Cherkulaev, I. V. Sazanovich, J. A. Weinstein, K. Porfyrakis, P. D. Beer, *J. Am. Chem. Soc.* **2018**, *140*, 1924-1936; d) Y. Xu, R. Kaur, B. Wang, M. B. Minameyer, S. Gsänger, B. Meyer, T. Drewello, D. M. Guldi, M. von Delius, *J. Am. Chem. Soc.* **2018**, *140*, 13413-13420; e) M. Wolf, A. Ogawa, M. Bechtold, M. Vonesch, J. A. Wytko, K. Oohara, S. Campidelli, T. Hayashi, D. M. Guldi, J. Weiss, *Chem. Sci.* **2019**, *10*, 3846-3853; f) S. Yang, C.-AX. Zhao, S. Crespi, X. Li, Q. Zhang, Z.-H. Zhang, J. Mei, H. Tian, D.-H. Qu, *Chem* **2021**, DOI: 10.1016/j.chempr.2021.02.019.
- [8] B. Delavaux-Nicot, H. Ben Aziza, I. Nierengarten, T. M. N. Trinh, E. Meichsner, M. Chessé, M. Holler, R. Abidi, E. Maisonhaute, J.-F. Nierengarten, *Chem. Eur. J.* **2018**, *24*, 133-140.
- [9] E. Meichsner, I. Nierengarten, M. Holler, M. Chessé, J.-F. Nierengarten, *Helv. Chim. Acta* **2018**, *101*, e1800059.
- [10] I. Nierengarten, E. Meichsner, M. Holler, P. M. Pieper, R. Deschenaux, B. Delavaux-Nicot, J.-F. Nierengarten, *Chem. Eur. J.* **2018**, *24*, 169-177.
- [11] I. Nierengarten, J.-F. Nierengarten, *ChemistryOpen* **2020**, *9*, 393-400.
- [12] The post-modification of pre-constructed [2]rotaxanes by stopper exchange reactions has been rarely used and only a few examples

- have been reported so far, see: a) S. J. Rowan, J. F. Stoddart, *J. Am. Chem. Soc.* **2000**, *122*, 164-165; b) S. J. Rowan, S. J. Cantrill, J. F. Stoddart, A. J. P. White, D. J. Williams, *Org. Lett.* **2000**, *2*, 759-762; c) R. J. Bordoli, S. M. Goldup, *J. Am. Chem. Soc.* **2014**, *136*, 4817-4820; d) D. W. Zehnder II, D. B. Smithrud, *Org. Lett.* **2001**, *3*, 2485-2487; e) N. Kihara, S. Motoda, T. Yokozawa, T. Takata, *Org. Lett.* **2005**, *7*, 1199-1202; f) J. S. Hannam, S. M. Lacy, D. A. Leigh, C. G. Saiz, A. M. Z. Slawin, S. G. Stichelell, *Angew. Chem. Int. Ed.* **2004**, *43*, 3260-3264; *Angew. Chem.* **2004**, *116*, 3322-3326; g) S. C. Rajappan, D. R. McCarthy, J. P. Campbell, J. B. Ferrel, M. Sharafi, O. Ambrozaite, J. Li, S. T. Schneebeli, *Angew. Chem. Int. Ed.* **2020**, *59*, 16668-16674; *Angew. Chem.* **2020**, *132*, 16811-16817; h) For a recent review on post-synthetic modification of interlocked molecules, see: P. Waelès, M. Gauthier, F. Coutrot, *Angew. Chem. Int. Ed.*, doi.org/10.1002/anie.202007496; *Angew. Chem.*, doi.org/10.1002/ange.202007596.
- [13] S.-Y. Han, Y.-A. Kim, *Tetrahedron* **2004**, 2447-2467.
- [14] S. R. Samanta, R. Cai, V. Percec, *Polym. Chem.* **2015**, *6*, 3259-3270.
- [15] R. Milev, A. Lopez-Pacheco, I. Nierengarten, T. M. N. Trinh, M. Holler, R. Deschenaux, J.-F. Nierengarten, *Eur. J. Org. Chem.* **2015**, 479-485.
- [16] M. Holler, T. Stoerkler, A. Louis, F. Fischer, J.-F. Nierengarten, *Eur. J. Org. Chem.* **2019**, 3401-3405.
- [17] For other examples of mechanochemical synthesis of MIMs, see: a) A. Orita, J. Okano, Y. Tawa, L. Jiang, J. Otera, *Angew. Chem. Int. Ed.* **2004**, *43*, 3724-3728; *Angew. Chem.* **2004**, *116*, 3810-3814; b) S. Y. Hsueh, K. W. Cheng, C.-C. Lai, S.-H. Chiu, *Angew. Chem. Int. Ed.* **2008**, *47*, 4436-4439; *Angew. Chem.* **2008**, *47*, 4508-4511; c) C.-C. Hsu, N. C. Chen, C.-C. Lai, Y.-H. Liu, S.-M. Peng, S.-H. Shiu, *Angew. Chem. Int. Ed.* **2008**, *47*, 7475-7478; *Angew. Chem.* **2008**, *47*, 7585-7588; d) C.-C. Hsu, C.-C. Lai, S.-H. Chiu, *Tetrahedron* **2009**, *65*, 2824-2829; e) P.-N. Chen, C.-C. Lai, S.-H. Chiu, *Org. Lett.* **2011**, *13*, 4660-4663; f) K.-D. Wu, Y.-H. Lin, C.-C. Lai, S.-H. Chiu, *Org. Lett.* **2014**, *16*, 1068-1071; g) H.-G. Li, G.-W. Wang, *J. Org. Chem.* **2017**, *82*, 6341-6348.
- [18] For reviews on supramolecular concepts in mechanochemistry, see: a) T. Friščić, *Chem. Soc. Rev.* **2012**, *41*, 3493-3510; b) H.-G. Li, L. Li, G.-W. Wang, *Current Org. Chem.* **2018**, *22*, 923-929; c) A. Inthasot, T.-Y. Tai, S.-H. Chiu, *J. Chin. Chem. Soc.* **2019**, *66*, 134-145.
- [19] For selected examples of covalent photoactive molecular devices combining C₆₀ with Bodipy dyes, see: a) R. Ziessel, B. D. Allen, D. B. Rewinska, A. Harriman, *Chem. Eur. J.* **2009**, *15*, 7382-7393; b) J.-Y. Liu, M. E. El-Khouly, S. Fukuzumi, D. K. P. Ng, *Chem. Asian J.* **2010**, *6*, 174-179; c) J. Iehl, J.-F. Nierengarten, A. Harriman, T. Bura, R. Ziessel, *J. Am. Chem. Soc.* **2012**, *134*, 988-998; d) V. Bandi, H. B. Gobeze, P. A. Karr, F. D'Souza, *J. Phys. Chem. C* **2014**, *118*, 18969-18982; e) C. B. KC, G. N. Lim, V. N. Nesterov, P. A. Karr, F. D'Souza, *Chem. Eur. J.* **2014**, *20*, 17100-17112.
- [20] A. Loudet, K. Burgess, *Chem. Rev.* **2007**, *107*, 4891-4932.
- [21] a) N. Armaroli, G. Accorsi, D. Felder, J.-F. Nierengarten, *Chem. Eur. J.* **2002**, *8*, 2314-2323; b) J.-F. Eckert, J.-F. Nicoud, J.-F. Nierengarten, S.-G. Liu, L. Echegoyen, F. Barigelletti, N. Armaroli, L. Ouali, V. Krasnikov, G. Hadzioannou, *J. Am. Chem. Soc.* **2000**, *122*, 7467-7479.
- [22] T. Lazarides, G. Charalambidis, A. Vuillamy, M. Réglie, E. Klontzas, G. Froudakis, S. Kuhri, D. M. Galdi, A. G. Coutsolelos, *Inorg. Chem.* **2011**, *50*, 8926-8936.
- [23] a) T. Ogoshi, K. Kitajima, T. Aoki, S. Fujinami, T.-a. Yamagishi, Y. Nakamoto, *J. Org. Chem.* **2010**, *75*, 3268-3273; b) M. Holler, N. Allenbach, J. Sonet, J.-F. Nierengarten, *Chem. Commun.* **2012**, 2576-2578.
- [24] A. N. Kursunlu, C. Baslak, *Tetrahedron Lett.* **2018**, *59*, 1958-1962.

Entry for the Table of Contents

Pillar[5]arene-containing [2]rotaxane building blocks bearing pentafluorophenyl ester functions have been prepared and stopper exchange reactions carried out to generate a wide range of [2]rotaxanes with amide, ester or thioester stoppers. Starting from a fullerene-functionalized pillar[5]arene derivative, this new strategy gave easy access to a photoactive [2]rotaxane incorporating a C₆₀ moiety and two Bodipy stoppers.

Institute and/or researcher Twitter usernames: @INC_CNRS - @Unistra - @LIMA_UMR7042 (Institutions) / @IwonaNieren - @nierengarten6 (Researchers).