Dispersal limitation by structures is more important than intermittent drying effects for metacommunity dynamics in a highly fragmented river network

Mailys Gauthier, Guillaume Le Goff, Bertrand Launay, Christophe J Douady, T. Datry

► To cite this version:

Mailys Gauthier, Guillaume Le Goff, Bertrand Launay, Christophe J Douady, T. Datry. Dispersal limitation by structures is more important than intermittent drying effects for metacommunity dynamics in a highly fragmented river network. *Freshwater Science*, 2021, 40 (2), pp.302 - 315. 10.1086/714376 . hal-03369180

HAL Id: hal-03369180

<https://hal.science/hal-03369180>

Submitted on 7 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial| 4.0 International License

Dispersal limitation by structures is more important than intermittent drying effects for metacommunity dynamics in a highly fragmented river network

Mailys Gauthier^{1,2,3}, Guillaume Le Goff^{1,4}, Bertrand Launay^{1,5}, Christophe J. Douady^{2,6}, and Thibault Datry^{1,7}

¹Institut National de la Recherche Agronomique (INRAE), UR-Riverly, Centre de Lyon-Villeurbanne, 5 rue de la Doua CS70077, 69626 Villeurbanne Cedex, France

²University of Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, UMR5023 LEHNA, F-69622 Villeurbanne, France

Abstract: In the Anthropocene, river networks are globally threatened by human activities, creating dramatic consequences for aquatic biodiversity. Permanent and temporary fragmentation by manmade structures and drying events, respectively, are both increasing in rivers worldwide. Although both of these fragmentation types can limit species dispersal, their individual and combined effects in shaping metacommunity dynamics at both local (site-level) and regional (network-level) scales have not been explored. Here, we examined whether processes structuring aquatic invertebrate metacommunities vary through time in response to flow variability in a river network affected by drying and severe fragmentation by manmade structures. We also compared the relative influences of permanent and temporary fragmentation on metacommunity dynamics and hypothesized that permanent fragmentation would be the primary determinant of metacommunity dynamics. We conducted an intensive sampling effort (30 sampling sites × 6 dates) and measured 12 local environmental variables and 4 spatial distances to assess the relative importance of niche- and dispersal-based processes on benthic invertebrate metacommunities across hydrological phases. Spatial distances considered here were overland, network, and 2 fragmentation distances integrating the permanent fragmentation by manmade structures and temporary fragmentation by drying. We used Mantel tests to identify relationships between community dissimilarities and environmental and spatial distance matrices. We identified temporal variability in metacommunity processes with a predominant role of dispersal and no effect of niche-based processes. Metacommunities were shaped primarily by permanent fragmentation, whereas we detected no effect of fragmentation by network-scale drying. Dispersal-based metacommunity processes varied over time, following the wet–dry cycles that characterize dynamic river networks. Our results suggest the importance of key hydrological phases that should be incorporated into conservation perspectives. In addition, we highlight the need to quantify context dependency in metacommunity studies to optimize biodiversity conservation strategies in river-network ecosystems.

Key words: fragmentation, metacommunity, intermittent rivers, temporal variability, manmade structures, drying, benthic invertebrates, dispersal, beta diversity

Ecosystem fragmentation, the division of habitats into smaller and isolated fragments separated by a matrix of human-transformed land cover, is a major threat to biodiversity in the Anthropocene era (Haddad et al. 2015, Crooks et al. 2017). Fragmentation decreases habitat availability and alters dispersal, i.e., the movements of organisms among local habitat patches within ecosystems (Leibold et al. 2004, Brown et al. 2011, Heino et al. 2017). Dispersal interacts with local

environmental selection to play a major role in promoting species coexistence at regional levels (Poff 1997, Leibold et al. 2004, Brown and Swan 2010, Sarremejane et al. 2020). Globally, human activities, including land-use change, dams, and extraction of surface and ground water, are modifying runoff patterns through catchments and altering river flow regimes (de Graaf et al. 2019, Reid et al. 2019). In recent decades, the speed of these changes is unprecedented, accelerating the

E-mail addresses: ³mailys.gauthier@inrae.fr; ⁴guillaume.legoff@inrae.fr; ⁵bertrand.launay@inrae.fr; ⁶christophe.douady@univ-lyon1.fr; ⁷thibault.datry@inrae.fr

DOI: 10.1086/714376. Received 25 June 2020; Accepted 29 December 2020; Published online XX Month 2021; Associate Editor, Bryan L. Brown. Freshwater Science. 2021. 40(2):000–000. © 2021 by The Society for Freshwater Science.

000

fragmentation and intensity of flow regulation of river networks in most parts of the world (Grill et al. 2019). These alterations threaten freshwater biodiversity, which is experiencing higher extinction rates than in other ecosystems (Reid et al. 2019).

Increased hydropower development in many countries has fragmented river networks through the construction of large and small structures, such as dams, weirs, and culverts (Zarfl et al. 2014). Additionally, the increasing frequency of droughts in recent decades has motivated the development of small water retention structures, which are often located within streams (Habets et al. 2014). These structures include small to very small impoundments with an impounded area $<0.1 \text{ km}^2$ and a volume $<0.2 \text{ hm}^3$ (Lehner et al. 2011). In spite of the low volume of stored water, these impoundments represent 99.5% of the total number of reservoirs worldwide and are the most common anthropogenic fragmentation structures in river networks (Lehner et al. 2011). Together with large dams, they constitute barriers that permanently fragment rivers and, thus, prevent dispersal of organisms within or across networks (Fuller et al. 2015).

Temporary fragmentation of river networks is also becoming increasingly common as a result of natural drying events caused by water abstraction and climate change (Datry et al. 2018, de Graaf et al. 2019). Drying is defined here as the cessation of flow or the complete disappearance of surface water in a given reach (Ruhí et al. 2017). Of the total length of river networks globally, an estimated $\frac{1}{2}$ is prone to drying, and this proportion is increasing in regions experiencing drying trends and over-abstraction (Döll and Schmied 2012). The local effects of drying on aquatic biodiversity are fairly well understood (e.g., Datry et al. 2017), but the effects of drying-induced fragmentation on aquatic biodiversity remain understudied at the network scale for biotic groups, including aquatic invertebrates (but see Cañedo-Argüelles et al. 2015, Crabot et al. 2020, Gauthier et al. 2020, Sarremejane et al. 2020).

Metacommunities, defined as sets of local communities connected by dispersal (Leibold et al. 2004), are also affected by both temporary and permanent fragmentation, which alters the dispersal of organisms among localities (Fagan 2002) and can change metacommunity dynamics (Cañedo-Argüelles et al. 2015). Many studies have used the biogeographical framework of distance decay relationships (Nekola and White 1999) to identify local, niche-based processes as the main determinants of freshwater metacommunity dynamics (Heino et al. 2015). Alternatively, metacommunities in fragmented freshwater systems can be structured by dispersal-based processes (Thrush et al. 2008, Brown et al. 2011, Sarremejane et al. 2017, Gauthier et al. 2020). However, whether this effect of fragmentation is stable or varies over time, notably according to flow variability and wet–dry cycles, remains unknown (Ruhí et al. 2017). During drying periods, local environmental conditions can change dramatically (e.g., increased temperature,

biotic interaction changes, decreased dissolved oxygen), which could favor niche-based processes. In contrast, during rewetting phases, dispersal could become the major determinant of metacommunity structure because of active and passive recolonization (Sarremejane et al. 2017). Characterizing temporal variability in metacommunity processes in river networks prone to drying will promote more efficient monitoring and management of their biodiversity.

The ways in which permanent and temporary fragmentation of river networks jointly shape aquatic metacommunity dynamics may have implications for biodiversity conservation but have not been explored. When manmade structures permanently limit aquatic organisms' dispersal, the effect of drying events on metacommunity dynamics could be reduced because the movement of organisms is already constrained by manmade structures and, thus, temporary dispersal limitation could be undetectable. In contrast, where drying extends over long distances (e.g., several km), dry sections could impose stronger dispersal barriers than manmade structures, particularly for strictly aquatic organisms. Therefore, contextualizing the combined effects of drying with manmade structures on riverine biodiversity is fundamental because focusing on one type of fragmentation or the other could lead to contrasting conservation strategies.

Here, we explored the temporal dynamics of aquatic invertebrate metacommunities in a river network prone to drying and severe fragmentation by manmade structures through the biogeographical distance decay lens. We first hypothesized that processes structuring metacommunity would change within a hydrological year (Fig. 1A). Accordingly, we predicted niche-based processes to be the predominant determinants of metacommunity dynamics during drying periods as shifts from lotic to lentic environmental conditions develop. Conversely, we predicted dispersal-based processes to be predominant upon flow resumption, when organisms recolonize previously dry habitats from perennial refuges in the network. We also hypothesized that permanent fragmentation by manmade structures would be the primary factor determining metacommunity dynamics in this network (Fig. 1B). Accordingly, we predicted stronger relationships between community dissimilarity and physical distances in reaches with only permanent fragmentation (i.e., manmade structures) compared with those including drying events.

METHODS

To test our hypotheses, we coupled an intensive sampling campaign in both time and space to a fine-scale description of permanent and temporary fragmentation patterns in the Thouaret river network, which is a sub-tributary of the Loire river in the Deux-Sèvres department in western France (Fig. 2A). We did this by characterizing environmental variables and sampling benthic invertebrates at river network sites and by

Figure 1. Theoretical contribution of dispersal-based and niche-based processes on metacommunity assembly in fragmented river systems across hydrological phases (A) and expected beta diversity relationships with permanent (i.e., artificial or natural barriers) and temporary (i.e., drying events) fragmentation in such systems (B).

identifying temporary (drying events) and permanent (man-made structures) fragmentation in the river network. We then used Mantel tests to address our 1st hypothesis by investigating metacommunity patterns across hydrological phases. We also used Mantel tests to address our 2nd hypothesis and assess the relative importance of permanent and temporary fragmentation on metacommunity dynamics.

Sampling design

Site selection We selected 30 sampling sites, including sites in 13 perennial and 17 intermittent reaches (Fig. 2B–D), to encompass spatial variability within the Thouaret net-

work and to control for the effects of confluences and man-made structures on community composition. The Thouaret's hydrological regime reflects the region's oceanic climate with rainy and cold winters (for the period 1958–2019: mean $\pm$ SD cumulative rainfall 233 ± 87 mm, mean temperature $5.2 \pm 1.4^\circ\text{C}$; Vidal et al. 2010) and hot and dry summers (mean cumulative rainfall 135 ± 59 mm, mean temperature $18.0 \pm 1.3^\circ\text{C}$). The impermeable granitic bedrock of the region results in flashy flood events and a lack of baseflow during summer. Hydrological variability is, thus, considerable, ranging between flooding and severe summer low flows and associated drying events (Datry et al. 2016c). The 96-km² Thouaret headwater network drains an area dominated by agriculture

Figure 2. The study river network, the Thouaret, situated in western France (A), during different hydrological phases (B–D), with site locations, manmade structures, and drying patch locations. Hydrological phases for which the entire catchment is flowing (i.e., wet [March, April], drying [June] and rewetting [December]) (B) and the dry phase: September (C) and October (D). Small water retention structures are shown in light blue. Manmade structures are only presented in panel B by black double lines on the network for the sake of clarity.

that includes pasture (35.4%), arable land (28.3%), complex cultivation patterns (23.9%), forest (10%), and urban areas (2.4%). Thousands of small water retention structures have been constructed for agriculture and public water supply in the past decades (exact installation dates are unknown), and some small lentic water bodies along and within the river network are used for recreation and aquaculture activities. These structures permanently fragment the network (Fig. 2B). In addition, dry reaches temporarily fragment the network during low-flow periods between August and October (Fig. 3).

From March to December 2018, we sampled each site 6×, if flowing, to characterize different hydrological phases. One sampling date was during the wet phase in March, 2 dates were during the drying phase in April and June, 2 dates were during the dry phase in September and October, and 1 date was in December during the rewetting phase (Fig. 3).

Characterization of environmental variables We measured 12 environmental variables at each sampling site on each sampling date. At each invertebrate sampling point (see below), we measured water depth (cm), wetted width (cm), flow velocity (cm/s, measured at $0.4 \times$ water depth with a Mini-water[®]20 flow meter; Schiltknecht Messtechnik, Gossau, Switzerland), and main and secondary substrate types. We classified substrate type into 14 classes (mud, silt, sand, gravel, pebbles, rocks, slab, litter, root tufts, wood, algae, helophyte, spermaphyte, and bryophyte). We obtained elevation (m), latitude, and longitude at each site with a GPSMAP[®] 62ST handheld navigator (Garmin[®] International, Olathe, Kansas) and measured specific conductivity ($\mu\text{S}/\text{cm}$) and temperature with an HQ14d meter and pH and dissolved oxygen percent-

age with an HQ40d multi meter (Hach[®], Loveland, Colorado). Along an upstream distance of $20 \times$ the wetted width of the sampling point, we visually assessed % canopy cover and quantified channel slope with a clinometer. We selected these environmental variables because they are useful in characterizing riffle-associated invertebrate community habitat (Karaouzas et al. 2019).

Benthic invertebrate sampling To address our hypotheses by exploring between-date differences in metacommunity patterns, we sampled organisms from riffle habitat, which is the 1st stream habitat type to dry during drying events (Arscott et al. 2010, Datry et al. 2014, Ruhí et al. 2017, Crabot et al. 2020). This sampling design was validated from a metacommunity perspective through preliminary analysis of 2 other datasets (see Tables S2, S3). We used a Surber sampler (area 0.1 m^2 , mesh size $500 \mu\text{m}$; Sefar, Switzerland) to collect samples, which we then preserved in 96% ethanol. We counted and identified organisms to the lowest practical taxonomic level with available identification keys (list of references for identification available on Zenodo, <https://doi.org/10.5281/zenodo.3569563>). We identified most insects, crustaceans, triclads, and all mollusks to species except for immature specimens lacking identification features. At a minimum, we identified Diptera to genus, sub-family, or tribe depending on family, Coleoptera, Hemiptera, Trichoptera, and Crustacea (Amphipoda) to genus, and Ostracoda and Copepoda to order. We identified Achaeta and Tricladida to genus and identified Nematoda, Acarina, Hydrozoa, and Oligochaeta to order or sub-order. We assigned each taxon to 1 of 2 dispersal categories according to Tachet et al. (2000):

Figure 3. Discharge time series for the year 2018 of a gauging station situated 55 km downstream of the studied Thouaret headwater. Hydrological phases are shown. Sampling dates are represented by vertical black arrows on the x -axis. Data were downloaded from the Hydro database (www.hydro.eaufrance.fr): gauging station L8213010.

1) strictly aquatic invertebrates that can only disperse in water (57 taxa), and 2) those with an aerial life stage (110 taxa), which can also disperse by flying or crawling.

Description of temporary and permanent fragmentation

To identify temporary fragmentation by drying, we quantified drying events by cross-referencing 3 different information sources. First, volunteer members of Carg'eau association (Collectif Associatif Régional pour la Gestion de l'EAU; www.eau-poitou-charentes.org) made visual observations of the Thouaret every 2 wk from June to October of 2018. Observations were classified as 1 of 4 states: 1) flowing, when water flowed across the whole channel width; 2) low flow, when <50% of the reach width was flowing; 3) no flow, when the reach was dry apart from disconnected pools; and 4) dry, when surface water was absent. Observations were made along the entire network whenever possible, depending on accessibility. Datry et al. (2016c) provide a detailed description of the observation methods. Second, within <4 d of the volunteers' observations, we visually assessed the same instream states at each sampling site and along each reach visited during the sampling campaigns. Finally, we installed Hobo® water presence loggers (Onset®, Bourne, Massachusetts) at each sampling site on the streambed in riffle heads. The loggers recorded the presence or absence of surface water every 2 h from March to December 2018. We used this water presence data to calculate the dry duration at each site, defined as the time in seconds without flowing water (i.e., no-flow and dry states). To characterize temporary fragmentation, we calculated the number of dry reaches, the mean and total length of dry reaches, and the percentage of the catchment, as in Datry et al. (2016c). Dry patch delimitations were integrated into a local database based on the larger Carthage network database (<https://geo.data.gouv.fr/fr/datasets/54917fd94fb1cd2fcb6f5d3295dbf33acaf1847e>). To characterize permanent fragmentation by manmade structures, we calculated density (no. of structures/km) and mean structure height for all structures in the network. All measurements related to manmade structures were based on a national Référentiel des Obstacles à l'Écoulement (ROE) database (<http://www.sandre.eaufrance.fr/atlas/srv/fre/catalog.search#/metadata/59057026-b40c-4cf9-9e3e-7296e0aa1a78>). We used QGIS (v3.10; QGIS Development Team, <https://qgis.org/en/site/>) modules *postres*, *postgis*, and *pgrouting* to calculate all network measurements for both temporary and permanent fragmentation.

Data analysis

Hypothesis 1: Temporal variation of metacommunity processes within a hydrological year To investigate metacommunity patterns across hydrological phases, we used Mantel tests (Legendre and Legendre 2012) to assess the relationships between community beta diversity and environmental and spatial distances between pairwise combinations

of sites on each sampling date. We created beta diversity matrices based on the Chao dissimilarity index to measure dissimilarity after $\log(x + 1)$ transformation of abundance data to better represent rare taxa (Cañedo-Argüelles et al. 2015) with the *vegdist* function in the *vegan* package (Oksanen et al. 2019) in R (version 1.3; R Project for Statistical Computing, Vienna, Austria). For each sampling date, we calculated beta diversity matrices for the whole community and assemblages of taxa in the 2 dispersal categories (i.e., strictly aquatic taxa and taxa with an aerial life stage). To represent niche-based processes, we constructed environmental distance matrices using Euclidean distances based on the 12 environmental variables and the dry duration, after variable standardization (mean $\pm$ 1SD = 0) with *vegdist*. To represent distance-based processes, we assessed 2 spatial distance matrices commonly used in stream metacommunity research (e.g., Sarremejane et al. 2017): an overland matrix based on Euclidean distances constructed from site GPS coordinates with *vegdist*, and a network matrix based on instream distances within the river network constructed using the theoretical hydrographic network described by Pella et al. (2012). For environmental distances, we used the Moran spectral randomization procedure to account for spatial patterns of the environment and to avoid inflation of correlations in Mantel tests (Crabot et al. 2019). We used the Holm correction to adjust *p*-values for multiple comparisons in Mantel tests (Heino and Tolonen 2017).

Hypothesis 2: Effects of permanent and temporary fragmentation on metacommunity structure To assess the importance of both permanent and temporary fragmentation on metacommunity structure, we used Mantel tests to assess relationships between the community dissimilarity matrix described above and 2 fragmentation distances. We used QGIS to calculate a permanent fragmentation matrix by summing the height of manmade structures (ROE database) in meters between pairwise combinations of sites because height can affect the ability of organisms to pass a structure (Fuller et al. 2015). Although this approach did not distinguish the effect of 1 large dam from that of several smaller structures, cumulative height can be more relevant than number of structures when studying network-scale dispersal (Perkin et al. 2015). We also used QGIS to calculate the temporary fragmentation matrix by summing the length in meters of dry patches observed in September between pairwise combinations of sites. We observed the maximum fragmentation by drying in September and, thus, we determined that performing Mantel tests on matrices that included only the final 3 sampling dates was more ecologically meaningful than including all sampling dates. We explored the correlation between the permanent and temporary fragmentation matrices to determine whether to use classic or partial (i.e., correcting for the influence of the other matrix) Mantel tests. We did not find a correlation between

the 2 fragmentation matrices (Mantel test, $r = 0.10$, $p = 0.21$), so we used classic Mantel tests to explore the influence of fragmentation type on community dissimilarities.

RESULTS

Environmental variability

We observed considerable spatial variability in environmental variables on each sampling date (see results in Table 1). The network included sites experiencing a wide range of hydrological conditions, including small and shallow to large and deep reaches (e.g., June wetted widths of 4.7–419.5 cm and depths of 0.6–25.0 cm) as well as standing to fast-flowing conditions (e.g., June flow velocity: 0–0.25 m/s). Physicochemical variables, except for pH, were also spatially variable, and canopy cover encompassed both unshaded and forested reaches. As is typical in riffle habitats, the main substrates were pebbles and gravel, across all sites and dates. The secondary substrate type was mainly composed of gravel, followed by sand and pebbles.

Permanent and temporary fragmentation of the Thouaret network

Both permanent and temporary fragmentation were observed throughout the Thouaret network. We recorded 32 manmade structures, corresponding to a density of 0.37 manmade structures/linear km (Table 2). Dams were the most abundant structures ($n = 15$), supporting water retention for agriculture and aquaculture purpose (60%) and recreation activities (40%). Other manmade structures included weirs ($n = 12$), bridge inverts ($n = 3$), and culverts ($n = 2$). Manmade structure heights ranged from 0.2 to 4.5 m (mean $\pm$ SD: 1.7 ± 1.3 m). We observed temporary fragmentation by dry reaches on both dry-phase sampling dates (September and October; Fig. 2C, D). There were 20 dry reaches in September and 14 in October, with a mean length of 20.6 and 20.3 km, respectively. Dry reaches represented 46.1 and 32.6% of the network in September and October, respectively. Drying duration ranged from 5 h to 3 mo (mean duration: 4.2 wk).

Invertebrate communities

Benthic invertebrate sampling yielded a high number of individuals with similar richness across sites but variable richness over the hydrologic year. We collected a total of 74,337 individuals from 167 taxa with a mean ($\pm$ SD) of 15,723 (± 3188) individuals/date (Table S1) and of 1226 (± 1146) individuals from 19 (± 6) taxa/sample. On average, 58% ($\pm 2.8\%$) of taxa had an aerial stage and 42% ($\pm 2.8\%$) were strictly aquatic. Communities were dominated by Oligochaeta (relative abundance: 32.3%), Tanytarsini (Diptera, 9.1), and Gammaridae (Amphipoda, 8.8%). Oligochaeta was also the most ubiquitous taxon, occurring at 92% of all sites, followed by Orthocladiinae (Diptera, 84.4%) and Tanytarsini (Diptera, 72.5%). Taxa richness was 21.2 (± 5.8) taxa/sample

across all sites and dates (Table S1) and were comparable at intermittent and perennial sites (Student's t -test, $t = -0.006$, $p = 0.99$). Highest average taxa richness was in September (26.9), which corresponded to the dry period when flowing reaches can act as refuges (Bonada et al. 2017) and when density of species is high because of decreased habitat area (Dewson et al. 2007).

Hypothesis 1: Temporal variation of processes influencing metacommunity dynamics

The processes determining aquatic invertebrate metacommunities varied during the hydrological year, supporting our 1st hypothesis; however, our prediction that niche-based processes would dominate during drying periods was not supported. Overland distances were associated with community dissimilarities during the dry phase (mean Mantel's $r = 0.24$, p -value: 0.03–0.04), whereas associations were lower during the rewetting (mean $r = 0.13$, p -value: 0.01–0.04) and wet ($r = 0.12$, p -value = 0.04) phases, and we found no relationships during the drying phase. We detected relationships between community dissimilarities and overland distances for strictly aquatic taxa but not for taxa with an aerial life stage (Fig. 4B, C, Table 3). Community dissimilarities were not associated with environmental or network distances on any date, indicating that, based on the 12 measured variables, niche-based processes were not important determinants of metacommunity dynamics in this network (Fig. 4A–C, Table 3).

Hypothesis 2: Effects of permanent and temporary fragmentation on metacommunity structure

During the dry phase, permanent fragmentation by manmade structures explained more variance in metacommunity structure than temporary fragmentation explained, supporting our 2nd hypothesis. Dissimilarities for the whole community and strictly aquatic taxa were positively correlated with the permanent fragmentation matrix (mean Mantel's $r = 0.20$, p -value: 0.01–0.04; Fig. 5A–C, Table 4). Before the dry phase (March, June), we observed positive relationships between strictly aquatic taxa dissimilarities and permanent fragmentation, suggesting a long-term effect of manmade structures (Table 4). In contrast, taxa with an aerial life stage had no detectable relationships with permanent fragmentation matrices. Community dissimilarities were not correlated with temporary fragmentation matrices.

DISCUSSION

River network fragmentation is one of the most important anthropogenic alterations threatening freshwater biodiversity (Reid et al. 2019). Fragmentation alters the dispersal of organisms within river networks and, thus, has cascading effects on metacommunity dynamics (Gauthier et al. 2020). However, to date, most studies have focused on the effects of permanent fragmentation by manmade structures, whereas

Table 1. Environmental variables measured for each sampling date. Mean ($\pm$ SD) are shown, with the range for each variable (minimum–maximum). Percentage for each substratum type is provided for the main and secondary substrata.

Environmental variable	Sampling Date					
	March	April	June	September	October	December
Elevation (m)			163 ± 19.2 (126–194)			
Slope (%)			1.3 ± 0.6 (0.5–3)			
Hydrologic phase	Wet	Drying	Drying	Dry	Dry	Rewetting
Depth (cm)	22.6 ± 10.7 (5–46)	16.9 ± 9.8 (4–45)	14.2 ± 6.1 (0.61–25)	6.0 ± 2.9 (2–12)	7.0 ± 3.2 (2–14)	16.4 ± 8.7 (3–44)
Wetted width (cm)	230.8 ± 163.6 (51.3–750)	179.7 ± 117.4 (45.4–429.1)	162.8 ± 111.5 (4.7–419.5)	116.9 ± 113.1 (30–457.3)	118.1 ± 104.5 (30.1–424.1)	191.2 ± 128.9 (30.1–532.3)
Flow velocity (m/s)	0.44 ± 0.19 (0.21–0.8)	0.4 ± 0.26 (0.1–1.22)	0.43 ± 0.19 (0.11–0.72)	0.1 ± 0.08 (0–0.25)	0.19 ± 0.15 (0.03–0.66)	0.56 ± 0.31 (0.02–1.15)
Temperature (°C)	9.9 ± 1.2 (7.1–11.3)	13.7 ± 2.5 (9–17.7)	18.9 ± 1.2 (16.3–20.7)	17.0 ± 0.9 (15.8–19)	14.2 ± 1.2 (12.4–16.6)	7.3 ± 0.7 (6–8.2)
Conductance (µS/cm)	225.9 ± 55.6 (96.3–366)	230.6 ± 41.5 (116.5–338)	245.2 ± 46.6 (177.6–373)	333.2 ± 129.2 (183.2–685)	347.0 ± 96.4 (192.8–537)	271.8 ± 76.7 (32.5–476)
Dissolved oxygen (%)	94.2 ± 6.3 (83.5–108.8)	87.9 ± 18.4 (10.26–102.9)	82.8 ± 7.8 (64.8–97.2)	43.5 ± 28.9 (0.6–89.1)	53.2 ± 22.5 (1.5–87.4)	88.3 ± 6.9 (62.3–96.6)
pH	7.42 ± 0.31 (6.81–8.18)	7.47 ± 0.22 (6.91–7.76)	7.5 ± 0.45 (6.86–8.5)	7.42 ± 0.17 (7.09–7.71)	7.17 ± 0.26 (6.39–7.61)	7.29 ± 0.31 (6.67–8.01)
Canopy cover (%)	0–25 to 25–50	0–25 to 25–50	0–25 to 75–100	25–50 to 75–100	0–25 to 75–100	0–25 to 75–100
Main substratum type (%)	Pebbles	Pebbles	Pebbles	Pebbles	Pebbles	Pebbles
	Gravels	Gravels	Gravels	Gravels	Gravels	Gravels
	Sand	Sand	Sand	Mud	Mud	Wood
	Slab	Mud	Mud	Litter	Sand	Helophytes
	Mud	Helophytes	Helophytes	Sand	Silt	Slab
		Litter			Litter	Sand
Secondary substratum type (%)	Gravels	Gravels	Gravels	Gravels	Gravels	Gravels
	Sand	Sand	Pebbles	Sand	Sand	Sand
	Helophytes	Pebbles	Root tufts	Pebbles	Pebbles	Wood
	Wood	Mud	Bryophytes	Wood	Mud	Mud
	Mud		Wood		Silt	Spermaphytes
	Algae				Litter	
	Root tufts					
	Slab					

Table 2. Descriptors of permanent and temporary fragmentation in the Thouaret network. For the temporary fragmentation, values for both dry phase sampling dates are provided. SD = standard deviation.

Fragmentation type	Variables	Value	
Permanent (manmade structures)	Number of structures	32	
	Density (number of structures/linear km)	0.37	
	Mean height of structures (m)	1.69 (± 1.34 SD)	
Temporary (dry reaches)		September	October
	Number of dry patches	20	14
	Average length of dry patch (km)	20.57	20.34
	Total length of dry reaches (km)	40.69	28.80
	Percentage of the catchment (%)	46.09	32.63

the consequences of temporary fragmentation of river networks by drying have only recently been explored (Sarremejane et al. 2017, Gauthier et al. 2020). These recent studies suggest that the temporary fragmentation of river networks leads to shifts in metacommunity processes over time. By repeatedly quantifying the associations between metacommunity and fragmentation patterns in a river network affected by both permanent and temporary fragmentation, our results confirmed that metacommunity processes can greatly vary over time, following wet–dry cycles and temporary fragmentation patterns at the network scale. We found that fragmentation by drying had little effect on metacommunity dynamics within a network subject to considerable permanent fragmentation by manmade structures, indicating that contextualization of the effects of fragmentation by drying is critical to protect and conserve biodiversity of river networks in the Anthropocene.

As we hypothesized, processes structuring the invertebrate metacommunity varied through time across hydro-

logical phases. This finding echoes recent conceptual (Datry et al. 2016a), empirical (Sarremejane et al. 2017), and methodological (Jabot et al. 2020) calls to incorporate temporal variability into the metacommunity framework. These studies also suggested that the relative importance of local and regional processes structuring metacommunities in dynamic river networks could vary substantially over time. Our metacommunity study, which, to our knowledge, is the 1st to describe the physical and biological model of a river network across multiple hydrological phases at such a fine spatiotemporal scale, confirmed that the relative roles of niche-based and dispersal-based metacommunity processes change over time. We suggest that this conclusion may apply to other dynamic freshwater, terrestrial, and marine ecosystems (e.g., Gerisch 2014, Dell et al. 2019).

Although the processes structuring metacommunities varied over time, our conceptual model predicting shifts between local niche-based processes and regional dispersal-based processes was only partially validated. First, we expected

Figure 4. Strength of the relationships (i.e., Mantel's r -values) between community dissimilarities and typical distances (environmental, overland, and network distances) used in metacommunity studies through the different sampling dates for the whole community (A), the community of taxa with aerial life stage (B), and strictly aquatic taxa (C). Relationships with p -values < 0.05 are shown with thicker outlines and an asterisk. r -values of all Mantel tests are available in Table 3.

Table 3. r -values from Mantel tests based on Chao dissimilarity index and typical distances used in metacommunity studies. Distances: Envi = environmental, Over = overland, Netw = network. Relationships with p -values <0.05 are shown in bold.

Sampling date (hydrologic phase)	Distance	Whole community (r)	Taxa with aerial life stage (r)	Taxa strictly aquatic (r)
March (wet)	Envi	−0.04	−0.03	0.05
	Over	0.05	−0.02	0.12
	Netw	−0.04	−0.02	−0.04
April (drying)	Envi	0.00	0.05	−0.06
	Over	−0.04	−0.03	−0.05
	Netw	−0.11	−0.12	−0.10
June (drying)	Envi	0.04	0.02	0.04
	Over	0.10	0.04	0.12
	Netw	0.00	0.03	−0.03
September (dry)	Envi	0.24	0.28	0.13
	Over	0.25	0.12	0.25
	Netw	−0.04	−0.14	0.04
October (dry)	Envi	−0.02	−0.04	−0.04
	Over	0.14	−0.03	0.22
	Netw	0.10	0.00	0.18
December (rewetting)	Envi	0.18	0.01	0.22
	Over	0.16	0.12	0.10
	Netw	0.01	0.04	−0.02

niche-based processes, as defined by the 12 locally measured environmental variables, to be influential during drying, when temperature, nutrient levels, and interspecific competition typically increase and when dissolved oxygen can dramatically decrease (Datry et al. 2016a). However, these environmental variables were not associated with metacommunity structures during any hydrological phase, indicating that niche-based processes were not associated with metacommunity dynamics. In the Thouaret network, although environmental variables varied substantially in space, they did not vary considerably between sampling dates and may not have reached ecologically relevant thresholds. For example, during the drying phase, recorded temperatures did not exceed 20.7°C at any site, which is far below the thermal threshold of most aquatic invertebrates (Quinn et al. 1994). Also, we sampled riffle habitat, but we did not sample pool habitat, which could experience greater change in physicochemical conditions during drying. Including pool habitat may have allowed us to detect a stronger role of niche-based processes. However, the limited influence of niche-based processes on riffle communities in this fragmented river network suggests that metacommunities in intermittent river networks may predominantly be controlled by dispersal. However, many aquatic taxa have adaptations promoting resistance to drying, including the production of desiccation-tolerant, dormant life stages (e.g., Stubbington and Datry 2013), and these adaptations may also partly explain the absence of relationships with local environmental conditions in this net-

work. Such strategies are conceptualized as enabling ‘temporal dispersal’ in metacommunities (e.g., Datry et al. 2016b, Wisnoski et al. 2019) because they promote the persistence of some species in local communities through time, regardless of whether the local environmental conditions define their preferred niche.

We expected niche-based processes to influence invertebrate metacommunity structure during drying phases; however, our result that dispersal was the main process structuring metacommunities in the Thouaret network, independent of hydrological phase, supports other recent studies proposing that metacommunities in dynamic river ecosystems are primarily structured by dispersal (Sarremejane et al. 2017, Gauthier et al. 2020). In intermittent river networks such as the Thouaret, aquatic communities experience local wet-dry cycles, which involve repeated extinction and recolonization events (Datry et al. 2016a). At the river network scale, recurrent recolonization through active and passive dispersal from nearby refuges is a major influence on the organization of biodiversity in space and time (Crabot et al. 2020). Indeed, in our study, community dissimilarities were exclusively explained by physical, overland distances on 4/6 sampling dates. However, these distances only explained a small proportion of total variation, which may reflect ecological drift (Silva et al. 2015), unmeasured environmental factors (e.g., pollutants; Jacobsen 1998), or fragmentation (see below).

In the Thouaret network, permanent fragmentation had a considerable effect on invertebrate dispersal, whereas the

Figure 5. Relationships between community dissimilarities (Chao dissimilarity index) and the 2 fragmentation matrices for September (A), October (B), and December (C) sampling dates and for each community type. Permanent and temporary fragmentation matrices are shown in black and gray circles, respectively. Correspondence of sampling date and hydrological phases: September and October: dry; December: rewetting. Only relationships with p -values < 0.05 are shown with corresponding exact r - and p -values indicated. r -values of all Mantel tests are available in Table 4.

effects of fragmentation by drying were either null or undetectable at the network scale. In contrast, observations by other studies of local-scale differences between perennial and intermittent sites reflect the key role of drying in struc-

turing aquatic communities (e.g., Leigh and Datry 2017, Ruhí et al. 2017), highlighting the effects of drying on riverine biodiversity as context dependent. Additionally, previous studies reporting strong effects of fragmentation by drying on

Table 4. *r*-values from Mantel tests based on Chao dissimilarity index and fragmentation distances used in this study. Perm = permanent fragmentation by manmade structures, Temp = temporary fragmentation by drying. Maximum temporary fragmentation by dry reaches was observed in September, 2018. Relationships with *p*-values <0.05 are shown in bold.

Sampling date (hydrologic phase)	Fragmentation type	Whole community (<i>r</i>)	Taxa with aerial life stage (<i>r</i>)	Taxa strictly aquatic (<i>r</i>)
March (wet)	Perm	0.10	0.05	0.14
	Temp	−0.18	−0.16	−0.13
April (drying)	Perm	0.09	0.07	0.05
	Temp	−0.27	−0.25	−0.21
June (drying)	Perm	0.13	0.07	0.17
	Temp	−0.10	−0.06	−0.12
September (dry)	Perm	0.18	0.00	0.21
	Temp	−0.15	−0.14	−0.07
October (dry)	Perm	0.21	0.062	0.27
	Temp	−0.02	0.01	−0.02
December (rewetting)	Perm	0.18	0.13	0.12
	Temp	−0.13	−0.04	−0.14

metacommunity dynamics were in systems subject to little fragmentation by manmade structures (e.g., Cañedo-Argüelles et al. 2015). In such cases, dry reaches limited aquatic invertebrate dispersal during dry phases, which cascaded through metacommunity dynamics. We found more, and stronger, distance decay relationships between community dissimilarities and physical distances accounting for permanent fragmentation compared with distances including the spatial extent of drying, which suggests that small manmade structures strongly alter invertebrate species dispersal within our river network, as reported for other river metacommunities (Katano et al. 2006, Gauthier et al. 2020) and metapopulations (Blanchet et al. 2010, Taylor et al. 2018). However, the relationships were not very strong, which could be because the high level of fragmentation led to high isolation and, thus, increased ecological drift and species turnover (Merritt et al. 2010).

Context dependency of the effects of river fragmentation were further illustrated by the differential responses to fragmentation by strictly aquatic invertebrate taxa as compared with those that have an aerial life stage, which were much less affected by fragmentation. This partly reflects the highly diverse dispersal modes and capacities of aquatic invertebrates (Heino et al. 2017), which lead to contrasting responses to fragmentation. For example, aquatic invertebrate taxa with an aerial life stage, such as mayflies, caddisflies, and non-biting midges, can colonize previously dry reaches by flying without being severely limited by manmade structures. In contrast, the dispersal of crustaceans, mollusks, and other strictly aquatic taxa was limited, as also has been observed for fishes (Van Looy et al. 2014). In addition to ongoing calls and efforts to better quantify the dispersal routes and capacities of aquatic invertebrates (Downes et al. 2017, Heino et al. 2017), assessing effective dispersal (i.e., dispersal

followed by successful reproduction, ensuring gene flow) by measuring genetic variation among local populations represents the next step toward understanding metacommunity dynamics in fragmented systems and informing conservation plans (Coulon et al. 2010).

Our study, which, to our knowledge, is the 1st to document the effects of both permanent and temporary fragmentation of riverine metacommunities, highlights the need to account for context dependency in metacommunity ecology, particularly for conservation purposes. For example, we observed no additive effects of permanent and temporary fragmentation in any hydrological phases or for any community type (results not shown), which contrasts with other fragmentation contexts in headwater streams of France (Gauthier et al. 2020). Our results also indicate the need to consider the complexity of the effects of drying in river ecology, particularly when exploring network-scale patterns and processes. Further research should explore the synergies and antagonisms between permanent and temporary fragmentation of river networks, which is critical for refining the management and conservation of the rivers of the Anthropocene. This type of study requires substantial effort for extensive field surveys and quantification of biodiversity through sample processing and invertebrate identification, which could present barriers to highly detailed network-scale studies, but novel approaches such as metabarcoding represent opportunities improve the efficiency of research into metacommunity ecology (Bush et al. 2020).

Implications for the management and conservation of river networks

Our results add to recent calls to recognize the temporal variability of metacommunity processes in biodiversity

research toward river conservation (Datry et al. 2016a, Sarremejane et al. 2017, Jabot et al. 2020). That such processes can vary over short timescales in river networks, and most likely in other dynamic ecosystems (e.g., Fernández et al. 2014, Wolkovich et al. 2014), has multiple implications for ecosystem management and conservation planning. For example, assessments of monitoring-site ecological status are mostly based on niche-based processes (Cid et al. 2020), and certain hydrological phases could be more relevant than others for accurate assessments. Identifying such key hydrological phases, as a parallel for keystone species or habitats (Kotliar 2000), will increase understanding of critical metacommunity processes, which will be crucial for improving the success of management practices. Also, conservation actions often need to promote hydrological connectivity during dispersal-limited phases. In cases where fragmentation is mostly due to drying, limiting surface and groundwater abstraction during typically dry summer months could ensure sufficient connectivity for species dispersal within drying river networks (de Graaf et al. 2019).

In the Anthropocene, river networks are experiencing increasing fragmentation by drying events and by the development of small water retention structures for irrigation and human needs (Reid et al. 2019). Previous studies have shown little or no effect of fragmentation by manmade structures on invertebrate communities compared with fish communities (Van Looy et al. 2014, Hwan and Carlson 2016), but our metacommunity approach indicates that small water retention structures can have severe effects on biodiversity patterns. More importantly, the cumulative effects of such structures on riverine biodiversity have not yet been investigated but could be substantial (Linares et al. 2019). As small water retention structures are increasingly promoted as an adaptation strategy to climate change (Habets et al. 2013), we argue that they might considerably alter metacommunity dynamics, resulting in riverine biodiversity loss. Further, determining tipping points of fragmentation levels on metacommunity dynamics is needed to allow prioritization of management efforts, notably with respect to restoration. For example, identifying structure characteristics (e.g., height, passability; Fuller et al. 2015) and their associated effects on metacommunities, as well as determining what fragmentation level is acceptable without altering dispersal among local communities, would be useful for prioritizing geographic locations for restoration and for updating permitting requirements before structures are built. In turn, generating such fragmentation–ecology relationships could reconcile water uses and biodiversity, as well as optimizing the conservation of river networks of the Anthropocene.

ACKNOWLEDGEMENTS

Author contributions: MG, GLF, and BL collected samples. MG, CJD, and TD conceived methodology design. GLF and BL conducted laboratory work. MG and TD conducted data analysis.

MG and TD led the writing of the manuscript, and all authors contributed to writing the manuscript.

This work was supported by the French Agency for Biodiversity (AFB) through the program n°26 entitled “Headwater biodiversity dynamics”. We thank the AFB members, in particular Claire Magand and Claire Bramard, for their help in selecting and expertise on Poitou-Charente rivers. We thank the Nouvelle-Aquitaine Regional Agency for Biodiversity, particularly Arnaud Vaudelet, and Hervé Pella and Jean-Philippe Vidal for their help on GIS and fishing federation data. We thank Julie Crabot, Christel Blot, and Benjamin Gerfand for help with sample collection and processing. We thank Maxence Forcellini, Julien Barnasson, and Jacques Mouthon for identification expertise on Trichoptera and mollusks. We thank Julie Crabot and Nuria Cid-Puey for useful comments on hypotheses and data analysis. This manuscript benefited from meaningful comments from Rachel Stubbington, Romain Sarremejane, Jani Heino, 2 anonymous reviewers, and the associate editor.

SUPPORTING INFORMATION

Mean value of abundance for each taxon and each sampling date (Table S1), supplementary analyses to test for the reliability of the study sampling design (Tables S2, S3) and outputs of non-parametric permutational multivariate analysis of variances testing differences in community composition between intermittent and perennial sites (Table S4) are available online.

LITERATURE CITED

- Arcott, D. B., S. Larned, M. R. Scarsbrook, and P. Lambert. 2010. Aquatic invertebrate community structure along an intermittence gradient: Selwyn River, New Zealand. *Journal of the North American Benthological Society* 29:530–545.
- Blanchet, S., O. Rey, R. Etienne, S. Lek, and G. Loot. 2010. Species-specific responses to landscape fragmentation: Implications for management strategies. *Evolutionary Applications* 3:291–304.
- Bonada, N., S. M. Carlson, T. Datry, D. S. Finn, C. Leigh, D. A. Lytle, M. T. Monaghan, and P. A. Tedesco. 2017. Genetic, evolutionary, and biogeographical processes in intermittent rivers and ephemeral streams. Pages 405–431 in T. Datry, N. Bonada, and A. Boulton (editors). *Intermittent rivers and ephemeral streams*. Academic Press, Cambridge, Massachusetts.
- Brown, B. L., and C. M. Swan. 2010. Dendritic network structure constrains metacommunity properties in riverine ecosystems. *Journal of Animal Ecology* 79:571–80.
- Brown, B. L., C. M. Swan, D. A. Auerbach, E. H. Campbell Grant, N. P. Hitt, K. O. Maloney, and C. Patrick. 2011. Metacommunity theory as a multispecies, multiscale framework for studying the influence of river network structure on riverine communities and ecosystems. *Journal of the North American Benthological Society* 30:310–327.
- Bush, A., W. A. Monk, Z. G. Compson, D. L. Peters, T. M. Porter, S. Shokralla, M. T. G. Wright, M. Hajibabaei, and D. J. Baird. 2020. DNA metabarcoding reveals metacommunity dynamics in a threatened boreal wetland wilderness. *Proceedings of the National Academy of Sciences* 117:8539–8545.
- Cañedo-Argüelles, M., K. S. Boersma, M. T. Bogan, J. D. Olden, I. Phillipsen, T. A. Schriever, D. A. Lytle, and R. Ladle. 2015. Dispersal strength determines meta-community structure in

- a dendritic riverine network. *Journal of Biogeography* 42:778–790.
- Cid, N., N. Bonada, J. Heino, M. Cañedo-Argüelles, J. Crabot, R. Sarremejane, J. Soininen, R. Stubbington, and T. Datry. 2020. A metacommunity approach to improve biological assessments in highly dynamic freshwater ecosystems. *BioScience* 70:427–438.
- Coulon, A., J. W. Fitzpatrick, R. Bowman, and I. J. Lovette. 2010. Effects of habitat fragmentation on effective dispersal of Florida scrub-jays. *Conservation Biology* 24:1080–1088.
- Crabot, J., S. Clappe, S. Dray, T. Datry, and R. McCrea. 2019. Testing the Mantel statistic with a spatially-constrained permutation procedure. *Methods in Ecology and Evolution* 10:532–540.
- Crabot, J., J. Heino, B. Launay, and T. Datry. 2020. Drying determines the temporal dynamics of stream invertebrate structural and functional beta diversity. *Ecography* 43:620–635.
- Crooks, K. R., C. L. Burdett, D. M. Theobald, S. R. B. King, M. Di Marco, C. Rondinini, and L. Boitani. 2017. Quantification of habitat fragmentation reveals extinction risk in terrestrial mammals. *Proceedings of the National Academy of Sciences* 114:7635–7640.
- Datry, T., N. Bonada, and A. J. Boulton. 2017. Intermittent rivers and ephemeral streams: Ecology and management. Academic Press, Cambridge, Massachusetts.
- Datry, T., N. Bonada, and J. Heino. 2016a. Towards understanding the organisation of metacommunities in highly dynamic ecological systems. *Oikos* 125:149–159.
- Datry, T., A. J. Boulton, N. Bonada, K. Fritz, C. Leigh, E. Sauquet, K. Tockner, B. Hugué, and C. N. Dahm. 2018. Flow intermittence and ecosystem services in rivers of the Anthropocene. *Journal of Applied Ecology* 55:353–364.
- Datry, T., K. Fritz, and C. Leigh. 2016b. Challenges, developments and perspectives in intermittent river ecology. *Freshwater Biology* 61:1171–1180.
- Datry, T., S. T. Larned, K. M. Fritz, M. T. Bogan, P. J. Wood, E. I. Meyer, and A. N. Santos. 2014. Broad-scale patterns of invertebrate richness and community composition in temporary rivers: Effects of flow intermittence. *Ecography* 37:94–104.
- Datry, T., H. Pella, C. Leigh, N. Bonada, and B. Hugué. 2016c. A landscape approach to advance intermittent river ecology. *Freshwater Biology* 61:1200–1213.
- de Graaf, I. E. M., T. Gleeson, L. P. H. Rens van Beek, E. H. Sutanudjaja, and M. F. P. Bierkens. 2019. Environmental flow limits to global groundwater pumping. *Nature* 574:90–94.
- Dell, J. E., D. M. Salcido, W. Lumpkin, L. A. Richards, S. M. Pokswinski, E. L. Loudermilk, J. J. O'Brien, and L. A. Dyer. 2019. Interaction diversity maintains resiliency in a frequently disturbed ecosystem. *Frontiers in Ecology and Evolution* 7:145.
- Dewson, Z. S., A. B. W. James, and R. G. Death. 2007. A review of the consequences of decreased flow for instream habitat and macroinvertebrates. *Journal of the North American Benthological Society* 26:401–415.
- Döll, P., and H. M. Schmied. 2012. How is the impact of climate change on river flow regimes related to the impact on mean annual runoff? A global-scale analysis. *Environmental Research Letters* 7:014037.
- Downes, B. J., J. Lancaster, A. Glaister, and W. D. Bovill. 2017. A fresh approach reveals how dispersal shapes metacommunity structure in a human-altered landscape. *Journal of Applied Ecology* 54:588–598.
- Fagan, W. F. 2002. Connectivity, fragmentation, and extinction risk in dendritic metapopulations. *Ecology* 83:3243–3249.
- Fernández, M. E., J. E. Gyenge, S. Varela, and M. de Urquiza. 2014. Effects of the time of drought occurrence within the growing season on growth and survival of *Pinus ponderosa* seedlings. *Trees* 28:745–756.
- Fuller, M. R., M. W. Doyle, and D. L. Strayer. 2015. Causes and consequences of habitat fragmentation in river networks. *Annals of the New York Academy of Sciences* 1355:31–51.
- Gauthier, M., B. Launay, G. Le Goff, H. Pella, C. J. Douady, and T. Datry. 2020. Fragmentation promotes the role of dispersal in determining 10 intermittent headwater stream metacommunities. *Freshwater Biology* 65:2169–2185.
- Gerisch, M. 2014. Non-random patterns of functional redundancy revealed in ground beetle communities facing an extreme flood event. *Functional Ecology* 28:1504–1512.
- Grill, G., B. Lehner, M. Thieme, B. Geenen, D. Tickner, F. Antonelli, S. Babu, P. Borrelli, L. Cheng, H. Crochetiere, H. Ehalt Macedo, R. Filgueiras, M. Goichot, J. Higgins, Z. Hogan, B. Lip, M. E. McClain, J. Meng, M. Mulligan, C. Nilsson, J. D. Olden, J. J. Opperman, P. Petry, C. Reidy Liermann, L. Saenz, S. Salinas-Rodriguez, P. Schelle, R. J. P. Schmitt, J. Snider, F. Tan, K. Tockner, P. H. Valdujo, A. van Soesbergen, and C. Zarfl. 2019. Mapping the world's free-flowing rivers. *Nature* 569:215–221.
- Habets, F., J. Boé, M. Déqué, A. Ducharne, S. Gascoin, A. Hachour, E. Martin, C. Pagé, E. Sauquet, L. Terray, D. Thiéry, L. Oudin, and P. Vienne. 2013. Impact of climate change on the hydrogeology of two basins in northern France. *Climatic Change* 121:771–785.
- Habets, F., E. Philippe, E. Martin, C. H. David, and F. Leseur. 2014. Small farm dams: Impact on river flows and sustainability in a context of climate change. *Hydrology and Earth System Sciences* 18:4207–4222.
- Haddad, N. M., L. A. Brudvig, J. Clobert, K. F. Davies, A. Gonzalez, R. D. Holt, T. E. Lovejoy, J. O. Sexton, M. P. Austin, C. D. Collins, W. M. Cook, E. I. Damschen, R. M. Ewers, B. L. Foster, C. N. Jenkins, A. J. King, W. F. Laurance, D. J. Levey, C. R. Margules, B. A. Melbourne, A. O. Nicholls, J. L. Orrock, D.-X. Song, and J. R. Townshend. 2015. Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science Advances* 1:e1500052.
- Heino, J., J. Alahuhta, T. Ala-Hulkko, H. Antikainen, L. M. Bini, N. Bonada, T. Datry, T. Erös, J. Hjort, O. Kotavaara, A. S. Melo, and J. Soininen. 2017. Integrating dispersal proxies in ecological and environmental research in the freshwater realm. *Environmental Reviews* 25:334–349.
- Heino, J., A. S. Melo, T. Siqueira, J. Soininen, S. Valanko, and L. M. Bini. 2015. Metacommunity organisation, spatial extent and dispersal in aquatic systems: Patterns, processes and prospects. *Freshwater Biology* 60:845–869.
- Heino, J., and K. T. Tolonen. 2017. Ecological drivers of multiple facets of beta diversity in a lentic macroinvertebrate metacommunity. *Limnology and Oceanography* 62:2431–2444.
- Hwan, J. L., and S. M. Carlson. 2016. Fragmentation of an intermittent stream during seasonal drought: Intra-annual and interannual patterns and biological consequences. *River Research and Applications* 32:856–870.
- Jabot, F., F. Laroche, F. Massol, F. Arthaud, J. Crabot, M. Dubart, S. Blanchet, F. Munoz, P. David, and T. Datry. 2020. Assessing

- metacommunity processes through signatures in spatiotemporal turnover of community composition. *Ecology Letters* 23:1330–1339.
- Jacobsen, D. 1998. The effect of organic pollution on the macroinvertebrate fauna of Ecuadorian highland streams. *Fundamental and Applied Limnology* 143:179–195.
- Karaouzas, I., C. Theodoropoulos, A. Vourka, K. Gritsalis, and N. T. Skoulidakis. 2019. Stream invertebrate communities are primarily shaped by hydrological factors and ultimately fine-tuned by local habitat conditions. *Science of the Total Environment* 665:290–299.
- Katano, O., T. Nakamura, S. Abe, S. Yamamoto, and Y. Baba. 2006. Comparison of fish communities between above- and below-dam sections of small streams: Barrier effect to diadromous fishes. *Journal of Fish Biology* 68:767–782.
- Kotliar, N. B. 2000. Application of the new keystone-species concept to prairie dogs: How well does it work? *Conservation Biology* 14:1715–1721.
- Legendre, P., and L. Legendre. 2012. *Numerical ecology*. 3rd edition. Elsevier, Amsterdam, The Netherlands.
- Lehner, B., C. R. Liermann, C. Revenga, C. Vörösmarty, B. Fekete, P. Crouzet, P. Doll, M. Endean, K. Frenken, J. Magome, C. Nilsson, J. C. Robertson, R. Rödel, N. Sindorf, and D. Wisser. 2011. High-resolution mapping of the world's reservoirs and dams for sustainable river-flow management. *Frontiers in Ecology and the Environment* 9:494–502.
- Leibold, M. A., M. Holyoak, N. Mouquet, P. Amarasekare, J. M. Chase, M. F. Hoopes, R. D. Holt, J. B. Shurin, R. Law, D. Tilman, M. Loreau, and A. Gonzalez. 2004. The metacommunity concept: A framework for multi-scale community ecology. *Ecology Letters* 7:601–613.
- Leigh, C., and T. Datry. 2017. Drying as a primary hydrological determinant of biodiversity in river systems: A broad-scale analysis. *Ecography* 40:487–499.
- Linares, M. S., W. Assis, R. R. de Castro Solar, R. P. Leitão, R. M. Hughes, and M. Callisto. 2019. Small hydropower dam alters the taxonomic composition of benthic macroinvertebrate assemblages in a neotropical river. *River Research and Applications* 35:725–735.
- Merritt, D. M., C. Nilsson, and R. Jansson. 2010. Consequences of propagule dispersal and river fragmentation for riparian plant community diversity and turnover. *Ecological Monographs* 80:609–626.
- Nekola, J. C., and P. S. White. 1999. The distance decay of similarity in biogeography and ecology. *Journal of Biogeography* 26:867–878.
- Oksanen, J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. Minchin, R. B. O'Hara, G. L. Simpson, P. Solymos, M. H. H. Stevens, E. Szoecs, and H. Wagner. 2019. *vegan*: Community ecology package. (Available from: <https://cran.r-project.org/web/packages/vegan/index.html>)
- Pella, H., J. Lejot, N. Lamouroux, and T. Snelder. 2012. Le réseau hydrographique théorique (RHT) français et ses attributs environnementaux. *Géomorphologie: relief, processus, environnement* 18:317–336.
- Perkin, J. S., K. B. Gido, A. R. Cooper, T. F. Turner, M. J. Osborne, E. R. Johnson, K. B. Mayes, and C. Nilsson. 2015. Fragmentation and dewatering transform Great Plains stream fish communities. *Ecological Monographs* 85:73–92.
- Poff, N. L. 1997. Landscape filters and species traits: Towards mechanistic understanding and prediction in stream ecology. *Journal of the North American Benthological Society* 16:391–409.
- Quinn, J. M., G. L. Steele, C. W. Hickey, and M. L. Vickers. 1994. Upper thermal tolerances of twelve New Zealand stream invertebrate species. *New Zealand Journal of Marine and Freshwater Research* 28:391–397.
- Reid, A. J., A. K. Carlson, I. F. Creed, E. J. Eliason, P. A. Gell, P. T. J. Johnson, K. A. Kidd, T. J. MacCormack, J. D. Olden, S. J. Ormerod, J. P. Smol, W. W. Taylor, K. Tockner, J. C. Vermaire, D. Dudgeon, and S. J. Cooke. 2019. Emerging threats and persistent conservation challenges for freshwater biodiversity. *Biological Reviews* 94:849–873.
- Ruhí, A., T. Datry, and J. L. Sabo. 2017. Interpreting beta-diversity components over time to conserve metacommunities in highly dynamic ecosystems. *Conservation Biology* 31:1459–1468.
- Sarremejane, R., M. Cañedo-Argüelles, N. Prat, H. Mykrä, T. Muotka, and N. Bonada. 2017. Do metacommunities vary through time? Intermittent rivers as model systems. *Journal of Biogeography* 44:2752–2763.
- Sarremejane, R., J. England, C. E. M. Sefton, S. Parry, M. Eastman, and R. Stubbington. 2020. Local and regional drivers influence how aquatic community diversity, resistance and resilience vary in response to drying. *Oikos* 129:1877–1890.
- Silva, J. L. A., A. F. Souza, J. G. Jardim, and B. T. Goto. 2015. Community assembly in harsh environments: The prevalence of ecological drift in the heath vegetation of South America. *Ecosphere* 6:1–18.
- Stubbington, R., and T. Datry. 2013. The macroinvertebrate seedbank promotes community persistence in temporary rivers across climate zones. *Freshwater Biology* 58:1202–1220.
- Tachet, H., P. Richoux, M. Bournaud, and P. Usseglio-Polatera. 2000. *Invertébrés d'eau douce: systématique, biologie, écologie*. CNRS Editions, Paris, France.
- Taylor, A. T., M. Papeş, and J. M. Long. 2018. Incorporating fragmentation and non-native species into distribution models to inform fluvial fish conservation. *Conservation Biology* 32:171–182.
- Thrush, S. F., J. Halliday, J. E. Hewitt, and A. M. Lohrer. 2008. The effects of habitat loss, fragmentation, and community homogenization on resilience in estuaries. *Ecological Applications* 18:12–21.
- Van Looy, K., T. Tormos, and Y. Souchon. 2014. Disentangling dam impacts in river networks. *Ecological Indicators* 37:10–20.
- Vidal, J. P., E. Martin, L. Franchistéguy, M. Baillon, and J. M. Soubeyroux. 2010. A 50-year high-resolution atmospheric reanalysis over France with the Safran system. *International Journal of Climatology* 30:1627–1644.
- Wisnoski, N. I., M. A. Leibold, and J. T. Lennon. 2019. Dormancy in metacommunities. *American Naturalist* 194:135–151.
- Wolkovich, E. M., B. I. Cook, K. K. McLauchlan, and T. J. Davies. 2014. Temporal ecology in the Anthropocene. *Ecology Letters* 17:1365–1379.
- Zarfl, C., A. E. Lumsdon, J. Berlekamp, L. Tydecks, and K. Tockner. 2014. A global boom in hydropower dam construction. *Aquatic Sciences* 77:161–170.

Freshwater Science

Author Fee Billing Information

Please fill in the box at the bottom of the form and return by fax (preferred) to +1.773.753.3616 or by mail to:
Cindy Garrett, The University of Chicago Press, Journals Division
1427 East 60th Street, Chicago, Illinois 60637

We cannot accept credit card information via e-mail. Prompt payment is required to avoid delays in publication of your article. **An author with an outstanding debt from a previous article cannot publish a new article until the original debt is settled.**

FEES

- Page charges: \$30/page for the first 13 typeset pages; \$50/page for every typeset page over 13
- Open access: \$400 (*Editor approval required **BEFORE** this form is submitted; Please email: fws.editor@freshwater-science.org*)
- Color figures in the printed edition \$150.00 per color figure
- Additional fees for special treatment of the manuscript may be assessed at the editor's discretion.

CHECKS/PURCHASE ORDERS

- Please list the vendor as The University of Chicago Press, 11030 Langley Avenue, Chicago, Illinois 60628.
- If the purchase order is sent separately from this form, it must be marked "confirming" and mailed to the address above. Please note: Purchase orders will not be processed without a manuscript number.
- All purchase orders must include the following information:
Name of journal, Issue date (month and year), Author's name, Amount of the invoice
- Make checks and purchase orders payable to The University of Chicago Press

ELECTRONIC TRANSFER INFORMATION

- Financial institution: The Northern Trust Bank, 50 South LaSalle Street, Chicago, Illinois 60603
- Account name: University of Chicago Press Account number: 2725681
- Routing number: 071000152 SWIFT #: CNORUS44

Article Information
Vol. _____ Issue ____ Month _____ Year _____ Manuscript No. _____ Total pages _____
Author(s) _____ Total color figures _____
Open Access charge _____ (<i>Editor approval required before this form is submitted; Please Email: fws.editor@freshwater-science.org</i>)
Article title _____

Payment Method
1. Institutional Purchase Order No. _____ attached () to come ()
2. Check ()
3. Electronic Fund Transfer ()
4. Credit Card: Visa () MasterCard () American Express () Discover ()
Name as it appears on card: _____
Card number: _____
Expiry date: _____ 3-4 digit security code: _____
Signature: _____
Phone number: _____ Email: _____
<i>Please Note: Invoice and credit card charges will appear from the Chicago Distribution Center or the University of Chicago Press</i>

Mandatory Address for Invoice
E-mail: _____
Mail: Name: _____
Address _____
City: _____ State/Province: _____
Post code: _____ Country: _____